

Title	First report of Pelargonium zonate spot virus from wild Brassicaceae plants in Japan
Author(s)	Kamitani, Mari; Nagano, Atsushi J.; Honjo, Mie N.; Kudoh, Hiroshi
Citation	Journal of General Plant Pathology (2017), 83(5): 329-332
Issue Date	2017-09
URL	http://hdl.handle.net/2433/234686
Right	This is a post-peer-review, pre-copyedit version of an article published in 'Journal of General Plant Pathology'. The final authenticated version is available online at: https://doi.org/10.1007/s10327-017-0727-6 / The full-text file will be made open to the public on 28 July 2018 in accordance with publisher's 'Terms and Conditions for Self-Archiving'. / This is not the published version. Please cite only the published version. この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。
Type	Journal Article
Textversion	author

1 First report of *Pelargonium zonate spot virus* from wild Brassicaceae plants in Japan

2

3 Mari Kamitani, MK, 1

4 Atsushi J. Nagano, AJN, 12

5 Mie N. Honjo, MNH, 1

6 Hiroshi Kudoh, HK, 1

7

8 1: Center for Ecological Research, Kyoto University, Hirano 2-509-3, Otsu 520-2113,

9 Japan

10 2: Faculty of Agriculture, Ryukoku University, Yokotani 1-5, Seta Oe-cho, Otsu 520-2914,

11 Japan

12

13 **Corresponding authors**

14 Mari Kamitani

15 kamitani@ecology.kyoto-u.ac.jp

16 Hiroshi Kudoh

17 kudoh@ecology.kyoto-u.ac.jp

18 TEL: +81-77-549-8200 FAX: +81-77-549-8201

19

20 **Total text pages: 7**

21

22 **Numbers of tables and figures: 1 table and 2 figures**

23

24

25

26 **Abstract**

27 *Pelargonium zonate spot virus* (PZSV) was identified from two wild Brassicaceae plant
28 species, *Arabidopsis halleri* and *Rorippa indica*, in central Japan using RNA-Seq and
29 reverse transcription polymerase chain reaction. The deduced amino acid sequences of
30 RNA-dependent RNA polymerase and coat protein were highly similar to those of
31 previously reported PZSV isolates, with 96.6%–98.2% and 93.7%–98.0% identity,
32 respectively. Mechanical inoculation revealed the pathogenicity of the PZSV isolate to
33 *Nicotiana benthamiana* and *Brassica oleracea*. To the best of our knowledge, this is the
34 first report of PZSV from Japan.

35

36 **Keywords**

37 *Pelargonium zonate spot virus*, *Arabidopsis halleri*, *Rorippa indica*, Japan, RNA-Seq,
38 *Brassica oleracea*

39

40

41

42

43

44

45

46 *Pelargonium zonate spot virus* (PZSV) belongs to the family *Bromoviridae* genus
47 *Anulavirus* and consists of three linear, positive-sense single-stranded RNAs (ssRNAs)
48 (Codoner & Elena, 2006). The three RNA segments, RNA-1, RNA-2, and RNA-3, encode
49 protein 1a, protein 2a (RNA-dependent RNA polymerase, RdRP), and two other proteins
50 (movement and coat proteins), respectively (Finetti-Sialer & Gallitelli, 2003). PZSV was
51 isolated from *Pelargonium zonale* (Quacquarelli and Gallitelli 1979 cited in Finetti-Sialer
52 and Gallitelli 2003) and has been reported to be a causal agent of tomato diseases in Italy
53 (Gallitelli, 1982), Spain (Luis-Arteaga & Cambra, 2000), France (Gebre-Selassie *et al.*,
54 2002), Israel (Lapidot *et al.*, 2010), Australia (Luo *et al.*, 2010), Argentina (Giolitti *et al.*,
55 2014) and USA (Liu & Sears, 2007). PZSV has also been shown to have a wider host
56 range, including plants from Solanaceae, Actinidiaceae, Brassicaceae, and Asteraceae
57 (reviewed in (Li *et al.*, 2016)). In this study, we report the first cases of PZSV infection
58 in wild Brassicaceae plants in Japan.

59 We made samplings in two localities where multiple Brassicaceae species co-occur,
60 i.e. Mino-gawa (Mino, Osaka Prefecture, Japan, alt. ca. 200 m, June 18, 2014) and
61 Gongen-dani (Taga, Shiga Prefecture, Japan, alt. ca. 320 m, June 28, 2014). The former
62 and the latter communities contained four [*Arabidopsis halleri* subsp. *gemmaifera* (*A.*
63 *halleri*, hereafter), *Rorippa indica*, *Cardamine scutata* and *C. occulta*] and six (*A. halleri*,
64 *R. indica*, *C. leucantha*, *C. impatiens*, *C. hirsuta* and *Arabis flagellosa*) Brassicaceae
65 species, respectively. At Mino-gawa, *A. halleri* grew by forming patches along the valley,
66 and we collected 17 leaves (Fig. 1a) from 17 plant patches (one sample/ patch) at 15–20-
67 m intervals within a sampling range of approximately 350 m in distance. At Gongen-dani,
68 we collected 9 leaves of *R. indica* (Fig. 1a) along the valley at 5–10-m intervals within a
69 sampling range of approximately 100 m. Immediately after sampling of cauline leaves,
70 each sample was immersed in 1.0 mL RNAlater (Life Technologies, CA, USA) to avoid

71 RNA degradation. Total RNA was extracted from each sample using 300–600 μ L (10
72 volumes) TRIzol reagent (Life Technologies, Carlsbad, CA, USA) according to the
73 manufacturer's instructions. We conducted RNA-Seq with selective depression of rRNA
74 by thermostable RNaseH (Morlan *et al.*, 2012, Nagano *et al.*, 2015) using a HiSeq 2500
75 instrument (Illumina, San Diego, CA, USA) at Macrogen Japan. Detailed procedures for
76 the RNA-Seq library preparation, mapping, and virus detection are described in a
77 previous study (Kamitani *et al.* 2016). We obtained 1.8 Gb (giga base) sequence data in
78 total that contained 8,132,772 and 9,820,064 reads from *R. indica* (9 samples) and *A.*
79 *halleri* (17 samples), respectively. We determined the infecting viruses by considering the
80 coverage of virus genome and the amount of reads which mapped on the virus genome.

81 PZSV was detected in five samples of *A. halleri* and four samples of *R. indica* (red
82 bold IDs in Fig. 1a and red circles in Fig. 1b). Some infected leaves showed chlorosis or
83 yellowing, but others did not show visible symptom at a leaf level (red bold IDs in Fig.
84 1a). Therefore, we could not judge whether symptom-like phenotypes of our samples had
85 been caused by the infection of PZSV. Eight of the nine putative infected samples
86 contained nearly the full length of the PZSV genome sequence, as indicated by the nearly
87 100% genome coverage by RNA-Seq reads (Fig. 1b). One sample of *A. halleri* showed
88 relatively low coverage (24%) which may be caused by low read number in this sample.
89 In addition to PZSV, *Turnip mosaic virus* (TuMV), *Cucumber mosaic virus* (CMV) and
90 *Brassica yellows virus* (BrYV) were detected from *A. halleri* at Mino-gawa. In 5 PZSV-
91 infected samples, two samples showed coinfections with TuMV (A11 in Fig. 1a) and with
92 TuMV and CMV (A8 in Fig. 1a), respectively. No other virus was detected from *R. indica*
93 at Gongen-dani.

94 To characterize PZSV sequences from each host species, consensus sequences of
95 RdRP, encoded in RNA-2, and coat protein (CP), encoded in RNA-3, were determined

96 using the RNA-Seq reads, respectively. The nucleotide and deduced amino acid (AA)
97 sequences from the two hosts were deposited in GenBank (accession numbers were
98 shown in Tables 1 and 2). The two AA sequences of RdRP and CP were similar (> 96%
99 identity and > 93% identity, respectively) to those of three previously reported PZSV
100 isolates (Tables 1 and 2). We found 35 polymorphic AA sites in RdRP across the five
101 sequences (Fig. S1).

102 To test the pathogenicity of the virus, mechanical inoculation was conducted using two
103 infected leaves of *R. indica* collected at the Gongen-dani site. Each leaf was crushed
104 completely in 500- μ L of 0.1M phosphate buffer (pH 7.4, consists of disodium hydrogen
105 phosphate and sodium dihydrogen phosphate) and inoculated onto *Nicotiana*
106 *benthamiana* using a carborundum (600 mesh). At 28–35 days post-inoculation (dpi), *N.*
107 *benthamiana* showed a susceptible phenotype with mild chlorosis and stunting relative to
108 the uninfected plant (Fig. 2a, b). The stunting symptom by the infection of PZSV has been
109 reported in *N. benthamiana* (Lapidot *et al.*, 2010). We extracted RNA from
110 inoculated/upper leaves of PZSV-inoculated or buffer (mock)-inoculated plants at 28 dpi.
111 RT-PCR was conducted to confirm the infections by amplifying RNA-3 genome fragment
112 of PZSV (403 bp in length). The RT reaction was conducted using a High-Capacity cDNA
113 Reverse Transcription Kit (Life Technologies) with random primers. PCR was conducted
114 using KOD -plus- (TOYOBO, Japan) with specific primers (forward, 5'-
115 AGATTTTCCGGGCTCTCTA-3' and reverse, 5'-
116 GTTCAACTGTTTTACCAGGATAG-3'). Based on the RNA 3 sequences from our study
117 sites, the sequence of the primers were modified from a previous study (Choi *et al.*, 2013).
118 The RT-PCR assay detected PZSV from both inoculated leaves and upper leaves (Fig. 2c),
119 suggesting that the virus caused systemic infection in *N. benthamiana*.

120 Mechanical inoculations were also conducted to test the pathogenicity of the virus in

121 *Solanum lycopersicum* (tomato, cultivar 'Momotaro' obtained from TAKII SEED, and
122 another 1 cultivar) and *Brassica oleracea* var. *capitata* (cabbage, cultivar 'Shoshu',
123 obtained from TAKII SEED). For all cultivars of *S. lycopersicum* and *B. oleracea*, in total
124 four plant individuals (two for inoculation and two for mock inoculation) were used for
125 the test. From 20 dpi and later, *B. oleracea* showed symptoms of chlorosis and occasional
126 ring patterns along the leaf veins (Fig. 3a) that were distinct from mock-inoculated
127 phenotypes (Fig. 3b). RT-PCR confirmed systemic PZSV infection in inoculated *B.*
128 *oleracea*, and we continually observed symptoms on newly expanded leaves. Although
129 further quantitative evaluation is required, we judged that PZSV has a potential risk to
130 reduce the cabbage yield. PZSV has been the causal agent of multiple disease outbreaks
131 in commercial tomato crops (Hanssen *et al.*, 2010), but infection was not detected either
132 of the two cultivars of *S. lycopersicum*. We also tested mechanical inoculations on two *A.*
133 *halleri* plants, but infection was not detected.

134 In this study, we identified PZSV from two natural Brassicaceae populations in Japan
135 and, to our knowledge, this is the first report of PZSV in Japan. PZSV has been previously
136 reported from Brassicaceae weeds, e.g. *Capsella bursa-pastoris* and *Diplotaxis eruroides*
137 in Europe (Finetti-Sialer & Gallitelli, 2003) and *Cakile maritima* in Australia (Luo *et al.*,
138 2010), but has not been reported in cabbage. Because we found PZSV pathogenicity to
139 one cultivar of cabbage, further studies are needed to determine whether the Japanese
140 isolates of PZSV in *A. halleri* and *R. indica* can be transmitted to agricultural crops. It has
141 been reported that the PZSV is transmitted from the Brassicaceae weed, *D. eruroides*, to
142 tomato via pollen grains carried by thrips (Vovlas *et al.*, 1989, cited in Finetti-Sialer &
143 Gallitelli, 2003). Our study indicated that surveys of virus infection in wild plants can
144 contribute to improve our knowledge of potential reservoirs of pathogens for crop plants.

145

146 **Acknowledgements**

147 We thank M. Sato and G. Atsumi for their comments. This work was supported by a
148 Grant-in-Aid for Scientific Research (S) (26221106) MEXT to HK, a Grant-in-Aid for
149 JSPS Fellows (15J00628) to MK, and the Platform for Supporting in Drug Discovery,
150 Informatics, and Structural Life Science from Japan AMED (PDIS0046).

151

152 **Figure legends**

153 **Fig. 1.** Detection of PZSV. **a.** Sampled leaves of *A. halleri* (Mino-gawa) and *R. indica*
154 (Gongen-dani). The sample IDs are listed above the photos. The red bold IDs correspond
155 to samples that were found to be infected by PZSV. Asterisks next to the sample IDs
156 indicate the mixed-infection of PZSV with TuMV (*), and with TuMV and CMV (**). **b.**
157 The amount of RNA-Seq reads mapped on the PZSV genome (vertical axis; virus reads
158 per million host reads) and the genome coverage (horizontal axis) plotted for each leaf
159 sample. The red circles indicate samples that were found to be infected by PZSV. The
160 black circles represent samples without any PZSV reads.

161 **Fig. 2.** Inoculation experiment and confirmation of PZSV infection in *N. benthamiana*. **a.**
162 Symptoms observed in *N. benthamiana* at 35 dpi in an inoculated plant (*right*) and a
163 mock-inoculated plant (*left*). **b.** Chlorosis observed in a leaf of *N. benthamiana* on 28 and
164 35 dpi. **c.** Detection of PZSV genome fragments by RT-PCR. The characters “I” and “U”
165 in the panel indicate inoculated leaves and upper leaves of the PZSV-inoculated plant,
166 respectively. “Mock-I” and “Mock-U” indicate the inoculated and upper leaves of the
167 mock (buffer)-inoculated plant, respectively.

168 **Fig. 3.** Inoculation of PZSV to *B. oleracea*. **a.** Symptoms observed in an upper leaf of the
169 PZSV-inoculated *B. oleracea*. **b.** An upper leaf of the mock-inoculated *B. oleracea*.

170

171

172 **References**

- 173 Choi H, Cho WK, Yu J, Lee JS & Kim KH (2013) Highly Specific Detection of Five Exotic
174 Quarantine Plant Viruses using RT-PCR. *Plant Pathol J* **29**: 99-104.
- 175 Codoner FM & Elena SF (2006) Evolutionary relationships among members of the
176 *Bromoviridae* deduced from whole proteome analysis. *Arch Virol* **151**: 299-307.
- 177 Finetti-Sialer M & Gallitelli D (2003) Complete nucleotide sequence of *Pelargonium zonate*
178 *spot virus* and its relationship with the family *Bromoviridae*. *J Gen Virol* **84**: 3143-3151.
- 179 Gallitelli D (1982) Properties of tomato isolate of *Pelargonium zonate spot virus*. *Ann Appl*
180 *Biol* **100**: 457-466.
- 181 Gebre-Selassie K, Delecolle B, Gognalons P, Dufour O, Gros C, Cotillon AC, Parrella G &
182 Marchoux G (2002) First report of an isolate of *Pelargonium zonate spot virus* in
183 commercial glasshouse tomato crops in Southeastern France. *Plant Disease* **86**: 1052.
- 184 Giolitti F, Bejerman N, Nome C, Visintin G, Breuil Sd & Lenardon S (2014) Biological and
185 molecular characterization of an isolate of *Pelargonium zonate spot virus* infecting
186 sunflower in Argentina. *J Plant Pathol* **96**: 189-194.
- 187 Hanssen IM, Lapidot M & Thomma BP (2010) Emerging viral diseases of tomato crops. *Mol*
188 *Plant Microbe Interact* **23**: 539-548.
- 189 Kamitani M, Nagano AJ, Honjo MN & Kudoh H (2016) RNA-Seq reveals virus-virus and
190 virus-plant interactions in nature. *FEMS Microbiol Ecol* **92**: doi: 10.1093/femsec/fiw1176.
- 191 Lapidot M, Guenoune-Gelbart D, Leibman D, Holdengreber V, Davidovitz M, Machbash Z,
192 Klieman-Shoval S, Cohen S & Gal-On A (2010) *Pelargonium zonate spot virus* is
193 transmitted vertically via seed and pollen in tomato. *Phytopathology* **100**: 798-804.
- 194 Li H, Zhang C, Luo H, Jones MGK, Sivasithamparam K, Koh S-H, Ong JWJ & Wylie SJ
195 (2016) Yellow tailflower mild mottle virus and *Pelargonium zonate spot virus* co-infect a

- 196 wild plant of red-striped tailflower in Australia. *Plant Pathol* **65**: 503–509.
- 197 Liu H & Sears JL (2007) First report of Pelargonium zonate spot virus from tomato in the
198 United States. *Plant Dis* **91**: 633.
- 199 Luis-Arteaga M & Cambra M (2000) First report of natural infection of greenhouse-grown
200 tomato and weed species by *Pelargonium zonate spot virus* in Spain. *Plant Dis* **84**: 807.
- 201 Luo H, Wylie SJ & Jones MG (2010) Identification of plant viruses using one-dimensional
202 gel electrophoresis and peptide mass fingerprints. *J Virol Methods* **165**: 297-301.
- 203 Morlan JD, Qu K & Sinicropi DV (2012) Selective Depletion of rRNA Enables Whole
204 Transcriptome Profiling of Archival Fixed Tissue. *PLoS ONE* **7**: e42882.
- 205 Nagano AJ, Honjo MN, Mihara M, Sato M & Kudoh H (2015) Detection of plant viruses in
206 natural environments by using RNA-Seq. *Methods Mol Biol* **1236**: 89-98.
- 207 Quacquarelli A, Gallitelli D (1979) [Three virus diseases of Pelargonium in Apulia].
208 *Phytopathol Mediterr* **18**, 61-70 (in Italian).
- 209 Tomlinson JA, Carter AL, Dale WT & Simpson CJ (1970) Weed plants as sources of
210 cucumber mosaic virus. *Ann Appl Biol* **66**: 11-16.
- 211 Vovlas C, Gallitelli D & Conti M (1989). Preliminary evidence for an unusual mode of
212 transmission in the ecology of *Pelargonium zonate spot virus* (PZSV). IVth International
213 Plant Virus Epidemiology Workshop, pp. 302–305 (Montpellier, France, 3–8 September
214 1989).
- 215

Fig. 2





Fig. 3



Tabel 1 Identity of RdRP amino acid sequences between PZSV isolates

	Accession in Genbank	<i>A. halleri</i> , Mino-gawa	<i>R. inidica</i> , Gongen-dani	Isolate tomato	Isolate SW13
<i>A. halleri</i> , Mino-gawa ^(a)	LC178561	-	-	-	-
<i>R. inidica</i> , Gongen-dani ^(a)	LC178560	0.983	-	-	-
Isolate tomato, Italy ^(b)	NC_003650.1	0.966	0.966	-	-
Isolate SW13, Australia ^(c)	KF790761.3	0.979	0.978	0.969	-
Isolate parana, Argentina ^(d)	JQ350739.1	0.978	0.982	0.970	0.979

(a) this study, (b) Gallitelli (1982), (c) Luo *et al.* (2010), (d) Giolitti *et al.* (2014)

Tabel 2 Identity of CP amino acid sequences between PZSV isolates

	Accession in Genbank	<i>A. halleri</i> , Mino-gawa	<i>R. inidica</i> , Gongen-dani	Isolate tomato	Isolate SW13
<i>A. halleri</i> , Mino-gawa ^(a)	LC269013	-	-	-	-
<i>R. inidica</i> , Gongen-dani ^(a)	LC269014	0.966	-	-	-
Isolate tomato, Italy ^(b)	NC_003651.1	0.937	0.961	-	-
Isolate SW13, Australia ^(c)	KF790762.4	0.975	0.980	0.956	-
Isolate parana, Argentina ^(d)	JQ350737.1	0.956	0.971	0.951	0.980

(a) this study, (b) Gallitelli (1982), (c) Luo *et al.* (2010), (d) Giolitti *et al.* (2014)




Fig. S1 Comparison of deduced amino acid (AA) sequences of RdRP obtained in this study and those reported in NCBI/Genbank (accession numbers in Table 1). Black shade indicates the major AA residue ($n \geq 2$) at the position. An asterisk under the sequence (AA position 110) indicates the position with two major AA (M and T).