

Title	Daily cumulative hip moment is associated with radiographic progression of secondary hip osteoarthritis
Author(s)	Tateuchi, H.; Koyama, Y.; Akiyama, H.; Goto, K.; So, K.; Kuroda, Y.; Ichihashi, N.
Citation	Osteoarthritis and Cartilage (2017), 25(8): 1291-1298
Issue Date	2017-8
URL	http://hdl.handle.net/2433/234976
Right	© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/ ; The full-text file will be made open to the public on 1 March 2018 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version. この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。
Type	Journal Article
Textversion	author

Title: Daily cumulative hip moment is associated with radiographic progression of secondary hip osteoarthritis

Authors: Hiroshige Tateuchi, PT, PhD^{1*}, Yumiko Koyama PT, MS¹, Haruhiko Akiyama, MD, PhD², Koji Goto, MD, PhD³, Kazutaka So, MD, PhD³, Yutaka Kuroda, MD, PhD³, Noriaki Ichihashi, PT, PhD¹

¹ Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto, Japan

² Department of Orthopaedic Surgery, School of Medicine, Gifu University, Gifu, Japan

³ Department of Orthopaedics Surgery, Graduate School of Medicine, Kyoto University, Kyoto, Japan

Grant: JSPS KAKENHI Grant-in-Aid for Scientific Research (C) Grant No. 24500578.

Competing interests: The work reported in this manuscript has not received financial support from any commercial source. There are no conflicts of interest to declare with regard to this study.

*** Corresponding author:**

Hiroshige Tateuchi, Ph.D.

Kyoto University

53 Kawara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

Tel; +81-75-751-3964: Fax; +81-75-751-3909

E-mail; tateuchi.hiroshige.8x@kyoto-u.ac.jp

Running title: Daily cumulative hip moment and hip OA progression

1 **ABSTRACT**

2

3 **Objective:**

4 To investigate whether higher daily cumulative hip moment at baseline is associated with
5 subsequent radiographic progression of hip osteoarthritis (OA) over 12 months.

6 **Design:**

7 Fifty patients with secondary hip OA, excluding patients with end-stage hip OA, participated in this
8 prospective cohort study. Joint space width (JSW) of the hip was measured at baseline and 12
9 months later. With radiographic progression of hip OA (> 0.5 mm/year in JSW) as dependent
10 variable (yes/no), univariable and multivariable logistic regression analyses were performed to
11 assess the association between load-related parameters during gait (i.e., peak hip moment, hip
12 moment impulse, and daily cumulative hip moment [product of hip moment impulse and mean
13 steps/day]) and hip OA progression with and without adjustment for age, body weight, and
14 minimum JSW.

15 **Results:**

16 Of the 50 patients (47.4 ± 10.7 years old), 21 (42.0%) were classified into the progression group.
17 The higher daily cumulative hip moment in the frontal plane at baseline was statistically
18 significantly associated with radiographic progression of hip OA (adjusted OR [95% CI], 1.34
19 [1.06–1.70]; $P = 0.013$). The higher daily cumulative hip moment in the sagittal plane was also
20 approaching significance in its association with hip OA progression (adjusted OR, 1.80 [0.99–
21 3.26]; $P = 0.052$).

22 **Conclusions:**

23 In the female patients with secondary hip OA, higher daily cumulative hip moment, particularly in
24 the frontal plane, was a predictor of radiographic progression of hip OA over 12 months. Reduction
25 in daily cumulative hip moment by modification in gait and physical activity may potentially slow
26 hip OA progression.

27

28

29 **Keywords:** Hip osteoarthritis, Gait, Biomechanics, cumulative joint moment

1 INTRODUCTION

2

3 Although progression of hip osteoarthritis (OA) seems to be multifactorial, genetic mutation¹,
4 higher age², female, narrower joint space width (JSW) and higher Kellgren and Lawrence score at
5 baseline^{2,3}, abnormal hip morphology such as hip dysplasia^{1,2,4,5}, atrophic bone response^{2,6}, and hip
6 pain³ are known potential risk factors for progression of hip OA. Especially for secondary hip OA,
7 which is more prevalent than primary OA⁷, abnormal hip morphology and malalignment between
8 acetabular and proximal femoral head play an important role in radiographic progression^{4,8}.

9

10 In knee OA, a mechanical factor (i.e., excessive knee adduction moment and moment impulse
11 during gait) has been identified as an important contributor to OA progression⁹⁻¹³. However, gait
12 biomechanics associated with progression of hip OA remain unknown. Extended exposure to heavy
13 physical work such as heavy lifting and standing can increase the risk for hip OA¹⁴, although it is
14 not known whether excessive load during gait is related to progression of hip OA. A recent
15 longitudinal study reported that patients with hip OA who later underwent total hip arthroplasty
16 (THA) had less hip extension moment and hip extension angle during gait at baseline compared to
17 those without surgery¹⁵. Although that study did not necessarily examine the causal relationship
18 between gait and radiographic progression since the decision of operation depends on multiple
19 factors, it highlights the need for investigation of the association between gait biomechanics and
20 progression of hip OA. However, the mechanical risk factor during gait for hip OA progression has
21 not been identified.

22

23 The external joint moment during gait can be used to estimate mechanical load since joint load
24 cannot be directly measured *in vivo* noninvasively. Hip contact force during gait can be predicted
25 from absolute hip joint moment in the three planes during the stance phase of gait^{16,17}. Peak joint
26 moment and joint moment impulse have been used as indicators of joint load^{11,13}. Peak joint
27 moment represents instantaneous load at a specific point during stance phase, and moment impulse
28 measures the total amount of load during stance phase by incorporating both load magnitude and
29 duration. Furthermore, total exposure to joint load during daily activities has been measured as

1 daily cumulative joint moment calculated as the product of the moment impulse during the stance
2 phase and the mean number of steps/day¹⁸. Daily cumulative moment may be particularly important,
3 as it was nearly doubled in the patients with knee OA compared with the healthy individuals¹⁹, and
4 daily cumulative hip moment was associated with JSW in patients with hip OA in cross-sectional
5 studies²⁰.

6

7 The purpose of this study was to evaluate the association between mechanical load during gait
8 at baseline and subsequent radiographic progression of hip OA over 12 months. Given that cartilage
9 degeneration depends on load magnitude and duration^{21,22}, it is possible that mechanical load
10 during gait, especially daily cumulative hip moment rather than the peak moment and moment
11 impulse, could critically influence degeneration of hip joint. We hypothesized that daily cumulative
12 hip moment at baseline is associated with radiographic progression of hip OA.

13

14

15 **PATIENTS AND METHODS**

16

17 **Patients**

18

19 In this prospective cohort study, non-surgical outpatients were selected in the Department of
20 Orthopaedic Surgery at Kyoto University Hospital. Patients with secondary hip OA aged 20 years
21 and older were recruited from April 2013 to March 2015. A total of 53 patients were eligible for
22 inclusion in our study, and were measured at baseline. Three patients were excluded from analysis
23 because of missing measurements 12 months later.

24

25 The inclusion criteria were as follows: 1) a diagnosis of preosteoarthritis (acetabular dysplasia
26 with no other abnormal radiographic findings) or early (slight joint space narrowing and abnormal
27 subchondral sclerosis) or advanced-stage (marked joint space narrowing with or without cysts or
28 sclerosis) hip OA, and 2) ability to walk without any assistive device in daily life. The exclusion
29 criteria were as follows: 1) patients with a baseline JSW of < 0.5 mm, as more than 0.5 mm/year in

1 JSW was defined as progression of hip OA; 2) a history of previous hip surgeries (e.g., osteotomy,
2 arthroplasty); and 3) neurologic, vascular, or other conditions that affect gait or activity of daily
3 living.

4
5 Although the candidates for our study included both males and females, our sample was biased
6 in gender (percentage of males: 7.1%), similar to previous reports on secondary hip OA (percentage
7 of males: 7.6–9.2%)^{8,23,24}. Therefore, only female patients were included in this study. Many of the
8 patients had bilateral hip OA, and the side on which the radiographic OA change was more severe
9 was used for analysis. All participants provided informed consent, and the protocol was approved
10 by the Ethics Committee of the Kyoto University Graduate School and Faculty of Medicine
11 (protocol identification number: E1683).

12 13 **Radiographic assessment**

14
15 A digital supine anteroposterior pelvic radiograph was obtained in a standardized manner by the
16 same skilled radiology technicians at baseline and approximately 12 months later. The influence of
17 position (supine versus standing) on the radiographic parameters of hip joint is discrepant^{25–27}.
18 However, radiographic parameters regarding hip dysplasia and joint space width differ little
19 between supine and standing anteroposterior radiographs^{25,27}. Therefore, to improve image quality,
20 we used radiograph in the supine position. Radiography at baseline was performed within 30 days
21 prior to gait analysis. To avoid unnecessary radiation exposure, we used radiographs taken for
22 general practice. From the radiograph, a single experienced examiner measured joint space width
23 (JSW) to assess degeneration, Sharp angle, lateral center edge (CE) angle, acetabular head index
24 (AHI), and acetabular roof obliquity (ARO) to assess morphologic abnormalities. These
25 measurements had high inter- and intrarater reliability^{28,29}, and are commonly used to diagnose
26 dysplasia and hip OA²⁹. Images were reviewed and measured on Centricity Enterprise Web, version
27 3.0 (GE Health care, Buckinghamshire, England). The JSW was measured at three locations, lateral
28 margin of the subchondral sclerotic line, apical transection of the weight-bearing surface by a
29 vertical line through the center of femoral head, and medial margin of the weight-bearing surface

1 bordering on the fovea, in 0.1 mm increments from an image magnified 4 times (Fig. 1). If the
2 minimum JSW was found aside from the 3 locations in the weight-bearing area, JSW of the
3 narrowest point was also recorded as a fourth measurement. According to previous research⁴,
4 minimum JSW was defined as the smallest of these 3 or 4 measurements. The intrarater reliability
5 [intraclass correlation (ICC) 1,1] of each radiographic measurement for 20 randomly selected
6 radiographs was 0.95 to 0.99.

7

8 To assess the change in JSW, films at baseline and approximately 12 months later were paired
9 by patients but blinded as to patient and sequence to the reader to avoid bias, as recommended²⁹.
10 All radiographic measurements were performed by the same examiner. Radiographic progression of
11 hip OA has been defined as a reduction of more than 0.5 mm in JSW based on minimum detectable
12 change (MDC) of the JSW^{30,31}. Although the MDC₉₅ (MDC at 95% confidence level) of the JSW in
13 the current study was 0.39 mm by using the formula ($MDC_{95} = \text{standard error of measurement} \times \sqrt{2}$
14 $\times 1.96$), we defined reduction of more than 0.5mm/year in JSW at any of the 3 or 4 locations as hip
15 OA progression.

16

17 (Fig. 1)

18

19 **Pain and functional assessment**

20

21 The average hip pain during daily life in the last 3 months was assessed on a 100-mm visual
22 analog scale (0 = no pain and 100 = the worst imaginable pain). The Harris hip score was recorded
23 to overview the functional status of the patient. Pain and functional assessment were conducted on
24 the day of gait analysis.

25

26 **Gait analysis**

27

28 Gait-related variables were recorded using an 8-camera Vicon motion system (Vicon Nexus;
29 Vicon Motion Systems Ltd. Oxford, England), at a sampling rate of 200 Hz with a fourth-order

1 Butterworth low-pass filter with a 6-Hz cut-off, and using force plates embedded flush with the
2 floor (Kistler Japan Co., Ltd. Tokyo, Japan), at a sampling rate of 1,000 Hz with a low-pass filter
3 (20 Hz). Patients were clothed in close-fitting shorts and T-shirts, and were asked to walk at a
4 self-selected speed without assistive devices. To closely match usual daily walking, patients were
5 given several practice trials before recording. The start position was adjusted so that participants
6 could step on the force plate naturally. At least 3 successful trials for each patient were recorded for
7 analysis.

8

9 Reflective markers were placed by a single experienced examiner. A total of 20 markers were
10 placed bilaterally on the anterior superior iliac spine, posterior superior iliac spine, superior aspect
11 of the greater trochanter, lateral femoral condyle, medial femoral condyle, lateral malleolus, medial
12 malleolus, heel, fifth metatarsal head, and first metatarsal head. The pelvic segment contained 4
13 markers placed at the bilateral anterior superior iliac spine and posterior superior iliac spine. The
14 thigh segment had 3 markers placed at the superior aspect of the greater trochanter and the medial
15 and lateral femoral condyles. The shank segment had 4 markers placed at the medial and lateral
16 femoral condyles and the medial and lateral malleoli.

17

18 We calculated 3-dimensional external joint moments of the hip using BodyBuilder software
19 (Vicon Motion Systems Ltd. Oxford, England). The joint center of the hip was determined by first
20 calculating a vector linking both greater trochanter markers. The joint center was then determined
21 at a point interpolated at a distance of 18% of the vector norm from each reflective marker of the
22 superior aspect of the greater trochanter along the vector³². The joint moment was calculated using
23 a link segment model in which segments were connected together at nodal points. To compute the
24 joint moment, coordinate data were added to the ground reaction force data, in which the position
25 of the center of mass, weight portion, and moment of inertia of each segment were used as
26 parameters. The peak external hip joint moment and hip joint moment impulse (area under the
27 moment-time curve), were calculated for stance phase in each of the three planes (Fig. 2). Although
28 the normalized value of the joint moment during gait is useful for group comparison, it can distract
29 attention from the actual load on the joint³³. Therefore, in the context of the purpose of evaluating

1 the association between mechanical load during gait and hip OA progression, non-normalized
2 values were used in moment peak and impulse according to a previous study¹⁹. Mean values of
3 gait-related variables from 3 trials were calculated and used for analysis.

4

5 (Fig. 2)

6

7 **Daily cumulative hip moment**

8

9 A pedometer (EX-500, Yamasa Tokei Co. Ltd., Tokyo, Japan) with validated accuracy was
10 given to all patients after being instructed in its use on the day of gait analysis^{34,35}. We confirmed
11 that the pedometers we used had good accuracy ($\pm 2.8\%$) when worn inside the pockets of the pants.
12 Patients were asked to wear the pedometer from the time of awaking until the time of sleeping, both
13 indoors and outdoors. The number of steps was recorded for 7 consecutive typical days within a
14 month from the day of gait analysis. The duration of extraordinary events such as illness or
15 traveling were excluded. We received the record of the number of steps via mail. Three to 5 days
16 are believed to be required to reliably assess habitual physical activity³⁶. However, we recorded
17 steps throughout the entire week in consideration of differences between individuals regarding the
18 balance of work days and non-work days within a week³⁷. Daily cumulative hip moment was
19 calculated as a product of the non-normalized hip moment impulse in each of the three planes and
20 the mean number of steps/day for the affected limb (number of steps/day divided by 2)²⁰.

21

22 **Statistical analysis**

23

24 SPSS version 19.0 (IBM Japan Ltd., Tokyo, Japan) was used for statistical analysis. Normality
25 of data was assessed using Shapiro-Wilk test. To test the hypothesis, univariable and multivariable
26 logistic regression analyses with likelihood ratio tests were used to identify predictors of hip OA
27 progression. The dependent variable was radiographic progression of hip OA (yes/no). Univariable
28 logistic regression was performed to estimate each odds ratio (OR) and accompanying 95%
29 confidence interval (CI). Multivariable logistic regression was performed to assess the association

1 between load-related parameters during gait (i.e., peak hip moment, hip moment impulse, and daily
2 cumulative hip moment) and radiographic progression of hip OA. Furthermore, as age, body weight,
3 and minimum JSW at baseline can be potential confounders^{2,20,38,39}, those 3 variables were included
4 in the multivariable model. Variables correlated at absolute coefficients > 0.7 were defined as
5 multicollinearity⁴⁰. A P value < 0.05 was considered statistical significant.

6
7

8 **RESULTS**

9

10 Baseline characteristics of patients are presented in Table I. Of the 50 patients, 21 (42.0%) were
11 classified into the progression group. Change in JSW in the progression group was 1.3 ± 0.8 mm.

12

13 In the univariable logistic regression analysis (Table II), higher daily cumulative hip moment in
14 the frontal plane at baseline was statistically significantly associated with progression of hip OA
15 (crude OR [95% CI], 1.23 [1.01 to 1.49]; $P = 0.038$; Fig 3). Minimum JSW (crude OR, 0.68 [0.45
16 to 1.03]; $P = 0.066$) and steps/day (crude OR, 1.26 [0.99 to 1.61]; $P = 0.062$) were also potential
17 predictors of hip OA progression.

18

19 In the multivariable analysis, higher daily cumulative hip moment in the frontal plane was
20 statistically significantly associated with radiographic hip OA progression even after adjustment for
21 age, body weight, and minimum JSW (adjusted OR, 1.34 [1.06 to 1.70]; $P = 0.013$; Table II). In
22 addition, higher daily cumulative hip moment in the sagittal plane was also approaching statistical
23 significance in its association with progression of hip OA (adjusted OR, 1.80 [0.99 to 3.26], $P =$
24 0.052 ; Table II). No statistically significant association was found between the peak and impulse of
25 the hip moment and hip OA progression.

26

27 There was no multicollinearity between variables. No outlier defined as its residual outside 3
28 standard deviations was found. Although only 21 patients were included in the progression group,
29 even the multivariable model (i.e., 4 independent variables) fulfilled the rule of a minimum of 5

1 events per variable⁴¹.

2

3 (Table I)

4 (Table II)

5 (Fig. 3)

6

7

8 **DISCUSSION**

9

10 The most important finding of this study was that higher daily cumulative hip moment at
11 baseline, particularly in the frontal plane, was a statistically significant independent predictor of hip
12 OA progression. The finding supports our hypothesis, and to our knowledge, this study is the first
13 to reveal an association between mechanical load during gait and radiographic progression of hip
14 OA. The ratio of the patients in progression group (42.0%) was nearly the same as the ratio (34.5%)
15 reported in a previous study although the study defined reduction of more than 0.6 mm/year in JSW
16 as progression⁴².

17

18 We included peak moment, moment impulse, and daily cumulative hip moment in load-related
19 parameters during gait, since each variable can estimate hip joint load during gait from different
20 aspects. Daily cumulative hip moment, rather than peak moment and moment impulse, was found
21 as a predictor of hip OA progression. This indicates that the physical activity during a day as well
22 as product of magnitude and duration of loading during a gait cycle must be considered for hip
23 loading. The mean value of steps/day in the progression group (i.e., $7,411 \pm 2,869$ steps/day) was
24 slightly higher than the age- and gender-specific standard value of steps/day in the same country
25 (i.e., $7,373 \pm 3,807$ steps/day)⁴³. Although it is not really known what type of stress causes cartilage
26 damage provoking joint degeneration, repetitive loading even at the same level as that during level
27 walking kill articular cartilage chondrocytes⁴⁴. Repetitive loading can cause microdamage to
28 accumulate; consequently, chondrocyte apoptosis can be induced, much like fatigue failure in
29 engineering materials. Patients with more steps/day may accelerate the progression of hip OA by

1 increased hip loading associated with excessive physical activity. Lifestyle changes through pacing
2 of physical activity would be needed for such patients as it has been regarded as one of the key
3 elements of non-pharmacological core management of OA⁴⁵.

4
5 In the 3-dimensional daily cumulative hip moments, that in the frontal plane was found as a
6 statistically significant predictor of hip OA progression, and that in the sagittal plane was also a
7 potential predictor of hip OA progression. In the hip moment impulses that compose cumulative hip
8 moment, that in the frontal plane was largest (66.7% of the total); subsequently, that in the sagittal
9 plane was large (25.7% of the total). Furthermore, change in hip contact force can be predicted by
10 the change in hip moments in the frontal and sagittal planes. The first peak of the hip contact force
11 during gait can be predicted well even by only the hip adduction moment, and combining the hip
12 adduction and flexion moments increased coefficients of determination at the second peak hip
13 contact force¹⁷. Therefore, it seems reasonable that daily cumulative hip moment, particularly in the
14 frontal plane, was the important factor among the load-related parameters during gait related to hip
15 OA progression. While body weight can directly affect joint moment during gait, daily cumulative
16 hip moment in the frontal plane remained in the multivariable logistic model even after adjustment
17 for body weight. This suggests the importance of change in gait pattern that increases the hip
18 moment impulse compared with increases in joint loading due to overweight. Daily cumulative hip
19 moment in the frontal plane is modifiable by gait modification (e.g., wide-based gait¹⁷ and lateral
20 trunk lean⁴⁶), which can reduce hip adduction moment, and/or avoid excessive physical activity.
21 Future studies should include a daily cumulative load modifying interventional trial to assess the
22 causal relationship between joint load and hip OA progression more closely.

23
24 In the previous cross-sectional study²⁰, higher daily cumulative hip moment was associated
25 with wider minimum JSW. However, in this cohort study, daily cumulative hip moment was higher
26 and minimum JSW was narrower in the progression group than in the no-progression group. These
27 findings seem contradictory; however, it would attribute to differences in dependent variables in the
28 statistical analysis, not contradict. Daily cumulative hip moment was associated with hip OA
29 progression even after adjustment for minimum JSW, which was also a potential predictor. It can be

1 interpreted that each of the higher daily cumulative hip moment and narrower minimum JSW was
2 an independent predictor when the hip OA progression was a dependent variable. The risk of
3 progression of hip OA may be particularly high in the patients with both higher daily cumulative
4 hip moment and narrower minimum JSW.

5
6 The finding that minimum JSW at baseline was a potential predictor of hip OA progression is
7 consistent with those of previous studies, where the narrower the JSW at baseline, the faster the
8 progression of hip OA³ and the higher the need for THA². This association can be explained by the
9 finding that subchondral sclerosis associated with JSW narrowing and cartilage degeneration results
10 in increased cartilage stress and pressure⁴⁷. Patients with less JSW at baseline would potentially
11 have tissue alterations which would hasten hip OA progression, and hip OA progression might have
12 already begun in those patients at baseline in this study.

13
14 Several limitations to this study should be noted. Because the daily cumulative hip moment
15 calculated in this study only reflects loading during steady waking, loading during other movement
16 such as lifting, standing, and stair climbing may have been underestimated. However, it is difficult
17 technically to measure the magnitude and duration of joint loading in daily life. Daily cumulative
18 hip moment can be estimated by using both 3-dimensional gait analysis systems and pedometers;
19 thus, it is difficult to measure the daily cumulative hip moment easily in clinical settings. More
20 time- and cost-effective sensors that can measure cumulative joint load need to be developed in the
21 future. The follow-up duration of 12 months was minimal. Although the yearly mean narrowing of
22 the hip JSW has been reported as a risk factor for hastening of THA⁴⁸, a longer follow-up would be
23 needed to establish the relationship between hip joint loading and degenerative change in articular
24 cartilage. Furthermore, the change in the daily cumulative hip moment in 12 months was not
25 measured; thus, which change in hip loading during gait affects radiographic change in hip joint is
26 not yet known. Patients with hip JSW of < 0.5 mm were excluded from this study. The findings of
27 this study could not be generalized to the relationship between gait biomechanics and hip OA
28 progression in patients with end-stage OA. In addition, as this study included patients with
29 secondary hip OA to reduce the heterogeneity of the study population, different predictors for hip

1 OA progression will be found in patients with primary hip OA, although primary hip OA is rare⁷.
2 Nevertheless, the secondary hip OA group may show heterogeneity in our study. Identification of
3 the predictor of hip OA progression might be necessary with respect to each subgroup of secondary
4 hip OA. Finally, we used a logistic regression analysis to estimate the adjusted OR because relative
5 risk could not be calculated in some of the independent variables. However, the OR underestimates
6 or overestimates the relative risk when the event being modeled is not rare (>10%)⁴⁹.

7

8 In conclusion, this study revealed that the higher daily cumulative hip moment at baseline,
9 particularly in the frontal plane, was associated with the radiographic progression of hip OA
10 defined by a > 0.5-mm cartilage thickness loss in 12 months. Our findings may help identify
11 patients with a higher risk of hip OA progression and clarify the target of intervention to slow hip
12 OA progression.

13

14

15 **Acknowledgements**

16

17 The authors would like to thank all staffs and patients for participation in this study. We would
18 also like to thank Junji Katsuhira, Rui Tsukagoshi, and Yoshihiro Fukumoto for their assistance in
19 acquisition of data and the kinematic and kinetic analysis.

20

21 **Author Contributions**

22

23 HT: concept and design, obtaining of funding, analysis and interpretation of the data, drafting of
24 the article, and final approval of the article. HT was the main investigator of this study, and
25 performed all of the measurements.

26 YK: acquisition of data, analysis and interpretation of data, critical revision of the article for
27 important intellectual content, and final approval of the article.

28 HA: provision of study patients, analysis and interpretation of data, critical revision of the
29 article for important intellectual content, and final approval of the article.

1 KG: provision of study patients, analysis and interpretation of data, critical revision of the
2 article for important intellectual content, and final approval of the article.

3 KS: provision of study patients, analysis and interpretation of data, critical revision of the
4 article for important intellectual content, and final approval of the article.

5 YK: provision of study patients, analysis and interpretation of data, critical revision of the
6 article for important intellectual content, and final approval of the article.

7 NI: concept and design, obtaining of funding, analysis and interpretation of data, critical
8 revision of the article for important intellectual content, and final approval of the article.

9

10 **Funding source**

11

12 This study was supported by the JSPS KAKENHI Grant-in-Aid for Scientific Research (C)
13 Grant No. 24500578.

14

15 **Competing interest**

16

17 There are no conflicts of interest to declare with regard to this study.

18

19

20 **References**

21

22 1. Sandell LJ. Etiology of osteoarthritis: genetics and synovial joint development. *Nat Rev*
23 *Rheumatol* 2012;8:77–89.

24 2. Lieveense AM, Bierma-Zeinstra SMA, Verhagen AP, Verhaar JA, Koes BW. Prognostic factors
25 of progress of hip osteoarthritis: A systematic review. *Arthritis Rheum* 2002;47:556–62.

26 3. Reijman M, Hazes JM, Pols HA, Bernsen RM, Koes BW, Bierma-Zeinstra SM. Role of
27 radiography in predicting progression of osteoarthritis of the hip: prospective cohort study.
28 *BMJ* 2005;330:1183–5.

29 4. Jacobsen S, Sonne-Holm S. Hip dysplasia: a significant risk factor for the development of hip

- 1 osteoarthritis. A cross-sectional survey. *Rheumatology* 2005;44:211–8.
- 2 5. Nicholls AS, Kiran A, Pollard TC, Hart DJ, Arden CP, Spector T, et al. The association between
3 hip morphology parameters and nineteen-year risk of end-stage osteoarthritis of the hip.
4 *Arthritis Rheum* 2011;63:3392–400.
- 5 6. Hochberg MC. Do risk factors for incident hip osteoarthritis (OA) differ from those for
6 progression of hip OA? *J Rheumatol Suppl* 2004;70:6–9.
- 7 7. Hoaglund FT. Primary osteoarthritis of the hip: A genetic disease caused by European genetic
8 variants. *J Bone Joint Surg Am* 2013;95:463–8.
- 9 8. Aoki H, Nagao Y, Ishii S, Masuda T, Beppu M. Acetabular and proximal femoral alignment in
10 patients with osteoarthritis of the dysplastic hip and its influence on the progression of disease.
11 *J Bone Joint Surg Br* 2010;92-B:1703–9.
- 12 9. Miyazaki T, Wada M, Kawahara H, Sato M, Baba H, Shimada S. Dynamic load at baseline can
13 predict radiographic disease progression in medial compartment knee osteoarthritis. *Ann*
14 *Rheum Dis* 2002;61:617–22.
- 15 10. Thorp LE, Sumner DR, Block JA, Moision KC, Shott S, Wimmer MA. Knee joint loading
16 differs in individuals with mild compared with moderate medial knee osteoarthritis. *Arthritis*
17 *Rheum* 2006;54:3842–9.
- 18 11. Bennell KL, Bowles KA, Wang Y, Cicuttini F, Davies-Tuck M, Hinman RS. Higher dynamic
19 medial knee load predicts greater cartilage loss over 12 months in medial knee osteoarthritis.
20 *Ann Rheum Dis* 2011;70:1770–4.
- 21 12. Chehab EF, Favre J, Erhart-Hledik JC, Andriacchi TP. Baseline knee adduction and flexion
22 moments during walking are both associated with 5 year cartilage changes in patients with
23 medial knee osteoarthritis. *Osteoarthritis Cartilage* 2014;22:1833–9
- 24 13. Chang AH, Moision KC, Chmiel JS, Eckstein F, Guermazi A, Prasad PV, et al. External knee
25 adduction and flexion moments during gait and medial tibiofemoral disease progression in
26 knee osteoarthritis. *Osteoarthritis Cartilage* 2015;23:1099–106.
- 27 14. Sulsky SI, Carlton L, Bochmann F, Ellegast R, Glitsch U, Hartmann B, et al. Epidemiological
28 evidence for work load as a risk factor for osteoarthritis of the hip: A systematic review.
29 *PlosOne* 2012;7:e31521.

- 1 15. Eitzen I, Fernandes L, Kallerud H, Nordsletten L, Knarr B, Risberg MA. Gait characteristics,
2 symptoms, and function in persons with hip osteoarthritis: A longitudinal study with 6 to 7
3 years of follow-up. *J Orthop Sports Phys Ther* 2015;45:539–49.
- 4 16. Foucher KC, Hurwitz DE, Wimmer MA. Relative importance of gait vs. joint positioning on
5 hip contact forces after total hip replacement. *J Orthop Res* 2009;27:1576–82.
- 6 17. Wesseling M, de Groot F, Meyer C, Corten K, Simon JP, Desloovere K, et al. Gait alterations
7 to effectively reduce hip contact forces. *J Orthop Res* 2015;33:1094–102.
- 8 18. Maly MR. Abnormal and cumulative loading in knee osteoarthritis. *Curr Opin Rheumatol*
9 2008;20:547–52.
- 10 19. Maly MR, Robbins SM, Stratford PW, Birmingham TB, Callaghan JP. Cumulative knee
11 adductor load distinguishes between healthy and osteoarthritic knees. A proof of principle study.
12 *Gait Posture* 2013;37:397–401.
- 13 20. Tateuchi H, Koyama Y, Tsukagoshi R, Kuroda Y, So K, Goto K, et al. Association of
14 radiographic degeneration and pain with daily cumulative hip loading in patients with
15 secondary hip osteoarthritis. *J Orthop Res* 2016, Epub ahead of print.
- 16 21. Chen CT, Bhargava M, Lin PM, Torzilli PA. Time, stress, and location dependent chondrocyte
17 death and collagen damage in cyclically loaded articular cartilage. *J Orthop Res* 2003;21:888–
18 98.
- 19 22. Roemhildt ML, Beynon BD, Gauthier AE, Gardner-Morse M, Ertem F, Badger GJ. Chronic in
20 vivo load alteration induces degenerative changes in the rat tibiofemoral joint. *Osteoarthritis*
21 *Cartilage* 2013;21:346–57.
- 22 23. Nakamura J, Oinuma K, Ohtori S, Watanabe A, Shigemura T, Sasho T, et al. Distribution of hip
23 pain in osteoarthritis patients secondary to developmental dysplasia of the hip. *Mod Rheumatol*
24 2013;23:119–24.
- 25 24. Okano K, Takaki M, Okazaki N, Shindo H. Bilateral incidence and severity of acetabular
26 dysplasia of the hip. *J Orthop Sci* 2008;13:401–4.
- 27 25. Pessis E, Chevrot A, Drapé JL, Leveque C, Sarazin L, Minoui A, et al. Study of the joint space
28 of the hip on supine and weight-bearing digital radiographs. *Clin Radiol*. 1999;54:528–32.
- 29 26. Troelsen A, Jacobsen S, Rømer L, Søballe K. Weightbearing anteroposterior pelvic radiographs

- 1 are recommended in DDH assessment. *Clin Orthop Relat Res* 2008;466:813–9.
- 2 27. Terjesen T, Gunderson RB. Reliability of radiographic parameters in adults with hip dysplasia.
3 *Skeletal Radiol* 2012;41:811–6.
- 4 28. Nelitz M, Guenther KP, Gunkel S, Puhl W. Reliability of radiological measurements in the
5 assessment of hip dysplasia in adults. *Br J Radiol* 1999;72:331–4.
- 6 29. Reijman M, Hazes JMW, Koes BW, Verhagen AP, Bierma-Zeinstra SM. Validity, reliability,
7 and applicability of seven definitions of hip osteoarthritis used in epidemiological studies: a
8 systematic appraisal. *Ann Rheum Dis* 2004;63:226–32.
- 9 30. Altman RD, Bloch DA, Dougados M, Hochberg M, Lohmander S, Pavelka K, et al.
10 Measurement of structural progression in osteoarthritis of the hip: the Barcelona consensus
11 group. *Osteoarthritis Cartilage* 2004;12:512–24.
- 12 31. Ornetti P, Brand K, Hellio-Le Graverand MP, Hochberg M, Hunter DJ, Kloppenburg M, et al.
13 OARSI-OMERACT definition of relevant radiological progression in hip/knee osteoarthritis.
14 *Osteoarthritis Cartilage* 2009;17:856–63.
- 15 32. Kito N, Shinkoda K, Yamasaki T, Kanemura N, Anan M, Okanishi N, et al. Contribution of
16 knee adduction moment impulse to pain and disability in Japanese women with medial knee
17 osteoarthritis. *Clin Biomech* 2010;25:914–9.
- 18 33. Browning RC, Kram R. Effects of obesity on the biomechanics of walking at different speeds.
19 *Med Sci Sports Exerc* 2007;39:1632–41.
- 20 34. Crouter SE, Schneider PL, Karabulut M, Bassett DR Jr. Validity of 10 electronic pedometers
21 for measuring steps, distance, and energy cost. *Med Sci Sports Exerc* 2003;35:1455–60.
- 22 35. Schneider PL, Crouter SE, Bassett DR Jr. Pedometer measures of free-living physical activity:
23 Comparison of 13 models. *Med Sci Sports Exerc* 2004;36:331–5.
- 24 36. Trost SG, Mciver KL, Pate RR. Conducting accelerometer-based activity assessments in
25 field-based research. *Med Sci Sports Exerc* 2005;37:S531–43.
- 26 37. Thorp LE, Orozco D, Block JA, Sumner DR, Wimmer MA. Activity levels in healthy older
27 adults: Implications for joint arthroplasty. *ISRN Orthop* 2012;1:pii 727950.
- 28 38. Leyton P, Muir K, Doherty S, Doherty M. Age and sex differences in hip joint space among
29 asymptomatic subjects without structural change: implications for epidemiologic studies.

- 1 Arthritis Rheum 2003;48:1041–6.
- 2 39. Recnik G, Kralj-Iglič V, Iglič A, Antolic V, Kramberger S, Rigler I, et al. The role of obesity,
3 biomechanical constitution of the pelvis and contact joint stress in progression of hip
4 osteoarthritis. *Osteoarthritis Cartilage* 2009;17:879–82.
- 5 40. Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, et al. Collinearity: a review of
6 methods to deal with it and a simulation study evaluating their performance. *Ecography*
7 2013;36:27–46.
- 8 41. Vittinghoff E, McCulloch CE. Relaxing the rule of ten events per variable in logistic and cox
9 regression. *Am J Epidemiol* 2007;165:710–8.
- 10 42. Chevalier X, Conrozier T, Gehrman M, Claudepierre P, Mathieu P, Unger S, et al. Tissue
11 inhibitor of metalloprotease-1 (TIMP-1) serum level may predict progression of hip
12 osteoarthritis. *Osteoarthritis Cartilage* 2001;9:300–7.
- 13 43. Inoue S, Ohya Y, Tudor-Locke C, Tanaka S, Yoshiike N, Shimomitsu T. Time trends for
14 step-determined physical activity among Japanese adults. *Med Sci Sports Exerc* 2011;43:1913–
15 9.
- 16 44. Clements KM, Bee ZC, Crossingham GV, Adams MA, Sharif M. How severe must repetitive
17 loading be to kill chondrocytes in articular cartilage? *Osteoarthritis Cartilage* 2001;9:499–507.
- 18 45. Fernandes L, Hagen KB, Bijlsma JW, Andreassen O, Christensen P, Conaghan PG, et al.
19 EULAR recommendations for the non-pharmacological core management of hip and knee
20 osteoarthritis. *Ann Rheum Dis* 2013;72:1125–35.
- 21 46. Hunt MA, Simic M, Hinman RS, Bennell KL, Wrigley TV. Feasibility of a gait retraining
22 strategy for reducing knee joint loading: increased trunk lean guided by real-time biofeedback.
23 *J Biomech* 2011;15:943–7.
- 24 47. Speirs AD, Beaulé PE, Ferguson SJ, Frei H. Stress distribution and consolidation in cartilage
25 constituents is influenced by cyclic loading and osteoarthritic degeneration. *J Biomech*
26 2014;47:2348–53.
- 27 48. Conrozier T, Jousseume CA, Mathieu P, Tron AM, Caton J, Bejui J, et al. Quantitative
28 measurement of joint space narrowing progression in hip osteoarthritis: a longitudinal
29 retrospective study of patients treated by total hip arthroplasty. *Br J Rheumatol* 1998;37:961–8.

- 1 49. Davies HTO, Crombie IK, Tavakoli M. Who can odds ratios mislead? *BMJ* 1998;316:989–91.

Table I. Baseline characteristics of study participants

(Footnotes for Table 1)

Values are reported as mean \pm standard deviation. VAS = visual analogue scale; JSW = joint space width; CE angle = center edge angle; AHI = acetabular head index; ARO = acetabular roof obliquity.

	All patients (n = 50)	No progression (n = 29)	Progression (n = 21)
Age, years	47.4 \pm 10.7	46.6 \pm 10.2	48.6 \pm 11.6
Weight, kg	55.2 \pm 10.2	54.2 \pm 9.8	56.5 \pm 10.9
Height, cm	156.9 \pm 5.6	157.5 \pm 6.8	156.1 \pm 3.5
Minimum JSW, mm	3.3 \pm 1.4	3.7 \pm 1.4	2.9 \pm 1.4
Pain (VAS), mm	42.0 \pm 27.5	37.7 \pm 1.4	47.9 \pm 26.4
Harris hip score (total 100 points)	86.9 \pm 9.9	87.9 \pm 8.7	85.6 \pm 11.4
Morphology parameters			
Sharp angle, degrees	45.0 \pm 6.5	45.6 \pm 7.4	44.1 \pm 4.8
CE angle, degrees	23.4 \pm 11.5	22.0 \pm 11.1	25.5 \pm 12.1
AHI, degrees	73.8 \pm 11.0	72.8 \pm 10.7	75.2 \pm 11.6
ARO, degrees	22.4 \pm 7.9	22.8 \pm 8.6	21.8 \pm 7.0
Gait-related parameters			
Gait speed, meters/seconds	1.1 \pm 0.2	1.2 \pm 0.2	1.1 \pm 0.1
Steps/day	6,596 \pm 2,552	6,005 \pm 2,157	7,411 \pm 2,869
Load-related parameters during gait			
Peak external hip moment, Nm			
Hip flexion moment	39.1 \pm 9.8	39.5 \pm 8.9	38.6 \pm 11.2
Hip extension moment	25.5 \pm 8.7	25.3 \pm 7.3	25.7 \pm 10.5
Hip adduction moment	57.9 \pm 16.0	56.9 \pm 16.2	59.3 \pm 16.0
Hip internal rotation moment	9.0 \pm 3.9	9.5 \pm 3.5	8.3 \pm 4.3
Hip external rotation moment	6.2 \pm 2.5	6.1 \pm 2.4	6.4 \pm 2.6
Hip moment impulse, Nm•seconds			
Sagittal plane	8.4 \pm 2.7	8.4 \pm 2.1	8.5 \pm 3.5
Frontal plane	22.7 \pm 7.4	21.4 \pm 7.3	24.5 \pm 7.2
Transversal plane	2.5 \pm 0.8	2.5 \pm 0.7	2.5 \pm 1.0
Cumulative hip joint moment, kNm•seconds			
Sagittal plane	27.6 \pm 13.4	25.2 \pm 10.3	30.8 \pm 16.6
Frontal plane	74.6 \pm 41.4	63.0 \pm 29.4	90.6 \pm 50.2
Transversal plane	8.4 \pm 5.4	7.6 \pm 3.7	9.5 \pm 7.0

Table II. Univariable and multivariable logistic regression predicting the progression of hip osteoarthritis (n = 50)
(Footnotes for Table 2)

OR = odds ratio; 95% CI = 95% confidence interval. VAS = visual analogue scale; JSW = joint space width; CE angle = center edge angle; AHI = acetabular head index; ARO = acetabular roof obliquity.

* Unit is 1,000 steps/day. † Adjusted for age, body weight, and minimum JSW.

	Crude OR (95% CI)	P value	Adjusted OR† (95% CI)	P value
Age, years	1.02 (0.97 to 1.08)	0.499	–	–
Weight, kg	1.02 (0.97 to 1.08)	0.426	–	–
Minimum JSW, mm	0.68 (0.45 to 1.03)	0.066	–	–
Pain (VAS), mm	1.01 (0.99 to 1.04)	0.198	–	–
Morphology parameters				
Sharp angle, degrees	0.96 (0.88 to 1.06)	0.407	–	–
CE angle, degrees	1.03 (0.98 to 1.08)	0.286	–	–
AHI, degrees	1.02 (0.97 to 1.08)	0.447	–	–
ARO, degrees	0.98 (0.92 to 1.06)	0.654	–	–
Gait-related variables				
Steps/day*	1.26 (0.99 to 1.61)	0.062	–	–
Gait speed, meters/seconds	0.13 (0.00 to 4.73)	0.268	–	–
Load-related parameters during gait				
Peak external hip moment, Nm				
Hip flexion moment	0.99 (0.94 to 1.05)	0.758	0.99 (0.93 to 1.05)	0.640
Hip extension moment	1.01 (0.94 to 1.07)	0.877	0.99 (0.92 to 1.06)	0.701
Hip adduction moment	1.01 (0.97 to 1.05)	0.597	0.99 (0.94 to 1.05)	0.760
Hip internal rotation moment	0.92 (0.78 to 1.07)	0.278	0.88 (0.73 to 1.05)	0.160
Hip external rotation moment	1.05 (0.84 to 1.33)	0.670	1.03 (0.80 to 1.34)	0.799
Hip moment impulse, Nm•seconds				
Sagittal plane	1.01 (0.82 to 1.24)	0.950	1.00 (0.80 to 1.26)	0.971
Frontal plane	1.06 (0.98 to 1.15)	0.155	1.09 (0.96 to 1.24)	0.190
Transversal plane	0.90 (0.45 to 1.80)	0.774	0.83 (0.36 to 1.91)	0.663
Cumulative hip joint moment, 10kNm•seconds				
Sagittal plane	1.39 (0.88 to 2.21)	0.159	1.80 (0.99 to 3.26)	0.052
Frontal plane	1.23 (1.01 to 1.49)	0.038	1.34 (1.06 to 1.70)	0.013
Transversal plane	2.01 (0.61 to 6.68)	0.253	2.93 (0.71 to 12.11)	0.253

1 **Figure legends**

2

3 **Fig 1.** The three measurement locations of the joint space width (JSW) of the hip joint. If the
4 minimum JSW was found aside from the three locations, it was recorded as a fourth measurement.

5

6 **Fig. 2.** A typical example of the hip joint moment curve in sagittal (thin black line), frontal (thick
7 black line), and transversal (grey line) plane during stance phase of walking. Positive values
8 indicate external hip flexion, adduction, and external rotation moment, respectively.

9

10 **Fig. 3.** Distribution of daily cumulative hip joint moment in the sagittal (A), frontal (B), and
11 transversal plane (C) in each of no progression group (white) and progression group (grey).
12 Boxplots with upper and lower bars showing maximum and minimum values. Upper, middle, and
13 lower lines in the box indicate 75th, 50th (median), and 25th centiles, respectively. The cross mark
14 in the box indicates the mean value.




Fig. 1.




Fig. 2.




Fig. 3.