



Title	Loss of Bacitracin Resistance Due to a Large Genomic Deletion among Bacillus anthracis Strains
Author(s)	Furuta, Yoshikazu; Harima, Hayato; Ito, Emiko; Maruyama, Fumito; Ohnishi, Naomi; Osaki, Ken; Ogawa, Hirohito; Squarre, David; Hang'ombe, Bernard Mudenda; Higashi, Hideaki
Citation	Msystems, 3(5), e00182-18 <a href="https://doi.org/10.1128/mSystems.00182-18">https://doi.org/10.1128/mSystems.00182-18</a>
Issue Date	2018-09
Doc URL	<a href="http://hdl.handle.net/2115/72165">http://hdl.handle.net/2115/72165</a>
Rights	Copyright © 2018 Furuta et al.
Rights(URL)	<a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>
Type	article
File Information	e00182-18.full.pdf



[Instructions for use](#)



# Loss of Bacitracin Resistance Due to a Large Genomic Deletion among *Bacillus anthracis* Strains

 Yoshikazu Furuta,<sup>a</sup> Hayato Harima,<sup>b</sup> Emiko Ito,<sup>a</sup>  Fumito Maruyama,<sup>c</sup> Naomi Ohnishi,<sup>a\*</sup> Ken Osaki,<sup>d\*</sup> Hirohito Ogawa,<sup>b\*</sup> David Squarre,<sup>e</sup> Bernard Mudenda Hang'ombe,<sup>f</sup> Hideaki Higashi<sup>a</sup>

<sup>a</sup>Division of Infection and Immunity, Research Center for Zoonosis Control, Hokkaido University, Sapporo, Japan

<sup>b</sup>Hokudai Center for Zoonosis Control in Zambia, Research Center for Zoonosis Control, Hokkaido University, Sapporo, Japan

<sup>c</sup>Department of Microbiology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

<sup>d</sup>TOMY Digital Biology Co., Ltd., Tokyo, Japan

<sup>e</sup>Department of National Parks and Wildlife, Chilanga, Zambia

<sup>f</sup>Department of Para-clinical Studies, School of Veterinary Medicine, University of Zambia, Lusaka, Zambia

**ABSTRACT** *Bacillus anthracis* is a Gram-positive endospore-forming bacterial species that causes anthrax in both humans and animals. In Zambia, anthrax cases are frequently reported in both livestock and wildlife, with occasional transmission to humans, causing serious public health problems in the country. To understand the genetic diversity of *B. anthracis* strains in Zambia, we sequenced and compared the genomic DNA of *B. anthracis* strains isolated across the country. Single nucleotide polymorphisms clustered these strains into three groups. Genome sequence comparisons revealed a large deletion in strains belonging to one of the groups, possibly due to unequal crossing over between a pair of rRNA operons. The deleted genomic region included genes conferring resistance to bacitracin, and the strains with the deletion were confirmed with loss of bacitracin resistance. Similar deletions between rRNA operons were also observed in a few *B. anthracis* strains phylogenetically distant from Zambian strains. The structure of bacitracin resistance genes flanked by rRNA operons was conserved only in members of the *Bacillus cereus* group. The diversity and genomic characteristics of *B. anthracis* strains determined in this study would help in the development of genetic markers and treatment of anthrax in Zambia.

**IMPORTANCE** Anthrax is caused by *Bacillus anthracis*, an endospore-forming soil bacterium. The genetic diversity of *B. anthracis* is known to be low compared with that of *Bacillus* species. In this study, we performed whole-genome sequencing of Zambian isolates of *B. anthracis* to understand the genetic diversity between closely related strains. Comparison of genomic sequences revealed that closely related strains were separated into three groups based on single nucleotide polymorphisms distributed throughout the genome. A large genomic deletion was detected in the region containing a bacitracin resistance gene cluster flanked by rRNA operons, resulting in the loss of bacitracin resistance. The structure of the deleted region, which was also conserved among species of the *Bacillus cereus* group, has the potential for both deletion and amplification and thus might be enabling the species to flexibly control the level of bacitracin resistance for adaptive evolution.

**KEYWORDS** *Bacillus anthracis*, *Bacillus cereus* group, antibiotic resistance, bacitracin, genome analysis, rRNA operon, unequal crossing over

*Bacillus anthracis* is an endospore-forming Gram-positive bacterium that causes anthrax, a worldwide zoonotic disease (1). Among animals, the bacterium mainly infects herbivores and causes anthrax in livestock such as cattle, sheep, and goats, as

Received 23 August 2018 Accepted 4 October 2018 Published 30 October 2018

**Citation** Furuta Y, Harima H, Ito E, Maruyama F, Ohnishi N, Osaki K, Ogawa H, Squarre D, Hang'ombe BM, Higashi H. 2018. Loss of bacitracin resistance due to a large genomic deletion among *Bacillus anthracis* strains. *mSystems* 3:e00182-18. <https://doi.org/10.1128/mSystems.00182-18>.

**Editor** Yu-Liang Yang, Agricultural Biotechnology Research Center

**Copyright** © 2018 Furuta et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Yoshikazu Furuta, [yfuruta@czc.hokudai.ac.jp](mailto:yfuruta@czc.hokudai.ac.jp).

\* Present address: Naomi Ohnishi, Japanese Foundation for Cancer Research, Tokyo, Japan; Ken Osaki, 10x Genomics, Pleasanton, California, USA; Hirohito Ogawa, Department of Virology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan.

well as in wildlife such as kudus, elephants, and hippopotamuses (2). Contact with infected animals or materials contaminated with *B. anthracis* results in the transmission of anthrax to humans. Two plasmids of *B. anthracis*, pXO1 and pXO2, carry genes producing virulence factors and capsule synthetases, respectively (3). The pXO1 plasmid harbors genes encoding a protective antigen, lethal factor, and edema factor, all of which contribute to virulence (4), whereas pXO2 harbors an operon encoding proteins for synthesis of a poly- $\gamma$ -D-glutamate capsule that increases the stability of cells in a natural environment (3).

In Zambia, anthrax cases are observed every year, and outbreaks occur once every few years. Anthrax cases occur in both livestock in communities and wildlife in national parks, with occasional cases of infection among humans (5, 6). Most recent cases of human infection and animal infection were observed in an outbreak in the eastern part of Zambia in 2011, with more than 500 cases of human infection and five deaths, following the occurrence of an anthrax outbreak among hippopotamuses (7).


Several genetic markers have been used for the phylogenetic analysis of *B. anthracis* strains (8), including amplified fragment length polymorphisms (AFLPs) (9), multilocus variable-number tandem repeats (10, 11), single nucleotide polymorphisms (SNPs) (12), canonical SNPs (13, 14), and whole-genome sequences (15), although the species is known to have low genetic diversity (16, 17). Recent analysis using high-throughput short read sequencers enabled comparison of genome-wide polymorphisms and revealed phylogenetic clusters and the distribution and migration history of species at a higher resolution (18, 19). The results of this whole-genome analysis are consistent with phylogenetic groups, previously defined as A, B, and C, and additionally revealed subgroups and diversity within these phylogenetic groups (20–22).

To assess the genetic diversity of *B. anthracis* strains in Zambia in detail, we performed whole-genome sequencing of *B. anthracis* strains collected from Zambia, including isolates from recent cases of anthrax in animals and humans. Phylogenetic analysis of these strains revealed three groups among closely related strains based on genome-wide SNPs. Despite the characteristics of few genome rearrangements in the species (23), a large genomic deletion within a pair of rRNA operons was detected among strains belonging to one of the three groups, resulting in the loss of bacitracin resistance. A similar genomic deletion was also observed in other *B. anthracis* strains, as well as in species of the *Bacillus cereus* group, suggesting that such a deletion may contribute to the flexible control of the level of bacitracin resistance and to the adaptive evolution of *B. anthracis* strains.

## RESULTS

**SNPs and a large genomic deletion in Zambian *B. anthracis* strains.** To elucidate the diversity in *B. anthracis* strains in Zambia, genomic DNA of 14 Zambian strains sampled from various locations and sources and over several years (Fig. 1A and Table 1) was sequenced using the Illumina MiSeq platform. Sequence reads from all 14 strains were mapped on the reference genome sequence of the *B. anthracis* Ames ancestor to detect SNPs. A total of only 883 SNPs were identified, which is consistent with the low genetic diversity of *B. anthracis* species (see Table S1 in the supplemental material). Of the 883 SNPs, 409 were detected in all 14 Zambian strains. The remaining 474 core SNPs were therefore used for phylogenetic analysis of the Zambian strains. The Zambian strains were clustered into three groups showing some relationship with the sampling locations; these three groups were named Zambia1, Zambia2a, and Zambia2b (Fig. 1B).

In addition to SNPs, other genomic characteristics differentiating the Zambian strains were searched across the whole-genome sequences, and a large genomic deletion was identified specifically among the Zambia1 strains. Mapping of short reads of Zambian strains on the genome sequence of the *B. anthracis* Ames ancestor revealed a stretch of approximately 5 kb of unmapped genomic region specifically among the Zambia1 strains (Fig. 2A; see Fig. S1 in the supplemental material). Deletion of this region in Zambia1 strains was further confirmed by sequencing one of the Zambia1



**FIG 1** Single nucleotide polymorphisms (SNPs) in *B. anthracis* strains isolated in Zambia. (A) Map of Zambia and sampling locations of *B. anthracis* strains. (B) Heat map of SNPs in each *B. anthracis* strain isolated in Zambia. Gray indicates the presence of SNPs. The genome sequence of the *B. anthracis* Ames ancestor strain was used as reference. The color of strains corresponds to the color of the sampling locations in panel A.

strains, CZC5, using the PacBio RS platform, which generated long reads spanning the deleted region (see Fig. S2 in the supplemental material).


The deleted region was flanked by rRNA operons, which exist as 11 copies per genome of typical *B. anthracis* strains (Fig. 2B). Most of the rRNA operons harbored 16S, 23S, and 5S rRNA genes with the same order, orientation, and nucleotide sequence, suggesting that the deletion was caused by unequal crossing over between two long direct repeats of rRNA operons (24) (Fig. 2C). The flanking pair of rRNA operons was annotated as *rrnG* and *rrnH*. We, therefore, named the deletion DelGH.

Five genes deleted by DelGH were annotated as encoding undecaprenyl-diphosphatase, subunits of an ABC transporter, and members of a two-component system (Fig. 2D). The undecaprenyl-diphosphatase gene has sequence similarity to *uppP* of *Escherichia coli* (56% amino acid identity), which is known to confer resistance to bacitracin, a dodecapeptide antibiotic produced by *Bacillus licheniformis* and *Bacillus*

**TABLE 1** *Bacillus anthracis* strains isolated in Zambia

Strain	Location <sup>a</sup>	Source	Yr	Read depth	Accession no.	Group
NL2016_1	L.V.	Soil	2016	723	DRR118317	Zambia1
NL2016_2	L.V.	Soil	2016	848	DRR118318	Zambia1
NL2016_3	L.V.	Soil	2016	695	DRR118319	Zambia1
CZC5	L.V.	Hippopotamus	2011	171	DRR014735	Zambia1
Han19	L.V.	Soil	2012	50	DRR125654	Zambia1
Mongu1	Mongu	Cattle	2012	69	DRR125653	Zambia2a
Mongu37	Mongu	Cattle	2013	161	DRR014741	Zambia2a
Human29	Chama	Human	2013	207	DRR014739	Zambia2b
Mox4	Chama	Hippopotamus	2013	138	DRR014740	Zambia2b
LZprevious	L.Z.	Elephant	2011	161	DRR014736	Zambia2b
LZ2013T	L.Z.	Elephant	2011	98	DRR014737	Zambia2b
LZ88	L.Z.	Elephant	2013	70	DRR125657	Zambia2b
LZ3	L.Z.	Elephant	2013	59	DRR125655	Zambia2b
LZ131001	L.Z.	Elephant	2013	65	DRR125656	Zambia2b

<sup>a</sup>L.V., Luangwa Valley; L.Z., Lower Zambezi.




**FIG 2** Large genomic deletion between rRNA operons in Zambia1 *B. anthracis* strains. (A) Sequence coverage of short reads of the CZC5 strain mapped to the reference genome sequence of the Ames ancestor. Arrows above the chart indicate open reading frames in the corresponding genomic regions of the Ames ancestor chromosome from bp 260000 to 300000. Arrows of different colors indicate protein coding regions (with the protein product in parentheses) or rRNA as follows: green, rRNA operons; blue, *uppP* (undecaprenyl-diphosphatase); pink, *bcrB* (ABC transporter permease); red, *bcrA* (ABC transporter ATP-binding protein); yellow, *bacS* (sensor histidine kinase); orange, *bacR* (DNA-binding response regulator); white, other genes. (B) Positions of 11 rRNA operons in the Ames ancestor strain. Genomic DNA is shown in linear form. The bottom panel magnifies the genomic region encompassing the rRNA operons A to H. rRNA operons G and H, which flank the deleted region, are indicated in red. (C) Schematic representation of the deletion via unequal crossing over between rRNA operons present as 4.7-kb-long direct repeats. (D) Deletion of the genomic region flanked by rRNA operons G and H. Genes are indicated by the number of the locus tag that in the full locus tag designation follows the prefix “GBAA\_RS” for the Ames ancestor or “BAZ\_” for CZC5.

*subtilis* (25). The ABC transporter genes and the two-component system genes also showed sequence similarity to *bcrAB* and *bacRS*, respectively, both of which sense bacitracin and regulate bacitracin resistance in *B. licheniformis* (26, 27). Hence, strains showing the deletion of these five genes were speculated to be more sensitive to bacitracin than strains without the deletion.

**Loss of bacitracin resistance by DelGH.** To confirm the influence of DelGH on bacitracin resistance, disk diffusion tests were conducted using representative Zambian isolates. Zambia1 strains showed an inhibitory zone of approximately 20 mm, whereas Zambia2a and Zambia2b strains showed no inhibitory zones (Fig. 3; see Fig. S3 in the supplemental material). This suggests that the deletion of genes conferring bacitracin resistance (DelGH) rendered the Zambia1 strains susceptible to bacitracin.

We further analyzed the contribution of each gene in the deleted region by strains constructed by artificially introducing DelGH to *B. anthracis* vaccination strain 34F2 (Table 2). First, the MICs of bacitracin were compared between the wild-type 34F2 strain and DelGH mutant (BYF10028). This showed an approximately 19-fold lower MIC in BYF10028 (Fig. 4A), supporting the results of the disk test for Zambian isolates. Next, BYF10028 was complemented with each gene or combination of genes in the deleted region to test the contribution of these genes to bacitracin resistance (Fig. 4B). Complementation of a single gene or a combination of ABC transporter and two-component system genes did not show any recovery of bacitracin resistance. Recovery of bacitracin resistance approximately to the level of the 34F2 strain was observed only with the complementation of a combination of *uppP* and both ABC transporter genes, suggesting that these three genes are required for bacitracin resistance.



**FIG 3** Bacitracin disk test of Zambian *B. anthracis* strains. (A) Mean diameter of the inhibition zone. The error bar indicates standard deviation ( $n = 3$ ). No inhibition zone was observed among Zambia2a and Zambia2b strains. (B) An example of bacitracin disk test of CZC5. (C) An example of bacitracin disk test of Mongu37.

**Occurrence of large genomic deletions during the evolution of *B. anthracis*.** To understand the phylogenetic position of Zambian strains among *B. anthracis* strains, genome-wide SNPs were compared with genome sequences of other non-Zambian *B. anthracis* strains (Fig. 5; see Table S2 in the supplemental material). The results showed that Zambian strains clustered in the same clade, together with the K3 strain isolated in South Africa.

We also searched for DelGH and genomic deletions between other pairs of rRNA operons in non-Zambian *B. anthracis* strains and identified such a deletion within a region flanked by different pairs of rRNA operons, *rrnF* and *rrnG*, which we here refer to as DelFG. DelFG resulted in the deletion of genes annotated as encoding class I S-adenosylmethionine (SAM)-dependent methyltransferase, RNA methyltransferase, and glycosyl transferase (see Fig. S4 in the supplemental material). All of these genes are involved in the modification of nucleotides, proteins, and membrane lipids. However, the targets and types of modification of these enzymes were unclear; thus, it is difficult to speculate about the influence of DelFG on the phenotype.

Among the non-Zambian *B. anthracis* strains, DelFG was observed in BA1015, BA1035, HYU01, RA3, and SVA11, whereas DelGH was observed in H9401 and PAK-1. These deletions were confirmed by mapping short reads of strains to the reference genome sequence of the Ames ancestor (see Fig. S5 in the supplemental material). Most of the strains with the same deletion grouped in a single clade: strains RA3, HYU01, SVA11, and BA1035 carrying the DelFG deletion formed one cluster, and Zambia1 strains carrying the DelGH deletion grouped in another clade. This implies the occurrence of single deletion event in a common ancestor, followed by vertical transmission of the deletion. Three strains, including H9401 and PAK-1 with DelGH and BA1015 with DelFG, were phylogenetically distant from both the DelFG and DelGH clades, suggesting that these deletions occurred at a low frequency but multiple times during the course of evolution of *B. anthracis*.

**Structure of a bacitracin resistance gene cluster in *Bacillus* species.** To address whether the deletion of the whole bacitracin resistance gene cluster possibly via unequal crossing over between rRNA operons occurred only in *B. anthracis* or could be found in other *Bacillus* species, we searched for homologs of genes within this cluster in genomic sequences of strains of *Bacillus* species (see Table S3 in the supplemental material).

The structure of the bacitracin resistance gene cluster flanked by rRNA operons was found only in strains of the *B. cereus* group, including *B. anthracis* (Fig. 6). More than



**TABLE 2** Strains and plasmids for bacitracin MIC measurement


Strain or plasmid	Description	Reference
Strains		
<i>E. coli</i>		
SM10	<i>thi thr leu tonA lacY supE recA</i> RP4-2-Tc::Mu (Km)	64
BYF684	SM10 harboring pYF684	This study
S17.1	<i>thi pro hsdR</i> RP4-2-Tc::Mu Km::Tn7 (Tp Sm)	64
BYF701	S17.1 harboring pRP1099	This study
BYF794	S17.1 harboring pYF794	This study
BYF796	S17.1 harboring pYF796	This study
BYF797	S17.1 harboring pYF797	This study
BYF798	S17.1 harboring pYF798	This study
BYF799	S17.1 harboring pYF799	This study
BYF800	S17.1 harboring pYF800	This study
BYF801	S17.1 harboring pYF801	This study
BYF802	S17.1 harboring pYF802	This study
BYF803	S17.1 harboring pYF803	This study
<i>B. anthracis</i>		
34F2	<i>B. anthracis</i> vaccine strain	62
BYF10028	34F2 with the region between <i>rrnG</i> and <i>rrnH</i> deleted (DelGH)	This study
BYF10031	BYF10028 harboring pYF794	This study
BYF10033	BYF10028 harboring pYF798	This study
BYF10034	BYF10028 harboring pYF799	This study
BYF10035	BYF10028 harboring pYF800	This study
BYF10036	BYF10028 harboring pYF801	This study
BYF10037	BYF10028 harboring pYF796	This study
BYF10038	BYF10028 harboring pYF797	This study
BYF10039	BYF10028 harboring pYF802	This study
BYF10040	BYF10028 harboring pYF803	This study
Plasmids		
pRP1028	Vector for introduction of I-SceI recognition site	61
pYF684	pRP1028 cloned with sequence around <i>bcrB</i> of 34F2	This study
pRP1099	Vector for introduction of the gene encoding I-SceI	61
pYF794	pRP1099 without I-SceI and AmCyan	This study
pYF796	pYF794 with cloned <i>uppP</i> , <i>bcrB</i> , and <i>bcrA</i> of 34F2	This study
pYF797	pYF794 with cloned <i>bcrB</i> and <i>bcrA</i> of 34F2	This study
pYF798	pYF794 with cloned <i>bcrA</i> of 34F2	This study
pYF799	pYF794 with cloned <i>bacS</i> and <i>bacR</i> of 34F2	This study
pYF800	pYF794 with cloned <i>bacR</i> of 34F2	This study
pYF801	pYF794 with cloned <i>uppP</i> of 34F2	This study
pYF802	pYF794 with cloned <i>bcrB</i> of 34F2	This study
pYF803	pYF794 with cloned <i>bacS</i> of 34F2	This study

60% of strains of each species of the *B. cereus* group were found with the gene cluster flanked by rRNA operons. Thus, it is likely that the bacitracin resistance gene operon was present in the common ancestor of the *B. cereus* group but was lost in some strains during the evolution of each species of the *B. cereus* group.

Other *Bacillus* species either did not possess the cluster or possessed it only at a very low frequency. For example, only two *Bacillus amyloliquefaciens* strains and one *B. subtilis* strain were found to possess the cluster, suggesting that these species acquired the cluster at a very low frequency. In addition, the *uppP* gene was substituted for with the *bcrC* gene homolog, whose product has the same activity of undecaprenyl-diphosphatase but is composed of different domain sequences (26, 28). Differences in the relative positions of the two-component system genes were also observed in *Bacillus vallismortis*, *B. amyloliquefaciens*, *Bacillus velezensis*, and *B. subtilis*. Taken together, the structure of bacitracin resistance gene cluster flanked by rRNA operons and the presence of *uppP* homolog within the cluster was specifically conserved among species of the *B. cereus* group.

## DISCUSSION

In this study, we conducted genome sequencing of Zambian isolates of *B. anthracis* to determine their genetic diversity. Results revealed three groups based on genome-




**FIG 4** Effect of the deletion between rRNA operons G and H (DelGH) on bacitracin resistance. (A) MICs of the 34F2 vaccine strain and the derivative strain introduced with DelGH. Plus and minus signs indicate the presence and absence of bacitracin resistance genes, respectively, on the chromosome of each strain. (B) MICs of strains with DelGH complemented with bacitracin resistance gene(s). Plus and minus signs indicate the presence and absence of bacitracin resistance genes, respectively, on complementing plasmids. All strains in this panel lack all the bacitracin resistance genes on their chromosome. The error bar indicates the standard deviation ( $n = 3$ ).

wide SNPs and the deletion of a large genomic region encompassing the bacitracin resistance genes, possibly via unequal crossing over between a pair of rRNA operons flanking the region. Experiments with natural isolates and artificially constructed strains of *B. anthracis* confirmed that the deletion of bacitracin resistance gene cluster results in the loss of bacitracin resistance. The deletion of this gene cluster was also identified in a few other non-Zambian *B. anthracis* strains, and the structure of this genomic region was highly conserved only in species of the *B. cereus* group.


All of the Zambian strains of *B. anthracis* in this study were classified in the A.Br.005/006 group, according to the canonical SNPs (see Table S4 in the supplemental material) and clustered with the K3 strain isolated in South Africa, according to the clustering based on genome-wide SNPs (Fig. 5). These results were consistent with previous works on Zambian *B. anthracis* strains (13, 22, 29). Within the closely related Zambian strains, genome-wide SNPs clustered the strains into three groups—Zambia1, Zambia2a, and Zambia2b—which showed some relationship with the sampling location. The genomic deletion resulted in a clear phenotypic difference between Zambia1 and the other groups, but not between the Zambia2a and Zambia2b groups. This information on the genetic diversity and phenotypic differences among *B. anthracis* strains would be useful for the development of genetic markers to differentiate among Zambian *B. anthracis* strains. Additionally, these markers would help in determining the strains' group in the event of an anthrax outbreak in Zambia in the future.





**FIG 5** Phylogenetic tree of Zambian and non-Zambian strains of *B. anthracis*. Zambian strains are indicated in red and bold letters. Strains with the DelFG deletion are indicated by "FG" in a white circle. Strains with the DelGH deletion are indicated by "GH" in a black circle.

The large genomic deletion DelGH included the bacitracin resistance gene cluster and resulted in a significant loss of bacitracin resistance, suggesting bacitracin as an alternative antibiotic when treating anthrax caused by Zambia1 strains. Bacitracin is a cyclic peptide antibiotic, which inhibits bacterial cell wall synthesis by sequestering undecaprenyl-pyrophosphate, an essential carrier lipid (25). Anthrax is usually treated with antibiotics such as benzylpenicillin, amoxicillin, doxycycline, and ciprofloxacin or ofloxacin (30, 31). To our knowledge, *B. anthracis* infections in animals and in humans are quite rarely treated with bacitracin in Zambia and in other countries. Reports of *B. anthracis* strains resistant to antibiotics are rare; however, some studies have shown that long-term treatment of *B. anthracis* *in vitro* with increasing amounts of antibiotics results in strains with higher MICs for various antibiotics, including fluoroquinolones



**FIG 6** Structure of the bacitracin resistance gene cluster in *Bacillus* species. The phylogenetic tree was constructed based on the 16S rRNA sequences of each species. Numbers following the names of strains indicate the number of strains of the species possessing the bacitracin resistance gene cluster out of the number of strains analyzed.

(32–36). Therefore, it is important to prepare for and suppress the emergence of antibiotic-resistant strains by having an alternative choice of antibiotics, although bacitracin treatment might be limited to cutaneous anthrax because of its nephrotoxic effect (37).

In *Bacillus*, two main gene clusters conferring bacitracin resistance are known: one of these clusters is present in species of the *B. cereus* group, including *B. anthracis*, which includes the genes *bacRS* and *bcrAB*, and the other cluster is present in other *Bacillus* species, such as *B. subtilis*, which contains *bceRSAB* (38, 39). Both gene clusters consist of two-component system (*bacRS* and *bceRS*) and ABC transporter (*bcrAB* and *bceAB*) genes. The cluster of *bacRS* and *bcrAB* usually possesses another gene with the function of the undecaprenyl-diphosphatase gene, *uppP* or *bcrC*, immediately adjacent to the four genes (Fig. 6). Genes with sequence similarity to *uppP* or *bcrC* are also found in the genome of species with *bceRSAB* but are usually not adjacent to the cluster. UppP and BcrC have the same activity as undecaprenyl-diphosphatase, but belong to a different protein superfamily (25, 40). It is difficult to determine how the two genes *uppP* and *bcrC* were swapped in the immediate vicinity of the *bacRS* and *bcrAB* cluster during the evolution of *Bacillus* species (Fig. 6).

The activity and regulation of proteins encoded by the *bacRS*, *bcrAB*, and *uppP* cluster have been studied less than those encoded by the *bceRSAB* cluster, which is present in the model organism *B. subtilis* (38, 39, 41–45). Undecaprenyl-diphosphatase activity of UppP has been shown to confer resistance to bacitracin when the *E. coli* homolog of *uppP* was overexpressed in *E. coli* cells (46); however, this was not the case in *B. anthracis* (Fig. 4B). We observed full recovery of bacitracin resistance only when UppP was expressed together with BcrA and BcrB in the strain with DelGH. In the *bceRSAB* cluster, it is hypothesized that the ABC transporter consists of BceA and BceB and functions as a flippase to flip the undecaprenyl-pyrophosphate and bacitracin complex to the outer leaflet of the membrane, thus, promoting the reaction of membrane proteins that utilize bacitracin as a substrate (43). If the ABC transporter that consists of BcrA and BcrB also possesses such flippase activity, the ABC transporter may promote the activity of UppP, which is also a membrane protein (25), thus, leading to a complete recovery of bacitracin resistance. On the other hand, BacR, the response regulator of the *bacRS* two-component system, has been shown to act as a negative regulator of the *bcrA* gene in *B. licheniformis* (27). This is consistent with our results showing that genes of the two-component system are not required for the recovery of bacitracin resistance in strains carrying the DelGH deletion (Fig. 4).

Deletion of the bacitracin resistance gene cluster was identified in several *B. anthracis* strains and among species of the *B. cereus* group. This deletion was most likely caused by unequal crossing over between a pair of rRNA operons flanking the gene cluster. Genome rearrangement due to unequal crossing over between direct repeats is a well-studied phenomenon in bacteria (47). This results not only in gene deletion but also gene amplification and an increase in the expression of genes in the amplified region (47). The rRNA operon has been shown to cause amplification of the flanking region in *Salmonella enterica* serovar Typhimurium and *E. coli* (48, 49). Such genome amplification has been suggested as a mechanism to increase the antibiotic resistance of bacteria, thus, increasing their ability to survive under high antibiotic pressure (50). For example, amplification of antibiotic resistance genes due to flanking insertion sequences approximately 1 kb in size has increased the resistance of *E. coli* to specific drugs (51–54). Decreased antibiotic pressure results in the reduction of copy numbers of the amplified region, again by unequal crossing over, to revert the copy number before amplification, thus maximizing the fitness of bacteria. The bacitracin resistance gene cluster in species of the *B. cereus* group may also have experienced both deletion and amplification to flexibly control the level of bacitracin resistance to survive in the environment with bacitracin secreted by different *Bacillus* species (25), although further experiments, such as culturing of *B. anthracis* strains *in vitro* under gradually increasing concentrations of bacitracin, are required to confirm the plasticity and contribution of the gene cluster.

In conclusion, our sequencing analysis of Zambian isolates of *B. anthracis* revealed three groups with different SNPs and large genomic deletions affecting bacitracin resistance. The pair of rRNA operons flanking the bacitracin resistance gene cluster are an ideal target for unequal crossing over, which not only leads to the deletion of the whole region but also has the potential for amplification of the region, to possibly enhance bacitracin resistance. Although further experimental confirmation is required, our data suggest that this gene composition contributes to the evolution of *B. anthracis* strains by controlling the bacitracin resistance phenotype for adaptation.

## MATERIALS AND METHODS

***B. anthracis* natural isolates.** *B. anthracis* strains were isolated from various hosts from different locations in Zambia in different years (Table 1). All of the Zambian isolates were handled at the BSL3 facility at the Hokudai Center for Zoonosis Control in Zambia. Samples from hippopotamuses (*Hippopotamus amphibius*), African elephants (*Loxodonta africana*), and cattle (*Bos taurus*) were collected from tissues and fluid of carcasses. Soil samples (approximately 25 mg) were collected from a depth of 15 cm. Isolation of *B. anthracis* strains from samples was performed as described previously (55). Briefly, 1 g of a specimen was suspended in 10 ml of sterilized saline and incubated at 75°C for 20 min to enrich for spores, followed by spreading the samples on brain heart infusion (BHI) agar plates with 10% (vol/vol) sheep blood agar. Colonies with a “medusa head” were identified as *B. anthracis*. These were inoculated in LB medium and incubated 37°C overnight on a shaker. The resulting cultures were stored at –80°C as glycerol stocks. For genomic DNA extraction, glycerol stocks were inoculated and streaked on BHI agar plates with 10% sheep blood and incubated at 37°C for 16 h. Colonies were harvested and used for genomic DNA extraction with the QIAamp PowerFecal DNA kit (Qiagen, Hilden, Germany).

A *B. anthracis* strain isolated from a human case of anthrax was provided by the School of Veterinary Medicine, University of Zambia, Zambia.

**Sequencing and genome analysis of Zambian strains.** For whole-genome sequencing, 10 ng of genomic DNA was used to prepare genomic sequencing libraries using the Nextera DNA library preparation kit (Illumina, San Diego, CA) and sequenced using the Illumina MiSeq platform (Illumina, San Diego, CA). Paired-end reads with a read length of 300 bp were obtained. These short reads were used for SNP detection, reconstruction of a consensus sequence, and detection of genomic deletion by mapping to the reference genome (chromosome, pXO1, and pXO2) of the *B. anthracis* Ames ancestor (RefSeq accession no. NC\_007530.2, NC\_007322.2, and NC\_007323.3, respectively) using Snippy v3.2 (<https://github.com/tseemann/snippy>), which maps reads using bwa mem (56) and performs variant calling using Freebayes (57) with default parameters.

To obtain long reads from strain CZC5, 5 µg of genomic DNA of CZC5 was used to construct a sequencing library with the SMRTbell Template Prep kit (Pacific Biosciences, Menlo Park, CA) and sequenced using the PacBio RS platform (Pacific Biosciences, Menlo Park, CA) with a 360-min movie time. This generated sequence reads of approximately 130 Mbp, with an average read length of 5,732 bp. *De novo* assembly was also conducted for CZC5 on the SMRT portal using the pipeline script RS\_HGAP\_Assembly. The assembled sequences were corrected with short reads, produced by the Illumina MiSeq platform as described above, by mapping the short reads to the assembled sequences using bwa mem with the default parameter, followed by using Pilon (58) with “–changes –fix snps,indels,gaps” options. This procedure resulted in changes of 19 single-nucleotide indels and one SNP.

For visualization of DelGH, long reads were mapped to the reference genome of the *B. anthracis* Ames ancestor and CZC5 using minimap2 v2.5 (59) with default parameters, followed by visualization with IGV v2.3.97 (60).

**Bacitracin disk test.** Glycerol stock of each *B. anthracis* strain was inoculated on a BHI plate with 10% (vol/vol) sheep blood and incubated at 37°C for 16 h. Bacterial colonies were harvested, resuspended in phosphate-buffered saline, and diluted to an optical density at 600 nm (OD<sub>600</sub>) of approximately 0.2, followed by plating of 300 µl on blood agar plates. A bacitracin disk (Sigma-Aldrich, St. Louis, MO; catalog no. 08382-50DISCS-F [0.04 U/disk]) was placed at the center of each plate and incubated at 37°C for 16 h. CFU were measured for the resuspended and diluted samples of each strain, and samples from each strain were confirmed to have less than a 10-fold difference in CFU.

**Strain construction.** Strains were constructed to measure the MIC values (Table 2) using the markerless allelic exchange strategy (61). To construct the strain with DelGH in *B. anthracis* 34F2 (62), genomic sequences (500 bp) flanking the *bcrB* gene were amplified by PCR using genomic DNA of 34F2 as a template and inserted in pRP1028 together with the recognition site of I-SceI using Gibson assembly (63). The constructed plasmid, pYF684, was transformed into *E. coli* SM10 cells (64) and introduced into the 34F2 strain by conjugation. The strain with the plasmid integrated into its genomic DNA was selected and conjugated with pRP1099 harboring the gene expressing I-SceI from *E. coli* S17.1 (64) to induce double-strand breaks (DSBs) in the genomic DNA of the *B. anthracis* intermediate strain. The DSBs promoted homologous recombination between rRNA operons G and H flanking the entire bacitracin resistance gene cluster rather than the introduced flanking sequence of the *bcrB* gene, which resulted in the 34F2 derivative strain with DelGH, BYF10028. Strains with the *bcrB* gene or other genes of the bacitracin resistance gene cluster deleted could not be constructed, probably because of much higher probability for homologous recombination between 4.7-kb rRNA operons than between 500-bp flanking sequence of the target gene. Introduction of DelGH in BYF10028 was confirmed by sequencing on the MinION platform using flowcell R9.5 and the Rapid barcode sequencing kit SQK-RBK001 (Oxford

Nanopore Technologies, Oxford Science Park, United Kingdom). Reads were mapped onto the reference genome sequence of the *B. anthracis* Ames ancestor strain and CZC5 strain using minimap2 v2.5 (59) with default parameters (see Fig. S6 in the supplemental material).

For the complementation of bacitracin resistance genes, genes were cloned under the control of their native promoter in pRP1099 with substituting genes encoding I-SceI and AmCyan. The resulting plasmids, pYF794 to 803, were transformed into *E. coli* S17.1 cells and introduced into BYF10028 by conjugation. Attempts to introduce complementation plasmid with four genes or more into *B. anthracis* cells were unsuccessful.

**Measurement of MIC for bacitracin.** Glycerol stocks of strains were inoculated on BHI agar plates with or without 20  $\mu\text{g}/\text{ml}$  kanamycin and incubated at 37°C overnight. A single colony was isolated, inoculated in BHI with or without 20  $\mu\text{g}/\text{ml}$  kanamycin, and incubated at 37°C on a shaker overnight. Ten microliters of overnight cultures was added to 980  $\mu\text{l}$  of BHI and then added with 10  $\mu\text{l}$  of a 2-fold serial dilution series of bacitracin solution (Sigma-Aldrich, St. Louis, MO; catalog no. B0125, lot no. 115M4104V, with a potency of 77,120 U/g) to have cultures in final bacitracin concentrations of 0, 4, 8, 16, 32, 64, 128, 256, and 512  $\mu\text{g}/\text{ml}$ . Cultures were incubated at 37°C on a shaker overnight, and the OD was measured at 595 nm using Multiskan FC (Thermo Fisher Scientific, Waltham, MA). The lowest concentration of bacitracin with an OD of  $<0.1$  was determined as the MIC.

**Comparison of whole-genome sequences of non-Zambian strains.** Genome sequences of non-Zambian *B. anthracis* strains registered as a complete assembly in RefSeq as of 18 July 2018 were used for the construction of a phylogenetic tree (Table S2). Genome sequences of the strains obtained in artificial evolution experiments were not used in this analysis to focus only on natural isolates (65).

Reconstructed consensus sequences of Zambian strains and whole-genome sequences of non-Zambian *B. anthracis* strains were aligned and subjected to phylogenetic analysis by Parsnp included in the Harvest suite (66). Whole-genome sequences were compared using Mauve (67, 68) to detect genomic deletions between rRNA operons. Genome maps were plotted by genoPlotR (69). Short sequence reads of strains PAK-1 (SRR2155549), BA1015 (SRR2175366), SVA11 (SRR974942), and BA1035 (SRR2174564) were obtained from the DNA Data Bank of Japan Sequence Read Archive (DRA) database and used to confirm the large genomic deletion by mapping.

**Detection of bacitracin resistance genes in *Bacillus* species.** Genome sequences of strains of *Bacillus* species that are registered with complete genome sequences were downloaded from the RefSeq database as of 18 July 2018 (Table S3). Genome sequences were searched for bacitracin resistance genes with tblastn (70) using amino acid sequences of the following accession numbers as queries: BcrC, NP\_391534.1; UppP, WP\_001280040.1; BcrB, WP\_000247643.1; BcrA, WP\_000074565.1; BacS, WP\_000686989.1; and BacR, WP\_000651978.1. The top-most hits according to amino acid percentage of identity with an alignment length covering  $>80\%$  of the query length and with  $>40\%$  sequence similarity were assumed as homologs. Strains with four or five of the query genes in tandem, allowing insertion of up to three genes, were determined to harbor the bacitracin resistance gene cluster. A phylogenetic tree was constructed using 16S rRNA sequence of each species in MEGA7 (71) by the neighbor-joining method with the Kimura two-parameter model (72).

**Accession number(s).** DNA sequences generated in this study were deposited in the DRA database under accession no. DRR118317 to DRR118319 and DRR125653 to DRR125657 for reads of Zambian isolates using the Illumina MiSeq platform (Illumina, San Diego, CA), DRR118320 for the CZC5 strain using the PacBio RS platform (Pacific Biosciences, Menlo Park, CA), and DRR147392 and DRR147393 for the 34F2 and BYF10028 strains, respectively, using the MinION platform (Oxford Nanopore Technologies, Oxford Science Park, United Kingdom). The complete whole-genome sequence of CZC5 was deposited in GenBank under the following accession numbers: chromosome, AP018443; pXO1, AP018444; and pXO2, AP018445.

## SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mSystems.00182-18>.

**FIG S1**, TIF file, 2.4 MB.

**FIG S2**, PDF file, 0.9 MB.

**FIG S3**, PDF file, 1.5 MB.

**FIG S4**, PDF file, 0.9 MB.

**FIG S5**, PDF file, 2.7 MB.

**FIG S6**, PDF file, 1.6 MB.

**TABLE S1**, PDF file, 0.2 MB.

**TABLE S2**, PDF file, 0.1 MB.

**TABLE S3**, PDF file, 0.1 MB.

**TABLE S4**, PDF file, 0.1 MB.

## ACKNOWLEDGMENTS

We thank Roger Plaut for providing strains and plasmids to construct knockout strains of *B. anthracis*, Miwako Kitazume for conducting experiments for genome sequencing using the PacBio RS, Aiko Ohnuma for helping with genome sequencing



using the Illumina MiSeq platform, and Evans Mulenga for helping with the bacitracin disk tests. We also thank Zambia Wildlife Authority and Department of National Parks and Wildlife for their support and provision of biological materials used in this study.

This work was supported by the Japan Prize Foundation to Y.F., the Akiyama Life Science Foundation to Y.F., Grants-in-Aid for Regional R&D Proposal-Based Program from Northern Advancement Center for Science & Technology of Hokkaido Japan to Y.F., the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID) to H.H., the Ministry of Education, Culture, Sports, Science and Technology in Japan or Japan Society for the Promotion of Science under Grants-in-Aid for Scientific Research (KAKENHI) to Y.F. (grant no. 18K14672) and F.M. (grant no. 16H05501), the Ichiro Kanehara Foundation Scholarship Grant for Research in Basic Medical Sciences and Medical Care to F.M., and a Senri Life Science Foundation Kishimoto Grant to F.M. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

## REFERENCES

- Turnbull PC, Hugh-Jones ME, Cosivi O. 1999. World Health Organization activities on anthrax surveillance and control. *J Appl Microbiol* 87: 318–320. <https://doi.org/10.1046/j.1365-2672.1999.00903.x>.
- Mock M, Fouet A. 2001. Anthrax. *Annu Rev Microbiol* 55:647–671. <https://doi.org/10.1146/annurev.micro.55.1.647>.
- Moayeri M, Leppla SH, Vrentas C, Pomerantsev AP, Liu S. 2015. Anthrax pathogenesis. *Annu Rev Microbiol* 69:185–208. <https://doi.org/10.1146/annurev-micro-091014-104523>.
- Pezard C, Berche P, Mock M. 1991. Contribution of individual toxin components to virulence of *Bacillus anthracis*. *Infect Immun* 59: 3472–3477.
- Siamudaala VM, Bwalya JM, Munang'andu HM, Sinyangwe PG, Banda F, Mweene AS, Takada A, Kida H. 2006. Ecology and epidemiology of anthrax in cattle and humans in Zambia. *Jpn J Vet Res* 54:15–23.
- Tuchili LM, Pandey GS, Sinyangwe PG, Kaji T. 1993. Anthrax in cattle, wildlife and humans in Zambia. *Vet Rec* 132:487. <https://doi.org/10.1136/vr.132.19.487-a>.
- Hang'ombe MB, Mwansa JC, Muwowa S, Mulenga P, Kapina M, Musenga E, Squarre D, Mataa L, Thomas SY, Ogawa H, Sawa H, Higashi H. 2012. Human-animal anthrax outbreak in the Luangwa Valley of Zambia in 2011. *Trop Doct* 42:136–139. <https://doi.org/10.1258/td.2012.110454>.
- Keim P, Van Ert MN, Pearson T, Vogler AJ, Huynh LY, Wagner DM. 2004. Anthrax molecular epidemiology and forensics: using the appropriate marker for different evolutionary scales. *Infect Genet Evol* 4:205–213. <https://doi.org/10.1016/j.meegid.2004.02.005>.
- Keim P, Kalif A, Schupp J, Hill K, Travis SE, Richmond K, Adair DM, Hugh-Jones M, Kuske CR, Jackson P. 1997. Molecular evolution and diversity in *Bacillus anthracis* as detected by amplified fragment length polymorphism markers. *J Bacteriol* 179:818–824. <https://doi.org/10.1128/jb.179.3.818-824.1997>.
- Keim P, Price LB, Klevytska AM, Smith KL, Schupp JM, Okinaka R, Jackson PJ, Hugh-Jones ME. 2000. Multiple-locus variable-number tandem repeat analysis reveals genetic relationships within *Bacillus anthracis*. *J Bacteriol* 182:2928–2936. <https://doi.org/10.1128/JB.182.10.2928-2936.2000>.
- Marston CK, Allen CA, Beaudry J, Price EP, Wolken SR, Pearson T, Keim P, Hoffmaster AR. 2011. Molecular epidemiology of anthrax cases associated with recreational use of animal hides and yarn in the United States. *PLoS One* 6:e28274. <https://doi.org/10.1371/journal.pone.0028274>.
- Kenefic LJ, Beaudry J, Trim C, Huynh L, Zanecki S, Matthews M, Schupp J, Van Ert M, Keim P. 2008. A high resolution four-locus multiplex single nucleotide repeat (SNR) genotyping system in *Bacillus anthracis*. *J Microbiol Methods* 73:269–272. <https://doi.org/10.1016/j.mimet.2007.11.014>.
- Van Ert MN, Easterday WR, Huynh LY, Okinaka RT, Hugh-Jones ME, Ravel J, Zanecki SR, Pearson T, Simonson TS, U'Ren JM, Kachur SM, Leadem-Dougherty RR, Rhoton SD, Zinser G, Farlow J, Coker PR, Smith KL, Wang B, Kenefic LJ, Fraser-Liggett CM, Wagner DM, Keim P. 2007. Global genetic population structure of *Bacillus anthracis*. *PLoS One* 2:e461. <https://doi.org/10.1371/journal.pone.0000461>.
- Simonson TS, Okinaka RT, Wang B, Easterday WR, Huynh L, U'Ren JM, Dukerich M, Zanecki SR, Kenefic LJ, Beaudry J, Schupp JM, Pearson T, Wagner DM, Hoffmaster A, Ravel J, Keim P. 2009. *Bacillus anthracis* in China and its relationship to worldwide lineages. *BMC Microbiol* 9:71. <https://doi.org/10.1186/1471-2180-9-71>.
- Pearson T, Busch JD, Ravel J, Read TD, Rhoton SD, U'Ren JM, Simonson TS, Kachur SM, Leadem RR, Cardon ML, Van Ert MN, Huynh LY, Fraser CM, Keim P. 2004. Phylogenetic discovery bias in *Bacillus anthracis* using single-nucleotide polymorphisms from whole-genome sequencing. *Proc Natl Acad Sci U S A* 101:13536–13541. <https://doi.org/10.1073/pnas.0403844101>.
- Price LB, Hugh-Jones M, Jackson PJ, Keim P. 1999. Genetic diversity in the protective antigen gene of *Bacillus anthracis*. *J Bacteriol* 181: 2358–2362.
- Sacchi CT, Whitney AM, Mayer LW, Morey R, Steigerwalt A, Boras A, Weyant RS, Popovic T. 2002. Sequencing of 16S rRNA gene: a rapid tool for identification of *Bacillus anthracis*. *Emerg Infect Dis* 8:1117–1123. <https://doi.org/10.3201/eid0810.020391>.
- Parkhill J, Wren BW. 2011. Bacterial epidemiology and biology—lessons from genome sequencing. *Genome Biol* 12:230. <https://doi.org/10.1186/gb-2011-12-10-230>.
- Pena-Gonzalez A, Rodriguez-R LM, Marston CK, Gee JE, Gulvik CA, Kolton CB, Saile E, Frace M, Hoffmaster AR, Konstantinidis KT. 2018. Genomic characterization and copy number variation of *Bacillus anthracis* plasmids pXO1 and pXO2 in a historical collection of 412 strains. *mSystems* 3:e00065-18. <https://doi.org/10.1128/mSystems.00065-18>.
- Derzelle S, Girault G, Kokotovic B, Angen Ø. 2015. Whole genome-sequencing and phylogenetic analysis of a historical collection of *Bacillus anthracis* strains from Danish cattle. *PLoS One* 10:e0134699. <https://doi.org/10.1371/journal.pone.0134699>.
- Girault G, Blouin Y, Vergnaud G, Derzelle S. 2014. High-throughput sequencing of *Bacillus anthracis* in France: investigating genome diversity and population structure using whole-genome SNP discovery. *BMC Genomics* 15:288. <https://doi.org/10.1186/1471-2164-15-288>.
- Vergnaud G, Girault G, Thierry S, Pourcel C, Madani N, Blouin Y. 2016. Comparison of French and worldwide *Bacillus anthracis* strains favors a recent, post-Columbian origin of the predominant North-American clade. *PLoS One* 11:e0146216. <https://doi.org/10.1371/journal.pone.0146216>.
- Keim P, Gruendike JM, Klevytska AM, Schupp JM, Challacombe J, Okinaka R. 2009. The genome and variation of *Bacillus anthracis*. *Mol Aspects Med* 30:397–405. <https://doi.org/10.1016/j.mam.2009.08.005>.
- Kuzminov A. 2011. Homologous recombination—experimental systems, analysis, and significance. *EcoSal Plus* 2011. <https://doi.org/10.1128/ecosalplus.7.2.6>.
- El Ghachi M, Bouhss A, Blanot D, Mengin-Lecreux D. 2004. The *bacA* gene of *Escherichia coli* encodes an undecaprenyl pyrophosphate phosphatase activity. *J Biol Chem* 279:30106–30113. <https://doi.org/10.1074/jbc.M401701200>.
- Podlessek Z, Comino A, Herzog-Velikonja B, Zgur-Bertok D, Komel R, Grabnar M. 1995. *Bacillus licheniformis* bacitracin-resistance ABC

- transporter: relationship to mammalian multidrug resistance. *Mol Microbiol* 16:969–976. <https://doi.org/10.1111/j.1365-2958.1995.tb02322.x>.
27. Neumüller AM, Konz D, Marahiel MA. 2001. The two-component regulatory system BacRS is associated with bacitracin 'self-resistance' of *Bacillus licheniformis* ATCC 10716. *Eur J Biochem* 268:3180–3189. <https://doi.org/10.1046/j.1432-1327.2001.02203.x>.
  28. Podlesek Z, Comino A, Herzog-Velikonja B, Grabnar M. 2000. The role of the bacitracin ABC transporter in bacitracin resistance and collateral detergent sensitivity. *FEMS Microbiol Lett* 188:103–106. <https://doi.org/10.1111/j.1574-6968.2000.tb09176.x>.
  29. Fasanella A, Serrecchia L, Chiaverini A, Garofolo G, Muuka GM, Mwambazi L. 2018. Use of canonical single nucleotide polymorphism (CanSNPs) to characterize *Bacillus anthracis* outbreak strains in Zambia between 1990 and 2014. *PeerJ* 6:e5270. <https://doi.org/10.7717/peerj.5270>.
  30. Centers for Disease Control and Prevention. 2001. Update: interim recommendations for antimicrobial prophylaxis for children and breastfeeding mothers and treatment of children with anthrax. *MMWR Morb Mortal Wkly Rep* 50:1014–1016.
  31. Inglesby TV, O'Toole T, Henderson DA, Bartlett JG, Ascher MS, Eitzen E, Friedlander AM, Gerberding J, Hauer J, Hughes J, McDade J, Osterholm MT, Parker G, Perl TM, Russell PK, Tonat K, Working Group on Civilian Biodefense. 2002. Anthrax as a biological weapon, 2002: updated recommendations for management. *JAMA* 287:2236–2252. <https://doi.org/10.1001/jama.287.17.2236>.
  32. Athamna A, Athamna M, Abu-Rashed N, Medlej B, Bast DJ, Rubinstein E. 2004. Selection of *Bacillus anthracis* isolates resistant to antibiotics. *J Antimicrob Chemother* 54:424–428. <https://doi.org/10.1093/jac/dkh258>.
  33. Brook I, Elliott TB, Pryor HI, Sautter TE, Gnade BT, Thakar JH, Knudson GB. 2001. In vitro resistance of *Bacillus anthracis* Sterne to doxycycline, macrolides and quinolones. *Int J Antimicrob Agents* 18:559–562. [https://doi.org/10.1016/S0924-8579\(01\)00464-2](https://doi.org/10.1016/S0924-8579(01)00464-2).
  34. Price LB, Vogler A, Pearson T, Busch JD, Schupp JM, Keim P. 2003. In vitro selection and characterization of *Bacillus anthracis* mutants with high-level resistance to ciprofloxacin. *Antimicrob Agents Chemother* 47:2362–2365. <https://doi.org/10.1128/AAC.47.7.2362-2365.2003>.
  35. Choe CH, Bouhaouala SS, Brook I, Elliott TB, Knudson GB. 2000. In vitro development of resistance to ofloxacin and doxycycline in *Bacillus anthracis* Sterne. *Antimicrob Agents Chemother* 44:1766. <https://doi.org/10.1128/AAC.44.6.1766-1766.2000>.
  36. Bryskier A. 2002. *Bacillus anthracis* and antibacterial agents. *Clin Microbiol Infect* 8:467–478. <https://doi.org/10.1046/j.1469-0691.2002.00527.x>.
  37. Michie AJ, Zintel HA. 1949. The nephrotoxicity of bacitracin in man. *Surgery* 26:626–632.
  38. Bernard R, Joseph P, Guiseppi A, Chippaux M, Denizot F. 2003. YtsCD and YwoA, two independent systems that confer bacitracin resistance to *Bacillus subtilis*. *FEMS Microbiol Lett* 228:93–97. [https://doi.org/10.1016/S0378-1097\(03\)00738-9](https://doi.org/10.1016/S0378-1097(03)00738-9).
  39. Ohki R, Giyanto, Tateno K, Masuyama W, Moriya S, Kobayashi K, Ogasawara N. 2003. The BceRS two-component regulatory system induces expression of the bacitracin transporter, BceAB, in *Bacillus subtilis*. *Mol Microbiol* 49:1135–1144. <https://doi.org/10.1046/j.1365-2958.2003.03653.x>.
  40. Bernard R, El Ghachi M, Mengin-Lecreux D, Chippaux M, Denizot F. 2005. BcrC from *Bacillus subtilis* acts as an undecaprenyl pyrophosphate phosphatase in bacitracin resistance. *J Biol Chem* 280:28852–28857. <https://doi.org/10.1074/jbc.M413750200>.
  41. Dintner S, Heermann R, Fang C, Jung K, Gebhard S. 2014. A sensory complex consisting of an ATP-binding cassette transporter and a two-component regulatory system controls bacitracin resistance in *Bacillus subtilis*. *J Biol Chem* 289:27899–27910. <https://doi.org/10.1074/jbc.M114.596221>.
  42. Radeck J, Gebhard S, Orchard PS, Kirchner M, Bauer S, Mascher T, Fritz G. 2016. Anatomy of the bacitracin resistance network in *Bacillus subtilis*. *Mol Microbiol* 100:607–620. <https://doi.org/10.1111/mmi.13336>.
  43. Bernard R, Guiseppi A, Chippaux M, Foglino M, Denizot F. 2007. Resistance to bacitracin in *Bacillus subtilis*: unexpected requirement of the BceAB ABC transporter in the control of expression of its own structural genes. *J Bacteriol* 189:8636–8642. <https://doi.org/10.1128/JB.01132-07>.
  44. Rietkötter E, Hoyer D, Mascher T. 2008. Bacitracin sensing in *Bacillus subtilis*. *Mol Microbiol* 68:768–785. <https://doi.org/10.1111/j.1365-2958.2008.06194.x>.
  45. Kingston AW, Zhao H, Cook GM, Helmann JD. 2014. Accumulation of heptaprenyl diphosphate sensitizes *Bacillus subtilis* to bacitracin: implications for the mechanism of resistance mediated by the BceAB transporter. *Mol Microbiol* 93:37–49. <https://doi.org/10.1111/mmi.12637>.
  46. Cain BD, Norton PJ, Eubanks W, Nick HS, Allen CM. 1993. Amplification of the *bacA* gene confers bacitracin resistance to *Escherichia coli*. *J Bacteriol* 175:3784–3789. <https://doi.org/10.1128/jb.175.12.3784-3789.1993>.
  47. Reams AB, Kofoid E, Kugelberg E, Roth JR. 2012. Multiple pathways of duplication formation with and without recombination (RecA) in *Salmonella enterica*. *Genetics* 192:397–415. <https://doi.org/10.1534/genetics.112.142570>.
  48. Anderson P, Roth J. 1981. Spontaneous tandem genetic duplications in *Salmonella typhimurium* arise by unequal recombination between rRNA (*rrm*) cistrons. *Proc Natl Acad Sci U S A* 78:3113–3117. <https://doi.org/10.1073/pnas.78.5.3113>.
  49. Hill CW, Harnish BW. 1981. Inversions between ribosomal RNA genes of *Escherichia coli*. *Proc Natl Acad Sci U S A* 78:7069–7072. <https://doi.org/10.1073/pnas.78.11.7069>.
  50. Sandegren L, Andersson DI. 2009. Bacterial gene amplification: implications for the evolution of antibiotic resistance. *Nat Rev Microbiol* 7:578–588. <https://doi.org/10.1038/nrmicro2174>.
  51. Nichols BP, Guay GG. 1989. Gene amplification contributes to sulfonamide resistance in *Escherichia coli*. *Antimicrob Agents Chemother* 33:2042–2048. <https://doi.org/10.1128/AAC.33.12.2042>.
  52. Nicoloff H, Perreten V, Levy SB. 2007. Increased genome instability in *Escherichia coli* lon mutants: relation to emergence of multiple-antibiotic-resistant (Mar) mutants caused by insertion sequence elements and large tandem genomic amplifications. *Antimicrob Agents Chemother* 51:1293–1303. <https://doi.org/10.1128/AAC.01128-06>.
  53. Nicoloff H, Perreten V, McMurry LM, Levy SB. 2006. Role for tandem duplication and *lon* protease in AcrAB-TolC-dependent multiple antibiotic resistance (Mar) in an *Escherichia coli* mutant without mutations in *marRAB* or *acrRAB*. *J Bacteriol* 188:4413–4423. <https://doi.org/10.1128/JB.01502-05>.
  54. Meyer J, Iida S. 1979. Amplification of chloramphenicol resistance transposons carried by phage P1Cm in *Escherichia coli*. *Mol Gen Genet* 176:209–219.
  55. Ogawa H, Ohnuma M, Squarre D, Mweene AS, Ezaki T, Fujikura D, Ohnishi N, Thomas Y, Hang'ombe BM, Higashi H. 2015. *Bacillus cereus* from the environment is genetically related to the highly pathogenic *Bacillus cereus* in Zambia. *J Vet Med Sci* 77:993–995. <https://doi.org/10.1292/jvms.15-0059>.
  56. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
  57. Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing. *arXiv arXiv1207 [q-bio.GN]*. <http://arXiv.org/abs/1207.3907v2>.
  58. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
  59. Li H. 2017. Minimap2: fast pairwise alignment for long nucleotide sequences. *arXiv arXiv1708 [q-bio.GN]*. <https://arXiv.org/abs/1708.01492>.
  60. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011. Integrative genomics viewer. *Nat Biotechnol* 29:24–26. <https://doi.org/10.1038/nbt.1754>.
  61. Plaut RD, Stibitz S. 2015. Improvements to a markerless allelic exchange system for *Bacillus anthracis*. *PLoS One* 10:e0142758. <https://doi.org/10.1371/journal.pone.0142758>.
  62. Sterne M. 1939. The use of anthrax vaccines prepared from avirulent (uncapsulated) variants of *Bacillus anthracis*. *Onderstepoort J Vet Sci Anim Industry* 13:307–312.
  63. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* 6:343–345. <https://doi.org/10.1038/nmeth.1318>.
  64. Simon R, Priefer U, Puhler A. 1983. A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in Gram negative bacteria. *Nat Biotechnol* 1:784–791. <https://doi.org/10.1038/nbt1183-784>.
  65. Molohon KJ, Blair PM, Park S, Doroghazi JR, Maxson T, Hershfield JR, Flatt KM, Schroeder NE, Ha T, Mitchell DA. 2016. Plantazolicin is an ultra-narrow spectrum antibiotic that targets the *Bacillus anthracis* mem-



- brane. *ACS Infect Dis* 2:207–220. <https://doi.org/10.1021/acinfecdis.5b00115>.
66. Treangen TJ, Ondov BD, Koren S, Phillippy AM. 2014. The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol* 15:524. <https://doi.org/10.1186/s13059-014-0524-x>.
67. Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.
68. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <https://doi.org/10.1371/journal.pone.0011147>.
69. Guy L, Kultima JR, Andersson SG. 2010. genoPlotR: comparative gene and genome visualization in R. *Bioinformatics* 26:2334–2335. <https://doi.org/10.1093/bioinformatics/btq413>.
70. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. <https://doi.org/10.1186/1471-2105-10-421>.
71. Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33:1870–1874. <https://doi.org/10.1093/molbev/msw054>.
72. Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>.