

Elsevier Editorial System(tm) for Fish and
Shellfish Immunology
Manuscript Draft

Manuscript Number:

Title: Responses of *Mytilus galloprovincialis* to challenge with the emerging marine pathogen *Vibrio coralliilyticus*

Article Type: Full Length Article

Keywords: bivalves; pathogenic vibrios; *Vibrio coralliilyticus*; Mediterranean mussels; hemocytes; immune response; embryos

Corresponding Author: Dr. teresa balbi,

Corresponding Author's Institution: University of Genova

First Author: teresa balbi

Order of Authors: teresa balbi; Manon Auguste; Katia Cortese; Michele Montagna; Alessio Borello; Carla Pruzzo; Luigi Vezzulli; Laura Canesi

Abstract: *Vibrio coralliilyticus* (V.c.) has emerged as a coral pathogen of concern throughout the Indo-Pacific reef. The interest towards understanding its ecology and pathogenic potential has increased since V.c. was shown to be strongly virulent also for other species; in particular, it represents a serious threat for bivalve aquaculture, being one of the most important emerging pathogen responsible for oyster larval mortalities worldwide. V.c. has a tightly regulated temperature-dependent virulence and it has been related to mass mortalities events of benthic invertebrates also in the temperate north-western Mediterranean Sea. However, no data are available on the effects of V.c. in the mussel *Mytilus galloprovincialis*, the most abundant aquacultured species in this area.

In this work, responses of *M. galloprovincialis* to challenge with V.c. (ATCC BAA-450) were investigated. In vitro, short term responses of mussel hemocytes were evaluated in terms of lysosomal membrane stability, bactericidal activity, lysozyme release, ROS and NO production, and ultrastructural changes, evaluated by TEM. In vivo, hemolymph parameters were measured in mussels challenged with V.c. at 24h p.i. Moreover, the effects of V.c. on mussel early embryo development (at 48 hpf) were evaluated. The results show that both in vitro and in vivo, mussels were unable to activate immune response towards V.c., and that challenge mainly induced lysosomal stress in the hemocytes. Moreover, V.c. showed a strong and concentration-dependent embryotoxicity. Overall, the results indicate that, although *M. galloprovincialis* is considered a resistant species to vibrio infections, the emerging pathogen V.c. can represent a potential threat to mussel aquaculture.

Suggested Reviewers: Miren Cajaraville
miren.p.cajaraville@ehu.es
expert on mussel responses to environmental stress

Daniela Ceccarelli
daniela.ceccarelli@wur.nl
expert on *Vibrio*

Maria Jose Figueras
mariajose.figueras@urv.cat
expert on Vibrio

Christine Paillard
christine.paillard@univ-brest.fr
expert on pathogen bivalve interactions

Elizabeth Dyrinda
E.A.Dyrinda@hw.ac.uk
expert on invertebrate immunology

Dear Prof. Hirono,

I send you the manuscript “Responses of *Mytilus galloprovincialis* to challenge with the emerging marine pathogen *Vibrio coralliilyticus*” to be considered for publication in Fish & Shellfish Immunology.

I thank you for your kind attention and look forward to hearing from you.

Best regards,

Teresa Balbi

Dept. of Earth, Environment and Life Sciences (DISTAV)

University of Genoa

Corso Europa 26

16132 – Genoa

Italy

Teresa.Balbi@unige.it

Highlights


- Evaluation of *Mytilus* responses to the emerging pathogen *V. coralliilyticus*
- In vitro, in vivo and early developmental effects were evaluated
- *V. coralliilyticus* induced lysosomal stress in the hemocytes
- No induction of the immune response
- Strong and concentration-dependent effects on embryo development

V. coralliilyticus



M. galloprovincialis
hemocytes

M. galloprovincialis
embryo



Responses of *Mytilus galloprovincialis* to challenge with the emerging marine pathogen *Vibrio coralliilyticus*

Teresa Balbi^{a,*}, Manon Auguste^a, Katia Cortese^b, Michele Montagna^a, Alessio Borello^a, Carla Pruzzo^a, Luigi Vezzulli^a, Laura Canesi^a

^a*Dept. of Earth, Environment and Life Sciences (DISTAV), University of Genoa, Italy;*

^b*Dept. of Experimental Medicine (DIMES), University of Genoa, Italy*

Keywords: bivalves; pathogenic vibrios; *Vibrio coralliilyticus*; Mediterranean mussels; hemocytes; immune response, embryos

Running title: Effects of V. coralliilyticus in Mytilus galloprovincialis

* Corresponding Author

Teresa Balbi
Dept. of Earth, Environment and Life Sciences (DISTAV)
University of Genoa
Corso Europa 26
16132 – Genoa
Italy

Teresa.Balbi@unige.it

1 **1. Introduction**

2 Marine bivalves, due to their filter-feeding habit, accumulate large numbers of bacteria from
3 the harvesting waters. Bivalves possess both cellular and humoral defence mechanisms that co-
4 operate to kill and eliminate infecting bacteria [1,2]. However, some bacteria can be pathogenic to
5 the bivalve host, in particular those belonging to the genus *Vibrio*. Pathogenic vibrios can mainly
6 affect larval stages of cultured bivalves, and are also involved in diseases of juveniles and adults [3-
7 5]. The *Vibrio* species with importance for bivalve hatcheries due to the known pathogenicity for
8 larvae and spat have been recently summarized [6]. These include species from the *Anguillarum*,
9 *Coralliilyticus*, *Harveyi*, *Orientalis*, *Pectenocida* and *Splendidus* clades.

10 *Vibrio coralliilyticus* has emerged as a coral pathogen of concern throughout the Indo-Pacific
11 reef [7,8]. The interest towards understanding its ecology and pathogenic potential has increased
12 since *V. coralliilyticus* was shown to be strongly virulent also for other species, such as unicellular
13 algae [9,10], flies [10,11], rainbow trout (*Oncorhynchus mykiss*) and larval brine shrimp (*Artemia*
14 spp.) [12]. Moreover, *V. coralliilyticus* represents a serious threat for bivalve aquaculture, being one
15 of the most important emerging pathogen responsible for oyster larval mortalities worldwide [13-
16 15]. *V. coralliilyticus* has been also associated with outbreaks of vibriosis in several other bivalve
17 species, such as hard clam (*Mercenaria mercenaria*), New Zealand green-lipped mussel (*Perna*
18 *canaliculus*), Atlantic bay scallop (*Argopecten irradians*) and naval shipworm (*Teredo navalis*). *V.*
19 *coralliilyticus* has a tightly regulated temperature-dependent virulence and it has been related to
20 mass mortality events of the purple gorgonian *Paramuricea clavata* in the temperate north-western
21 Mediterranean Sea [16]. However, no information is available of the effects of *V. coralliilyticus* in
22 the Mediterranean mussel *Mytilus galloprovincialis*, which represents the most abundant
23 aquacultured species in this area.

24 Although *Mytilus* spp., including *M. galloprovincialis*, is particularly resistant to bacterial
25 infections, it shows a remarkable specificity of the immune response towards different *Vibrio* spp.
26 and strains, as demonstrated both *in vitro* and *in vivo* studies in adults [2,17,18]. In contrast, little

27 information is available on the possible vibrio pathogens affecting *Mytilus* embryo development
28 [19].

29 In this work, data are presented on responses of *M. galloprovincialis* to challenge with the
30 emerging marine pathogen *V. coralliilyticus*. *In vitro*, short term responses of mussel hemocytes to
31 *V. coralliilyticus* were evaluated in terms of lysosomal membrane stability (LMS), bactericidal
32 activity, extracellular lysozyme release, ROS and NO production. The effects on hemocyte
33 morphology were also investigated by TEM. *In vivo*, hemocyte LMS, ROS production and serum
34 lysozyme activity were measured in mussels challenged with *V. coralliilyticus* at 24 h post-
35 injection. Moreover, the effects of *V. coralliilyticus* on mussel early embryo development (at 48 h
36 post fertilization) were evaluated.

37

38 **2. Methods**

39 **2.1 Mussels and bacteria**

40 Mussels (*Mytilus galloprovincialis* Lam), 4-5 cm long, were purchased from an aquaculture
41 farm (Arborea-OR, Italy) in October 2017 and kept for 1 day in static tanks containing aerated
42 artificial sea water (ASW), salinity 36 ppt (1 L/mussel) at 18°C. Hemolymph was extracted from
43 the posterior adductor muscle using a sterile 1 mL syringe with an 18 G1/2" needle. With the needle
44 removed, hemolymph was filtered through a sterile gauze and pooled in 50 mL Falcon tubes at
45 18°C. Hemolymph serum was obtained by centrifugation of whole hemolymph at 100 x g for 10
46 min, and the supernatant was sterilized through a 0.22 µm-pore filter.

47 *V. coralliilyticus* ATCC BAA-450 and *V. coralliilyticus* TAV24 (isolated from diseased
48 *Paramuricea clavata* colonies [16]) were cultured in Zobell Marine Broth 2216 (Difco
49 Laboratories) at 20°C under static conditions; after overnight growth, cells were harvested by
50 centrifugation (4500 x g, 10 min), washed three times with artificial seawater (ASW), and
51 resuspended to an Abs₆₀₀ = 1 (about 10⁸ CFU/mL). Thiosulfate Citrate Bile Salts Sucrose (TCBS)
52 Agar (Conda Lab, Spain) was used throughout the experiments.

53

54 **2.2 *In vitro* challenge of *Mytilus* hemocytes with *V. coralliilyticus***

55 Hemocyte monolayers were prepared as previously described [18,20] and incubated at 18°C
56 with suspensions of *V. coralliilyticus* suitably diluted in hemolymph serum at different
57 concentrations (5×10^5 , 5×10^6 , 5×10^7 CFU/mL), for different periods of times, depending on the
58 endpoint measured. Untreated hemocyte samples in serum were run in parallel. All experiments
59 were performed in triplicate.

60

61 **2.2.1 Determination of lysosomal membrane stability-LMS**

62 Lysosomal membrane stability-LMS in hemocyte monolayers was evaluated by the NRRT
63 assay as previously described [17,18,20]. Hemocyte monolayers on glass slides were pre-incubated
64 for 30 min with different concentrations of *V. coralliilyticus*. Hemocyte monolayers were washed
65 out and incubated with 20 µL of a neutral red (NR) solution (final concentration 40 µg/mL from a
66 stock solution of NR 20 mg/mL DMSO-dimethylsulfoxide). After 15 min, excess dye was washed
67 out, 20 µL of ASW was added, and slides were sealed with a coverslip. Every 15 min, slides were
68 examined under optical microscope and the percentage of cells showing loss of dye from lysosomes
69 in each field was evaluated. For each time point, 10 fields were randomly observed, each containing
70 8-10 cells. The endpoint of the assay was defined as the time at which 50% of the cells showed sign
71 of lysosomal leaking, i.e. the cytosol becoming red and the cells rounded. All incubations were
72 carried out at 18°C.

73 For comparison, LMS was evaluated using the Mediterranean strain *V. coralliilyticus* TAV24
74 [16] in the same experimental conditions as described above.

75

76 **2.2.2 Bactericidal activity**

77 Bactericidal activity was evaluated as previously described [20,21]. Hemocyte monolayers
78 were incubated with different concentrations of *V. coralliilyticus* at 18°C, for different periods of

79 time (60-90 min). Immediately after the inoculum (T = 0) and after 60 and 90 min of incubation,
80 supernatants were collected and hemocytes were lysed by adding filter sterilized ASW
81 supplemented with 0.05% Triton x-100 and by 10 s agitation. Supernatants and lysates were pooled
82 and tenfold serial diluted in ASW. Aliquots (100 μ L) of the diluted samples were plated onto TCBS
83 Agar. After overnight incubation at 20°C, the number of colony-forming units (CFU) per hemocyte
84 monolayer (representing live, culturable bacteria) was determined. Percentages of killing were
85 determined in comparison to values obtained at zero time. The number of CFU in control
86 hemocytes never exceeded 0.1% of those enumerated in experimental samples.

87

88 **2.2.3 Lysozyme release, ROS and NO production**

89 For these endpoints, hemocytes were incubated with suspensions of *V. coralliilyticus* in serum
90 at 5×10^6 CFU/mL. Lysosomal enzyme release by mussel hemocytes was evaluated by measuring
91 lysozyme activity in the extracellular medium as previously described [20]. Lysozyme activity in
92 aliquots of serum of control hemocytes and hemocytes incubated *V. coralliilyticus* for different
93 periods of time (from 5 to 30 min), was determined spectrophotometrically at 450 nm using a
94 suspension of *Micrococcus lysodeikticus* (15 mg/100 mL in 66 mM phosphate buffer, pH 6.4). Data
95 are expressed as percentage of control values.

96 Extracellular generation of reactive oxygen species (ROS) was measured by the reduction of
97 cytochrome c as previously described [18]. Aliquots of hemocyte suspension were incubated for 30
98 min with cytochrome c solution (75 mM ferricytochrome c in TBS), with or without *V.*
99 *coralliilyticus*. Cytochrome c in TBS was used as a blank. Samples were read at 550 nm and the
100 results expressed as changes in OD per mg protein.

101 Nitric oxide (NO) production was evaluated as described previously [18] by the Griess
102 reaction, which quantifies the nitrite (NO_2^-) content of supernatants. Aliquots of hemocyte
103 suspensions were incubated at 18°C with or without bacterial suspension of *V. coralliilyticus* for 2
104 h. After the incubation, samples were frozen and stored at -80°C until analysis. Before analysis,

105 samples were thawed and centrifuged (12000 x g for 30 min at 4°C). Aliquots of supernatants were
106 incubated for 10 min in the dark with 1% (w/v) sulphanilamide in 5% H₃PO₄ and 0.1% (w/v) N-(1-
107 naphthyl)-ethylenediamine dihydrochloride. Samples were read at 540 nm, and the molar
108 concentration of NO₂⁻ in the sample was calculated from standard curves generated using known
109 concentrations of sodium nitrite. Data are expressed as nitrite accumulation per protein content,
110 determined according to the bicinchoninic acid (BCA) method using bovine serum albumin (BSA)
111 as a standard.

112

113 **2.2.4 Transmission electron microscopy (TEM)**

114 TEM of mussel hemocytes was carried out as previously described [20]. Hemocyte monolayers
115 were seeded on glass chamber slides for 20 min at 18°C (Lab-Tek, Nunc, 177380), and incubated
116 with *V. coralliilyticus* (5 x 10⁶ CFU/mL in hemolymph serum) for 5, 15 and 30 min. Samples were
117 washed out with 0.1 M cacodylate buffer in ASW and fixed in 0.1 M cacodylate buffer in ASW
118 containing 2.5% glutaraldehyde in ASW, for 1 h at room temperature. The cells were postfixed in
119 1% osmium tetroxide in ASW for 10 min and 1% uranyl acetate in ASW for 1 h. Subsequently,
120 samples were dehydrated through a graded ethanol series and embedded in epoxy resin (Poly-Bed;
121 Polysciences, Inc., Warrington, PA) overnight at 60°C. About 50 cells per sample were observed by
122 F20 Tecnai electron microscope (Philips, Eindhoven, The Netherlands), and representative images
123 were taken with an Eagle CCD camera and iTEM software and processed with Adobe Photoshop
124 CS2.3.2.

125

126 **2.3 *In vivo* challenge of adult mussels with *V. coralliilyticus***

127 Mussels were kept for 24 h in static tanks containing aerated artificial sea water (1 L/mussel)
128 at 18°C. Mussels were *in vivo* challenged by injection of live *V. coralliilyticus* into the posterior
129 adductor muscle, as previously described [20], with 50 µL of a bacterial suspension containing 1 x
130 10⁸ CFU/mL in PBS-NaCl (in order to obtain a nominal concentration of 5 x 10⁶ CFU/mussel).

131 Control mussels were injected with PBS-NaCl. After challenge, mussels were returned to sea water.
132 At 24 h post injection (p.i.), hemolymph was collected from the posterior adductor muscle of 4
133 pools of 4 mussels each. No mortality was observed during the experiments.

134 At 24 h p.i. in hemolymph samples from control and vibrio-injected mussels, hemocyte LMS
135 and ROS production, soluble lysozyme activity, as well as bacterial counts, evaluated as number of
136 CFU/mL of whole hemolymph were determined as described above.

137

138 **2.4 Effects of *V. coralliilyticus* on embryo development**

139 Sexually mature mussels (*M. galloprovincialis* Lam.), purchased from an aquaculture farm in
140 the Ligurian Sea (La Spezia, Italy) between November and March, were transferred to the
141 laboratory and acclimatized in static tanks containing aerated artificial sea water [22], pH 7.9-8.1,
142 36 ppt salinity (1 L/animal), at $18 \pm 1^\circ\text{C}$. Mussels were utilized within 2 days for gamete collection.
143 When mussels beginning to spontaneously spawn were observed, each individual was immediately
144 placed in a 250 mL beaker containing 200 mL of aerated ASW until complete gamete emission.
145 After spawning, mussels were removed from beakers and sperms and eggs were sieved through 50
146 mm and 100 mm meshes, respectively, to remove impurities. Egg quality (shape, size) and sperm
147 motility were checked using an inverted microscope. For each experiment, eggs and sperm from
148 two individuals were selected and counted to give a single pairing. Eggs were fertilized with an
149 egg:sperm ratio 1:10 in polystyrene 96-microwell plates (Costar, Corning Incorporate, NY, USA).
150 After 30 min fertilization success (n. fertilized eggs/n. total eggs x 100) was verified by
151 microscopical observation (>85%).

152 The 48-h embryotoxicity assay [22] was carried out in 96-microwell plates as described by
153 [23]. Aliquots of 20 μL of suspensions of *V. coralliilyticus* (obtained from a 10^7 CFU/mL stock
154 suspension), suitably diluted in ASW, were added to fertilized eggs in each microwell to reach the
155 nominal final concentrations (10^1 , 10^2 , 10^3 , 10^4 , 10^5 , 10^6 CFU/mL) in a 200 μL volume. At each
156 dilution step, all suspensions were immediately vortexed prior to use. Microplates were gently

157 stirred for 1 min, and then incubated at $18 \pm 1^\circ\text{C}$ for 48 h, with a 16 h:8 h light:dark photoperiod.
158 All the following procedures were carried out following [22]. At the end of the incubation time,
159 samples were fixed with buffered formalin (4%). All larvae in each well were examined by optical
160 and/or phase contrast microscopy using an inverted Olympus IX53 microscope (Olympus, Milano,
161 Italy) at 40X, equipped with a CCD UC30 camera and a digital image acquisition software
162 (cellSens Entry). Observations were carried out by an operator blind to the experimental conditions.
163 A larva was considered normal when the shell was D-shaped (straight hinge) and the mantle did not
164 protrude out of the shell, and malformed if had not reached the stage typical for 48 hpf (trochophore
165 or earlier stages) or when some developmental defects were observed (concave, malformed or
166 damaged shell, protruding mantle). The acceptability of test results was based on controls for a
167 percentage of normal D-shell stage larvae $>75\%$ [22]. Moreover, in each sample the percentage of
168 malformed D-veligers, imature veligers, and trocophorae was evaluated.

169

170 **2.5 Data analysis**

171 The results are the mean \pm SD of at least 4 experiments and analyses, unless otherwise
172 indicated, performed in triplicate. Statistical analysis was performed by ANOVA followed by
173 Tukey's post hoc test or by Mann-Whitney U test using the GraphPad InStat software.

174 Embriotoxicity test data, representing the mean \pm SD of 4 independent experiments, carried out
175 in 6 replicate samples in 96-microwell plates, were analyzed by ANOVA plus Tukey's post test.
176 The EC_{50} was defined as the concentration causing 50% reduction in the embryogenesis success,
177 and their 95% confidence intervals (CI) were calculated by PRISM 5 software (GraphPad Prism 5
178 software package, GraphPad Inc.).

179

180 **3. Results**

181 **3.1 Effects of *in vitro* challenge with *V. coralliilyticus* on hemocyte functional parameters**

182 Lysosomal membrane stability: as shown in Fig. 1A, incubation with *V. coralliilyticus* ATCC
183 BAA-450 for 30 min induced a dose-dependent decrease in hemocyte LMS, evaluated by the
184 NRRT assay, with respect to controls. The lowest concentration tested (5×10^5 CFU/mL) was
185 ineffective, while a moderate decrease was observed at 5×10^6 CFU/mL (-25%; $p < 0.05$). At the
186 highest concentration (5×10^7 CFU/mL) lysosomal membranes were completely destabilized (-
187 98%; $p < 0.01$). Interestingly, similar results were obtained with the Mediterranean strain *V.*
188 *coralliilyticus* TAV24 (Fig. S1).

189 Bactericidal activity: the capacity of mussel hemocytes to kill *V. coralliilyticus* ATCC BAA-
190 450 was investigated using a bactericidal assay that evaluates the number of live, culturable bacteria
191 at different times of incubation (Fig. 1B). Hemocytes were incubated with *V. coralliilyticus*, at the
192 same concentrations utilized in the LMS assay, and the results are reported as % of killed bacteria
193 with respect to the inoculum. The results clearly show a dose-dependent bactericidal activity
194 towards *V. coralliilyticus*. At 5×10^5 CFU/mL, *V. coralliilyticus* was efficiently killed by mussel
195 hemocytes (from 40% at 60 min to 65% at 90 min). A lower percentage of killing was observed at
196 the concentration of 5×10^6 CFU/mL (20% at both 60 and 90 min). At the highest concentration
197 tested (5×10^7 CFU/mL) no significant bactericidal activity was recorded (less than 10% at 90 min).

198 On the basis of these results, subsequent experiments to evaluate other immune parameters
199 were carried out using a concentration of bacteria of 5×10^6 CFU/mL, and the results are reported
200 in Fig. 2. Immediately after addition of ATCC BAA-450 bacteria, a significant increase in
201 extracellular lysozyme activity was observed with respect to controls (+37%, $p < 0.05$). No
202 differences were measured at subsequent times of incubation. *V. coralliilyticus* ATCC BAA-450 did
203 not affect extracellular ROS production (B) or nitrite accumulation (C) after 30 min and 2 h,
204 respectively.

205

206 **3.2 Effects of *in vitro* challenge with *V. coralliilyticus* on hemocyte ultrastructure**

207 The effects of challenge with *V. coralliilyticus* ATCC BAA-450 (5×10^6 CFU/mL) on the
208 morphology of mussel hemocytes were observed by TEM at different times of incubation (5, 15 and
209 30 min) and representative images are reported in Fig. 3. Fig. 3A shows *V. coralliilyticus* ATCC
210 BAA-450 before the addition to the hemocytes. A control hemocyte is shown in Fig. 3B; as
211 previously reported [18,20], in hemocyte monolayers control cells are mainly represented by
212 granulocytes, whose cytoplasm is filled by small intracellular granules of different electron
213 densities.

214 *V. coralliilyticus* induced morphological changes in the hemocytes at the plasma membrane
215 and cytoplasmic level as soon as 5 min from addition. Some cells formed irregular pseudopodial
216 extensions (Fig. 3C), while others showed a more flattened shape, with the cell membrane lining
217 portions of empty cytoplasm (Fig. 3D). Different ultrastructural changes were more evident at 15
218 min post-infection. In addition to the formation of long pseudopodia, *V. coralliilyticus* mainly
219 affected the intracellular vacuolar system, as shown by the appearance of enlarged vacuoles of
220 heterogeneous content, empty vacuoles, or vacuoles containing granular material (Fig. 3E and 3F).
221 At 30 min, large electron dense vacuoles of heterogeneous content were observed, suggesting
222 lysosomal fusion events, together with empty vacuoles and irregular plasma membrane surfaces
223 (Fig. 3G and 3H). No bacterial internalization was observed at any time of incubation (not shown).

224

225 **3.3 Effects of *in vivo* challenge with *V. coralliilyticus* on hemolymph parameters**

226 Mussels were injected with *V. coralliilyticus* ATCC BAA-450 in order to reach a nominal
227 concentration of 5×10^6 CFU/mL hemolymph and samples were collected after 24 h p.i. Hemocyte
228 LMS, serum lysozyme activity and ROS production were evaluated, as well as bacterial cell counts
229 in whole hemolymph samples. The results show that *in vivo* challenge with *V. coralliilyticus* lead to
230 a moderate but significant decrease in LMS at 24 h p.i. (-23%; $p < 0.05$) (Fig. 4A), comparable to
231 that observed in *in vitro* experiments. No increases in serum lysozyme activity (Fig. 4B) and
232 hemocyte ROS production (Fig.4C) were observed; interestingly, the basal levels of ROS were even

233 reduced with respect to controls (-24%, $p < 0.05$). Finally, in *V. coralliilyticus*-injected mussels,
234 *Vibrio* counts were significantly higher (about 7-folds; $p < 0.01$) in hemolymph collected at 24 h p.i.,
235 compared to those in hemolymph collected immediately after infection (T=0) (Fig. 4D), indicating
236 bacterial growth.

237

238 **3.4 Effects of *V. coralliilyticus* on embryo development**

239 Fertilized eggs were exposed to different concentrations (from 10^1 to 10^6 CFU/mL) of *V.*
240 *coralliilyticus* ATCC BAA-450 in 96-microwell plates, and the percentage of normal D-larvae was
241 evaluated after 48 hpf. The results, reported in Fig. 5, show that *V. coralliilyticus* significantly
242 affected normal larval development, with an EC_{50} value of 5.045×10^3 CFU/mL (4.599 - 5.492,
243 95% CI) (Fig. 5A). The percentage of normal D-larvae was significantly reduced from the lowest
244 concentration tested (from -30% vs controls at 10 CFU/mL) and a dose-dependent effect was
245 observed at increasing concentrations, up to a complete impairment of normal D-larvae
246 development at 10^6 CFU/mL (-92.8%).

247 When the type of effect caused by bacterial challenge was evaluated (Fig. 5B) *V. coralliilyticus*
248 induced a progressive increase in the percentage of malformed embryos. At the highest
249 concentration tested (10^6 CFU/mL), the presence of trocophorae/immature D-veligers was also
250 observed, indicating developmental arrest. In Fig. 5C representative images of control embryos and
251 embryos exposed to different concentrations of *V. coralliilyticus* are reported.

252

253 **4. Discussion**

254 The present work represents the first investigation on the responses of *M. galloprovincialis* to
255 challenge with the emerging marine pathogen *V. coralliilyticus*. To this aim, the reference ATCC
256 BAA-450 strain isolated from bleached corals near Zanzibar [7], was utilized. *In vitro* experiments
257 were carried out in the presence of hemolymph serum, in order to simulate the *in vivo* conditions,
258 taking into account also the possible role of soluble hemolymph components, and functional

259 responses of *M. galloprovincialis* hemocytes were evaluated. The results show that challenge with
260 *V. coralliilyticus* induced a dose-dependent lysosomal membrane destabilization that was inversely
261 correlated with bactericidal activity. In particular, whereas at the lowest vibrio concentration tested,
262 hemocytes, in the absence of lysosomal stress, were able to efficiently kill bacteria, at the highest
263 concentration tested (10^7 CFU/mL) *V. coralliilyticus* was cytotoxic, and no bactericidal activity was
264 observed. Interestingly, from these data *V. coralliilyticus* appears to be more virulent to *M.*
265 *galloprovincialis* hemocytes with respect to other *Vibrio* species and strains tested in the same
266 experimental conditions (*V. splendidus*, *V. aestuarianus*, *V. anguillarum*, *V. tapetis*, *V. cholerae*) [2
267 and references quoted therein].

268 Other functional responses were evaluated at intermediate *Vibrio* concentrations (10^6
269 CFU/mL), when moderate lysosomal destabilization in hemocytes and some bactericidal activity
270 were observed. The results indicate no activation of immune parameters, except for an extremely
271 rapid extracellular lysozyme release, observed immediately after addition of *V. coralliilyticus*. In
272 these conditions, TEM analysis of hemocytes showed that as soon as after 5 min incubation, cell
273 membranes lining empty portions of cytoplasm were present, thus indicating possible
274 degranulation. Moreover, after 30 min incubation *V. coralliilyticus* induced lysosomal fusion
275 events, in line with LMS data. However, no vibrio internalization was observed. Overall, TEM
276 observations confirm the results of functional parameters and indicate that *in vitro* challenge with *V.*
277 *coralliilyticus* does not result in intracellular degradation of bacteria. The limited bactericidal
278 activity observed in these conditions (about 20%) may be probably related to the rapid extracellular
279 degranulation of hydrolytic enzymes.

280 The effects of *V. coralliilyticus* were also investigated *in vivo*, in hemolymph from injected
281 mussels sampled after 24 h p.i. In these conditions, challenge with *V. coralliilyticus* induced a
282 significant decrease in hemocyte LMS, but did not result in activation of immune parameters, thus
283 confirming the *in vitro* data. In addition, vibrio challenge even reduced basal ROS production.

284 Accordingly, the results indicate that *V. coralliilyticus* can grow within mussel hemolymph, as
285 shown by the large increase in *Vibrio* counts registered in whole hemolymph samples at 24 h p.i.

286 *V. coralliilyticus* also affected mussel embryo development, inducing a dose-dependent
287 decrease in the percentage of normal D-veligers at 48 hpf, with an EC₅₀ of 5.045 x 10³ CFU/mL.
288 Interestingly, the effect was significant from the lowest concentration tested (10 CFU/mL),
289 approximately corresponding to a ratio of 2 CFU/40 embryos in each well.

290 Challenge with *V. coralliilyticus* resulted in embryo malformations at all the concentrations
291 tested. Moreover, at higher concentrations, the presence of trocophorae/immature D-veligers was
292 observed, indicating that *V. coralliilyticus* could also induce a delay in development. In all
293 experimental conditions, erratic closing of the valves, velum detachment, and bacterial swarming
294 around the embryos were observed, which are clear signs of disease in the larvae [4,24,25]. In both
295 Eastern and Pacific oyster larvae, challenge with *V. coralliilyticus* ATCC BAA-450 for 6 days
296 resulted in mortalities with LD₅₀ of 2.1 and 4 x 10⁴ CFU/mL, respectively [15]. In *C. gigas*, *V.*
297 *coralliilyticus* also induced a wide range of physiological, enzymatic, biochemical and molecular
298 changes [14]. However, oyster data were obtained in 1-2 weeks old larvae. The results here reported
299 represent the first data on the effects of *V. coralliilyticus* on early developmental stages of bivalves.
300 In *M. galloprovincialis*, immune capacities arise during mussel development as early as the
301 trocophorae stage (24 hpf). At this developmental stage, gene expression has contributions of
302 maternal origin, but stimulation induces the expression of immune-related genes [26]. However, the
303 present results show that mussel early embryos are particularly sensitive to *V. coralliilyticus*, and
304 indicate that they are unable to mount a defence response towards this pathogen.

305 *V. coralliilyticus* possess several virulence mechanisms, including powerful extracellular
306 enzymes that have been linked to direct lysis of coral tissue [8]. Several authors demonstrated that
307 the virulence of some strains is associated with the production of toxins, mainly extracellular
308 metalloprotease (VtpA) and hemolysin (VthA) [27-30]. Furthermore, coral diseases not only
309 depend on the presence of *Vibrio* pathogens and their virulence level, but are also the result of

310 complex interactions between the expression of different bacterial virulence factors and an increase
311 of seawater temperature or other environmental stresses, as well as the physiological and immune
312 status of the coral host [31].

313 *Vibrio* species are strongly thermodependent. In particular, for the reference strain of *V.*
314 *coralliilyticus* ATCC BAA-450 a direct temperature regulation of multiple virulence mechanisms
315 has been demonstrated at 27°C [32]. *V. coralliilyticus* is able to invade and to lyse the tissue of the
316 coral *Pocillopora damicornis*, one of the most affected organisms, at temperatures higher than
317 27°C, while in a temperature range between 24°C and 26°C it kills the symbiotic algae of the coral
318 [9]. At temperature below 24°C is totally avirulent [9,33]. In the present work, all experiments were
319 carried out at the constant temperature of 18°C, in order to ensure the health and immune status of
320 the mussels. However, even in these conditions, both adult and embryos of *M. galloprovincialis* are
321 apparently unable to mount an efficient immune response towards *V. coralliilyticus*. This results in
322 lysosomal stress in the hemocytes both *in vitro* and *in vivo*, in bacterial growth in the hemolymph of
323 adult mussels challenged *in vivo*, and in malformations in early embryos. Recent data indicate that
324 *in vivo* challenge of the New Zealand Greenshell Mussel *Perna canaliculus* with a *V.*
325 *coralliilyticus/neptunius*-like isolate induced perturbations of the immune system, oxidative stress,
326 inflammation and metabolic changes at 6 days p.i. [34]. Overall, these findings provide a further
327 insight into the pathogenic effects of *V. coralliilyticus* in mussels.

328 In a global warming scenario, an increase in the seawater temperature could promote the
329 proliferation and the potential disease outbreaks associated with *Vibrio* pathogens also in mussels.
330 This is of particular concern in temperate regions such as the Mediterranean sea, where the relative
331 increase in seawater temperature seems to be higher than in tropical areas [31]. Mediterranean
332 strains of *V. coralliilyticus* have been isolated from diseased *P. clavata* colonies collected at
333 Tavolara island (Sardinia, Italy) [16]. Among these, the most virulent strain is TAV24, recently
334 identified as a new genotype of *V. coralliilyticus* by MLST and *vcpA* gene sequencing analyses
335 [35]. The results here reported indicate that the *in vitro* effects of the TAV24 strain on hemocyte

336 lysosomal membrane stability were comparable with those of the reference strain. The responses of
337 *M. galloprovincialis* to challenge with the highly virulent Mediterranean strain require further
338 investigation. Despite the fact that *V. coralliilyticus* appears to be a global bivalve pathogen, there is
339 limited information about its pathogenicity, infection mechanism and/or disease mitigation. These
340 studies will contribute to understand the potential threat of this vibrio to bivalve aquaculture in the
341 Mediterranean.

342

343

344 **Acknowledgments**

345 This work was funded by the EU project H2020 VIVALDI (Preventing and Mitigating Farmed
346 Bivalve Diseases), contract n° 678589.

347 The authors wish to thank Dr. Rita Fabbri for her invaluable technical assistance and Dr.
348 Giovanni Tassistro for bacterial cultures.

349

350 **References**

- 351 [1] L. Canesi, M. Gavioli, C. Pruzzo, G. Gallo, Bacteria-hemocyte interactions and phagocytosis in
352 marine bivalves, *Microsc. Res. Tech.* 57 (2002) 469-476.
- 353 [2] L. Canesi, C. Pruzzo, Specificity of innate immunity in bivalves: a lesson from bacteria, in: L.
354 Ballarin, M. Cammarata (Eds.), *Lessons in Immunity: from Single-cell Organisms to*
355 *Mammals*, Elsevier Inc, 2016, pp. 79-91.
- 356 [3] C. Paillard, F. Le Roux, J.J. Borrego, Bacterial disease in marine bivalves, a review of recent
357 studies: trends and evolution, *Aquat. Living Resour.* 17 (2004) 477-498.
- 358 [4] R. Beaz-Hidalgo, S. Balboa, J.L. Romalde, M.J. Figueras, Diversity and pathogenicity of
359 *Vibrio* species in cultured bivalve molluscs, *Environ. Microbiol. Rep.* 2 (2010) 34-43.
- 360 [5] M.A. Travers, K. Boettcher Miller, A. Roque, C.S. Friedman, Bacterial diseases in marine
361 bivalves, *J. Invertebr. Pathol.* 131 (2015) 11-31.

- 362 [6] J. Dubert, J.L. Barja, J.L. Romalde, New insights into pathogenic *Vibrios* affecting bivalves in
363 hatcheries: Present and future prospects, *Front. Microbiol.* 8 (2017) 762.
- 364 [7] Y. Ben-Haim, E. Rosenberg, A novel *Vibrio* sp. pathogen of the coral *Pocillopora damicornis*,
365 *Mar. Biol.* 141 (2002) 47-55.
- 366 [8] M. Sussman, B.L. Willis, S. Victor, D.G. Bourne, Coral pathogens identified for white
367 syndrome (WS) epizootics in the Indo-Pacific, *Plos ONE* 3 (2008) e2393.
- 368 [9] Y. Ben-Haim, M. Zicherman-Keren, E. Rosenberg, Temperature-regulated bleaching and lysis
369 of the coral *Pocillopora damicornis* by the novel pathogen *Vibrio coralliilyticus*, *Appl.*
370 *Environ. Microbiol.* 69 (2003) 4236-4242.
- 371 [10] E. de Oliveira Santos, N. Alves Jr, G.M. Dias, A.M. Mazotto, A. Vermelho, G.J. Vora, B.
372 Wilson, V.H. Beltran, D.G. Bourne, F. Le Roux, F.L. Thompson, Genomic and proteomic
373 analyses of the coral pathogen *Vibrio coralliilyticus* reveal a diverse virulence repertoire, *ISME*
374 *J.* 5 (2011) 1471-1483.
- 375 [11] N. Alves Jr, O.S. Neto, B.S. Silva, R.L. De Moura, R.B. Francini-Filho, C. Barreira e Castro,
376 R. Paranhos, B.C. Bitner-Mathé, R.H. Kruger, A.C. Vicente, C.C. Thompson, F.L. Thompson,
377 Diversity and pathogenic potential of vibrios isolated from Abrolhos Bank corals, *Environ.*
378 *Microbiol. Rep.* 2 (2010) 90-95.
- 379 [12] B. Austin, D. Austin, R. Sutherland, F. Thompson, J. Swings, Pathogenicity of vibrios to
380 rainbow trout (*Oncorhynchus mykiss*, Walbaum) and *Artemia* nauplii, *Environ. Microbiol.* 7
381 (2005) 1488-1495.
- 382 [13] A. Kesarcodi-Watson, P. Miner, J.L. Nicolas, R. Robert, Protective effect of four potential
383 probiotics against pathogen-challenge of the larvae of three bivalves: Pacific oyster
384 (*Crassostrea gigas*), flat oyster (*Ostrea edulis*) and scallop (*Pecten maximus*), *Aquaculture*
385 344-349 (2012) 29-34.

- 386 [14] B. Genard, P. Miner, J.L. Nicolas, D. Moraga, P. Boudry, F. Pernet, R. Tremblay, Integrative
387 study of physiological changes associated with bacterial infection in Pacific oyster larvae, Plos
388 ONE 8 (2013) e64534.
- 389 [15] G.P. Richards, M.A. Watson, D.S. Needleman, K.M. Church, C.C Häse, Mortalities of Eastern
390 and Pacific oyster larvae caused by the pathogens *Vibrio coralliilyticus* and *Vibrio tubiashii*,
391 Appl. Environ. Microbiol. 81 (2015) 292-297.
- 392 [16] L. Vezzulli, M. Previati, C. Pruzzo, A. Marchese, D.G. Bourne, C. Cerrano, the VibrioSea
393 Consortium, *Vibrio* infections triggering mass mortality events in a warming Mediterranean
394 Sea, Environ. Microbiol. 12 (2010) 2007-2019.
- 395 [17] C. Ciacci, A. Manti, B. Canonico, R. Campana, G. Camisassi, W. Baffone, L. Canesi,
396 Responses of *Mytilus galloprovincialis* hemocytes to environmental strains of *Vibrio*
397 *parahaemolyticus*, *Vibrio alginolyticus*, *Vibrio vulnificus*, Fish Shellfish Immunol. 65 (2017)
398 80-87.
- 399 [18] T. Balbi, K. Cortese, C. Ciacci, G. Bellese, L. Vezzulli, C. Pruzzo, L. Canesi, Autophagic
400 processes in *Mytilus galloprovincialis* hemocytes: Effects of *Vibrio tapetis*, Fish Shellfish
401 Immunol. 73 (2018) 66-74.
- 402 [19] C. Anguiano-Beltrán, M.L. Lizárraga-Partida, R. Searcy-Bernal, Effect of *Vibrio alginolyticus*
403 on larval survival of the blue mussel *Mytilus galloprovincialis*, Dis. Aquat. Org. 59 (2004) 119-
404 123.
- 405 [20] T. Balbi, R. Fabbri, K. Cortese, A. Smerilli, C. Ciacci, C. Grande, L. Vezzulli, C. Pruzzo, L.
406 Canesi, Interactions between *Mytilus galloprovincialis* hemocytes and the bivalve pathogens
407 *Vibrio aestuarianus* 01/032 and *Vibrio splendidus* LGP32, Fish Shellfish Immunol. 35 (2013)
408 1906-1915.
- 409 [21] L. Canesi, C. Pruzzo, R. Tarsi, G. Gallo, Surface interactions between *Escherichia coli* and
410 hemocytes of the mediterranean mussel *Mytilus galloprovincialis* Lam. leading to efficient
411 bacterial clearance, Appl. Environ. Microbiol. 67 (2001) 464-468.

- 412 [22] ASTM, Standard guide for conducting static acute toxicity tests starting with embryos of four
413 species of saltwater bivalve molluscs, <http://dx.doi.org/10.1520/E0724-98>, 2004.
- 414 [23] R. Fabbri, M. Montagna, T. Balbi, E. Raffo, F. Palumbo, L. Canesi, Adaptation of the bivalve
415 embryotoxicity assay for the high throughput screening of emerging contaminants in *Mytilus*
416 *galloprovincialis*, Mar. Environ. Res. 99 (2014) 1-8.
- 417 [24] S. Prado, J.L. Romalde, J. Montes, J.L. Barja, Pathogenic bacteria isolated from disease
418 outbreaks in shellfish hatcheries. First description of *Vibrio neptunius* as an oyster pathogen,
419 Dis. Aquat. Org. 67 (2005) 209-215.
- 420 [25] R. Rojas, C.D. Miranda, R. Opazo, J. Romero, Characterization and pathogenicity of *Vibrio*
421 *splendidus* strains associated with massive mortalities of commercial hatchery-reared larvae of
422 scallop *Argopecten purpuratus* (Lamarck, 1819), J. Invertebr. Pathol. 124 (2015) 61-69.
- 423 [26] P. Balseiro, R. Moreira, R. Chamorro, A. Figueras, B. Novoa, Immune responses during the
424 larval stages of *Mytilus galloprovincialis*: metamorphosis alters immunocompetence, body
425 shape and behavior, Fish Shellfish Immunol. 3 (2013) 438-447.
- 426 [27] M.H. Kothary, R.B. Delston, S.K. Curtis, B.A. McCardell, B.D. Tall, Purification and
427 characterization of vulnificolysin-like cytotoxin produced by *Vibrio tubiashii*, Appl. Environ.
428 Microbiol. 67 (2001) 3707-3711.
- 429 [28] R.B. Delston, M.H. Kothary, K.A. Shangraw, B.D. Tall, Isolation and characterization of zinc-
430 containing metalloprotease expressed by *Vibrio tubiashii*, Can. J. Microbiol. 49 (2003) 525-
431 529.
- 432 [29] H. Hasegawa, E.J. Lind, M.A. Boin, C.C. Häse, The extracellular metalloprotease of *Vibrio*
433 *tubiashii* is a major virulence factor for Pacific oyster (*Crassostrea gigas*) larvae, Appl.
434 Environ. Microbiol. 74 (2008) 4101-4110.
- 435 [30] H. Hasegawa, C.C. Häse, TetR-type transcriptional regulator VtpR functions as a global
436 regulator in *Vibrio tubiashii*, Appl. Environ. Microbiol. 75 (2009) 7602-7609.

- 437 [31] E. Rubio-Portillo, J.F. Gago, M. Martínez-García, L. Vezzulli, R. Rosselló-Móra, J. Antón,
438 A.A. Ramos-Esplá, *Vibrio* communities in scleractinian corals differ according to health status
439 and geographic location in the Mediterranean Sea, *Syst. Appl. Microbiol.* 41 (2018) 131-138.
- 440 [32] N.E. Kimes, C.J. Grim, W.R. Johnson, N.A. Hasan, B.D. Tall, M.H. Kothary, H. Kiss, A.C.
441 Munk, R. Tapia, L. Green, C. Detter, D.C. Bruce, T.S. Brettin, R.R. Colwell, P.J. Morris,
442 Temperature regulation of virulence factors in the pathogen *Vibrio coralliilyticus*, *ISME J.* 6
443 (2012) 835-846.
- 444 [33] J. Tout, N. Siboni, L.F. Messer, M. Garren, R. Stocker, N.S. Webster, P.J. Ralph, J.R.
445 Seymour, Increased seawater temperature increases the abundance and alters the structure of
446 natural *Vibrio* populations associated with the coral *Pocillopora damicornis*, *Front. Microbiol.*
447 6 (2015) 432.
- 448 [34] T.V. Nguyen, A.C. Alfaro, T. Young, S. Ravi, F. Merien, Metabolomics study of immune
449 responses of New Zealand Greenshell™ mussels (*Perna canaliculus*) infected with pathogenic
450 *Vibrio* sp., *Mar. Biotechnol.* 20 (2018) 396-409.
- 451 [35] B. Wilson, A. Muirhead, M. Bazanella, C. Huete-Stauffer, L. Vezzulli, D.G. Bourne, An
452 improved detection and quantification method for the coral pathogen *Vibrio coralliilyticus*,
453 *Plos ONE* 812 (2013) e81800.

454

455

456 **Figure legends.**

457 **Fig. 1 - *In vitro* effects of *V. coralliilyticus* on lysosomal membrane stability-LMS and**
458 **bactericidal activity.**

459 A) Hemocyte monolayers were treated with different concentrations (5×10^5 , 5×10^6 , 5×10^7
460 CFU/mL) of *V. coralliilyticus* for 30 min and LMS was evaluated as described in Methods. Data,
461 expressed as percent values with respect to controls and representing the mean \pm SD of 4

462 experiments in triplicate, were analysed by ANOVA followed by Tukey's post hoc test (* = $p < 0.05$;
463 ** = $p < 0.01$).

464 B) Hemocytes were incubated for different periods of time (60-90 min) with *V. coralliilyticus*, at
465 the same concentrations utilized in the LMS assay, and the number of viable, cultivable bacteria
466 (CFU) per monolayer was evaluated. Percentages of killing were determined in comparison to
467 values obtained at zero time.

468

469 **Fig. 2 - *In vitro* effects of *V. coralliilyticus* on functional parameters of *Mytilus* hemocytes.**

470 Lysosomal enzyme release (A), extracellular ROS production (B) and NO accumulation (C) were
471 evaluated after incubation with *V. coralliilyticus* (*V.c.*) at 5×10^6 CFU/mL in hemolymph serum.
472 Data are the mean \pm SD of at least 4 experiments performed in triplicate. Statistical analysis was
473 performed by ANOVA followed by Tukey's post hoc test (* = $p < 0.05$).

474

475 **Fig. 3 - Early *in vitro* effects of *V. coralliilyticus* on the ultrastructure of mussel hemocytes**
476 **evaluated by TEM.**

477 Representative images of A) *V. coralliilyticus* before addition to the hemocytes; B) Control
478 hemocyte; C-H), hemocytes incubated with *V. coralliilyticus* (5×10^6 CFU/mL) for 5 min (C-D), 15
479 min (E-F) and 30 min (G-H). As soon as after 5 min incubation, the formation of irregular
480 pseudopodial extensions and membrane vesicles was observed (C); moreover, many cells showed a
481 more adherent, flattened shape, with the cell membrane lining empty portions of cytoplasm
482 (arrowhead in D) and vesicles. At 15 min, single long filopodia were formed, as well as many
483 vacuoles with heterogeneous content (E), empty vesicles and vesicles with granular material (F). At
484 30 min, large electron dense vacuoles of heterogeneous content were observed, suggesting
485 lysosomal fusion events (G and enlargement in H), together with empty vacuoles, cytoplasmic
486 disorganization and irregular plasma membrane surfaces (arrowhead in H). No intracellular bacteria
487 were observed.

488 Scale bars: A) 1 μm ; B-H) 5 μm .

489

490 **Fig. 4 - *In vivo* effects of *V. coralliilyticus* on hemolymph parameters of *Mytilus* hemocytes.**

491 Hemocyte lysosomal membrane stability-LMS (A), serum lysozyme activity (B), ROS production
492 (C) and bacterial cell counts (D) were evaluated in hemolymph sampled from mussels challenged
493 with *V. coralliilyticus* (*V.c.*) at 24 h p.i.. Data are the mean \pm SD of at least 4 experiments performed
494 in triplicate. Statistical analysis was performed by ANOVA followed by Tukey's post hoc test (* =
495 $p < 0.05$; ** = $p < 0.01$).

496

497 **Fig. 5 - Effects of different concentrations of *V. coralliilyticus* on *M. galloprovincialis* normal
498 larval development in the 48 h embryotoxicity assay.**

499 A) Percentage of normal D-shaped larvae with respect to controls. B) Percentage of normal D-
500 veliger (dark grey), malformed D-veliger (light grey), pre-veligers (white) and trocophorae (black)
501 in each experimental condition. Data represent the mean \pm SD of 4 experiments carried out in 96-
502 multiwell plates (6 replicate wells for each sample). C) Representative images of control embryos
503 and embryos exposed to different concentrations of *V. coralliilyticus*, showing progressive shell
504 malformations, including asymmetric valvae, irregular hinges, externalized velum and, at the
505 highest concentration of bacteria, immature embryos. Bacteria swarming around larvae can be
506 observed at increasing concentrations.

Figure1
[Click here to download high resolution image](#)





Figure2

[Click here to download high resolution image](#)




Figure 3

[Click here to download high resolution image](#)




Figure4

[Click here to download high resolution image](#)






Figure 5

[Click here to download high resolution image](#)





Supplementary material for online publication only

[Click here to download Supplementary material for online publication only: Supplementary Information.doc](#)