



## Original article

Scand J Work Environ Health 2015;41(3):234-246

doi:10.5271/sjweh.3493

### Patterns of biomechanical demands are associated with musculoskeletal pain in the beginning of professional life: a population-based study

by [Lourenço S](#), [Araújo F](#), [Severo M](#), [Cunha Miranda L](#), [Carnide F](#), [Lucas R](#)

Patterns of highly physically demanding work-related tasks were associated with neck/shoulder pain and the presence and severity of upper-/lower-back pain. The beginning of professional life may be regarded as a key timeframe to quantify the preventive potential of interventions on work-related causes of neck, shoulder, and back pain.

**Affiliation:** EPIUnit – Institute of Public Health of the University of Porto, Rua das Taipas, 135-139, 4050-600 Porto, Portugal. [slourenco@med.up.pt](mailto:slourenco@med.up.pt)

Refers to the following texts of the Journal: [2013;39\(4\):321-426](#)  
[2001;27\(6\):0](#) [1999;25\(2\):0](#) [1990;16 suppl 1:1-79](#)

**Key terms:** [biomechanical demand](#); [biomechanical exposure](#); [musculoskeletal pain](#); [occupation](#); [pattern of biomechanical demand](#); [population-based study](#); [professional life](#); [work-related biomechanical exposure](#); [young adult](#)

This article in PubMed: [www.ncbi.nlm.nih.gov/pubmed/25808241](http://www.ncbi.nlm.nih.gov/pubmed/25808241)

### Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).



This work is licensed under a [Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/).

## Patterns of biomechanical demands are associated with musculoskeletal pain in the beginning of professional life: a population-based study

by Sara Lourenço, MSc,<sup>1</sup> Fábio Araújo, MSc,<sup>1</sup> Milton Severo, PhD,<sup>1,2,3</sup> Luís Cunha Miranda, MD,<sup>4</sup> Filomena Carnide, PhD,<sup>5</sup> Raquel Lucas PhD<sup>1,2</sup>

Lourenço S, Araújo F, Severo M, Cunha Miranda L, Carnide F, Lucas R. Patterns of biomechanical demands are associated with musculoskeletal pain in the beginning of professional life: a population-based study. Scand J Work Environ Health. 2015;41(3):234–246. doi:10.5271/sjweh.3493

**Objectives** This study aimed to describe patterns of occupational biomechanical demands in the beginning of professional life and to quantify their association with the presence and intensity of regional musculoskeletal pain.

**Methods** Cross-sectional data from 21-year-old participants were collected during the third wave of the EPITeen cohort study (N=1733, 37.5% were workers). Ten different work-related biomechanical tasks were characterized. Latent class analysis was conducted to identify empirical patterns of occupational biomechanical demands. The presence and intensity of regional musculoskeletal pain in the previous year were also evaluated.

**Results** Four patterns of occupational biomechanical demands were found: “low demands”, “sitting demands”, “repetitive and asymmetric demands”, and “high and vibrational demands”. When compared to workers with low demands, those with repetitive and asymmetric demands or high and vibrational demands presented 80% higher adjusted odds ratio (OR<sub>adj</sub>) of reporting neck/shoulder pain. High and vibrational demands occupations were significantly associated with upper-/lower-back pain in comparison to low demands [OR<sub>adj</sub> 1.80, 95% confidence interval (95% CI) 1.09–2.96]. In addition, workers with sitting demands were more likely to report any or severe upper-/lower-back pain [OR<sub>adj</sub> 1.56 (95% CI 0.99–2.45) and 1.66 (95% CI 1.03–2.66), respectively] when compared to those with low demands.

**Conclusions** Patterns of high work-related physical demands were associated with the presence of neck/shoulder pain and severity of upper-/lower-back pain. This emphasizes that even short-term biomechanical exposures at the workplace may be involved in the etiology of musculoskeletal complaints.

**Key terms** biomechanical exposure; occupation; work-related biomechanical exposure; young adult.

Musculoskeletal conditions are the most common cause of severe long-term pain and physical incapacity, representing 17% of the total years lived with disability among the young adult population worldwide (1, 2). Even though these diseases are multifactorial, work-related factors account for 30–40% of musculoskeletal complaints among active workers (3). In turn, musculoskeletal pain has been pointed as the main cause of work disability, and it is the most frequent reason for long-term sick leave and early retirement (4, 5).

Occupational exposures such as external loads, organizational factors and psychosocial work environment are strong determinants of musculoskeletal pain (6). Among those, physical factors such as heavy loads, bending or twisting, vibrations and awkward postures are the strongest predictors of multisite musculoskeletal pain, probably due to the direct mechanical tissue overload induced on joints (7–9).

Most research on the biomechanical etiology of work-related musculoskeletal conditions has focused

<sup>1</sup> EPIUnit – Institute of Public Health, University of Porto, Porto, Portugal.

<sup>2</sup> Department of Clinical Epidemiology, Predictive Medicine and Public Health, University of Porto Medical School, Porto, Portugal.

<sup>3</sup> Department of Medical Education and Simulation, University of Porto Medical School, Porto, Portugal.

<sup>4</sup> Portuguese Institute of Rheumatology, Lisboa, Portugal.

<sup>5</sup> Faculty of Human Kinetics, University of Lisbon, Cruz Quebrada – Dafundo, Portugal.

Correspondence to: Sara Lourenço, EPIUnit – Institute of Public Health of the University of Porto, Rua das Taipas, 135-139, 4050-600 Porto, Portugal. [E-mail: slourenco@med.up.pt]

on adults of heterogeneous age ranges (10). In such context, the well-documented healthy worker bias is expected to arise since workers with musculoskeletal complaints are more likely to have retired early or to be moving towards less demanding mechanical tasks. In addition, changes throughout life in the individual's work-related biomechanical demands lead to increasing misclassification of cumulative exposure with advancing stages of professional life.

Recent studies among newly-employed workers have shown that even short-term exposures to mechanical loads contribute significantly to the etiology of musculoskeletal pain, which suggests that investigating occupational health since the beginning of professional life may guide the timing of preventive strategies (11–14). However, research in early stages of employment remains fairly scarce, as well as restricted to a limited number of anatomical regions and specific occupational groups.

Work activities are commonly characterized by a narrow spectrum of co-occurring biomechanical demands (15) that cluster within workers and create different patterns of physical exposures in the occupational setting (16). Most previous evidence has emerged from independent analysis of isolated mechanical tasks, disregarding the potential synergistic effects of multiple combinations of distinct biomechanical factors in determining regional musculoskeletal pain (17). Such synergistic effect on low-back pain was indeed suggested in a 2011 study of students with traineeship-related occupations (18). Currently, still little is known about the association between the effective beginning of professional life – with the diversity of exposures as they occur in the current labor market – and the occurrence of pain in the musculoskeletal system as a whole.

Therefore, using data collected in a population-based sample of 21-year-old adults, we aimed to: (i) describe patterns of occupational biomechanical demands; and (ii) assess whether exposure to different patterns of work-related biomechanical demands have an effect on the presence and intensity of regional musculoskeletal pain in the beginning of professional life.

## Study population and methods

### Participants

In this study, we use cross-sectional data collected in 2011–2013 from young adults as part of the Epidemiological Health Investigation of Teenagers in Porto (EPI-Teen). The EPITeen study was designed as a prospective cohort first assembled during the 2003/2004 school year, when we approached all public and private schools in Porto (Portugal) providing teaching to adolescents born

in 1990. In the first wave, we identified 2787 eligible adolescents, of whom 2159 agreed to participate (77.5% baseline participation). In 2007/2008 (second wave), the initially recruited sample was re-evaluated and 783 students born in the same year but who moved to Porto after 2003/2004 were additionally recruited to the cohort. Sampling procedures and detailed methods have already been described (19).

During 2011–2013, 1761 young (21 years old) adults (of the 2942 eligible for follow-up) attended the third evaluation wave on which the present paper is exclusively based. We excluded 23 subjects due to neurodevelopmental or other illness (eg, autism spectrum disorders) and five due to missing data on work status. A final sample of 1733 individuals was included in the present analysis, considering both 1083 participants who were non-workers (more than 80% full-time students) and 650 workers (appendix 1: [http://www.sjweh.fi/data\\_repository.php](http://www.sjweh.fi/data_repository.php)).

Participants included (N=1733) were similar to those lost to follow-up regarding sex, body mass index and smoking behavior at 13 years of age. However, subjects attending follow-up were more sedentary (mostly sitting during leisure time: 31.1% versus 24.5%,  $P=0.012$ ) and had higher parental education (mean schooling years: 11.4 versus 9.2,  $P<0.001$ ).

The Ethics Committee of University Hospital of São João (Porto) approved the study protocol, which was carried out in accordance with the principles of the Declaration of Helsinki. Written informed consent was obtained from all participants.

### Data collection

**Occupational biomechanical characteristics.** Exposures to ten different work-related biomechanical demands (sitting posture, computer use, whole body vibrations, handling vibration tools, precision demands, repetitive tasks, overhead work, kneeling work, bending or rotation movements, and manual materials handling) were assessed among all participants who had worked for at least one month in the previous year.

The average fractions of a regular working day that were spent in sitting position, using the computer, exposed to whole body vibrations, and handling vibration tools were measured through a visual analogue scale, ranging from 0–100%. The absolute number of weekly hours of exposure to those mechanical demands was computed by multiplying the percentage of time performing each task by the daily number of working hours (full-time=8 hours, part-time=4 hours, less than a part-time job=2 hours), assuming a constant average of five working days per week. Participants were then grouped as above or below the sample median of weekly hours exposed to each biomechanical demand.

Workers were also asked to quantify the frequency of precision demands, repetitive tasks, overhead work, kneeling work, bending or rotation movements, and manual materials handling using Likert scales ranging from 0=never to 5=every day. Participants were classified as exposed to each biomechanical factor if it occurred at least one day per week. In particular, “bending or rotation movements” and “manual materials handling” were assessed using two items each. Since both items were intended to measure the same underlying construct, three categories of exposure to these biomechanical tasks were defined (both absent; one present; and both present) for the purpose of latent class analysis.

The remaining time of the working day not accounted for by any of the ten abovementioned biomechanical demands was classified as unexposed time.

**Regional musculoskeletal pain.** Regional musculoskeletal pain was assessed using the Nordic Musculoskeletal Questionnaire (20), which has been validated in the Portuguese population (21). This questionnaire evaluates the presence of musculoskeletal pain during the preceding 12 months in nine different anatomical regions: neck, shoulders, elbows, wrists/hands, upper back, lower back, hips/thighs/buttocks, knees and ankles/feet. In order to minimize potential misclassification and considering the expected effect of work-related demands on adjacent musculoskeletal regions, we aggregated anatomical sites into four major groups: neck and shoulders, upper limb (elbows and wrists/hands), upper and lower back, and lower limb (hips/thighs/buttocks, knees and ankles/feet). Moderate-to-severe pain was considered present if the reported intensity was >40 millimeters in the visual analogue scale (22). If more than one painful site was simultaneously present within aggregated regions, we chose the highest pain intensity. Additionally, the total number of painful anatomical sites was computed and varied between 0–9.

**Covariates.** Based on published evidence, anthropometrics and lifestyles (23), socioeconomic context (24), and psychosocial characteristics of the occupation (25) were considered *a priori* as confounders of the relation between occupational biomechanical demands and regional musculoskeletal pain.

Anthropometric parameters were obtained while subjects stood barefoot in light indoor clothing. Body mass index was then calculated as the weight (kilograms) divided by the square of height (meters) and individuals were grouped according to the World Health Organization classification: underweight (<18.5 kg/m<sup>2</sup>), normal weight (18.5–24.9 kg/m<sup>2</sup>), overweight (25.0–29.9 kg/m<sup>2</sup>) or obese (≥30 kg/m<sup>2</sup>). Duration and intensity of leisure time physical activity were also collected with participants classifying their activity

as: sitting most of the time, standing and/or walking (but not running) or very active. Frequency of smoking was gathered and young adults were grouped as never smokers, ex-smokers or current smokers. Educational level has been shown to be the most accurate indicator to assess the effect of socioeconomic position on health outcomes in the Portuguese population (26, 27). Accordingly, formal education was recorded as completed years of schooling and subjects were grouped according to the International Standard Classification of Education as ≤9, 10–12, or ≥13 years of education. Among workers, occupational psychosocial context was assessed using the self-administered Job Content Questionnaire (28) and participants’ jobs were categorized into one of the four quadrants of Karasek’s Job Strain Model: (i) low strain, high decision latitude and low psychological demands; (ii) passive, low decision latitude and low psychological demands; (iii) active, high decision latitude and high psychological demands; and (iv) high strain, low decision latitude and high psychological demands (29).

Additionally, overall exertion of work-related tasks was measured using the Borg scale (range: 6–20) (30) and individuals were categorized as having slight exertion (smallest scores), moderate exertion (13–14 points) or heavy exertion occupations (highest scores). Occupations were grouped according to the International Standard Classification of Occupations-88 (31) as low and high-skilled manual (armed forces, skilled agricultural and fishery workers, craft and related trade workers, plant and machine operators and assemblers and elementary occupations), low-skilled clerical (service workers and shop/market sales workers), intermediate-skilled clerical (clerks) and high-skilled clerical (legislators, senior officials, managers, professionals, technicians and associated professionals).

## Statistical analysis

**Patterns of occupational biomechanical demands.** Latent class analysis was used to uncover distinct groups of homogenous individuals from our sample in order to create patterns of biomechanical demands. This approach considers that the performance of an individual in a set of items is explained by a categorical latent (unmeasured) variable with K classes (32). Interpretation of the model is usually based on item profiles in each category, obtained from the probabilities of each subject endorsing each response item, conditional on class membership.

In this study, the number of latent classes (also referred to as patterns) was selected considering the models with the smallest Bayesian information criterion (BIC) and according to the Lo-Mendell-Rubin likelihood ratio test that quantifies the likelihood that the data can be described by a model with one-less class. Starting from one single class and increasing one class

at each step, the best solution was identified when the increase in the number of classes did not lead to a further decrease in the BIC. Based on previous knowledge regarding the data necessary to accurately measure the biomechanical exposure in the workplace (30, 33, 34), the ten occupational biomechanical factors described above (items) were used in the latent class analysis fitted using MPlus (V.5.2; Muthen & Muthen, Los Angeles, California, USA). Although the majority of participants had complete information for all items, 16.6% had missing values in one single item and 4.5% in two or more. As by default MPlus uses the full information maximum likelihood to estimate the latent class analysis model, individuals with missing values were not excluded, ie, they were classified into one of the classes using information from the remaining items.

To assess the criterion validity of the final class solution, we tested the relation between patterns of occupational biomechanical demands and education, work-related overall exertion, work type and occupational psychosocial context, ie, variables expected to vary between classes but not used to determine the class solution itself. The statistical significance of differences between classes was estimated using chi-square test, and post-hoc analyses were computed to determine which classes made a substantial contribution to the overall significance of the chi-square test (adjusted standardized residuals with absolute values >1.96 were considered relevant).

*Occupational biomechanical patterns and musculoskeletal pain.* Sample characteristics are presented as counts and proportions for all categorical variables. In order to quantify the associations between patterns of occupational biomechanical demands and regional musculoskeletal pain (presence and intensity), logistic regressions were performed using two different conceptual approaches: (i) demands versus non-workers: to assess the potential effect of each pattern of work-related demands versus no demands, we considered the whole sample and compared each of the four groups of workers with non-workers (the latter were 80% students and were considered unexposed to work-related biomechanical demands); (ii) high versus low work demands: to assess the potential effect of different physically-demanding patterns versus low demands, we considered only workers and compared each of the three groups with physically-demanding patterns with the low-demands group (the latter less exposed to work-related biomechanical demands). Using the same conceptual approaches, multinomial logistic regressions were also performed having multisite musculoskeletal pain as outcome.

Crude and adjusted odds ratios (OR and OR<sub>adj</sub>) with 95% confidence intervals (95% CI) were computed. For the demands versus non-workers approach, estimates

were adjusted for sex, body mass index, physical activity, smoking and education. In the high versus low work demands approach, in which confounding by occupational psychosocial context may occur, associations were additionally adjusted for job strain (Karasek's Job Strain Model).

Statistical analyses other than latent class analysis were performed using Stata® version 11.2 for Windows (Stata Corp LP, College Station, Texas, USA).

## Results

Table 1 summarizes the characteristics of EPITeen participants. Most subjects were female, had healthy weight, spent most of the time standing or walking in their leisure activities, were non-smokers, and had ≥13 schooling years. Among workers (37.5% of the whole sample), 51.4% were low-skilled clerical, 69.1% had been working ≤1 year, and 49.0% had a full-time job. Near a quarter of workers perceived their occupation as highly strained.

### Patterns of occupational biomechanical demands

As for occupational biomechanical patterns, BIC reached its optimum value at four or five classes, but the Lo-Mendell-Rubin likelihood ratio test of model fit confirmed that four classes were enough to describe the observed data (P=0.291). The average latent class probabilities (most likely latent class membership) showed high discrimination between classes, varying between 0.86–0.94. The final class solution was characterized by the following item profiles, which we named as follows: “low demands” assigned low probability to all items (eg, security guards and staff, receptionists, stewardesses and sales promoters); “sitting demands” was mainly characterized by sitting posture and computer use (>95%; eg, administrative staff, call-center operators, teachers and lawyers); “repetitive and asymmetric demands” presented high probability of repetitive tasks, bending or rotation movements and manual materials handling and intermediate probability (near 50%) of including overhead and kneeling work (eg, shop staff, market sales staff, and hairdressers); and “high and vibrational demands” allocated the highest probability to all items (except sitting and computer use) with a maximum probability of exposure to whole-body vibrations and vibration tools handling (eg, mechanics, electricians, warehouse staff and farmers) (table 2 and figure 1).

The criterion validity of the model is presented in table 3. High formal education was more frequent among workers with low-demands occupations (56.6%) and less frequent among high- and vibrational-demands

**Table 1.** Anthropometrics, lifestyles and socioeconomic characteristics of participants of the EPITeen cohort according to work status. Sample size is not constant due to missing data in body mass index (N=19), physical activity (N=2), smoking (N=2), education (N=1), work-related overall exertion (N=7), work type (N=68), work duration (N=65) and working time (N=41).

	Non-workers (N=1083)		Workers (N=650)	
	N	%	N	%
Sex				
Female	561	51.8	333	51.2
Male	522	48.2	317	48.8
Body mass index				
Underweight	67	6.3	34	5.3
Normal weight	766	71.6	430	66.8
Overweight	182	17.0	141	21.9
Obese	55	5.1	39	6.0
Physical activity				
Sitting most of the time	383	35.4	190	29.3
Standing and/or walking most of the time	505	46.7	275	42.4
Very active most of the time	194	17.9	184	28.3
Smoking				
Never smokers	315	29.1	173	26.7
Ex-smokers	371	34.3	237	36.6
Current smokers	397	36.6	238	36.7
Education				
≤9 years	62	5.7	77	11.8
10–12 years	180	16.7	284	43.7
≥13 years	840	77.6	289	44.5
Work-related overall exertion				
Slight exertion	.	.	224	34.8
Moderate exertion	.	.	256	39.8
Heavy exertion	.	.	163	25.4
Work type				
Low and high-skilled manual	.	.	77	13.2
Low-skilled clerical	.	.	299	51.4
Intermediate-skilled clerical	.	.	60	10.3
High-skilled clerical	.	.	146	25.1
Work duration				
<6 months	.	.	169	28.9
6–12 months	.	.	235	40.2
13–18 months	.	.	57	9.7
>18 months	.	.	124	21.2
Working time				
Full-time	.	.	298	48.9
Part-time (≥15 hours/week)	.	.	214	35.2
Part-time (<15 hours/week)	.	.	97	15.9
Occupational psychosocial context				
Low strain	.	.	153	23.5
Passive	.	.	129	19.8
Active	.	.	144	22.2
High strain	.	.	157	24.2
Missing	.	.	67	10.3

occupations (27.1%,  $P<0.001$ ). While workers with low- (46.1%) or sitting- (48.6%) demands occupations considered more frequently their work-related overall exertion as slight, workers with repetitive- and asymmetric- (13.7%) or high- and vibrational- (23.4%) demands occupations were those less frequently reporting their job exertion as slight ( $P<0.001$ ). Sitting demands (7.1%) were less frequently observed in manual jobs, whereas manual workers more frequently reported high- and vibrational-demands occupations

**Table 2.** Marginal percentage of subjects with each occupational biomechanical demand in each assigned latent class (pattern) to predict class membership among 21-year-old young adults (N=650).

	Demands (%)				
	Total	Low <sup>a</sup>	Sitting	Repetitive & asym-metric	High & vibra-tional
Total	28.0	28.2	21.7	22.2	
Sitting posture <sup>b</sup>					
≤4.2 hours/week	50.3	76.4	4.4	81.4	48.5
>4.2 hours/week	49.7	23.6	95.6	18.6	51.5
Computer use <sup>b</sup>					
≤5.0 hours/week	50.1	85.8	1.8	73.7	46.6
>5.0 hours/week	49.9	14.2	98.2	26.3	53.4
Whole body vibrations <sup>b</sup>					
≤0.8 hours/week	52.1	69.4	55.7	72.8	0.0
>0.8 hours/week	47.9	30.6	44.3	27.2	100
Vibration tools <sup>b</sup>					
≤0.4 hours/week	54.3	73.7	53.4	80.7	0.0
>0.4 hours/week	45.7	26.3	46.6	19.3	100
Precision work					
Less than weekly	84.9	98.3	91.2	80.9	62.4
At least weekly	15.1	1.7	8.8	19.1	37.6
Repetitive work					
Less than weekly	37.5	73.3	44.2	10.6	9.1
At least weekly	62.5	26.7	55.8	89.4	90.9
Overhead work					
Less than weekly	71.5	97.7	97.1	48.1	25.3
At least weekly	28.5	2.3	2.9	51.9	74.7
Kneeling work					
Less than weekly	74.4	98.2	98.6	47.1	37.3
At least weekly	25.6	1.8	1.4	52.9	62.7
Bending or rotation					
Less than weekly (both items)	53.8	79.7	89.5	11.7	13.4
One item present at least weekly	15.1	17.5	8.8	19.0	16.6
Both items present at least weekly	31.1	2.8	1.8	69.3	70.0
Manual materials handling					
Less than weekly (both items)	61.8	75.0	92.5	34.3	30.9
One item present at least weekly	16.6	15.9	3.9	32.3	18.4
Both items present at least weekly	21.6	9.1	3.5	33.4	50.7


<sup>a</sup> Workers with "low demands" spent, on average (per week), 3.4 hours sitting, 3.0 hours using the computer, 2.6 hours exposed to whole body vibrations and 1.7 hours handling vibration tools.

<sup>b</sup> The cut-offs were based on the median of weekly hours spent in each occupational biomechanical demand.

(24.8%,  $P<0.001$ ). Occupations with low (17.6%) or sitting (18.6%) demands were more frequently reported as highly strained while repetitive and asymmetric (36.2%) and high- and vibrational- (27.8%) demands occupations were less frequently considered to be highly strained ( $P<0.001$ ).

#### Patterns of occupational biomechanical demands and regional musculoskeletal pain

*Demands versus non-workers (workers versus non-workers).* Workers whose occupation was characterized by repetitive and asymmetric and high and vibrational demands were significantly more likely to report neck/



**Figure 1.** Marginal percentage of subjects with each occupational biomechanical demand in each assigned latent class (pattern) to predict class membership in 21-year-old young adults.

**Table 3.** Associations between patterns of occupational biomechanical demands and socioeconomic and occupational parameters in 21-year-old young adults. Sample size is not constant due to missing data in work-related overall exertion (N=7) and work type (N=68).

	Demands								P-value
	Low		Sitting		Repetitive & asymmetric		High & vibrational		
	N	%	N	%	N	%	N	%	
Total	182	28.0	183	28.2	141	21.7	144	22.2	
Education									
≤9 years	15	8.2	10	5.5 <sup>a</sup>	22	15.6	30	20.8 <sup>a</sup>	<0.001
10–12 years	64	35.2 <sup>a</sup>	85	46.4	60	42.6	75	52.1 <sup>a</sup>	
≥13 years	103	56.6 <sup>a</sup>	88	48.1	59	41.8	39	27.1 <sup>a</sup>	
Work-related overall exertion									
Slight exertion	83	46.1 <sup>a</sup>	89	48.6 <sup>a</sup>	19	13.7 <sup>a</sup>	33	23.4 <sup>a</sup>	<0.001
Moderate exertion	70	38.9	56	30.6 <sup>a</sup>	67	48.2 <sup>a</sup>	63	44.7	
Heavy exertion	27	15.0 <sup>a</sup>	38	20.8	53	38.1 <sup>a</sup>	45	31.9 <sup>a</sup>	
Work type									
Low and high-skilled manual	19	11.6	12	7.1 <sup>a</sup>	15	12.2	31	24.8 <sup>a</sup>	<0.001
Low-skilled clerical	103	62.8 <sup>a</sup>	47	27.6 <sup>a</sup>	82	66.7 <sup>a</sup>	67	53.6	
Intermediate-skilled clerical	9	5.5 <sup>a</sup>	39	22.9 <sup>a</sup>	7	5.7	5	4.0 <sup>a</sup>	
High-skilled clerical	33	20.1	72	42.4 <sup>a</sup>	19	15.4 <sup>a</sup>	22	17.6 <sup>a</sup>	
Occupational psychosocial context									
Low strain	48	26.4	51	27.9	26	18.4	28	19.4	<0.001
Passive	51	28.0 <sup>a</sup>	38	20.8	17	12.1 <sup>a</sup>	23	16.0	
Active	33	18.1	44	24.0	30	21.3	37	25.7	
High strain	32	17.6 <sup>a</sup>	34	18.6 <sup>a</sup>	51	36.2 <sup>a</sup>	40	27.8	
Missing	18	9.9	16	8.7	17	12.1	16	11.1	

<sup>a</sup> Substantial contribution to the overall significance of the chi-square test using the adjusted standardized residuals test.

**Table 4a.** Associations between patterns of occupational biomechanical demands and presence of regional musculoskeletal pain among 21-year-old young adults. [OR=odds ratio; 95% CI=95% confidence interval].

	Subjects reporting pain		Regional musculoskeletal pain							
			All participants <sup>a</sup>				Only workers <sup>b</sup>			
			Crude (N=1733)		Adjusted (N=1710)		Crude (N=650)		Adjusted (N=642)	
N	%	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	
Neck and shoulders										
Non-workers	484	44.7								
Workers (demands)										
Low	74	40.7	0.85	0.62–1.17	0.87	0.62–1.20				
Sitting	89	48.6	1.17	0.86–1.60	1.23	0.89–1.71	1.38	0.91–2.09	1.45	0.94–2.23
Repetitive & asymmetric	75	53.2	1.41	0.99–2.00	1.54	1.06–2.22	1.66	1.06–2.58	1.81	1.13–2.90
High & vibrational	76	52.8	1.38	0.98–1.96	1.56	1.08–2.27	1.63	1.05–2.54	1.82	1.14–2.90
Upper limb										
Non-workers	268	24.7								
Workers (demands)										
Low	46	25.3	1.03	0.72–1.48	1.03	0.71–1.48				
Sitting	50	27.3	1.14	0.80–1.63	1.10	0.76–1.59	1.11	0.70–1.77	1.05	0.64–1.70
Repetitive & asymmetric	49	34.8	1.62	1.12–2.35	1.57	1.06–2.32	1.57	0.97–2.55	1.56	0.94–2.60
High & vibrational	41	28.5	1.21	0.82–1.78	1.20	0.79–1.80	1.18	0.72–1.93	1.26	0.75–2.12
Upper and lower back										
Non-workers	601	55.5								
Workers (demands)										
Low	106	58.2	1.12	0.81–1.54	1.08	0.78–1.51				
Sitting	124	67.8	1.69	1.21–2.35	1.70	1.20–2.42	1.51	0.98–2.31	1.56	0.99–2.45
Repetitive & asymmetric	92	65.2	1.51	1.04–2.17	1.37	0.93–2.02	1.35	0.85–2.12	1.27	0.78–2.05
High & vibrational	102	70.8	1.95	1.33–2.84	1.95	1.30–2.92	1.74	1.09–2.77	1.80	1.09–2.96
Lower limb										
Non-workers	462	42.7								
Workers (demands)										
Low	82	45.1	1.10	0.80–1.51	1.07	0.78–1.48				
Sitting	73	39.9	0.89	0.65–1.23	0.83	0.60–1.16	0.81	0.53–1.23	0.80	0.52–1.23
Repetitive & asymmetric	61	43.3	1.02	0.72–1.46	0.98	0.68–1.42	0.93	0.60–1.45	0.91	0.57–1.44
High & vibrational	58	40.3	0.91	0.64–1.29	0.90	0.62–1.31	0.82	0.53–1.28	0.82	0.51–1.31

<sup>a</sup> The reference category comprised the young non-worker adults; multiple logistic regressions consider adjustment for sex, body mass index, physical activity, smoking and education.

<sup>b</sup> The reference category comprised the young adult workers with occupational low demands; multiple logistic regressions consider adjustment for sex, body mass index, physical activity, smoking, education and occupational psychosocial context.

shoulder pain when compared to non-workers [respectively, OR<sub>adj</sub> 1.54 (95% CI 1.06–2.22) and 1.56 (95% CI 1.08–2.27)]. Sitting and high and vibrational demands were significantly associated with the report of upper-/lower-back pain when compared to non-workers: OR<sub>adj</sub> 1.70 (95% CI 1.20–2.42) and 1.95 (95% CI 1.30–2.92), respectively. Sitting, repetitive and asymmetric, and high and vibrational demands were significantly associated with moderate-to-severe upper-/lower-back pain with OR<sub>adj</sub> ranging between 1.56 (95% CI 1.05–2.33) and 1.62 (95% CI 1.14–2.31) (tables 4a and 4b). As observed in figure 2, workers whose occupation was characterized by high and vibrational demands were more likely than non-workers to report pain in at least one anatomical site, and this effect was stronger as we increased the threshold of number of painful anatomical regions: OR<sub>adj1</sub> 2.21 (95% CI 1.08–4.50), OR<sub>adj2</sub> 2.75 (95% CI 1.34–5.62), and OR<sub>adj≥3</sub> 2.81 (95% CI 1.43–5.52).

*High versus low work demands (workers with physically-demanding patterns versus low demands).* As presented in tables 4a and 4b, workers exposed to occupations with repetitive and asymmetric and high and vibrational demands were more likely to report neck/shoulder pain than those with low demands [OR<sub>adj</sub> 1.81 (95% CI 1.13–2.90) and 1.82 (95% CI 1.14–2.90), respectively]. High and vibrational demands were significantly associated to the presence of upper-/lower-back pain when compared to low demands (OR<sub>adj</sub> 1.80, 95% CI 1.09–2.96). Workers with sitting demands were more likely to report any or moderate-to-severe upper-/lower-back pain than those with low demands: OR<sub>adj</sub> 1.56 (95% CI 0.99–2.45) and 1.66 (95% CI 1.03–2.66) respectively. High and vibrational demands were associated with pain in at least one anatomical site when compared to low demands, and the magnitude of associations was stronger as the threshold of number of painful anatomical regions increased: OR<sub>adj1</sub> 2.56 (95% CI 1.08–6.08), OR<sub>adj2</sub> 2.67 (95% CI 1.13–6.34), OR<sub>adj≥3</sub> 2.85 (95% CI 1.26–6.42) (figure 2).



**Table 4b.** Associations between patterns of occupational biomechanical demands and moderate to severe regional musculoskeletal pain in 21-year-old young adults. [VAS=visual analogue scale; OR=odds ratio; 95% CI=95% confidence interval].

	Subjects reporting pain		Moderate to severe regional musculoskeletal pain (VAS >40 millimetres)							
			All participants <sup>a</sup>				Only workers <sup>b</sup>			
			Crude (N=1733)		Adjusted (N=1710)		Crude (N=650)		Adjusted (N=642)	
N	%	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	
Neck and shoulders										
Non-workers	172	15.9								
Workers (demands)										
Low	29	15.9	1.00	0.65–1.54	1.00	0.64–1.54				
Sitting	34	18.6	1.21	0.80–1.81	1.17	0.77–1.79	1.20	0.70–2.07	1.24	0.71–2.18
Repetitive & asymmetric	35	24.8	1.75	1.15–2.65	1.56	1.01–2.42	1.74	1.00–3.02	1.61	0.90–2.88
High & vibrational	28	19.4	1.28	0.82–1.99	1.21	0.75–1.93	1.27	0.72–2.26	1.27	0.70–2.33
Upper limb										
Non-workers	106	9.8								
Workers (demands)										
Low	13	7.1	0.71	0.39–1.29	0.68	0.37–1.26				
Sitting	14	7.7	0.76	0.43–1.36	0.66	0.36–1.22	1.08	0.49–2.36	0.92	0.40–2.09
Repetitive & asymmetric	19	13.5	1.44	0.85–2.42	1.18	0.67–2.06	2.02	0.96–4.26	1.57	0.71–3.46
High & vibrational	19	13.2	1.40	0.83–2.36	1.24	0.71–2.16	1.98	0.94–4.15	1.78	0.81–3.88
Upper and lower back										
Non-workers	270	24.9								
Workers (demands)										
Low	48	26.4	1.08	0.75–1.54	0.98	0.67–1.43				
Sitting	68	37.2	1.78	1.28–2.48	1.62	1.14–2.31	1.65	1.06–2.58	1.66	1.03–2.66
Repetitive & asymmetric	56	39.7	1.98	1.38–2.86	1.60	1.09–2.37	1.84	1.15–2.95	1.60	0.96–2.66
High & vibrational	53	36.8	1.75	1.22–2.53	1.56	1.05–2.33	1.63	1.01–2.61	1.60	0.96–2.66
Lower limb										
Non-workers	210	19.4								
Workers (demands)										
Low	46	25.3	1.41	0.97–2.03	1.29	0.88–1.89				
Sitting	45	24.6	1.36	0.94–1.96	1.19	0.81–1.76	0.96	0.60–1.55	0.93	0.57–1.53
Repetitive & asymmetric	41	29.1	1.70	1.15–2.53	1.58	1.04–2.39	1.21	0.74–1.99	1.21	0.72–2.05
High & vibrational	33	22.9	1.24	0.81–1.87	1.14	0.73–1.78	0.88	0.53–1.47	0.90	0.53–1.55

<sup>a</sup> The reference category comprised the young non-worker adults; multiple logistic regressions consider adjustment for sex, body mass index, physical activity, smoking and education.

<sup>b</sup> The reference category comprised the young adult workers with occupational low demands; multiple logistic regressions consider adjustment for sex, body mass index, physical activity, smoking, education and occupational psychosocial context.

## Discussion

In this community-based study of young adults, we identified four clearly distinct patterns of exposure to occupational biomechanical demands. Even in the early stages of employment, repetitive and asymmetric and high and vibrational demands were associated with the presence of neck/shoulder pain. Patterns of increased biomechanical demands, especially sitting and high and vibrational demands, showed higher risk of any or severe upper-/lower-back pain. Moreover, the odds of multisite pain increased with growing biomechanical workplace demands.

In the beginning of professional life, work-related demands may affect health outcomes not only through a direct biomechanical effect on body structures, but also because labor market entry is a stressor in itself (18). Our comparison between subjects with each exposure pattern and non-workers intended to measure the cumulative effect of being employed plus having a certain pattern

of demands versus being unexposed to employment-related mechanical demands (ie, background propensity for pain in the source population). The comparison with the low demands group (exposed to employment but less exposed to mechanical demands) aimed to isolate the effects of the demands themselves. Despite this conceptual difference, associations remained very similar regardless of reference group, suggesting that meaningful differences are attributable to demanding biomechanical exposures rather than employment status by itself.

Even short-term exposures to repetitive and asymmetric or high and vibrational demands were associated with neck/shoulder pain in this investigation. Previous studies in samples of newly-employed workers have shown that repetitiveness, sustained trapezius muscle activity, and high force demands are significant predictors of neck and shoulder pain (11, 35, 36). Additionally, high and vibrational demands were also associated with upper-/lower-back pain. Occupational mechanical

load – lifting and pushing, pulling heavy weights or overhead tasks – has also been pointed out as playing a major role in the onset of low-back pain (14), even among adolescents with regular or irregular exposures to physically-demanding jobs (18).

In addition to the well-documented fact that work-related musculoskeletal pain can be caused by fibrotic and structural changes from tissue injury and consequent micro-trauma (37–39), work-related physically-demanding tasks have also been shown to be related to the occurrence of monotonic muscle contractions that may cause pain (40, 41). This is expected to be even more pronounced among young workers since their willingness to comply with all the work-related tasks in a short period of time may lead to involuntary muscle contraction in an uncoordinated way without relaxing pauses, which may ultimately result in regional pain. (42). Concordantly, our findings supported that short-term exposures to physically-demanding patterns of occupational biomechanical demands may have been sufficient to cause meaningful permanent or transient disturbance to neck/shoulder and upper/lower back structures.

Another site-specific harmful effect was observed regarding the pattern of sitting demands and the presence of any or severe upper-/lower-back pain. Due to the low activation of lumbar muscles while sitting, load is absorbed by viscoelastic structures and transmitted directly to the spine, causing muscle contractions and/or structural damage and consequent severe back pain (43). Nevertheless, controversy still exists regarding the impact of occupational sitting posture on back pain (17), since extremely high doses of exposure are supposedly needed to affect spinal structures (44). The detection of this effect in such early stages of employment may be the result of our empirical characterization of demands that captured a more realistic pattern in which computer use and sitting posture co-occur. We believe that this better represents overall sitting-related demands to the detriment of traditional measurements of time spent sitting as a single and independent exposure. An additional explanation is the potential independent effect of workplace rules and obligations, which are frequently interpreted as rigid and inflexible in the beginning of professional life and may lead to excessive periods of time spent uninterruptedly sitting (45).

Another relevant result of our study was that the risk of multisite pain increased with growing occupational biomechanical demands. Particularly, for high and vibrational demands occupations, associations were stronger as we increased the threshold of the number of painful anatomical regions, which is in agreement with previous evidence in long-term workers (46, 47). These findings suggest that, from the earliest stages of employment, high demands may rapidly produce musculoskeletal pain in more than one body region (particularly in the neck/

shoulder and upper-/lower-back areas, ie, the regions most frequently reported as painful). Adding to that, in the specific case of multisite pain, it is likely that the overall psychosocial context of the occupation contributes to the reported symptoms in addition to the potential effects of biomechanical exposures. However, we do not believe that the former was a major confounder here since our estimates were adjusted for job strain.


With scarce exceptions, we did not observe significant associations between patterns of biomechanical demands and musculoskeletal pain in the upper or lower limbs. The literature has been inconsistent in detecting an influence of work-related physical demands on limb pain in early stages of professional life (12, 13, 48) and, when associations have been found, they are commonly much weaker than those estimated for shoulders or back regions. In such early stages of professional life, it seems that the duration of exposure to adverse biomechanical demands is too short to produce tissue damage above the biological threshold for musculoskeletal pain.

To the best of our knowledge, this study was a pioneer in using a population-based approach to estimate the associations between empirically-defined patterns of occupational biomechanical demands and musculoskeletal pain among young adults in the beginning of professional life. Our population-based approach adds a scenario of a wide range of real-world occupations performed in a decade where the typology of the early stages of employment has changed dramatically (49). This updates previous knowledge and supports the generalization of our results to other young adult populations.

Additionally, this study aimed to identify patterns using an extensive characterization of interrelated biomechanical demands that may function as markers of early development of pain and, consequently, may be the target for intervention since the beginning of the professional life. Our latent class approach provided an innovative, pragmatic and likely more valid characterization of the exposure since it considers the co-occurrence of different but closely linked work-related biomechanical tasks (8).

Furthermore, we evaluated not only the presence of pain, but also severity and its co-occurrence in different anatomical regions. We believe that considering different pain outcomes (anatomical diversity, intensity and number of painful regions) allowed us not only to study the effect of occupational biomechanical demands across a wide spectrum of pain severity stages, but also to validate our findings regarding the anatomical plausibility of exposure-outcome relations, since the strength and statistical significance of associations were consistent across different outcomes.

Nevertheless, some methodological concerns should be addressed. Individuals comprising the non-workers



**Figure 2.** Adjusted associations between patterns of occupational biomechanical demands and multisite musculoskeletal pain among 21-year-old young adults. <sup>a</sup> The comparison group comprised the young adult non-workers without regional musculoskeletal pain (adjusted for sex, body mass index, physical activity, smoking and education). <sup>b</sup> The comparison group comprised the young adult workers in the occupational pattern of low demands without regional musculoskeletal pain (adjusted for sex, body mass index, physical activity, smoking, education and occupational psychosocial context).

reference group were not totally “unexposed” to biomechanical demands, but we did not collect detailed demands in this group. Nevertheless, a rapid assessment of physical exertion showed that nearly three quarters of non-workers reported to spend most of the day sitting (data not shown). In line with this, low-demand workers were exposed to all types of biomechanical demands, but all of them with low duration and frequency. It is likely that the remaining time (not accounted for by any of the biomechanical demands considered) was spent standing or walking without any other simultaneous physical demand, which in our perspective qualifies as unexposed time. This assumption was supported by the jobs included (vigilantes and security staff, receptionists, stewardesses and promoters), but we did not collect information on the duration/frequency of periods with no (or nearly null) biomechanical demands, ie we characterized exposure to a set of frequent and work-related demanding exposures and assumed that the remaining working time was unexposed.

As workers were at the beginning of their professional lives, exposure to occupational biomechanical demands was assumed to be reasonably short. This assumption was corroborated by our observation that the majority of participants had been working for  $\leq 12$  months (nearly 70%) and the maximum duration of work was 72 months.

Due to its cross-sectional design, conclusions on the temporality can be argued but not proven and prudence is needed when inferring causality from the associations found in this investigation. Firstly, although EPITeen participants have been followed since early adoles-

cence, data on musculoskeletal pain were not collected before 21 years of age. Therefore, we were not able to distinguish between acute and chronic complaints. Our methodological option of not assessing pain history cross-sectionally was related to the high likelihood of differential recall of past pain according to current complaints (50). As musculoskeletal complaints often begin in early stages of life (51), psychosocial and lifestyle pathways other than occupational exposures are plausible to have caused pain onset (52). Within this framework, we cannot exclude that our findings were the result of alternative pathways of musculoskeletal pain development throughout the life course, where occupational biomechanical demands may only partly explain musculoskeletal complaints. To deal with this, we adjusted estimates for other documented psychosocial causes of pain, which in our view have minimized potential confounding. Secondly, the healthy worker effect, through which musculoskeletal symptoms may themselves determine the choice or persistence of a specific less demanding occupation, cannot be excluded. However, it seems unlikely that pain has led a substantial proportion of individuals to radically change their exposure status before the age of 21 years. In fact, we believe it to be more likely that biomechanical factors have preceded pain rather than earlier pain has determined the selection of a specific less-demanding occupation. Thirdly, as exposures and outcomes were collected retrospectively, individuals with pain symptoms may have recalled occupational exposures as more frequent and/or demanding than those who did not present such complaints. Nevertheless, in this study, data have been

collected as part of an extensive protocol that assessed a wide spectrum of health-related information and subjects were not expected to be particularly aware of the main hypothesis addressed in this particular study.

Regional musculoskeletal complaints were assessed using the widely accepted standardized recall period of 12 months. Since our sample included only young adult workers, a relevant percentage of those had been employed for <1 year: 28.9% for <6 months, 21.2% for 6–9 months and 5.0% for 10–11 months. Therefore, recall times for exposure and outcome did not completely overlap in this fraction of the sample, ie, musculoskeletal pain might have referred to an unexposed period with regard to occupational biomechanical demands. Sensitivity analyses were performed by restricting estimates to individuals that had been working for  $\geq 1$  year and similar conclusions were supported by the results (appendix 2, [http://www.sjweh.fi/data\\_repository.php](http://www.sjweh.fi/data_repository.php)), despite the reduced statistical power to identify significant differences between groups.

Finally, the extent to which any cohort represents the source population throughout follow-up is likely to decrease over time because of differential losses to follow-up. Participants reassessed at 21 years were reasonably similar to those lost to follow-up in a wide number of characteristics measured in the baseline evaluation, but attrition was significantly higher among individuals with low social position. Despite that, we do not believe that the underrepresentation of young adults with social disadvantage has substantially biased our estimates since we found no associations between parental education and musculoskeletal pain.

In conclusion, we found four clearly distinct patterns of biomechanical demands in the general population of young workers. Even in early stages of employment, physically-demanding patterns at the workplace were associated with the presence of neck/shoulder pain and with the presence and severity of upper-/lower-back pain. Our study emphasizes that even short-term biomechanical exposures at the workplace may be involved in the etiology of musculoskeletal complaints.

### Acknowledgments

This work was supported by the Portuguese Foundation for Science and Technology (FCT, references: FCOMP-01-0124-FEDER-015750, FCT – PTDC/SAU-EPI/115254/2009). Sara Lourenço gratefully acknowledges a doctoral grant from the Portuguese Foundation for Science and Technology (FCT; reference SFRH/BD/77965/2011). We also gratefully acknowledge Dr. Cristina Mesquita for kindly providing the Portuguese version of the Nordic Musculoskeletal Questionnaire

used in this study. All authors declare that there are no conflicts of interest.

### References

1. Woolf AD, Akesson K. Understanding the burden of musculoskeletal conditions. The burden is huge and not reflected in national health priorities. *BMJ*. 2001;322:1079–80. <http://dx.doi.org/10.1136/bmj.322.7294.1079>.
2. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2197–223. [http://dx.doi.org/10.1016/S0140-6736\(12\)61689-4](http://dx.doi.org/10.1016/S0140-6736(12)61689-4).
3. European Foundation for the Improvement of Living and Working Conditions. 4th European working conditions survey. Luxembourg: Office for Official Publications of the European Communities: 2007.
4. Lidgren L, Smolen J, Bentley G, Delmas P, Woolf A, Åkesson K, et al. European action towards better musculoskeletal health: a public health strategy to reduce the burden of musculoskeletal conditions. Lund, Sweden: The Bone & Joint Decade Department of Orthopedics University Hospital; 2005.
5. Morse T, Dillon C, Kenta-Bibi E, Weber J, Diva U, Warren N, et al. Trends in work-related musculoskeletal disorder reports by year, type, and industrial sector: a capture-recapture analysis. *Am J Ind Med*. 2005;48:40–9. <http://dx.doi.org/10.1002/ajim.20182>.
6. Neupane S, Virtanen P, Leino-Arjas P, Miranda H, Siukola A, Nygard CH. Multi-site pain and working conditions as predictors of work ability in a 4-year follow-up among food industry employees. *Eur J Pain*. 2013;17:444–51. <http://dx.doi.org/10.1002/j.1532-2149.2012.00198.x>.
7. Bernard BP. Musculoskeletal disorders and workplace factors: a critical review of epidemiologic evidence for work-related musculoskeletal disorders of the neck, the upper-limb, and low back. Cincinnati, Ohio: U.S. Department of Health and Human Services (National Institute of Occupational Safety and Health): 1997.
8. Punnett L, Wegman DH. Work-related musculoskeletal disorders: the epidemiologic evidence and the debate. *J Electromyogr Kinesiol*. 2004;14:13–23. <http://dx.doi.org/10.1016/j.jelekin.2003.09.015>.
9. Mayer J, Kraus T, Ochsmann E. Longitudinal evidence for the association between work-related physical exposures and neck and/or shoulder complaints: a systematic review. *Int Arch Occup Environ Health*. 2012;85:587–603. <http://dx.doi.org/10.1007/s00420-011-0701-0>.
10. Bang Christensen K, Lund T, Labriola M, Villadsen E, Bultmann U. The fraction of long-term sickness absence attributable to work environmental factors: prospective results from the Danish Work Environment Cohort Study. *Occup Environ Med*. 2007;64:487–9. <http://dx.doi.org/10.1136>

- [oem.2006.028563](http://oem.2006.028563).
11. Harkness EF, Macfarlane GJ, Nahit ES, Silman AJ, McBeth J. Mechanical and psychosocial factors predict new onset shoulder pain: a prospective cohort study of newly employed workers. *Occup Environ Med.* 2003;60:850–7. <http://dx.doi.org/10.1136/oem.60.11.850>.
  12. Nahit ES, Taylor S, Hunt IM, Silman AJ, Macfarlane GJ. Predicting the onset of forearm pain: a prospective study across 12 occupational groups. *Arthritis Rheum.* 2003;49:519–25. <http://dx.doi.org/10.1002/art.11202>.
  13. Jones GT, Harkness EF, Nahit ES, McBeth J, Silman AJ, Macfarlane GJ. Predicting the onset of knee pain: results from a 2-year prospective study of new workers. *Ann Rheum Dis.* 2007;66:400–6. <http://dx.doi.org/10.1136/ard.2006.057570>.
  14. Harkness EF, Macfarlane GJ, Nahit ES, Silman AJ, McBeth J. Risk factors for new-onset low back pain amongst cohorts of newly employed workers. *Rheumatology (Oxford).* 2003;42:959–68. <http://dx.doi.org/10.1093/rheumatology/keg265>.
  15. Garg A, Kapellusch JM. Applications of biomechanics for prevention of work-related musculoskeletal disorders. *Ergonomics.* 2009;52:36–59. <http://dx.doi.org/10.1080/00140130802480794>.
  16. Radwin RG, Marras WS, Lavender SA. Biomechanical aspects of work-related musculoskeletal disorders. *Theor Issues Ergon.* 2001;2:153–217. <http://dx.doi.org/10.1080/14639220110102044>.
  17. Lis AM, Black KM, Korn H, Nordin M. Association between sitting and occupational LBP. *Eur Spine J.* 2007;16:283–98. <http://dx.doi.org/10.1007/s00586-006-0143-7>.
  18. Mikkonen P, Viikari-Juntura E, Remes J, Pienimäki T, Solovieva S, Taimela S, et al. Physical workload and risk of low back pain in adolescence. *Occup Environ Med.* 2012;69:284–90. <http://dx.doi.org/10.1136/oemed-2011-100200>.
  19. Ramos E, Barros H. Family and school determinants of overweight in 13-year-old Portuguese adolescents. *Acta Paediatr.* 2007;96:281–6. <http://dx.doi.org/10.1111/j.1651-2227.2007.00107.x>.
  20. Kuorinka I, Jonsson B, Kilbom A, Vinterberg H, Biering-Sorensen F, Andersson G, et al. Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Appl Ergon.* 1987;18:233–7. [http://dx.doi.org/10.1016/0003-6870\(87\)90010-X](http://dx.doi.org/10.1016/0003-6870(87)90010-X).
  21. Mesquita CC, Ribeiro JC, Moreira P. Portuguese version of the standardized Nordic musculoskeletal questionnaire: Cross cultural and reliability. *J Public Health (Oxf).* 2010;18:461–6. <http://dx.doi.org/10.1007/s10389-010-0331-0>.
  22. Breivik H, Borchgrevink PC, Allen SM, Rosseland LA, Romundstad L, Hals EK, et al. Assessment of pain. *Br J Anaesth.* 2008;101:17–24. <http://dx.doi.org/10.1093/bja/aen103>.
  23. Nilsen TI, Holtermann A, Mork PJ. Physical exercise, body mass index, and risk of chronic pain in the low back and neck/shoulders: longitudinal data from the Nord-Trøndelag Health Study. *Am J Epidemiol.* 2011;174:267–73. <http://dx.doi.org/10.1093/aje/kwr087>.
  24. Poleshuck EL, Green CR. Socioeconomic disadvantage and pain. *Pain.* 2008;136:235–8. <http://dx.doi.org/10.1016/j.pain.2008.04.003>.
  25. Torp S, Riise T, Moen BE. The impact of psychosocial work factors on musculoskeletal pain: a prospective study. *J Occup Environ Med.* 2001;43:120–6. <http://dx.doi.org/10.1097/00043764-200102000-00010>.
  26. Alves L, Azevedo A, Silva S, Barros H. Socioeconomic inequalities in the prevalence of nine established cardiovascular risk factors in a southern European population. *PLoS One.* 2012;7:e37158. <http://dx.doi.org/10.1371/journal.pone.0037158>.
  27. Bastos J, Peleteiro B, Barros R, Alves L, Severo M, de Fatima Pina M, et al. Sociodemographic determinants of prevalence and incidence of *Helicobacter pylori* infection in Portuguese adults. *Helicobacter.* 2013;18:413–22. <http://dx.doi.org/10.1111/hel.12061>.
  28. Karasek R, Brisson C, Kawakami N, Houtman I, Bongers P, Amick B. The Job Content Questionnaire (JCQ): an instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol.* 1998;3:322–55. <http://dx.doi.org/10.1037/1076-8998.3.4.322>.
  29. Karasek RA. Job Demands, Job Decision Latitude, and Mental Strain - Implications for Job Redesign. *Admin Sci Quart.* 1979;24:285–308. <http://dx.doi.org/10.2307/2392498>.
  30. Borg G. Psychophysical scaling with applications in physical work and the perception of exertion. *Scand J Work Environ Health.* 1990;16 Suppl 1:55–8. <http://dx.doi.org/10.5271/sjweh.1815>.
  31. International Labour Office. International Standard Classification of Occupations (ISCO-88): structure, group definitions and correspondence tables. Geneva: 2012.
  32. Bartholomew DJ, Knott M. Latent variable models and factor analysis. 2nd ed. London, New York: Arnold, Oxford University Press; 1999.
  33. Holmann S, Klimmer F, Schmidt KH, Kylian H. Validation of a questionnaire for assessing physical work load. *Scand J Work Environ Health.* 1999;25:105–14. <http://dx.doi.org/10.5271/sjweh.412>.
  34. Occhipinti E. OCRA: a concise index for the assessment of exposure to repetitive movements of the upper limbs. *Ergonomics.* 1998;41:1290–311. <http://dx.doi.org/10.1080/001401398186315>.
  35. Hakkanen M, Viikari-Juntura E, Martikainen R. Incidence of musculoskeletal disorders among newly employed manufacturing workers. *Scand J Work Environ Health.* 2001;27:381–7. <http://dx.doi.org/10.5271/sjweh.630>.
  36. Hanvold TN, Waersted M, Mengshoel AM, Bjertness E, Stigum H, Twisk J, et al. The effect of work-related sustained trapezius muscle activity on the development of neck and shoulder pain among young adults. *Scand J Work Environ Health.* 2013;39:390–400. <http://dx.doi.org/10.5271/sjweh.3357>.
  37. Barbe MF, Barr AE. Inflammation and the pathophysiology of work-related musculoskeletal disorders. *Brain Behav Immun.*

- 2006;20:423–9. <http://dx.doi.org/10.1016/j.bbi.2006.03.001>.
38. Andersen JH, Haahr JP, Frost P. Risk factors for more severe regional musculoskeletal symptoms: a two-year prospective study of a general working population. *Arthritis Rheum.* 2007;56:1355–64. <http://dx.doi.org/10.1002/art.22513>.
39. Pope MH, Goh KL, Magnusson ML. Spine ergonomics. *Annu Rev Biomed Eng.* 2002;4:49–68. <http://dx.doi.org/10.1146/annurev.bioeng.4.092101.122107>.
40. Hostens I, Ramon H. Assessment of muscle fatigue in low level monotonous task performance during car driving. *J Electromyogr Kinesiol.* 2005;15:266–74. <http://dx.doi.org/10.1016/j.jelekin.2004.08.002>.
41. Sbriccoli P, Yousuf K, Kupershtein I, Solomonow M, Zhou BH, Zhu MP, et al. Static load repetition is a risk factor in the development of lumbar cumulative musculoskeletal disorder. *Spine (Phila Pa 1976).* 2004;29:2643–53. <http://dx.doi.org/10.1097/01.brs.0000146052.44581.5f>.
42. Hughes LE, Babski-Reeves K, Smith-Jackson T. Effects of psychosocial and individual factors on physiological risk factors for upper extremity musculoskeletal disorders while typing. *Ergonomics.* 2007;50:261–74. <http://dx.doi.org/10.1080/00140130601049378>.
43. Morl F, Bradl I. Lumbar posture and muscular activity while sitting during office work. *J Electromyogr Kinesiol.* 2013;23:362–8. <http://dx.doi.org/10.1016/j.jelekin.2012.10.002>.
44. Mork PJ, Westgaard RH. Back posture and low back muscle activity in female computer workers: a field study. *Clin Biomech (Bristol, Avon).* 2009;24:169–75. <http://dx.doi.org/10.1016/j.clinbiomech.2008.11.001>.
45. Van Nieuwenhuysse A, Fatkhutdinova L, Verbeke G, Pirenne D, Johannik K, Somville PR, et al. Risk factors for first-ever low back pain among workers in their first employment. *Occup Med (Lond).* 2004;54:513–9. <http://dx.doi.org/10.1093/occmed/kqh091>.
46. Solidaki E, Chatzi L, Bitsios P, Coggon D, Palmer KT, Kogevas M. Risk factors for new onset and persistence of multi-site musculoskeletal pain in a longitudinal study of workers in Crete. *Occup Environ Med.* 2013;70:29–34. <http://dx.doi.org/10.1136/oemed-2012-100689>.
47. Solidaki E, Chatzi L, Bitsios P, Markatzi I, Plana E, Castro F, et al. Work-related and psychological determinants of multisite musculoskeletal pain. *Scand J Work Environ Health.* 2010;36:54–61. <http://dx.doi.org/10.5271/sjweh.2884>.
48. Bonde JP, Mikkelsen S, Andersen JH, Fallentin N, Baelum J, Svendsen SW, et al. Understanding work related musculoskeletal pain: does repetitive work cause stress symptoms? *Occup Environ Med.* 2005;62:41–8. <http://dx.doi.org/10.1136/oem.2003.011296>.
49. European Commission. *New skills and jobs in Europe: pathways towards full employment.* Belgium: Brussels; 2012.
50. Linton SJ, Melin L. The accuracy of remembering chronic pain. *Pain.* 1982;13:28–5. [http://dx.doi.org/10.1016/0304-3959\(82\)90017-3](http://dx.doi.org/10.1016/0304-3959(82)90017-3).
51. O’Sullivan PB, Beales DJ, Smith AJ, Straker LM. Low back pain in 17 year olds has substantial impact and represents an important public health disorder: a cross-sectional study. *BMC Public Health.* 2012;12:100. <http://dx.doi.org/10.1186/1471-2458-12-100>.
52. Jussila L, Paananen M, Nayha S, Taimela S, Tammelin T, Auvinen J, et al. Psychosocial and lifestyle correlates of musculoskeletal pain patterns in adolescence: a 2-year follow-up study. *Eur J Pain.* 2014;18:139–46. <http://dx.doi.org/10.1002/j.1532-2149.2013.00353.x>.

Received for publication: 14 November 2014