

Vegetation in the vicinity of the Toolik Field Station, Alaska

Donald A. Walker and Hilmar A. Maier
Alaska Geobotany Center, Institute of Arctic Biology, University of Alaska Fairbanks

Funded by: U.S. National Science Foundation grant no. ARC-0225517

Suggested citation: Walker, D.A. and H.A. Maier. 2008. Vegetation in the vicinity of the Toolik Field Station, Alaska. Biological Papers of the University of Alaska, No. 28. Institute of Arctic Biology, Fairbanks, Alaska.

ISSN: 0568-8604, No. 28 ISBN: 978-0-9767525-2-3

The maps presented are components of the Arctic Geobotanical Atlas (<http://www.ArcticAtlas.org/>).

© 2008, Institute of Arctic Biology, Biological Papers of the University of Alaska, University of Alaska Fairbanks, PO Box 757000, Fairbanks, AK 99775-7000. UAF is an AA/E/O employer and educational institution.

Explanation

Overview

This publication contains a group of vegetation maps at three scales in the vicinity of the Toolik Field Station, Alaska, which is an arctic research facility run by the Institute of Arctic Biology at the University of Alaska Fairbanks. The maps are intended to support research at the field station. The front side of this map sheet contains a vegetation map and ancillary maps of a 751-km² region surrounding the upper Kuparuk River watershed, including the Toolik Lake and the Creek research areas, as well as portions of the Dalton Highway and Trans-Alaska Pipeline from the northern end of Galbraith Lake to Slope Mountain. The reverse side shows more detailed vegetation maps of the 20-km² research area centered on Toolik Lake and a 12-km² intensive research grid on the south side of Toolik Lake (red rectangles on Map A). All the maps are part of a hierarchical geographic information system (GIS) and the Web-based Arctic Geobotanical Atlas (<http://www.arcticatlas.org/>). The atlas also includes other map themes for all three areas and a previously published hierarchy of maps of the Inuvait Creek area (Walker et al. 1989; Walker and Walker 1996) (black rectangles on Map A). Photos and explanations of the geobotanical mapping units and the supporting field data and metadata can also be found on the website.

Maps of the Upper Kuparuk River Region

The upper Kuparuk River region has terrain typical of the Southern Foothills of the Brooks Range, including landscapes affected by three major glacial events. Map A shows the vegetation of the upper Kuparuk River region at 1:63,360 scale. Other maps show a false-color infrared satellite image (Map B), glacial geology (Map C), surficial geomorphology (Map D) and Normalized Difference Vegetation Index (NDVI) biomass (Map E) - all at 1:225,000 scale. Maps A and C were derived from a geobotanical map of the region. The base map for the geobotanical map was a 1:25,000-scale black-and-white orthophoto-topographic map that was prepared especially for this project by Xerox Corp., Denver, CO in 1994 from stereo pairs of 160,000-scale, 9 x 9-inch color-infrared aerial photographs that were obtained by NASA in 1982. The base map was prepared without ground-control points, but was registered as closely as possible to the 1:63,360 USGS map of the region. Vegetation and other geobotanical features were mapped by photo-interpretation onto 1:25,000-scale enlargements of the 1982 NASA aerial photographs. The minimum mapping unit was approximately 0.6 ha (1/8" at 1:25,000-scale). No formal accuracy assessment was performed, but 320 of the map polygons representing 3.2% of the total map polygons, and about 16% of the total map area were checked on the ground during helicopter-assisted transects in 1994. Geobotanical variables coded for each map polygon include: primary vegetation, secondary vegetation, tertiary vegetation, landform, surface deposit, primary surficial geomorphology and secondary surficial geomorphology. Secondary and tertiary types are subdominant types that cover more than 30% of a map polygon. The geobotanical map was made using methods and legends specially developed for northern Alaska (Walker et al. 1980, 1986, 1989). The GIS was developed following the integrated terrain-unt mapping approach (Dangermond and Hamden 1990). The resulting geobotanical maps were presented at conferences in 1996 (e.g., Walker et al. 1996) but remained unpublished until now. In 2007 the map boundaries were modified to register with a recent digital elevation model (DEM) of the Kuparuk River region (Nolan 2003) and the 1989 Systeme Probatoire d'Observation de la Terre (SPOT) image of the region (Map B). The legends were also modified to better fit the hierarchy of other maps in the Arctic Geobotanical Atlas.

Map A: Vegetation

The vegetation of the region was studied and mapped as part of the Arctic Long-Term Ecological Research (LTER) project at Toolik Lake and the Department of Energy R4D (Response, Resistance, Resilience and Recovery of vegetation from Disturbance) project at Inuvait Creek (Walker et al. 1994, Walker and Walker 1996). Fifty-seven plant communities and land-cover types were recognized during the mapping of the upper Kuparuk River region and are designated by the numeric GIS codes in the second column of the legend. These were grouped into the 14 physiognomic map units shown on Map A, which are comparable with the Circumpolar Arctic Vegetation Map (CAVM Team 2003) and the Alaska Arctic Tundra Vegetation Map (Raynolds et al. 2005).

Map B: False-Color Infrared Satellite Image

The French SPOT satellite data (20-m resolution) were obtained on 28 July 1989 and provides a view of the mapped region from space. The false-color infrared image shows more densely vegetated areas as brighter red tones. When compared with the glacial geology map (Map C), the older Sagavanirktok-age glacial landscapes have few lakes and redder tones indicating more dense vegetation, and the younger Itkillik-age glacial surfaces that have more lakes and grayer colors. The image data were also used to produce the NDVI/biomass map (Map E).

Map C: Glacial Geology

This map shows a simplified version of Thomas Hamilton's glacial geology map of the upper Kuparuk River region (Hamilton 2003). The glacial history of the region affects a wide variety of landscape and ecosystem properties, including topographic variation, abundance of lakes, plant production, soil carbon, spectral reflectance, biodiversity, trace-gas fluxes and heat flux of these landscapes. Glacial deposits within the upper Kuparuk River region are assigned to Sagavanirktok (middle Pleistocene, about 780-125 kya), Itkillik I (late Pleistocene, about 120-50 kya) and Itkillik II (late Pleistocene, about 25-11.5 kya) glaciations of the central Brooks Range glacial succession (Hamilton 2003). The legend units are arranged approximately from oldest to youngest.

Map D: Surficial Geomorphology

The surfaces of the landscapes in the Toolik Lake region have been modified by a variety of geomorphological processes including alluviation (movement of material by water), colluviation (movement of material by gravity) and periglacial processes (freezing and permafrost-related phenomena). Many of the surface forms have been described for the Inuvait Creek region (Walker and Walker 1996). Common surficial geomorphological features within the mapped area include sorted and nonsorted circles (frost boils), turf hummocks, gelifluction lobes and terraces, water tracks, high- and low-centered ice-wedge polygons, wetland features (strangmoor, aligned hummocks, palsas) and thermokarst features.

Map E: NDVI and Plant Biomass

NDVI is an index of vegetation greenness that can be linked to plant biomass and other biophysical properties of the vegetation, such as CO₂ and photosynthesis. The NDVI = (NIR - R)/(NIR + R), where NIR and R are the spectral reflectance values of the near-infrared (790-890 nm) and red (610-680 nm) bands, respectively. This map is derived from the same SPOT data as Map B. It is modified from an earlier version (Shippert et al. 1995) using more recent biomass information (Walker et al. 2008 in press). Water and barrens are generally displayed as black. Dry tundra and sparsely vegetated areas are displayed in gray. Vegetation density increases with darker shades of green.

Literature Cited

CAVM Team. 2003. Circumpolar Arctic Vegetation Map. Conservation of Arctic Flora and Fauna (CAFF) Map No. 1, U.S. Fish and Wildlife Service, Anchorage, AK.
Dangermond, J. and E. Hamden. 1990. Map data standardization: A methodology for integrating thematic cartographic data before automation. ARC News 12:16-19.
Hamilton, T. D. 2003. Surficial geology of the Dalton Highway (Itkillik-Sagavanirktok rivers) area, southern Arctic foothills, Alaska. Professional Report 121. Alaska Division of Geological and Geophysical Surveys, Fairbanks, AK.
Nolan, M. 2003. Distribution of the Star3i DEM of the Kuparuk River watershed (on CD ROM), Joint Office of Scientific Support Boulder, CO.
Raynolds, M. K., D. A. Walker and H. A. Maier. 2005. Alaska Arctic Vegetation Map, Scale 1:4,000,000. Conservation of Arctic Flora and Fauna (CAFF) Map No. 2, U.S. Fish and Wildlife Service, Anchorage, AK.
Shippert, M. M., D. A. Walker, N. A. Auerbach and B. E. Lewis. 1995. Biomass and leaf-area index maps derived from SPOT images for Toolik Lake and Inuvait Creek areas, Alaska. Polar Record 31:147-154.
Walker, D. A., N. A. Auerbach, L. R. Lentz, S. V. Muller and M. D. Walker. 1996. A hierarchical GIS for studies of process, pattern and scale in arctic ecosystems: The Arctic System Science River Study, Kuparuk River Basin, Alaska. Poster presented at the Second Circumpolar Arctic Vegetation Mapping Workshop, Arendal, Norway, May 20-23, 1996.
Walker, D. A., E. Binman, B. M. Evans, N. D. Lederer, E. Nordstrand and P. J. Webber. 1989. Terrain, vegetation and landscape evolution of the R-4D research site, Brooks Range Foothills, Alaska. Holartic Ecology 12:238-261.
Walker, D. A., et al. 2008. Arctic patterned-ground ecosystems: A synthesis of field studies and models along a North American Arctic Transect. J. Geophys. Res. doi:10.1029/2007JG009504, in press.
Walker, D. A., K. R. Everett, P. J. Webber and J. Brown. 1980. Geobotanical atlas of the Prudhoe Bay Region, Alaska. CRRRL Report 80-14, U.S. Army Cold Regions Research and Engineering Laboratory.
Walker, D. A. and M. D. Walker. 1996. Terrain and vegetation of the Inuvait Creek Watershed. Pages 73-108 in J. E. Reynolds and J. D. Tenhunen, editors. Landscape Function: Implications for Ecosystem Disturbance, a Case Study in Arctic Tundra. Springer-Verlag, New York.
Walker, D. A., P. J. Webber, M. D. Walker, N. D. Lederer, R. H. Meenan and E. A. Nordstrand. 1986. Use of geobotanical maps and automated mapping techniques to examine cumulative impacts in the Prudhoe Bay Oilfield, Alaska. Environmental Conservation 13:149-160.
Walker, M. D., D. A. Walker and N. A. Auerbach. 1994. Plant communities of a tussock tundra landscape in the Brooks Range Foothills, Alaska. Journal of Vegetation Science 5:843-866.





A: Upper Kuparuk River Region Vegetation

Physiognomy	Plant Communities (GIS codes)	Typical Microsites	Area (ha)	% of Map
Barren				
1. Anthropogenic barren	Roads, gravel pads and pipeline pads. (Disturbances are overlaid on pre-disturbance vegetation, so area of anthropogenic barrens are not included in the total area of the map.)	Barren roads, gravel construction pads, airstrips and gravel mines.	(270)	(0.4)
2. Lichens on rocks	Lichen communities on rocks, including <i>Cetraria nigricans-Rhizocarpon geographicum</i> (12).	Bedrock and seric blockfields.	1,009	1.3
3. Partially vegetated barren	Complexes of vegetation with rock or soil on scree slopes (11), river gravels and other barrens (14), partially vegetated alpine areas (13) and areas dominated by nonsorted circles (15). Dominant plant communities include: <i>Saxifraga oppositifolia-Saxifraga eschscholtzii</i> (131); <i>Epilobium latifolium-Castilleja caudata</i> (141); revegetated gravel pads with <i>Festuca rubra</i> (142); <i>Anethula juratzkana-Juncus biglumis</i> (15).	Partially vegetated barrens.	1,805	2.4
Moist graminoid tundra				
4. Tussock-sedge, dwarf-shrub, moss tundra	Moist acidic tussock-tundra complexes dominated by graminoids (31, 311). Dominant plant communities include <i>Eriophorum vaginatum-Sphagnum</i> (31) and <i>Carex bigelowii-Sphagnum</i> (3111).	Mesic to subhygric, stable, acidic (pH < 5.5) sites with shallow to moderate snow. Flat areas and gentle slopes.	29,020	38.7
5. Nontussock-sedge, dwarf-shrub, moss tundra	Moist nonacidic tundra complexes (32). Dominant plant communities include: <i>Carex bigelowii-Dryas integrifolia</i> (321), <i>Carex bigelowii-Dryas integrifolia</i> (subtypes <i>Equisetum arvense</i> (322) and <i>Salix glauca</i> (324)), <i>Eriophorum vaginatum-Tomentypnum nitens</i> (323) and <i>Carex bigelowii-Sphagnum</i> , subtype <i>Cassiope tetragona</i> (3113). Also includes a few other graminoid-dominated communities, including: <i>Festuca altaica-Poa glauca</i> (disturbed thermokarst areas) (325), <i>Deschampsia caespitosa-Carex saxatilis</i> (drained lakes) (326), <i>Carex podocarpa-Salix chamosensis</i> (snowy streambanks) (515).	Mesic to subhygric, nonacidic (pH > 5.5) sites with shallow to moderate snow. Flat areas and gentle slopes. Also miscellaneous graminoid-dominated sites including deep-snow stream margins, landslides and some rocky drained-lake basins.	13,153	17.5
Wet graminoid tundra and water				
6. Sedge, moss tundra (poor fens)	Nutrient-poor fen wetland complexes (41). Dominant plant communities include: <i>Eriophorum angustifolium-Sphagnum orientale</i> (412) and <i>Eriophorum angustifolium-Sphagnum</i> (413). Raised microsites: <i>Sphagnum lenense-Salix fuscescens</i> (411).	Subhygric to hydric, acidic (pH < 4.5). Poor fens, meadows in colluvial basins.	1,934	2.6
7. Sedge, moss tundra (fens)	Nutrient-rich fen wetland complexes (4, 42). Dominant plant communities include: Lower microsites: <i>Carex aquatilis-Carex chodorhiza</i> (422) and <i>Eriophorum angustifolium-Carex aquatilis</i> (425). Raised microsites: <i>Trichophorum caespitosum-Tomentypnum nitens</i> (421) and <i>Carex bigelowii-Tomentypnum nitens</i> .	Subhygric to hydric, nonacidic (pH > 4.5). Water tracks, stream margins, fens, flarks on solifluction slopes.	1,160	1.5
8. Water and herbaceous marsh	Unvegetated water (6) and/or aquatic vegetation in lakes and streams. Dominant plant communities include: <i>Arctophila fulva-Hippuris vulgaris</i> and <i>Spartanium hyperboreum-Hippuris vulgaris</i> (43).	Hydric. Lakes, ponds and streams.	1,593	2.1
Prostrate-shrub tundra				
9. Prostrate dwarf-shrub, forb, fruticose-lichen tundra (acidic)	Dry acidic tundra complexes (21). Dominant plant communities include: <i>Dryas octopetala-Sedagella sibirica</i> (211), <i>Arctous alpina-Salix phlebotypha</i> (212); and lichen tundra <i>Cladonia arbuscula-Stereocaulon tomentosum</i> (215).	Xeric to xeromesic, acidic, wind blown to shallow winter snow cover. Ridge tops, exposed slopes, dry river terraces.	5,818	7.8
10. Prostrate dwarf-shrub, sedge, forb, fruticose-lichen tundra (nonacidic)	Dry nonacidic tundra complexes. <i>Dryas integrifolia-Oxytropis nigrescens</i> (24) and <i>Dryas integrifolia-Asragalus umbellatus</i> (22).	Xeromesic to mesic, nonacidic. Includes a wide variety of drier nonacidic habitats, including stable river terraces and nonsorted stripes on slopes.	895	1.2
11. Hemi-prostrate dwarf-shrub, fruticose-lichen tundra	Deeper snowbed complexes (23). Dominant plant communities include: <i>Cassiope tetragona-Carex microchaeta</i> (231), <i>Cassiope tetragona-Dryas integrifolia</i> (232); and <i>Salix rotundifolia-Saxonia uncinatus</i> (233).	Subseric to mesic, acidic to nonacidic, with deep snow. Snowbeds.	2,164	2.9
12. Hemi-prostrate and prostrate dwarf-shrub, forb, moss, fruticose-lichen tundra	Dry tundra with shallow snowbeds mainly on stripe complexes. Dominant plant communities include: <i>Cassiope tetragona-Calluna vulgaris inopiana</i> (214). Also includes dry areas with hemi-prostrate dwarf birch <i>Betula nana-Hieracium alpina</i> (213).	Subseric to mesic, acidic to nonacidic, somewhat deeper-snow areas. Depressions on acidic ridge crests, dry glacial till and outwash; nonsorted stripes.	2,116	2.8
Erect-shrub tundra				
13. Dwarf- to low-shrub, sedge, moss tundra	Moist acidic tundra complexes dominated by shrubs. Includes shrubby tussock tundra and mainly dwarf-shrub tundra areas (3112). Dominant plant communities include: <i>Betula nana-Eriophorum vaginatum</i> (312) and <i>Salix pulchra-Carex bigelowii</i> (no code). Also acidic shrub tundra dominated by dwarf birch or willows. Dominant plant communities include: <i>Betula nana-Rubus chamaemorus</i> and <i>Salix pulchra-Sphagnum</i> (513).	Mesic to subhygric, moderate snow. Includes a wide variety of habitats with dwarf shrubs, including wet lower slopes, margins of upland water tracks, palsas and high-centered polygons.	9,045	12.1
14. Low to tall shrublands	Shrublands dominated by low and tall shrubs (5) including: 1. Shrublands along streams and water tracks dominated by diamond-leaf willow (<i>Salix pulchra</i>). Dominant plant communities include: <i>Salix pulchra-Eriophorum angustifolium</i> (511) and <i>Salix pulchra-Calluna vulgaris canadensis</i> (514). 2. Shrublands in riparian complexes (51) dominated by feltleaf willow (<i>S. alaxensis</i>) and lamate willow (<i>S. richardsonii</i>) (512); includes tall shrublands (5121) and low shrublands (5122). 3. Upland shrublands dominated by glaucous willow (<i>Salix glauca</i>) or alder (<i>Alnus crispa</i>) (52).	Mesic to subhygric, often with deep snow. Stream margins, upland water tracks and south-facing slopes.	5,344	7.1

Total (excluding roads and pads) 75,056 ha 100%



Property of Ermer E. Rasmuson Library University of Alaska Fairbanks



Source: SPOT Imagery, 28 July 1989
Source: simplified from Hamilton, 2003
Source: see vegetation map above, 1994
Source: SPOT Imagery, 28 July 1989

Explanation for the maps of the Toolik Lake Area and the Toolik Lake Grid




Figure 1. Location of the Toolik Lake Area (large red rectangle) and Toolik Lake Grid (small red rectangle) within the upper Kuparuk River region.

Vegetation of the Toolik Lake Area

Map F is located near the western boundary of Map A (displayed on front) and encloses a 20-km² area surrounding Toolik Lake that stretches from the Dalton Highway on the east to Jade Mountain on the west (large red rectangle in Fig. 1). It includes the Toolik Field Station, the old Toolik Lake pipeline construction camp gravel pad and airstrip on the northeast side of the lake and the primary terrestrial research areas to the south, west and east sides of the lake, as well as several smaller research lakes in the immediate vicinity of Toolik Lake. The area contains surfaces with irregular topography that were glaciated during the Late Pleistocene (Fig. 2 and 3).

Map F portrays the physiognomy of the dominant plant communities in each mapped polygon. Fifty-one landscape types (GIS codes are in parentheses in the second column of the legend) were recognized in the field (minimum mapping unit approximately 250 m²). These were later grouped into the 14 physiognomic vegetation units on the map, which correspond to the same units on the 1:63,360-scale map of the upper Kuparuk River region (Map A).

Vegetation of the Toolik Lake Grid

Map G focuses on the 1.2-km² research grid on the south side of Toolik Lake (red rectangle on Map F and small red rectangle in Fig. 1). This area is one of the principal intensive research areas at the Toolik Lake Field Station. It includes many experimental research sites where long-term observations and experiments are being conducted, including the greenhouse and snow-fence experiments (Fig. 6-9). The grid was constructed in 1989 to provide geographic referencing for experimental plots and to provide a sampling scheme for periodic measurements of snow, active layer and plant communities.

Sixty-five plant communities were recognized (minimum mapping unit approximately 2.5 m²) in the field (GIS codes are in the second column of the legend) and were then grouped into the 24 units appearing on the map. The vegetation units are primarily at the plant-community level (compared to the physiognomic level for the maps of the Upper Kuparuk River Region and the Toolik Lake Area). Several of the dominant plant communities in the Toolik Lake area are shown in the photos (Fig. 10-16). Details of the methods for both maps, sources for aerial photos, orthophoto topographic map, cross-reference to the Braun-Blanquet system, plant community names (Walker et al. 1994) and other information are on the Arctic Geobotanical Atlas website, <http://www.arcticatlas.org/>.

Typical Plant Communities



Figure 10. Sagavarnikok-age glacial surface near Innavik Creek. The vegetation is tussock tundra (*Eriophorum vaginatum-Sphagnum* spp.), the most common plant community on old, stable, acidic landscapes in the region. This is the dominant plant community in unit four on Maps A, F and G.



Figure 11. Blockfield with *Cetraria nigricans-Rhizocarpon geographicum*, unit two on Map A, F and G.



Figure 12. Close-up of *Carex bigelowii-Dryas integrifolia*, typical subtype (401, 403), *Tomentypnum nitens-Carex bigelowii*, *Salix glauca* subtype (320).




Figure 13. Fen with *Carex aquatilis-C. chordeorrhiza*, a major component of unit seven on Map A and F, and unit nine on Map G.



Figure 14. Dry south-facing slope on kame with *Dryas octopetala-Selaginella sibirica*, unit nine on Map A, F, and unit 13 on Map G.



Figure 15. Deep, late-melting snow with *Salix rotundifolia* (at stake). Dark-colored vegetation above the stake is *Cassiope tetragona-Dryas integrifolia*, a common component of unit 11 on Map A and F and unit 17 on Map G.




Figure 16. Well-developed water track with *Salix pulchra-Eriophorum angustifolium*, a common component of map unit 14 on Map A and Map F and unit 23 on Map G.

All photos are by D.A. Walker except figures 6 and 7 which are courtesy of the Arctic LTER website (http://ecosystems.mbl.edu/ARC/terrest/maps_photos/index.html).

F: Toolik Lake Area Vegetation



Physiognomy	Plant Communities (GIS codes)	Typical Microsites	Area (ha)	% of Map
Barren				
1. Barren	Unvegetated (91, 101).	Unvegetated natural and anthropogenic barrens.	23.8	1.2
2. Lichens on rocks	Lichen communities on rocks, including <i>Cetraria nigricans-Rhizocarpon geographicum</i> (92).	Xeric blockfields, glacial erratics.	3.9	0.2
3. Partially vegetated barrens and revegetated disturbed areas	Revegetated gravel pads (e.g., <i>Festuca rubra</i> or <i>Salix alaxensis</i> 102).	Partially vegetated disturbed barrens on gravel pads, abandoned roads, bulldozed areas.	24.9	1.2
Moist graminoid tundra				
4. Tussock sedge, dwarf-shrub, moss tundra	Moist acidic tussock tundra complexes dominated by graminoids. Dominant plant communities include: <i>Eriophorum vaginatum-Sphagnum</i> (41) and <i>Carex bigelowii-Sphagnum</i> (no code).	Mesic to subhygric, acidic, shallow to moderate snow. Stable slopes. Some areas on steeper slopes with solifluction are dominated by Bigelow sedge (<i>Carex bigelowii</i>) (no code).	605.1	29.8
5. Nontussock sedge, dwarf-shrub, moss tundra	Moist nonacidic tundra complexes. Dominant plant communities include: <i>Carex bigelowii-Dryas integrifolia</i> (42) and other subtypes of this unit (e.g., <i>Salix glauca</i> (33), <i>Equisetum arvense</i> and <i>Cassiope tetragona</i> (no codes)). Includes some miscellaneous graminoid communities mostly on disturbed areas, such as <i>Deschampsia caespitosa</i> (45); <i>Rumex arcticus-Carex saxatilis</i> (75) <i>Salix chamissonis-Carex aquatilis</i> (65); <i>Ranunculus pedatifidus-Poa glauca</i> (104).	Mesic to subhygric, circumneutral, shallow to moderate snow. Solifluction areas and somewhat unstable slopes (42), mainly on tillik II glacial surfaces. Some south-facing slopes have scattered glaucous willow (<i>Salix glauca</i>) (33). Also includes some miscellaneous graminoid-dominated sites: deep-snow stream margins (65), landslides, some rocky drained lake basins (45, 75) and animal dens (104).	306.8	15.1
Wet graminoid tundra and water				
6. Sedge, moss tundra (poor fens)	Nutrient-poor fen wetland complexes. Dominant plant communities include: <i>Eriophorum angustifolium-Carex aquatilis</i> (82), <i>Eriophorum angustifolium-Carex aquatilis</i> (82), <i>Carex aquatilis-Scirpium scopioides</i> (74).	Subhygric to hydric, acidic (pH < 4.5). Wet meadows, poor fens in colluvial basins - mainly on older (tillik I) glacial surfaces.	7.8	0.4
7. Sedge, moss tundra (fens)	Nutrient-rich fen wetland complexes. Dominant plant communities include: <i>Eriophorum angustifolium-Carex aquatilis</i> (no code); <i>Eriophorum angustifolium-Carex aquatilis</i> (82), <i>Carex aquatilis-Scirpium scopioides</i> (74).	Subhygric to hydric, minerotrophic (pH > 4.5). Water tracks, stream margins, fens, flarks on solifluction slopes - mainly on younger (tillik II) glacial surfaces.	105.6	5.2
8. Water and herbaceous marsh	Unvegetated water (84); graminoid marsh <i>Arctophila fulva</i> (81) and <i>Spartanium hyperboreum-Hippuris vulgaris</i> (83).	Lakes, ponds and streams; aquatic vegetation in some protected sites.	196.5	9.7

Typical landscapes in the Upper Kuparuk River region:



Figure 2. View looking northeast from Jade Mountain across an Itkillik II glacial landscape with numerous glacial lakes, kames and kettles. This landscape is much more vegetatively complex than the Sagavarnikok-age glacial surfaces (Fig. 3).



Figure 3. View looking southeast across the headwaters of Innavik Creek into the Philip Smith Mountains of the Brooks Range. Vegetation is typical of the Sagavarnikok-age glacial surfaces, which cover large portions of Map A but do not occur on the terrain shown in Maps F and G.




Figure 4. Alpine area on limestone on Peak 1376 in the southeast corner of Map A, looking south into the valley of the Sagavarnikok River. The dominant vegetation is *Dryas integrifolia-Oxytropis nigrescens* (unit 10 on Map A).



Figure 5. Streamside vegetation along the inlet stream to Toolik Lake. The tallest shrubs are *Salix alaxensis*. Low shrubs along the far bank are a mix of *Betula nana* and *Salix pulchra*. The dominant vegetation unit along the stream is low to tall shrublands, unit 14 on Maps A and F, and unit 24 on Map G.

Total 2027.6 ha 100%

G: Toolik Lake Grid Vegetation



Plant Communities (GIS codes)	Description (physiognomy and typical microsite)	Area (ha)	% of Map
Barren			
1. Barren (901)	Unvegetated natural and anthropogenic barrens.	0.16	0.1
2. <i>Cetraria nigricans-Rhizocarpon geographicum</i> (902)	Lichen communities on rocks. Xeric blockfields, glacial erratics.	0.11	0.1
3. <i>Festuca rubra</i> (903); <i>Salix alaxensis</i> (904); <i>Eriophorum latifolium</i> (905); <i>Juncus biglomis-Luzula arctica</i> (no code)	Partially revegetated areas. Gravel pads (903), river gravels (904, 905) and nonsorted cirques (no code).	0.23	0.2
Moist graminoid tundra			
4. <i>Eriophorum vaginatum-Sphagnum</i> (406,407); <i>Carex bigelowii-Sphagnum</i> (404, 405)	Tussock sedge, dwarf-shrub, moss tundra (tussock tundra, moist acidic tundra). Mesic to subhygric, acidic, shallow to moderate snow, stable. This unit is the zonal vegetation on fine-grained substrates with ice-rich permafrost (406, 407). Some areas on steeper slopes with solifluction are dominated by Bigelow sedge (<i>Carex bigelowii</i>) (404, 405).	65.92	54.2
5. <i>Carex bigelowii-Dryas integrifolia</i> , typical subtype (401, 403); <i>Tomentypnum nitens-Carex bigelowii</i> , <i>Salix glauca</i> subtype (320)	Nontussock sedge, dwarf-shrub, moss tundra (moist nonacidic tundra). Mesic to subhygric, nonacidic (pH > 5.5), shallow to moderate snow. Solifluction areas and somewhat unstable slopes (401,403). Some south-facing slopes have scattered glaucous willow (<i>Salix glauca</i>) (320).	7.04	5.8
6. <i>Carex bigelowii-Dryas integrifolia</i> , <i>Equisetum arvense</i> subtype (402); <i>Tomentypnum nitens-Carex bigelowii</i> , <i>Carex aquatilis</i> subtype (410)	Nontussock sedge, prostrate dwarf-shrub, horsetail, moss tundra (wetter subtypes of moist nonacidic tundra, often with abundant horsetails). Mesic to subhygric, nonacidic, moderate snow. Seepage areas below snowbeds with abundant horsetails (<i>Equisetum arvense</i>) (402) or aquatic sedge (<i>C. aquatilis</i>) in wetter areas (410).	2.22	1.8
7. <i>Carex bigelowii-Dryas integrifolia</i> , <i>Cassiope tetragona</i> subtype (208); or other miscellaneous graminoid plant communities, including <i>Ranunculus pedatifidus-Poa glauca</i> (106); <i>Salix chamissonis-Carex podocarpa</i> (408); <i>Festuca aluticola-Artemisia arctica</i> (412); <i>Poa glauca-Eriophorum latifolium</i> (109)	Sedge, hemi-prostrate dwarf shrub, moss tundra (moist nonacidic tundra in deep snow accumulation areas). Mesic to subhygric, mostly nonacidic, moderate to deep snow. Inter-stripe areas in nonsorted stripe complexes on upper hill-slopes with moderate to deep snow and abundant Lapland heather (<i>Cassiope tetragona</i>). This unit also includes several miscellaneous graminoid, dwarf-shrub, forb communities that cover small areas, including animal dens (106), deep-snow stream and lake margins (106), dry snow accumulation areas (412) and stream banks (109).	1.64	1.3
Wet graminoid tundra and water			
8. <i>Tricophorum caespitosum-Tomentypnum nitens</i> (409), <i>Salix fuscescens-Sphagnum lenense</i> (411,508), <i>Carex aquatilis-Sphagnum warnstorffii</i> (506)	Sedge, prostrate dwarf-shrub, moss tundra. Hygric to subhygric. Hummocks, strags and raised microsites in fens (409) and poor fens in wet meadows and colluvial-basins (411, 508) and mossy colluvial basin margins (506).	0.32	0.3
9. <i>Eriophorum angustifolium-Carex aquatilis</i> , typical subtype (501), <i>Carex chordeorrhiza</i> subtype (502, 505), <i>Drepanolabus revolvens</i> subtype (503), <i>Carex saxatilis-Carex aquatilis</i> (504) and <i>Calligonum giganteum-Drepanolabus revolvens</i> (507)	Sedge, moss tundra in fens with flowing water. Subhygric to hydric. Lower microsites in colluvial basins, water tracks and stream margins (501, 502, 503, 505), wet pools on solifluction slopes (507).	2.30	1.9
10. Unvegetated water (602)	Hydric. Streams, lakes, ponds.	5.34	4.4
11. <i>Spartanium hyperboreum-Hippuris vulgaris</i> (603, 604); <i>Arctophila fulva</i> (601)	Herbaceous marsh. Hydric. Water to 1-m deep in lakes and ponds.	0.33	0.3

Total 121.66 ha 100%

Research within the Toolik Lake Grid:



Figure 6. Long-term experiments within the Toolik Lake grid, aerial view showing boardwalks, greenhouses (white structures) and shadehouses (black structures).



Figure 7. Experimental greenhouse with the side and top opened to show enhanced growth due to added warmth. Pre-treatment shrubs were same height as vegetation in the foreground.



Figure 8. Snowfence experiment within the Toolik Lake Grid, summer view showing fence, snow-depth monitoring stakes, and small open-top greenhouses.



Figure 9. Winter view of snowfence experimental area showing the drift that forms behind the fence. Tall stakes are the same as striped stakes in Figure 8.

Acknowledgements

Numerous people have contributed in major ways to the field work, map production and analysis of these maps, including Nancy Auerbach, Andrew Balsler, Edie Barbour, Tom Hamilton, Julie Knudsen, Nan Lederer, Leanne Lestak, Martha Raynolds and Marilyn Walker. Funding for the mapping came from the Arctic LTER project, the DOE R4D program and NSF grants ARC-0225517, ARC-0455541 and ARC-0531180.

