

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

## Sheep Liver Cytosolic Aldehyde

Dehydrogenase;

A fresh perspective

A thesis presented in partial fulfilment of the requirements for the degree of

Master of Science in Chemistry

at

Massey University New Zealand

Richard Malcolm Sime 1995

#### Abstract

The pre-steady-state mechanism of aldehyde dehydrogenase has been further investigated using synthesised deuterated 4-*trans*-(N,N-dimethylamino) cinnamaldehyde as a substrate.

Reporter groups of the active site of ALDH have indicated the presence of a divalent or trivalent metal electrophile, shown in chapter 3 as being either Fe<sup>(II)</sup> or Fe<sup>(III)</sup>.

Studies of the spectral properties of NADH bound to aldehyde dehydrogenase have revealed the presence of at least two spectrally different enzyme-NADH species. The consequences of this information are important in interpretation of the kinetic data and understanding apparently contradictory experimental results from different research workers.

The steady-state kinetics of ALDH have been further investigated.

A sensitive substrate for use in enzyme immunoassays has been designed and synthesised. The preliminary kinetic behaviour observed using this substrate has been studied with three enzymes.

Aldehyde dehydrogenase has been used as a model system for studying the effects of electromagnetic radiation on biological systems.

### Acknowledgements

I wish to thank my supervisors, Len Blackwell for his enthusiasm for science, from who I developed a lively interest in enzyme immunoassays, and Paul Buckley for his teaching and advice during this study.

I would also like to thank Jeremy Hill and Geoffery Barnes who also worked with me in an almost supervisory role, and were a source of much inspiration. Thanks are extended to all members of Chemistry-Biochemistry for helpful comments and also Bernardette Kilsby of the Department of Plant Biology and Biotechnology for useful references.

Finally I wish to thank my wife Lis, for her patience and encouragement.

### Table of contents

Introduction ......1

#### Chapter 1

#### Kinetic Isotope Effect on Acyl-Enzyme Formation During Oxidation of 4-*trans*-(N,N-dimethylamino) Cinnamaldehyde by Aldehyde Dehydrogenase

1.1 Introduction	3
1.1.1 4-trans-(N,N-dimethylamino) cinnamaldehyde as a	
substrate for aldehyde dehydrogenase	3
1.1.2 Investigation of the pre-steady state reaction of	
ALDH with DACA and its deuterated analogue as	
substrates	8
1.1.3 Kinetic isotope effects	9
1.2. Materials	10
1.3. Methods	11
1.3.1 Preparation of sheep liver cytosolic aldehyde	
dehydrogenase.	11
1.3.2 Enzyme assays	13
1.3.3 Protein determinations	13
1.3.4 Spectroscopic determinations	14
1.3.5 Computer simulations	14
1.3.6 Pre-steady state studies of the formation of the 464	
nm ALDH 4-trans-(N,N-dimethylamino) cinnamoyl -	
cysteine moiety from deuterated DACA.	14
1.3.7 Synthetic schemes for the preparation of deuterated	

DACA	14
1.4 Results and Discussion	18
1.4 .1 Attempts to synthesise deuterated DACA	18
1.4.2 Synthesis of 1-[2H]-4-trans-N,N-dimethyl	
cinnamaldehyde by acid catalysed condensation of N,N-	
dimethyl benzaldehyde with ethanal	18
1.4.3 Isotope effects on the pre-steady-state burst during	
the formation of the DACA acyl-enzyme intermediate	26
1.4.4 Kinetic modeling of the isotope effects on the pre-	
steady-state burst during the formation of the DACA	
acyl-enzyme intermediate	35
1.4.5 Ethanamoyl 4-trans-(N,N-dimethylamino)	
cinnamoate	44
1.4.6 Conclusion	48

# Formation of 4-*trans*-(N,N-dimethylamino cinnamaldehyde complexes as models for the acyl intermediate

2.1 Introduction	.9
2.2 Materials and Methods	51
2.2.1 Formation of chromophore complexes	1
2.2.2 Determination of difference spectra	3
2.2.3 Formation of the ALDH furylacryloyl cysteine	
moiety using thioNAD5	6
2.3 Results	6
2.3.1 DACA complexes formed with non-metal	
electrophiles5	6
2.3.2 DACA complexes formed with metal electrophiles 6	0
2.3.3 Furylacrolein complexes formed with metal and	
non-metal electrophiles6	4
2.3.4 Absorption spectra of the ALDH furylacryloyl	
cysteine moiety6	5
2.4 Discussion	7
2.4.1 Introduction to the discussion	7

2.4.2 Basic concepts in colour chemistry
2.4.3 Calculation of the approximate maximum possible
red-shift for a DACA moiety
2.4.4 Consequences for DACA aldehyde dehydrogenase
interactions74
2.4.5 Modelling experiments 77
2.4.6 Biological examples of red-shifted spectra induced
by enzymes not containing metal electrophiles
2.4.7 Model experiments for the non-metal electrophiles 84
2.4.8 Possible improvements for modelling non-metal
electrophiles90
2.4.9 Metal electrophiles
2.4.10 Comparison of the ALDH 4-trans-(N,N-
dimethylamino) cinnamoyl - cysteine moiety and Alcohol
dehydrogenase - DACA intermediates
2.5 Conclusion

#### Effects of metal ions on ALDH

3.1 Introduction	. 98
3.2 Materials	. 99
3.3 Methods	. 99
3.2.1 Determination of the number of magnesium binding	
sites using dialysis	. 99
3.2.2 Effect of calcium, magnesium and zinc ions on the	
ALDH activity at pH 5.22	. 100
3.2.3 Determination of the Metal ion content of highly	
purified ALDH	. 102
3.2.4 Examination of the iron profile of ALDH from the	
pH gradient column	. 103
3.2.5 Time dependent effects of EDTA	. 103
3.2.6 The pH dependence of EDTA inhibition	. 103
3.3 Results	. 104
3.3.1 Number of metal binding sites	. 104

3.3.2 Effect of Calcium Magnesium and Zinc on ALDH	
activity at pH 5.2210	)5
3.3.3 Iron, copper and zinc determination of highly	
purified ALDH using atomic absorption11	10
3.3.4 Examination of the iron profile of ALDH from the	
pH gradient column11	10
3.3.5 Time dependent effects of EDTA 11	12
3.3.6 pH dependence of the ALDH inactivation by	
EDTA11	13
3.4 Discussion	14
3.4.1 Effects of metal ions on ALDH11	4
3.4.2 Effects of EDTA on the ALDH activity 11	16
3.4.3 What is the active site metal electrophile?	17
3.5 Conclusion	17

# The Interactions of NADH on the ALDH Pathway

4.1 Introduction	. 119
4.2 Materials	. 120
4.3 Methods	. 121
4.3.1 Spectral properties of NADH bound to ALDH	. 121
4.3.2 Effect of pyrophosphate on ALDH activity	. 122
4.3.3 Activation and inhibition of ALDH by NAD	. 122
4.4 Results and Discussion	. 123
4.4.1 Spectral properties of NADH bound to ALDH	. 123
4.4.2 Probable cause of the ENADH red-shifted spectra	. 127
4.4.3 Calculation of the equilibrium constant for the two	
ENADH species	. 128
4.4.4 Effects of propanal on the ENADH species	. 132
4.4.5 The 364 nm species in the kinetic pathway	. 135
4.4.6 Effect of the acyl species on the absorption and	
fluorescence lag phases.	. 144
4.4.7 Relationship between decreased amplitudes and the	

observed rate	146
4.4.8 Explanation for the difference results observed by	
the two research groups	150
4.4.9 Effects of pyrophosphate on activity of ALDH	151
4.4.10. Steady-state kinetics	. 154
4.5 Conclusion	. 155

# Synthesis of a sensitive substrate for use in homogeneous enzyme immunoassays

5.1. Introduction
5.1.2 Desirable features when designing a substrate for a
homogeneous EIA 158
5.2 Materials
5.3 Methods
5.3.1 Synthesis of 4-((bis 4-N,N-dimethylaminophenyl)
methylinium ethanoate) phenol 160
5.3.2 Synthesis of 4-((bis 4-N,N-dimethylaminophenyl)
methylinium ethanoate) phenyl ethanoate
5.2.3 Measurement of the rate of hydrolysis of DMPE
with rabbit liver esterase and porcine liver esterase
5.3.4 Measurement of the rate of hydrolysis of DMPE
with ALDH
5.3.5 Inhibition of the DMPE esterase reaction by ALDH
4- trans (N,N-dimethylamino) cinnamoyl - cysteine
moiety
5.4 Results and Discussion
5.4.1 Synthesis of dDMP 165
5.4.2 Synthesis of DMPE
5.4.3 The comparison of the spectral properties of DMPE
with other structurally related compounds
5.4.4 Measurement of the rate of hydrolysis of DMPE
with rabbit liver esterase and porcine liver esterase
5.4.5 Measurement of the rate of hydrolysis of DMPE

with ALDH	175
5.4.6 Inhibition of the DMPE esterase reaction by ALD	Н
4-trans (N,N-dimethylamino) cinnamoyl - cysteine	
moiety.	176
5.4.7 Long term stability of DMPE	177
5.4.8 Safety of DMPE	179
5.4.9 Potential of DMPE in homogenous enzyme	
immunoassays	180
5.5 Conclusion	181

#### ALDH as biological model for the effects of electromagnetic fields

6.1 Introduction	182
6.1 Methods	184
6.3 Results and Discussion	185
6.4 Conclusion	190

## List of Tables

Table 1.4.4.1 Kinetic parameters for simulation of scheme	
1.4.4.4	39
Table 1.4.4.2 Summary of kinetic parameters obtained by	
computer fits to scheme 1.4.4.4.	40
Table 1.4.4.3 Kinetic parameters for simulation of scheme	
1.4.4.5 (B)	42
Table 1.4.4.4 Summary of kinetic parameters obtained by	
computer fits to scheme 1.4.4.5 (B).	43
Table 2.2.1.1 DACA complexing solutions	52
Table 2.2.1.2 Furylacrolein complexing solutions	
Table 2.3.1.1 Summary of the red-shifts induced in DACA by	
non-metal complexing agents	57
Table 2.3.2.1 Summary of the red-shifts induced by metal	
complexing agents in DACA	64
Table 2.3.3.1 Summary of the red-shifts induced by metal and	
non-metal complexing agents in furylacrolein	65
Table 2.4.3.1	73
Table 2.4.4.1 Stabilities of the ALDH 4-trans-(N,N-	
dimethylamino) cinnamoyl - cysteine moieties	76
Table 2.4.5.1 Comparative hypsochromicity of hydrogen, sulphur	
and oxygen.	79
In Figure 2.4.6.2 the complexing conditions were	82
Table 2.4.6.1 Calculated red-shifts induced by non-metallo	
enzymes in a 4-trans-(N,N-dimethylamino) cinnamoyl moiety	83
Table 2.4.7.1 Comparison of modelling techniques	87
Table 2.4.10.1 Red-shifts induced in DACA by human and horse	
liver alcohol dehydrogenase	95

Table 2.4.10.1 Comparison of aldehyde dehydrogenase and alcohol dehydrogenase.	96
Table 4.4.3.1 Calculation of the approximate equilibrium constant of the ENADH species	130
Table 4.4.3.2 Calculation of the approximate equilibrium constant of the ENADH species in different buffers Table 4.4.4.1 Correlation between ALDH-NADH-acyl and	130
ALDH-NADH-aldehyde Table 4.4.5.1 Parameters used for the NADH burst simulations	
Table 5.4.3.1 Spectral properties of tri-aryl carbonium dyes   Table 5.4.4.1 Measurement of kcat for porcine   esterase and rabbit esterase	

Table 6.3.1.1 Effects of parallel combinations of a.c. and d.c. magnetic fields on the activity of ALDH.....

#### Introduction

Sheep liver cytosolic aldehyde dehydrogenase is an enzyme which is involved in the metabolism of aldehdyes. It has been suggested that aldehyde dehydrogenase has a multifunction purpose, its broad substrate specificity allowing it to oxidise a wide variety of aldehydes to the corresponding carboxylic acids. Hence the physiological role of aldehyde dehydrogenase appears to be that of a general detoxifier (Weiner, 1989).

Aldehyde dehydrogenase was first purified by Kathy Kitson (nee Crow, thesis), and she has continued her involvement in the improvement enzyme quality thoughout. Kathy Kitson has been predominantly researching the physiological factors that control the rate alcohol metabolism, of which aldehyde dehydrogenase is one (Crow and Hardman, 1989).

The steady state and pre-steady state kinetics of sheep liver aldehyde dehydrogenase was first studied by MacGibbon (thesis). The significant outcomes of this work was that the product NADH was shown to be released from ALDH in a biphasic manner. The biphasic release of NADH was found to be the rate limiting process for many aldehyde substrates. MacGibbon also showed that ALDH could catalyse the hydrolysis of unstable esters. Later a group at Hull University, under the leadership of Mark Dickinson also started research on this enzyme. It was found that NADH could reduce the ALDH-acyl species (the reverse reaction) formed by reaction of the enzyme active site nucleophile with an anhydride. This group contributed much pre-steady state and steady information (Hart and Dickinson, 1982)

The rate of acyl enzyme hydrolysis (for many substrates) using proton release was predominantly studied by Bennett (thesis). Bennett also studied the effect of magnesium ions on the enzyme, magnesium ions being available in abundance in vivo (Soman et al, 1970). Motion (thesis) studied the kinetic effects of pH, buffers and furthered the studies on the reverse reaction. Dunn and Buckley (1982,1985), studied the formation of a coloured quassi-stable ALDH-acyl species formed as an intermediate in the oxidation of 4-trans-(N,N-dimethylamino) cinnamaldehyde. Jeremy Hill examined the enzyme structural equilibrium between the tetrameric state and the dissociated state (Buckley et al., 1991).

Trevor Kitson made the first stable ALDH-acyl species (Kitson T., 1989) which was found to have been incorporated with the residue cysteine 302 (Kitson et al.,1991). In conjunction with Kerry Loomes, Kitson studied the effects of a variety of sulfhydral modifiers on the kinetics of aldehyde dehydrogenase (Kitson T., (1989,b)).

A major focus of the research work presented in this thesis is to further increase the kinetic understanding of aldehyde dehydrogenase.