

Journal of Melittology

Bee Biology, Ecology, Evolution, & Systematics

The latest buzz in bee biology

No. 66, pp. 1–20

3 April 2017

The native bee fauna of the Palouse Prairie (Hymenoptera: Apoidea)

Paul Rhoades¹, Terry Griswold², Harold Ikerd², Lisette Waits³,
Nilsa A. Bosque-Pérez¹, & Sanford D. Eigenbrode¹

Abstract. While synoptic collections provide data on the range and general composition of the North American bee fauna, bee communities associated with specific habitats are largely uncharacterized. This report describes the community of native bees currently found in remnant fragments of the Palouse Prairie of northern Idaho and southeastern Washington State. Native bees were collected using standardized collection techniques including blue vane traps, colored pan traps and aerial netting. More than 13,000 individuals were collected, representing at least 174 species and 36 morphospecies in 29 genera. These data provide the most thorough characterization of the bee fauna of this vulnerable ecosystem, as well as community level information on bee species of unknown conservation status. These results are relevant to regional conservation efforts and, more broadly, are representative of conditions in fragmented grasslands surrounded by intense agriculture, a common global land use pattern of conservation concern.

INTRODUCTION

By 2005, cultivated systems covered one quarter of Earth's terrestrial surface (Sarikhan *et al.*, 2005). This habitat loss is responsible for worldwide reductions in species richness and diversity of many taxa including bees (Foley *et al.*, 2005; Brown & Paxton, 2009; Senapathi *et al.*, 2015). Temperate grasslands, such as the Palouse Prairie, are greatly impacted by anthropogenic land use change, with more than half of all temperate grassland, shrubland or savannah converted to agricultural or urban use

¹ Department of Plant, Soil and Entomological Sciences, University of Idaho, 875 Perimeter Drive, MS 2339, Moscow, Idaho 83844 (paul.r.rhoades@gmail.com; nbosque@uidaho.edu; sanforde@uidaho.edu).

² USDA-ARS Pollinating Insects Research Unit, BNR Room 261, Dept. Biology UMC 5310 Utah State University Logan, Utah 84322-5310 (terry.griswold@ars.usda.gov; hikerd@gmail.com).

³ Department of Fish and Wildlife Sciences, University of Idaho, 875 Perimeter Drive, Moscow, Idaho 83844 (lwaits@uidaho.edu).

doi: <http://dx.doi.org/10.17161/jom.v0i66.5703>

(White *et al.*, 2000). Habitat loss is associated with pollinator declines (Vanbergen, 2013) and can exacerbate reductions in bee species richness and abundance caused by pesticides (Park *et al.*, 2015). Additionally, fragmentation caused by habitat loss can impact remaining isolated populations through inbreeding (Zayed, 2004; Zayed & Packer, 2005; Darvill *et al.*, 2006; Ellis *et al.*, 2006); inability of small habitat fragments to support populations (Lennartsson, 2002); and through degradation of the remaining habitat, where depauperate bee communities inadequately pollinate necessary forage plants (Fontaine *et al.*, 2005).

Habitat loss and fragmentation functions in conjunction with disease, invasive plant spread, and pesticide use to cause bee declines (Brown *et al.*, 2002; Vanbergen, 2013; Goulson *et al.*, 2015, but see Winfree *et al.*, 2007). However, the extent and magnitude of native bee decline in North America remains unclear (NRC, 2007) despite the important role native bee pollinators play in agricultural production and ecosystem health (Ashman *et al.*, 2004; Klein *et al.*, 2007). The degree of bee species decline can be difficult to resolve because baseline data necessary to identify species of concern is lacking in many cases and the conservation status of most native bee species remains unknown (Meffe *et al.*, 1998; NRC, 2007; Goulson *et al.*, 2008). Yet, reductions in bee species' range and abundance have been documented throughout the world (Biesmeijer *et al.*, 2006; Fitzpatrick *et al.*, 2007; Potts *et al.*, 2010; Cameron *et al.*, 2011). While there have been some efforts to quantify bee species decline using museum specimens (Bartomeus *et al.*, 2013; Scheper *et al.*, 2014), systematic surveys of bee fauna presence and abundance are lacking in most parts of the world, including in the Palouse Prairie in northern Idaho and adjacent eastern Washington.

The Palouse Prairie is a discrete component of the Pacific Northwest bunchgrass biome, differentiated by its distinctive soils and topography (Tisdale, 1982). It is considered a subsection within Bailey's ecoregions (Bailey, 1995), a subregion in Omernik's ecoregions (1987), and a unit in Ertter and Moseley's floristic regions of Idaho (1992). It is bounded by the arid channeled scablands of central Washington to the west, the canyon grasslands adjacent to the Snake and Clearwater Rivers to south and southeast, and the forests of the Selkirk and Bitterroot Mountains to the north and east. The Palouse Prairie was continuous habitat across this region until the late 1800s when agricultural conversion began. Now approximately 1% of the Palouse Prairie remains (Black *et al.*, 1998), and so the ecosystem could be considered 'Critically Endangered' using the criteria of Keith *et al.* (2013). The remaining fragments are small (most less than 2 ha) with high perimeter-to-area ratios, located disproportionately along streams or on land too rocky or steep to farm (Looney & Eigenbrode, 2012). Although fragmented and surrounded by intensive agriculture, the Palouse Prairie still supports a species rich community of vascular plants, including rare and threatened plant species like *Silene spaldingii* S. Watson, *Symphytotrichum jessicae* (Piper) G.L. Nesom, *Astragalus arrectus* A. Gray, and *Calochortus nitidus* Douglas (Daubenmire, 1942; Lichthardt & Moseley, 1997; Hanson *et al.*, 2008; Davis, 2015). The native earthworm *Driloleirus americanus* Smith still persists in the Palouse (Sánchez-de León & Johnson-Maynard, 2008). Weevils (20 species), darkling beetles (five species), and scarab beetles (six species) present in Palouse Prairie fragments and adjacent agricultural fields have been characterized (Hatten *et al.*, 2004, 2007), and all eight regional species of carrion beetles are found in Palouse Prairie fragments (Looney *et al.*, 2004). However, native bee communities on the Palouse remain uncharacterized, a situation common throughout North America (NRC, 2007).

A compilation of historical records lists 257 bee species present in the Palouse

ecoregion (Bailey, 1995), among the highest of all ecoregions in the Columbia Basin (Tepedino & Griswold, 1995). However, these data were compiled from many different sources using a variety of collection methods, so the relative abundance of species in this assemblage is unknown. The objectives of this study were to: 1) provide a comprehensive species list of bee fauna of the Palouse Prairie, 2) assess the relative abundance of bee species, and 3) identify range expansions or new state records for bee species.

MATERIAL AND METHODS

Bee collection occurred at 32 sites on 29 fragments of the Palouse Prairie (Fig. 1) between May and July in 2012 and 2013. We chose to constrain sampling to these months to coincide with the period of highest species richness of plants in flower and the greatest abundance of active bees. The great majority (73%) of bee specimens recorded in the Global Biodiversity Information Facility (hereafter, GBIF) from the Palouse region were collected between May and July (GBIF, 2015). Only seven species recorded in GBIF have records of occurrence in the Palouse that do not overlap these three months. Of these seven species, five are represented by a single specimen in the GBIF database (GBIF, 2015). So, while this sampling scheme may miss some of the early spring and late summer species this was the most efficient use of collector time. Each site was sampled four times in each year, at sampling intervals of approximately three weeks. Sampling location within the fragment was determined by generating a random point within each prairie fragment at least 10 meters from the fragment edge, when possible, using the Create Random Points tool in ArcMap 10.0 (ESRI, Redlands, CA). If the sampling location fell within a thicket of shrubs or small trees, which would inhibit trap placement, the sampling location was moved 5 meters beyond the nearest edge of the thicket. Multiple methods of bee sampling were employed to maximize detection of the existing fauna: pan traps, blue vane traps, and aerial netting. Pan traps have been extensively used in standardized bee sampling regimes, but are known to have bias in bee capture, recovering Halictinae and *Perdita* Smith at greater rates than the genera *Anthidium* Fabricius, *Colletes* Latreille, and *Epeolus* Latreille, as compared with netting in the same locations (Wilson *et al.*, 2008). Blue vane traps were used so we could better compare results with the only other systematic bee collection effort in Pacific Northwest bunchgrass prairie, which was performed using only blue vane traps (Kimoto *et al.*, 2012). Blue vane traps filled with soapy water (Springstar Inc., Woodinville, WA) (Stephen & Rao, 2007) were hung about one meter off the ground on a bamboo tripod at the randomly determined sampling location. Three colored pan traps (3.25 oz. soufflé cups, Solo model #p325w-0007) filled with soapy water, one each of fluorescent yellow, fluorescent blue, and white, were set three meters apart in a transect leading away from the blue vane trap on a random heading. Pan trap colors were randomized within each transect. Traps were left open for 24 hours. Finally, an aerial net was used to collect bees from flowers within 50 meters of the random point for 5 minutes at the time of trap placement and again at removal for a total of 80 minutes of net collection at each site over the 2 years of sampling.

Sampling was only initiated on mostly sunny days with highs above 16°C, but quickly changing weather during the spring and early summer in this region meant some light rain fell during the 24 hours traps were left open. The average high temperature for sampling days was 22.7°C in 2012 and 27°C in 2013; the average low temperature was 5.4°C in 2012 and 6.2°C in 2013; 1.16 cm of precipitation fell over four




Figure 1. Map of sampling locations and Palouse Prairie fragments.

sampling days in 2012 and 0.15 cm fell over two days in 2013; the largest daily rainfall total on a sampling day was 0.71 cm in 2012 and 0.1 cm in 2013.

Netted bees were kept frozen before processing. Bees collected in blue vane traps or pan traps were rinsed in ethanol and then placed in a Whirl-Pak bag (Nasco, Fort Atkinson, WI) and covered with ethanol for temporary storage. Bees stored in ethanol were then washed and dried before further processing (methods adapted from Droege, 2009). All bees were pinned in the first year of collection. In the second year, very common and easily identifiable species including *Agapostemon angelicus* Cockerell, *A. virescens* (Fabricius), *A. femoratus* Crawford, and *Halictus tripartitus* Cockerell were identified without pinning to save time and resources. Additionally, because

species-level identification of *Lasioglossum* Curtis belonging to the *Hemihalictus* series (Michener, 2007) was not possible, they were counted then stored without pinning. Individuals in other genera not identified to species were damaged or, rarely, males. Voucher specimens reside in the William F. Barr Insect Museum at the University of Idaho and the U.S. National Pollinating Insects Collection, USDA Bee Biology and Systematics Laboratory, housed on the campus of Utah State University.

To determine historical records of bee occurrence in the Palouse region, data from GBIF were downloaded and used in conjunction with raw data used in a report on the bees of the Upper Columbia River basin (Tepedino & Griswold, 1995), obtained from the authors. Only records falling within the Palouse ecoregion (Omernik, 1987) were used.

The taxonomy of *Bombus* Latreille is relatively stable presently and historically, so good information on bumble bee community composition is readily available where similar information for other genera is not. There are four instances where *Bombus* community data can be compared to the Palouse: 1) A 2003 survey of *Bombus* was performed in the Palouse Prairie, reflecting recent community composition (Hatten *et al.*, 2013); and 2) a GBIF-derived dataset with 1675 records of *Bombus* occurrence when limited to pre-2000 records (1805–1999) within the Palouse, reflecting historical community composition [the preponderance of post-1999 records in the GBIF database were from the Hatten *et al.* (2013) study]; 3) a recent survey of native bees on the Zumwalt prairie (Kimoto *et al.*, 2012); and 4) *Bombus* community data extracted from a bee study of the nearby Okanogan National Forest (Wilson *et al.*, 2010). Bray-Curtis dissimilarity was calculated using the vegan package in R (Oksanen *et al.*, 2015; R Core Development Team, 2015), among these four datasets to evaluate: 1) the similarity of the Palouse *Bombus* community through time, and 2) the similarity of the contemporary Palouse *Bombus* community to nearby habitats. All community data were normalized to account for differing sampling regimes.

Incidence-based rarefaction without replacement was performed using EstimateS (Colwell *et al.*, 2012) to evaluate the number of non-*Hemihalictus* bee species that remain undetected in the study area. *Hemihalictus* Cockerell were excluded because of the large number of unidentified individuals in this group. Estimated species richness was extrapolated to twice the total number of collected non-*Hemihalictus* individuals.

RESULTS

Over two years of sampling, 13,293 bees were collected, comprising 174 species and 36 morphospecies in five families and 29 genera (Appendix). Rarefaction analysis indicates the total number of trappable, non-*Hemihalictus* bees was 253 \pm 22 species (Fig. 2).

The Halictidae were the most abundant family, comprising more than 64% of all collected bees (Appendix), followed by the Apidae (16%) and Megachilidae (11%). The most abundant species also belonged to Halictidae: *H. tripartitus* made up 10.2% of total collected bees, *A. virescens* (6.5%), *A. angelicus* (4.6%), and *L. sisymbrii* (Cockerell) (4.0%). The most abundant genus was the halictid *Lasioglossum*, comprising 37% of all collected bees, with the *Hemihalictus* series making up nearly 75% of collected *Lasioglossum*. *Halictus* Latreille (15%), *Agapostemon* Guérin-Ménéville (11%), the megachilid genus *Osmia* Panzer (8%), and the andrenid genus *Andrena* Fabricius (7%) were also abundant (Appendix).

The most speciose families were Megachilidae (64 species, 2 morphospecies) and




Figure 2. Extrapolated rarefaction curve with 95% confidence intervals based on all collected non-*Hemihalictus* bees. Vertical line indicates the actual number of collected non-*Hemihalictus* bees.

Apidae (52 species, 21 morphospecies), followed by Andrenidae (28 species) and Halictidae (23 species, 13 morphospecies) (Appendix). It is important to note that the *Hemihalictus* series, with 28% of collected individuals, were not identified to species. A similar study in a nearby Pacific Northwest bunchgrass system (Kimoto *et al.*, 2012), detected as many as 38 morphospecies within the *Hemihalictus* series. If species richness within the *Hemihalictus* series is comparable in the Palouse Prairie, it would nearly make the Halictidae the most speciose family. Colletidae were poorly represented (8 species) and Melittidae were absent. The most speciose genera include *Osmia* (Megachilidae, 33 species), *Andrena* (Andrenidae, 26 species), *Nomada* Scopoli (Apidae, 2 species and 17 morphospecies), *Bombus* (Apidae, 16 species), and *Lasioglossum* (14+ species) (Appendix).

DISCUSSION

This is the first thorough examination of the wild bee fauna in the Palouse Prairie. We noted several first records and range expansions. These data provide a baseline of presence and abundance of prairie-inhabiting bee species which will be useful in evaluating declines or range contractions of wild bees in the western United States. As Kimoto *et al.* (2012) noted, the Pacific Northwest bunchgrass ecosystem supports a rich community of wild bees. By utilizing a more diverse array of trapping methods and by identifying more individuals to species rather than morphospecies, we were able to more fully characterize the community of bees inhabiting bunchgrass prairie.

Remarkably, bee richness in the Palouse is greater than recorded for most studies in the extensive tallgrass prairie (Table 1). This may be due in part to sampling effort,

Table 1. Studies of prairie or grassland inhabiting bees.

Habitat	Years Collecting	Specimens Collected	Species Detected	Sampling Method	Sampling Period	Citation
Iowa tallgrass prairie	2	3566	86	pan trap	May–August	Davis <i>et al.</i> , 2008
grasslands near Boulder, CO	5	5207	104	pan trap & net	May–August	Kearns & Oliveras, 2009
Minnesota tallgrass prairie	3	3702	127	net	May–September	Reed, 1995
Illinois tallgrass prairie	1	4622	111	malaise trap, pan trap, & vane trap	June–October	Geroff <i>et al.</i> , 2014
Iowa tallgrass prairie	1	1149	73	pan trap & net	June–August	Hendrix <i>et al.</i> , 2010
Iowa tallgrass prairie & ruderal areas	1	582	56	pan traps & nets	June–August	Kwaiser & Hendrix, 2008
Wyoming shortgrass prairie	2	—	200	net	May–August	Tepedino & Stanton, 1981
Zumwalt bunchgrass prairie	2	9158	211	blue vane trap	June–August	Kimoto <i>et al.</i> , 2012
Palouse bunchgrass prairie	2	13,241	210	vane trap, pan trap, & net	May–July	this report

since a larger number of collected individuals will yield more detected species. Additionally, a wider variety of collection methods could have increased the bee fauna sampled, as trap type can affect the taxa detected (Geroff *et al.*, 2014). Finally, the Palouse and Zumwalt regions both have a more diverse array of habitats nearby. The Palouse Prairie is surrounded by forest, sagebrush steppe, and arid grasslands. Many detected taxa could be adapted to more mesic or arid environments, only marginally present on the Palouse.

Historic records from GBIF and Tepedino & Griswold (1995) list 273 positively identified species (*i.e.*, not morphospecies or generic-level determinations) of native bees in the Palouse region. Of the 174 positively identified species we collected, 117 species were previously noted as denizens of the Palouse and 57 species were not formerly observed in the region. Species previously unknown to the area are primarily in *Osmia* (12 species), *Andrena* (7 species), *Megachile* Latreille (5 species), and *Eucera* Scopoli (5 species). Discrepancies between the list of species historically present and the list of species observed in our study are stark. We observed less than half of historically present species and nearly a third of the species we identified were not previously observed in the Palouse region. This highlights the difficulty of fully characterizing the composition of bee communities which tend to be dominated by rare species.

Indeed, despite insect collection data stretching back more than a century, eleven

Table 2. New state records or range expansions for bee species collected in Palouse Prairie fragments.

Species	New record		Previously documented range
	ID	WA	
<i>Andrena fuscicauda</i>	X		Within and coastward of Sierra Nevada and Cascade mountains. Rarely found in central OR and NV.
<i>Andrena semipunctata</i>	X		Arid southwestern US and within and coastward of Sierra Nevada and Cascade mountains. Rarely found in eastern WA.
<i>Andrena shoshoni</i>		X	Very rare. Observed in SD and WY.
<i>Andrena aff. waldmerei</i>	X	X	Southern CA, more rarely found in northern CA.
<i>Anthophora affabilis</i>		X	Arid southwestern US, north to OR and ID.
<i>Melissodes plumosa</i>	X		Within and coastward of Sierra Nevada and Cascade mountains. One specimen observed in ND.
<i>Hylaeus granulatus</i>	X	X	CA to CO, north to OR.
<i>Osmia aglaia</i>		X	CA and OR.
<i>Osmia thysanisca</i>		X	OR, WY, and CA.
<i>Osmia trifoliama</i>	X		Within and coastward of Sierra Nevada and Cascade mountains.
<i>Stelis interrupta</i>	X		Southwest US, Cascade mountains in OR.
<i>Megachile snowi</i>			TX to CA. North to southern ID.
<i>Osmia raritatis</i>			CA to CO. North to Cascade mountains and southern ID.

species were recovered that are new records for Washington, Idaho, or both (Table 2), and range expansions were noted for two species, including *Megachile snowi* Mitchell (Table 2). However, *M. snowi* was recently elevated to the rank of species, being previously regarded as a subspecies of *Megachile mendica* Cresson (Bzdyk, 2012), for which there are records in the Palouse region, so it is not clear if this species has been previously observed in the Palouse region or not.

Since historic data for non-*Bombus* bee species are sparse, it is difficult to determine if once common species have disappeared from the region. However, *Osmia lignaria* Say was consistently found in the Palouse region by various collectors between 1905 and 1991 (GBIF, 2015), and was the most commonly detected species of *Osmia* during this period. While we did not recover this species in 2012 or 2013, *O. lignaria* is active very early in the spring and the preponderance (60%) of records in GBIF were collected in March or April, before sampling commenced in the present study. So, our failure to detect this species may be because we began sampling too late in the year. While we did fail to detect other species previously found on the Palouse, the paucity of historical data makes it impossible to say if this was due to our times of sampling, the rarity of these species, or their actual absence.

Unlike other bee taxa, data on bumble bee species is complete enough to make statements about alterations to the community over time and space. Despite differing trapping methods, Bray-Curtis dissimilarity analysis shows the three *Bombus* datasets from the Palouse Prairie (historic net collections, 2002 and 2003 pitfall traps, contemporary mixed methods) to be more similar to one another than to the Zumwalt Prairie

Table 3. Bray-Curtis dissimilarity matrix for *Bombus* community data from the Palouse Prairie (this report), Okanogan National Forest (Wilson *et al.*, 2010), the Zumwalt Prairie (Kimoto *et al.*, 2012), the Palouse Prairie from 2002 and 2003 (Hatten *et al.*, 2013), and Palouse Prairie data collected prior to 2001 (GBIF, 2015).

	Current Palouse	Okanogan	Zumwalt	Palouse 2002,2003
Okanogan	72.8	—		
Zumwalt	63.5	43.3	—	
Palouse 2002,2003	55.5	87.6	78.4	—
Historic Palouse	36.6	62.0	60.7	68.1

or the Okanogan National Forest, suggesting the Palouse Prairie has a distinctive *Bombus* community not shared by similar, nearby systems (Table 3). Within the Palouse, a few species of *Bombus* have either declined in abundance or disappeared from the region entirely. *Bombus occidentalis* Greene was once common in the Inland Northwest but is now rare in the region (Stephen, 1957; Rao & Stephen, 2007; Rao *et al.*, 2011; Rhoades *et al.*, 2016). GBIF data for the Palouse region shows 292 *B. occidentalis* collected between 1888 and 1997, forming about 16% of all pre-2000 *Bombus* occurrences recorded in GBIF for the Palouse region. *Bombus occidentalis* was present in our study, but at lower rates than is evident in the historical data (2.9% of total *Bombus*), mirroring trends found throughout its range (Cameron *et al.*, 2011). Additionally, one species of bumble bee listed as vulnerable [*B. morrisoni* Cresson (Hatfield *et al.*, 2014)] and one listed as critically endangered [*B. suckleyi* Greene (Hatfield *et al.*, 2015)] were not observed in this survey, despite likely past records of occurrence (GBIF, 2015).

The Palouse Prairie is a unique region that has been heavily impacted through fragmentation and habitat loss caused by conversion to agriculture (Donovan *et al.*, 2009). This study adds bees to this list of distinctive and diverse Palouse fauna and contributes to our limited but growing knowledge of the bees of the inland Northwest.

ACKNOWLEDGEMENTS

Thanks to Michael Orr, Karen Wright, Skyler Burrows, Jamie Strange, Jonathan Koch, and Zach Portman for help with identification and insect curation. This work was funded through an NSF-IGERT grant to the University of Idaho.

REFERENCES

- Ashman, T.L., T.M. Knight, J.A. Steets, P. Amarasekare, M. Burd, D.R. Campbell, M.R. Dudash, M.O. Johnston, S.J. Mazer, R.J. Mitchell, M.T. Morgan, & W.G. Wilson. 2004. Pollen limitation of plant reproduction: Ecological and evolutionary causes and consequences. *Ecology* 85(9): 2408–2421.
- Bailey, R.G. 1995. Descriptions of the ecoregions of the United States [2nd Revision]. *U.S. Department of Agriculture, Miscellaneous Publication* 1391: 1–77.
- Bartomeus, I., J.S. Ascher, J. Gibbs, B.N. Danforth, D.L. Wagner, S.M. Hedtke, & R. Winfree. 2013. Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proceedings of the National Academy of Sciences, USA* 110(12): 4656–4660.
- Biesmeijer, J.C., S.P.M. Roberts, M. Reemer, R. Ohlemüller, M. Edwards, T. Peeters, A.P. Schafers, S.G. Potts, R. Kleukers, C.D. Thomas, J. Settele, & W.E. Kunin. 2006. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* 313(5785): 351–354.

- Black, A.E., E. Strand, P. Morgan, J.M. Scott, R.G. Wright, & C. Watson. 1998. Biodiversity and land-use history of the Palouse Bioregion: Pre-European to present. In: Sisk, T.D. (Ed.), *Perspectives on the Land Use History of North America: A Context for Understanding Our Changing Environment*: 85–99. U.S. Geological Survey, Biological Resources Division, Biological Science Report USGS/BRD/BSR–1998–0003; Washington, D.C.; 104 pp.
- Brown, B.J., R.J. Mitchell, & S.A. Graham. 2002. Competition for pollination between an invasive species (purple loosestrife) and a native congener. *Ecology* 83(8): 2328–2336.
- Brown, M.J.F., & R.J. Paxton. 2009. The conservation of bees: A global perspective. *Apidologie* 40(3): 410–416.
- Bzdyk, E.L. 2012. A revision of the *Megachile* subgenus *Litomegachile* Mitchell with an illustrated key and description of a new species (Hymenoptera, Megachilidae, Megachilini). *ZooKeys* 221: 31–61.
- Cameron, S.A., J.D. Lozier, J.P. Strange, J.B. Koch, N. Cordes, L.F. Solter, & T.L. Griswold. 2011. Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences, USA* 108(2): 662–667.
- Colwell, R.K., A. Chao, N.J. Gotelli, S.Y. Lin, C.X. Mao, R.L. Chazdon, & J.T. Longino. 2012. Models and estimators linking individual-based and sample-based rarefaction, extrapolation and comparison of assemblages. *Journal of Plant Ecology* 5(1): 3–21.
- Darvill, B., J.S. Ellis, G.C. Lye, & D. Goulson. 2006. Population structure and inbreeding in a rare and declining bumblebee, *Bombus muscorum* (Hymenoptera: Apidae). *Molecular Ecology* 15(3): 601–611.
- Daubenmire, R.F. 1942. An ecological study of the vegetation of southeastern Washington and adjacent Idaho. *Ecological Monographs* 12(1): 53–79.
- Davis, C. 2015. *Biodiversity and Culturally Significant Plants of the Palouse Prairie*. Ph.D. Dissertation, University of Idaho; Moscow, ID; 128 pp.
- Davis, J.D., S.D. Hendrix, D.M. Debinski, & C.J. Hemsley. 2008. Butterfly, bee and forb community composition and cross-taxon incongruence in tallgrass prairie fragments. *Journal of Insect Conservation* 12(1): 69–79.
- Donovan, S.M., C. Looney, T. Hanson, Y. Sánchez de León, J.D. Wulhorst, S.D. Eigenbrode, M. Jennings, J. Johnson-Maynard, & N.A. Bosque-Pérez. 2009. Reconciling social and biological needs in an endangered ecosystem: The Palouse as a model for bioregional planning. *Ecology and Society* 14(1): 9 [1–24].
- Droege, S. 2009. *The Very Handy Bee Manual: How to Catch and Identify Bees and Manage a Collection*. 73 pp. [<http://www.pwrc.usgs.gov/nativebees/Handy%20Bee%20Manual/Handy%20Bee%20Manual.pdf>; last accessed 20 February 2016].
- Ellis, J.S., M.E. Knight, B. Darvill, & D. Goulson. 2006. Extremely low effective population sizes, genetic structuring and reduced genetic diversity in a threatened bumblebee species, *Bombus sylvarum* (Hymenoptera: Apidae). *Molecular Ecology* 15(14): 4375–4386.
- Ertter, B., & B. Moseley. 1992. Floristic regions of Idaho. *Journal of the Idaho Academy of Science* 28(2): 57–70.
- Fitzpatrick, Ú., T.E. Murray, R.J. Paxton, J. Breen, D. Cotton, V. Santorum, & M.J.F. Brown. 2007. Rarity and decline in bumblebees – a test of causes and correlates in the Irish fauna. *Biological Conservation* 136(2): 185–194.
- Foley, J.A., R. DeFries, G.P. Asner, C. Barford, G. Bonan, S.R. Carpenter, F.S. Chapin, M.T. Coe, G.C. Daily, H.K. Gibbs, J.H. Helkowski, T. Holloway, E.A. Howard, C.J. Kucharik, C. Monfreda, J.A. Patz, I.C. Prentice, N. Ramankutty, & P.K. Snyder. 2005. Global consequences of land use. *Science* 309(5734): 570–574.
- Fontaine, C., I. Dajoz, J. Meriguet, & M. Loreau. 2005. Functional diversity of plant–pollinator interaction webs enhances the persistence of plant communities. *PLoS Biology* 4(1): e1 [129–135].
- Geroff, R.K., J. Gibbs, & K.W. McCravy. 2014. Assessing bee (Hymenoptera: Apoidea) diversity of an Illinois restored tallgrass prairie: Methodology and conservation considerations. *Journal of Insect Conservation* 18(5): 951–964.
- Goulson, D., G.C. Lye, & B. Darvill. 2008. Decline and conservation of bumble bees. *Annual Re-*

- view of *Entomology* 53: 191–208.
- Goulson, D., E. Nicholls, C. Botías, & E.L. Rotheray. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* 347(6229): 1255957 1–9.
- Hanson, T., Y. Sánchez-de León, J. Johnson-Maynard, & S. Brunsfeld. 2008. Influence of soil and site characteristics on Palouse Prairie plant communities. *Western North American Naturalist* 68(2): 231–240.
- Hatfield, R., S. Jepsen, R. Thorp, L. Richardson, & S. Colla. 2014. *Bombus morrisoni*. The IUCN Red List of Threatened Species 2014: e.T44937666A69004519. [<http://dx.doi.org/10.2305/IUCN.UK.2014-3.RLTS.T44937666A69004519.en> ; last accessed 15 March 2017].
- Hatfield, R., S. Jepsen, R. Thorp, L. Richardson, & S. Colla. 2015. *Bombus suckleyi*. The IUCN Red List of Threatened Species 2015: e.T44937699A46440241. [<http://dx.doi.org/10.2305/IUCN.UK.2015-2.RLTS.T44937699A46440241.en> ; last accessed 15 March 2017].
- Hatten, T.D., S.D. Eigenbrode, N.A. Bosque-Pérez, S. Gebbie, F. Merickel, & C. Looney. 2004. Influence of matrix elements on prairie-inhabiting Curculionidae, Tenebrionidae, and Scarabaeidae in the Palouse. *Proceedings of the North American Prairie Conferences* 19: 101–108.
- Hatten, T.D., N.A. Bosque-Pérez, J.R. Labonte, S.O. Guy, & S.D. Eigenbrode. 2007. Effects of tillage on the activity density and biological diversity of carabid beetles in spring and winter crops. *Environmental Entomology* 36(2): 356–368.
- Hatten, T.D., C. Looney, J.P. Strange, & N.A. Bosque-Pérez. 2013. Bumble bee fauna of Palouse Prairie: Survey of native bee pollinators in a fragmented ecosystem. *Journal of Insect Science* 13(26): 1–19.
- Hendrix, S.D., K.S. Kwaiser, & S.B. Heard. 2010. Bee communities (Hymenoptera: Apoidea) of small Iowa hill prairies are as diverse and rich as those of large prairie preserves. *Biodiversity and Conservation* 19(6): 1699–1709.
- Hurd, P.D., Jr. 1979. Superfamily Apoidea. In: Krombein, K.V., P.D. Hurd, Jr., D.R. Smith, & B.D. Burks (Eds.), *Catalog of Hymenoptera in America North of Mexico, Vol. 2, Apocrita (Aculeata): 1741–2209*. Smithsonian Institution Press; Washington, D.C.; xvi+2209 pp.
- Kearns, C.A., & D.M. Oliveras. 2009. Environmental factors affecting bee diversity in urban and remote grassland plots in Boulder, Colorado. *Journal of Insect Conservation* 13(6): 655–665.
- Keith, D.A., J.P. Rodríguez, K.M. Rodríguez-Clark, E. Nicholson, K. Aapala, A. Alonso, M. Assmussen, S. Bachman, A. Basset, E.G. Barrow, J.S. Benson, M.J. Bishop, R. Bonifacio, T.M. Brooks, M.A. Burgman, P. Comer, F.A. Comin, F. Essl, D. Faber-Langendoen, P.G. Fairweather, R.J. Holdaway, M. Jennings, R.T. Kingsford, R.E. Lester, R.M. Nally, M.A. McCarthy, J. Moat, M.A. Oliveira-Miranda, P. Pisanu, B. Poulin, T. J. Regan, U. Rieckan, M.D. Spalding, & S. Zambrano-Martínez. 2013. Scientific foundations for an IUCN Red List of ecosystems. *PLoS ONE* 8(5): e62111 [1–25].
- Kimoto, C., S.J. DeBano, R.W. Thorp, S. Rao, & W.P. Stephen. 2012. Investigating temporal patterns of a native bee community in a remnant North American bunchgrass prairie using blue vane traps. *Journal of Insect Science* 12(1): 108 1–5.
- Klein, A.M., B.E. Vaissière, J.H. Cane, I. Steffan-Dewenter, S.A. Cunningham, C. Kremen, & T. Tscharntke. 2007. Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society, Series B, Biological Sciences* 274(1608): 303–313.
- Kwaiser, K.S., & S.D. Hendrix. 2008. Diversity and abundance of bees (Hymenoptera: Apiiformes) in native and ruderal grasslands of agriculturally dominated landscapes. *Agriculture, Ecosystems & Environment* 124(3): 200–204.
- Lennartsson, T. 2002. Extinction thresholds and disrupted plant-pollinator interactions in fragmented plant populations. *Ecology* 83(11): 3060–3072.
- Lichthardt, J., & R.K. Moseley. 1997. Status and conservation of the Palouse grassland in Idaho. Idaho Department of Fish and Game and US Fish and Wildlife Service. [https://www.tag.idaho.gov/ifwis/idnhp/cdc_pdf/palou97.pdf; last accessed 20 February 2016]
- Looney, C., & S.D. Eigenbrode. 2012. Characteristics and distribution of Palouse Prairie remnants: implications for conservation planning. *Natural Areas Journal* 32(1): 75–85.
- Looney, C., B. Caldwell, T. Hatten, C. Lorion, & S. Eigenbrode. 2004. Potential habitat factors influencing carrion beetle communities of Palouse Prairie remnants. *Proceedings of the North*

- American Prairie Conferences* 19: 117–121.
- Meffe, G.K., A.W. Gordon, P. Bernhardt, R. Bitner, A. Burquez, S. Buchmann, J. Cane, P.A. Cox, V. Dalton, P. Feinsinger, M. Ingram, D. Inouye, C.E. Jones, K. Kennedy, P.G. Kevan, H. Koo-powitz, R. Medellin, S. Medellin-Morales, & G.P. Nabhan. 1998. The potential consequences of pollinator declines on the conservation of biodiversity and stability of food crop yields. *Conservation Biology* 12(1): 8–17.
- NRC. 2007. *Status of Pollinators in North America*. The National Academies Press; Washington, D.C.; 322 pp.
- Oksanen, J., F.G. Blanchet, R. Kindt, P. Legendre, P.R. Minchin, R.B. O'Hara, G.L. Simpson, P. Solymos, M.H. Stevens, H. Wagner. 2015. Package “vegan.” Community Ecology Package, Version 2–2. [https://cran.r-project.org/web/packages/vegan/index.html; last accessed 15 March 2017]
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. *Annals of the Association of American Geographers* 77(1): 118–125.
- Park, M.G., E.J. Blitzer, J. Gibbs, J.E. Losey, & B.N. Danforth. 2015. Negative effects of pesticides on wild bee communities can be buffered by landscape context. *Proceedings of the Royal Society of London, Series B, Biological Sciences* 282(1809): 2015.0299 [1–9].
- Potts, S.G., J.C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, & W.E. Kunin. 2010. Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology & Evolution* 25(6): 345–353.
- R Core Team. 2015. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing; Vienna, Austria. [https://www.R-project.org/; last accessed 15 March 2017]
- Rao, S., & W.P. Stephen. 2007. *Bombus (Bombus) occidentalis* (Hymenoptera: Apiformes): In decline or recovery. *Pan-Pacific Entomologist* 83(4): 360–362.
- Rao, S., W.P. Stephen, C. Kimoto, & S.J. DeBano. 2011. The status of the “Red-Listed” *Bombus occidentalis* (Hymenoptera: Apiformes) in northeastern Oregon. *Northwest Science* 85(1): 64–67.
- Reed, C.C. 1995. Insects surveyed on flowers in native and reconstructed prairies (Minnesota). *Restoration and Management Notes* 13(2): 210–213.
- Rhoades, P.R., J.B. Koch, L.P. Waits, J.P. Strange, & S.D. Eigenbrode. 2016. Evidence for *Bombus occidentalis* (Hymenoptera: Apidae) populations in the Olympic Peninsula, the Palouse Prairie, and forests of northern Idaho. *Journal of Insect Science* 16(1): 20 [1–5].
- Ribble, D.W. 1974. A revision of the bees of the genus *Andrena* of the Western Hemisphere sub-genus *Scaphandrena*. *Transactions of the American Entomological Society* 100(2): 101–189.
- Sánchez-de León, Y., & J. Johnson-Maynard. 2008. Dominance of an invasive earthworm in native and non-native grassland ecosystems. *Biological Invasions* 11(6): 1393–1401.
- Sarukhan, J., A. Whyte, R. Hassan, R. Scholes, N. Ash, S.T. Carpenter, P.L. Pingali, E.M. Bennett, M.B. Zurek, K. Chopra, R. Leemans, P. Kimar, H. Simons, D. Capistrano, C.K. Samper, M.J. Lee, & C. Raudsepp-Hearne. 2005. *Millennium Ecosystem Assessment: Ecosystems and Human Well-Being*. Island Press; Washington D.C.; x+137 pp.
- Scheper, J., M. Reemer, R. van Kats, W.A. Ozinga, G.T.J. van der Linden, J.H.J. Schaminée, H. Siepel, & D. Kleijn. 2014. Museum specimens reveal loss of pollen host plants as key factor driving wild bee decline in The Netherlands. *Proceedings of the National Academy of Sciences, USA* 111(49): 17552–17557.
- Senapathi, D., L.G. Carvalheiro, J.C. Biesmeijer, C.A. Dodson, R. L. Evans, M. Mc Kerchar, R.D. Morton, E.D. Moss, S.P.M. Roberts, W.E. Kunin, & S.G. Potts. 2015. The impact of over 80 years of land cover changes on bee and wasp pollinator communities in England. *Proceedings of the Royal Society, Series B, Biological Sciences* 282(1806): 20150294 1-8.
- Stephen, W.P. 1957. *Bumble Bees of Western America (Hymenoptera: Apoidea)*. Oregon State College, Agricultural Experiment Station; Corvallis, OR; 163 pp.
- Stephen, W.P., & S. Rao. 2007. Sampling native bees in proximity to a highly competitive food resource (Hymenoptera: Apiformes). *Journal of the Kansas Entomological Society* 80(4): 369–376.
- Tepedino, V.J., & N.L. Stanton. 1981. Diversity and competition in bee-plant communities on short-grass prairie. *Oikos* 36(1): 35–44.

- Tepedino, V.J., & T.L. Griswold. 1995. *The Bees of the Columbia Basin*. Final report, USDA Forest Service; Portland, OR; 212 pp.
- Tisdale, E.W. 1982. Grasslands of western North America: The Pacific Northwest bunchgrass. In: Tisdale, E.W., A. McLean, & T.E. Baker (Eds.), *Grassland Ecology and Classification Symposium*: 223–245. British Columbia Ministry of Forests; Victoria, Canada; x+351 pp.
- Vanbergen, A.J. 2013. Threats to an ecosystem service: Pressures on pollinators. *Frontiers in Ecology and the Environment* 11(5): 251–259.
- White, R.P., S. Murray, M. Rohweder, S.D. Prince, & K.M. Thompson. 2000. *Grassland Ecosystems*. World Resources Institute; Washington, D.C.; xi+69 pp.
- Wilson, J.S., T. Griswold, & O.J. Messinger. 2008. Sampling bee communities (Hymenoptera: Apiformes) in a desert landscape: Are pan traps sufficient? *Journal of the Kansas Entomological Society* 81(3): 288–300.
- Wilson, J.S., L.E. Wilson, L.D. Loftis, & T. Griswold. 2010. The montane bee fauna of North Central Washington, USA, with floral associations. *Western North American Naturalist* 70(2): 198–207.
- Winfrey, R., T. Griswold, & C. Kremen. 2007. Effect of human disturbance on bee communities in a forested ecosystem. *Conservation Biology* 21(1): 213–223.
- Zayed, A. 2004. Effective population size in Hymenoptera with complementary sex determination. *Heredity* 93(6): 627–630.
- Zayed, A., & L. Packer. 2005. Complementary sex determination substantially increases extinction proneness of haplodiploid populations. *Proceedings of the National Academy of Sciences, USA* 102(30): 10742–10746.

Appendix. List of all collected bee species and their abundance in each month in which collection occurred.

Species	2012			2013			Total
	May	June	July	May	June	July	
ANDRENIDAE	322	233	48	193	204	26	1026
<i>Andrena angustitarsata</i> Viereck	13	2		18	20		53
<i>Andrena astragali</i> Viereck & Cockerell	2						2
<i>Andrena candida</i> Smith	4	1					5
<i>Andrena chlorogaster</i> Viereck				3			3
<i>Andrena cressonii infasciata</i> Lanham	6	8	2	4	3		23
<i>Andrena fuscicauda</i> Viereck ¹	2			3			5
<i>Andrena hemileuca</i> Viereck	12			3			15
<i>Andrena hippotes</i> Robertson	4						4
<i>Andrena melanochoa</i> Cockerell	10	20	1	2	15	1	49
<i>Andrena merriami</i> Cockerell	2			4	3		9
<i>Andrena microchlora</i> Cockerell	39	13		16	5		73
<i>Andrena nigrihirta</i> (Ashmead)				8	25		33
<i>Andrena nigrocaerulea</i> Cockerell	112	62	3	72	45		294
<i>Andrena nivalis</i> Smith	13	6		12	13		44
<i>Andrena pallidifovea</i> Viereck	3	22	2		18		45
<i>Andrena piperi</i> Viereck				1	1		2
<i>Andrena prunorum</i> Cockerell	11	42	11	26	18	15	123
<i>Andrena semipunctata</i> Cockerell ¹		3	1		3		7
<i>Andrena shoshoni</i> Ribble ²				3			3
<i>Andrena sola</i> Viereck	6	2	1	2	7		18
<i>Andrena</i> spp.	13						13
<i>Andrena subtilis</i> Smith	3	8	2	2	2		17
<i>Andrena thaspiae</i> Graenicher		1	1	2	3	2	9
<i>Andrena topazana</i> Cockerell		2	2				4
<i>Andrena trevoris</i> Cockerell			3			5	8
<i>Andrena vierecki</i> Cockerell	1	1	1			1	4
<i>Andrena aff. waldmeri</i> LaBerge & Bouseman ^{1,2}	1	1		2			4
<i>Panurginus atriceps</i> (Cresson)	61	37	16	10	23	2	149
<i>Panurginus</i> spp.	4	1					5
<i>Perdita</i> sp.		1					1
<i>Perdita wyomingensis</i> Cockerell			2				2
APIDAE	450	366	195	305	378	606	2281
<i>Anthophora affabilis</i> Cresson ²					1		1
<i>Anthophora bombooides</i> Kirby	1		5		5	1	12
<i>Anthophora edwardsii</i> Cresson	1			1			2

Appendix. Continued.

Species	2012			2013			Total
	May	June	July	May	June	July	
<i>Anthophora occidentalis</i> Cresson						1	1
<i>Anthophora pacifica</i> Cresson				8			8
<i>Anthophora porterae</i> Cockerell				1			1
<i>Anthophora terminalis</i> Cresson					2	1	3
<i>Anthophora urbana</i> Cresson			1		2	17	20
<i>Anthophora ursina</i> Cresson	2	1	1	3	4		11
<i>Apis mellifera</i> Linnaeus	21	36	23	3	13	9	105
<i>Bombus appositus</i> Cresson	13	4	3	4	13	12	49
<i>Bombus bifarius</i> Cresson	69	18	4	2	1	1	95
<i>Bombus californicus</i> Smith	1	2	2	2	3	2	12
<i>Bombus centralis</i> Cresson	25	4	4	17	7	9	66
<i>Bombus fernaldae</i> (Franklin)				1			1
<i>Bombus fervidus</i> (Fabricius)	27	9	5	48	17	16	122
<i>Bombus flavifrons</i> Cresson	2					2	4
<i>Bombus griseocollis</i> (DeGeer)		1				7	8
<i>Bombus huntii</i> Greene	7	2		3	1	1	14
<i>Bombus insularis</i> (Smith)	12	5	1	1			19
<i>Bombus melanopygus</i> Nylander	1						1
<i>Bombus mixtus</i> Cresson	12		3	1		2	18
<i>Bombus nevadensis</i> Cresson	71	6	8	28	22	25	160
<i>Bombus occidentalis</i> Greene	9		1	1	1		12
<i>Bombus rufocinctus</i> Cresson	41	37	26	18	27	21	170
<i>Bombus</i> sp.						1	1
<i>Bombus vagans</i> Smith	1	1	1				3
<i>Ceratina acantha</i> Provancher	18	45	20	30	22	17	152
<i>Ceratina nanula</i> Cockerell	71	118	14	78	62	92	435
<i>Ceratina pacifica</i> H.S. Smith	3	4	34	8	18	82	149
<i>Ceratina</i> spp.	5					2	7
<i>Diadasia enavata</i> (Cresson)						7	7
<i>Diadasia nigrifrons</i> (Cresson)		1	2				3
<i>Epeolus minimus</i> (Robertson)			1				1
<i>Epeolus</i> sp.						1	1
<i>Eucera actiosa</i> (Cresson)	2	3			7		12
<i>Eucera delphinii</i> (Timberlake)	1			4			5
<i>Eucera edwardsii</i> (Cresson)		9	4	4	21		38
<i>Eucera frater</i> (Cresson)	6	23	16	13	103	50	211
<i>Eucera hurdi</i> (Timberlake)	4	1		1	2		8

Appendix. Continued.

Species	2012			2013			Total
	May	June	July	May	June	July	
<i>Eucera</i> spp.						3	3
<i>Habropoda cineraria</i> (Smith)	1						1
<i>Melecta pacifica</i> Cresson	2	1		1	1		5
<i>Melissodes</i> aff. <i>plumosa</i> LaBerge						4	4
<i>Melissodes agilis</i> Cresson			1				1
<i>Melissodes communis</i> Cresson			1		1	29	31
<i>Melissodes lupina</i> Cresson					1	90	91
<i>Melissodes menuachus</i> Cresson			1				1
<i>Melissodes metenua</i> Cockerell						20	20
<i>Melissodes microsticta</i> Cockerell			3		5	20	28
<i>Melissodes pallidisignata</i> Cockerell						1	1
<i>Melissodes plumosa</i> LaBerge ¹						20	20
<i>Melissodes rivalis</i> Cresson			2		1	12	15
<i>Melissodes</i> spp.						13	13
<i>Melissodes</i> sp. 1						3	3
<i>Melissodes</i> sp. 2						1	1
<i>Melissodes</i> sp. 3						1	1
<i>Melissodes</i> sp. 4						1	1
<i>Melissodes verbesinarum</i> Cockerell			2			6	8
<i>Nomada edwardsii</i> Cresson	1	1					2
<i>Nomada hemphilli</i> Cockerell				2			2
<i>Nomada</i> spp.	5	7		7	2	1	22
<i>Nomada</i> sp. 1	1	1		3			5
<i>Nomada</i> sp. 2	1	3		4	3		11
<i>Nomada</i> sp. 3	1	1					2
<i>Nomada</i> sp. 4	1			3			4
<i>Nomada</i> sp. 5		5					5
<i>Nomada</i> sp. 6					2		2
<i>Nomada</i> sp. 7	3						3
<i>Nomada</i> sp. 8					1		1
<i>Nomada</i> sp. 9					1		1
<i>Nomada</i> sp. 10				1			1
<i>Nomada</i> sp. 11	2				1		3
<i>Nomada</i> sp. 12	5	3		2	3		13
<i>Nomada</i> sp. 13					1		1
<i>Nomada</i> sp. 14	1						1
<i>Nomada</i> sp. 15		1					1

Appendix. Continued.

Species	2012			2013			Total
	May	June	July	May	June	July	
<i>Nomada</i> sp. 16				1	1		2
<i>Nomada</i> sp. 17				1			1
<i>Triepeolus heterurus</i> (Cockerell & Sandhouse)						2	2
Colletidae	0	13	6	1	16	3	36
<i>Colletes fulgidus</i> Swenk		1			2		
<i>Hylaeus affinis</i> (Smith)		2	1		1	1	5
<i>Hylaeus conspicuus</i> (Metz)					1		1
<i>Hylaeus granulatus</i> (Metz) ^{1,2}		10	2	1	8	1	22
<i>Hylaeus modestus</i> Say					1	1	2
<i>Hylaeus</i> spp.			2		1		3
<i>Hylaeus verticalis</i> (Cresson)			1				1
<i>Hylaeus wootoni</i> (Cockerell)					2		2
Halictidae	1635	1721	1030	1365	1722	975	8448
<i>Agapostemon angelicus</i> Cockerell	50	145	88	24	192	98	597
<i>Agapostemon coloradinus</i> (Vachal)	3				1	3	7
<i>Agapostemon femoratus</i> Crawford				7	11	4	22
<i>Agapostemon virescens</i> (Fabricius)	35	136	219	25	219	225	859
<i>Halictus confusus</i> Smith	4	16					20
<i>Halictus farinosus</i> Smith	11	9	11	19	17	3	70
<i>Halictus ligatus</i> Smith	46	21	6	13	29	10	125
<i>Halictus rubicundus</i> (Christ)	45	51	22	170	78	7	373
<i>Halictus</i> spp.	12	1		11	1		25
<i>Halictus tripartitus</i> Cockerell	259	213	113	324	276	172	1357
<i>Lasioglossum anhypops</i> McGinley	1	2					3
<i>Lasioglossum athabascense</i> (Sandhouse)	4		2		1	1	8
<i>Lasioglossum colatum</i> (Vachal)	3	16	14	5	14	1	53
<i>Lasioglossum Hemihalictus</i> series	815	661	365	694	652	328	3515
<i>Lasioglossum egregium</i> (Vachal)	16	90	45	3	51	41	246
<i>Lasioglossum mellipes</i> (Crawford)				2	7		9
<i>Lasioglossum olympiae</i> (Cockerell)	16	29	4	2	5		56
<i>Lasioglossum ovaliceps</i> (Cockerell)		1			1		2
<i>Lasioglossum pacificum</i> (Cockerell)		4	7	8	6	5	30
<i>Lasioglossum paraforbesii</i> McGinley						2	2
<i>Lasioglossum sisymbrii</i> (Cockerell)	64	238	81	27	91	24	525
<i>Lasioglossum</i> spp.	172	4	1				177
<i>Lasioglossum titusi</i> (Crawford)	50	54	27	13	18	40	202

Appendix. Continued.

Species	2012			2013			Total
	May	June	July	May	June	July	
<i>Lasioglossum trizonatum</i> (Cresson)	3	3	1	4	30	2	43
<i>Lasioglossum zonulum</i> (Smith)		5	12	8	4		29
<i>Sphecodes</i> spp.			1			7	8
<i>Sphecodes</i> sp. 1	2	1			1		4
<i>Sphecodes</i> sp. 2	3	1	1	2			7
<i>Sphecodes</i> sp. 3	2	1					3
<i>Sphecodes</i> sp. 4		1	1		1		3
<i>Sphecodes</i> sp. 5		1	1				2
<i>Sphecodes</i> sp. 6						1	1
<i>Sphecodes</i> sp. 7					1		1
<i>Sphecodes</i> sp. 8			2				2
<i>Sphecodes</i> sp. 9		1					1
<i>Sphecodes</i> sp. 10	1	4	1	1	6		13
<i>Sphecodes</i> sp. 11	8	11	5	3	8	1	36
<i>Sphecodes</i> sp. 12					1		1
<i>Sphecodes</i> sp. 13		1					1
Megachilidae	215	385	198	180	287	234	1510
<i>Anthidium manicatum</i> (Linnaeus) ³						1	1
<i>Anthidium utahense</i> Swenk			1		1	1	3
<i>Atoposmia copelandica</i> (Cockerell)					1		1
<i>Dianthidium curvatum</i> (Smith)						3	3
<i>Dianthidium subparvum</i> Swenk			2		2	4	8
<i>Heriades carinatus</i> Cresson			1			5	6
<i>Hoplitis albifrons argentifrons</i> (Cresson)		3	3		1	4	11
<i>Hoplitis fulgida</i> (Cresson)		8	4	3	13		28
<i>Hoplitis grinnelli</i> (Cockerell)	1	6	2		3	1	13
<i>Hoplitis hypocrita</i> (Cockerell)	3	3	1	7	6		20
<i>Hoplitis producta</i> (Cresson)	1	29	37	5	26	15	113
<i>Hoplitis sambuci</i> Titus			1		1	4	6
<i>Megachile apicalis</i> Spinola ³						2	2
<i>Megachile brevis</i> Say				1	2	4	7
<i>Megachile gemula</i> Cresson		1					1
<i>Megachile gentilis</i> Cresson					2	6	8
<i>Megachile lippiae</i> Cockerell						1	1
<i>Megachile</i> sp.						1	1
<i>Megachile melanophaea</i> Smith						2	2
<i>Megachile mellitarsis</i> Cresson	1						1

Appendix. Continued.

Species	2012			2013			Total
	May	June	July	May	June	July	
<i>Megachile montivaga</i> Cresson		12	12		9	24	57
<i>Megachile parallela</i> Smith						2	2
<i>Megachile perihirta</i> Cockerell		3	9		13	65	90
<i>Megachile relativa</i> Cresson						1	1
<i>Megachile rotundata</i> (Fabricius) ³						4	4
<i>Megachile snowi</i> Mitchell						1	1
<i>Osmia</i> spp.	22		1				34
<i>Osmia aglaia</i> Sandhouse ²	1						1
<i>Osmia albolateralis</i> Cockerell	3	2	2	5	26	15	53
<i>Osmia atrocyanea</i> Cockerell	20	19	3	15	22	4	83
<i>Osmia bakeri</i> Sandhouse	1			1			2
<i>Osmia brevis</i> Cresson	2	10					12
<i>Osmia bruneri</i> Cockerell	8	5			7	2	22
<i>Osmia caerulescens</i> (Linnaeus) ³		8	12		8	1	29
<i>Osmia californica</i> Cresson	3		1	1	1		6
<i>Osmia calla</i> Cockerell			4	1	5	4	14
<i>Osmia cobaltina</i> Cresson	1	1					2
<i>Osmia coloradensis</i> Cresson	3	3		8		1	15
<i>Osmia densa</i> Cresson	6	5		25	9	1	46
<i>Osmia dolerosa</i> Sandhouse					2		2
<i>Osmia giliarum</i> Cockerell	6	10	2	1		1	20
<i>Osmia grindeliae</i> Cockerell					1		1
<i>Osmia integra</i> Cresson					1		1
<i>Osmia iridis</i> Cockerell & Titus	1	3		5			9
<i>Osmia kincaidii</i> Cockerell	10	16	5	7	7	3	48
<i>Osmia marginipennis</i> Cresson			1				1
<i>Osmia nemoris</i> Sandhouse			1	4	2		7
<i>Osmia nigrifrons</i> Cresson		1		1			2
<i>Osmia paradisiaca</i> Sandhouse			1		2	3	6
<i>Osmia proxima</i> Cresson	4	11	7	3	3		28
<i>Osmia pusilla</i> Cresson	30	44	13	2	10	14	113
<i>Osmia raritatis</i> Michener	1			1	1		3
<i>Osmia simillima</i> Smith	5	6	1	1	2	2	17
<i>Osmia texana</i> Cresson					1		1
<i>Osmia thyanisca</i> Michener ²	1						1
<i>Osmia trevoris</i> Cockerell	73	167	62	77	81	29	489
<i>Osmia trifoliama</i> Sandhouse ¹		4		4	3	1	12

Appendix. Completed.

Species	2012			2013			Total
	May	June	July	May	June	July	
<i>Osmia tristella</i> Cockerell			2		2		4
<i>Osmia unca</i> Michener	2						2
<i>Osmia vandykei</i> Sandhouse	1			2			3
<i>Stelis holocyanea</i> (Cockerell)		1			5		6
<i>Stelis interrupta</i> Cresson ¹			3				3
<i>Stelis montana</i> Cresson	5	3	1				9
<i>Stelis monticola</i> Cresson			1		1	1	3
<i>Stelis</i> sp.					1		1
<i>Stelis</i> sp. 1		1	1				2
<i>Stelis</i> sp. 2					1	1	2
<i>Stelis</i> sp. 3			1		1		2
<i>Stelis submarginata</i> Cresson					2		2

¹ New state record for Idaho.

² New state record for Washington.

³ Exotic.



Journal of Melittology

A Journal of Bee Biology, Ecology, Evolution, & Systematics

The *Journal of Melittology* is an international, open access journal that seeks to rapidly disseminate the results of research conducted on bees (Apoidea: Anthophila) in their broadest sense. Our mission is to promote the understanding and conservation of wild and managed bees and to facilitate communication and collaboration among researchers and the public worldwide. The *Journal* covers all aspects of bee research including but not limited to: anatomy, behavioral ecology, biodiversity, biogeography, chemical ecology, comparative morphology, conservation, cultural aspects, cytogenetics, ecology, ethnobiology, history, identification (keys), invasion ecology, management, melittopalynology, molecular ecology, neurobiology, occurrence data, paleontology, parasitism, phenology, phylogeny, physiology, pollination biology, sociobiology, systematics, and taxonomy.

The *Journal of Melittology* was established at the University of Kansas through the efforts of Michael S. Engel, Victor H. Gonzalez, Ismael A. Hinojosa-Díaz, and Charles D. Michener in 2013 and each article is published as its own number, with issues appearing online as soon as they are ready. Papers are composed using Microsoft Word® and Adobe InDesign® in Lawrence, Kansas, USA.

Editor-in-Chief

Michael S. Engel
University of Kansas

Assistant Editors

Victor H. Gonzalez
University of Kansas

Ismael A. Hinojosa-Díaz
Universidad Nacional Autónoma de México

Journal of Melittology is registered in ZooBank (www.zoobank.org), and archived at the University of Kansas and in Portico (www.portico.org).

<http://journals.ku.edu/melittology>
ISSN 2325-4467