

USE OF PROBIOTICS AS GROWTH PROMOTERS AND IMMUNOSTIMULATORS IN FINGERLINGS OF CYPRINID FISH SPECIES

Ljubojević D.^{1}, Živkov-Baloš M.¹, Mihaljev Ž.¹, Jakšić S.¹, Dorđević V.², Radosavljević V.³, Ćirković M.¹*

¹ Scientific Veterinary Institute „Novi Sad“, Rumenački put 20, 21000 Novi Sad, Republic of Serbia

² Institute of Meat Hygiene and Technology, Kačanskog 13, 11000 Belgrade, Republic of Serbia

³ Scientific Institute of Veterinary Science of Serbia, Vojvode Toze 14, 11000 Belgrade, Republic of Serbia

*Correspondence: ljubojevic.ljubojevicd.dragana@gmail.com; dragana@niv.ns.ac.rs

Abstract

Intensive aquaculture production has required the development of an individual's resistance to disease rather than depending upon antibiotics or chemotherapeutics. The role of gastrointestinal microflora in disease resistance has been established in many fish species, which has led to the concept of manipulating gastrointestinal microflora for better health management. A number of studies has been conducted in different fish species with various useful microorganisms called 'probiotics' to amplify gastrointestinal microflora to fight against various infectious diseases. Probiotics are beneficial microorganisms which protect the host from diseases. Probiotic protection can be achieved by various mechanisms. Most probiotics used in aquaculture belong to the lactic acid bacteria, the genus *Bacillus*, the photosynthetic bacteria, the yeast, notwithstanding other genera and species have also been used. The immunostimulatory effect of probiotics has been established in many fish species, but their direct involvement in the immune response is not well established. It has also been proven that the application of probiotics in aquaculture has beneficial effects on growth of fish as well as on the environment. At present, data about the efficacy of probiotics in commercial aquaculture of Serbia is still lacking. This review discusses mainly the studies and applications about effects, problems and perspectives of probiotics used in fingerlings of cyprinid fish species, and highlights immunostimulatory effects and growth promotion effects of commercial probiotic products. In the present paper the results that show positive influence of probiotics in cyprinides nutrition on production performance and immune system are summarized. Special accent is given to criteria for proper selection of probiotics in cyprinides production.

Key words: *cyprinids, growth promoters, immune system, probiotics*

Introduction

In the several past years, the usage of some antibiotics has been forbidden by several countries due to serious environmental hazards and various adverse effects in many fish species (Gatesoupe, 2007). Furthermore, resistance of pathogen could increase which often leads to spreading of disease, especially under some stressful culture conditions or if antibiotics are used in lower doses for economical reasons (Suzer et al., 2008). Functional additives, like probiotics, represent a new concept on aquaculture. Probiotics have been

increasingly used in aquaculture after their first admission (Kozasa, 1986). Food and Agriculture Organization (FAO) and World Health Organization (WHO) defined probiotics as live micro-organisms that when administered in appropriate amount confer a health benefit on the host (FAO/WHO, 2001). Probiotics supplemented in fish feed resulted in growth performances and feed utilization better than that of the control basal feed, suggesting that the addition of probiotics reduced the culture cost of cyprinid fish species (Jovanović et al., 2011; Miščević et al., 2011 and 2012). In addition, the application of probiotics in aquaculture as the environment friendly treatments was also increasing rapidly (Gatesoupe, 1999). The immunostimulatory effect of probiotics has been also observed in many studies on different fish species (Salinas et al., 2008). Probiotics should also be considered as potentially useful for the control of fish diseases.

At present, data about the efficacy of probiotics in commercial aquaculture of Serbia are still lacking. This review discusses mainly the studies about applications, effects, problems and perspectives of probiotics used in fingerlings of cyprinid fish species, and highlights immunostimulatory effects and growth promotion effects of commercial probiotic products. In the present paper the results that show positive influence of probiotics in cyprinides nutrition on production performance and immunological system are summarized. Special accent is given to criteria for proper selection of probiotics in cyprinides production.

Probiotics

Probiotics represent components of a micro-organism that is beneficial to the health of the host, so they include certain bacteria strains and yeasts that are not adverse on continued use for a longer period of time (Irianto and Austin, 2002). The mode of action is that good and useful microbes multiply in order to compete with the harmful ones, thus suppressing their growth (Sahu et al., 2008). Wang et al. (2008) suggested that the definition of a probiotic in aquaculture should include the addition of live naturally occurring bacteria to ponds in which fish live. Most probiotics used in aquaculture belong to the genus *Bacillus*, *Lactobacillus* sp., *Bifidobacterium* sp., *Vibrio* sp., *Saccharomyces* sp., *Enterococcus* sp., the photosynthetic bacteria, the yeast, (Kumar et al., 2006) notwithstanding other genera and species that have also been used. Fuller (1987) noted that a probiotic should provide actual benefit to the host, be capable to survive in the digestive tract, be suitable for commercialization and be stable and viable for long-term storage conditions.

Probiotics as growth promoters in fingerlings of cyprinid fish species

Growth process in fish depends on numerous factors (Azaza et al., 2008; Deane and Woo, 2009). One of the most important factors on growth and health of fish is feed supply (Lall and Lewis-McCrea, 2007). According to DeVrese and Marteau (2007), mechanism and function of probiotics effect depend principally on the interactions between probiotic species and microbiota of the host or with immuno competent cell of the intestinal mucous. The growth of fish might have occurred by probiotics interaction that increased minerals absorption (Tewe et al., 1999). Jovanović et al. (2011), Miščević et al. (2011 and 2012) observed that the adding commercial probiotics in the diets for common carp fingerlings improve weight of fish. Results obtained by Faramarzi et al. (2011) indicate that the fingerlings of common carp fed with diets with probiotics supplemented exhibited greater growth than those fed with the control diet. Of the four probiotics treatments in the mentioned trial, the 40% protein diet supplemented with yeast showed the best growth

performance and feed efficiency, suggesting that yeast is an appropriate growth stimulating additive in common carp cultivation. Sahandi et al. (2012) also reported the similar results that demonstrate the effect of *Bacillus* spp on length of silver carp and increased fish growth. Effect of probiotics for common carp (*Cyprinus carpio* L.) and some other warm-water fish species based on growth performance and feed utilization (feed conversion ratio, FCR) were investigated also by Wang and Xu, 2006; Wang et al., 2008; Lara-Flores et al., 2010). The above mentioned authors established that the use of probiotics improve growth performance and show the growth promoting effect. All the probiotics supplemented diets resulted in growth performance and feed utilization better than that of the control diets, suggesting that the addition of probiotics reduced the culture cost of common carp (*Cyprinus carpio*). The use of probiotics was suggested as a tool for improving the growth performance and increasing feed digestibility (Sahandi et al., 2012; Ljubojević et al., 2013). Increase of the weight of fish under the probiotics consumption is not completely understood, but according to Lilly and Stillwell (1965) probiotics improve the gastrointestinal microbial populations and then increase enzyme secretion (Suzer et al., 2008) and consequently increase utilization of nutrients and protein anabolism as the length and weight gain occurs in fish.

Probiotics as immunostimulators in fingerlings of cyprinid fish species

Probiotics are used as dietary supplementations in aquaculture and their role in intestinal microbial balance, growth, nutrition, health status and resistance against infectious agents are already established (Gatesoupe, 1999; Ljubojević et al., 2013). Although, some reports are available on the activation of the immune response by dietary supplementation with probiotics in fish (Aly et al., 2008) their direct involvement in the immune response is not well established (Geijtenbeek and Gringhuis, 2009). Nayak et al. (2007) determined the effect of dietary supplementation with the probiotics bacterium, *Bacillus subtilis*, vitamin C in the form of ascorbyl polyphosphate and their combination on the immune response and disease resistance of Indian major carps. The non-specific immune system can be stimulated by probiotics (Balcázar et al., 2006). In the study conducted by Kumar et al. (2006), an increased growth rate was observed in *L. rohita* (Indian carp) fed diet containing *B. subtilis* compared with control. Furthermore, the survival rate after challenge with *A. hydrophila* was significantly higher in the treatment group compared with the control. Administration of probiotics enhances the survival of carp infected with *A. hydrophila* (Selvaraj et al., 2005). The high rates of establishment of bacterium in the gastro-intestinal tract of fish treated with *B. subtilis* have suppressed the *A. hydrophila* infection, which ultimately resulted in the higher survival (Kumar et al., 2006).

Beneficial effects of probiotics on the environment

Probiotics could help in improvement of the water quality in fish ponds (Panigrahi and Azad, 2007). This is due to the capability of the probiotic bacteria to participate in the turnover of organic nutrients in the ponds, to reduce ammonia, nitrite and total nitrogen levels in water and thus improve the chemical quality of water (Zhang et al., 2013).

Proper selection of probiotics in cyprinides production

For proper selection of probiotics it is essential to understand the mechanisms of their action and to define selection criteria for potential probiotics (Klaenhammer and Kullen,

1999). According to Sahu et al. (2008) methods to select probiotic bacteria for use in the aquaculture should include the following steps: collection of background information, acquisition of putative probiotics, screening of putative probiotics, evaluation of pathogenicity and survival test, *in vivo* evaluation, and evaluation of their effects in rearing conditions.

Future perspectives

Generally, the study of probiotics in fish production of Serbia is still in its early stage, and there have been no commercial probiotics products produced in Serbia so far. In the future, probiotics will gain more popularity in aquaculture of the Republic of Serbia, and the application areas will be expanded. Thus, quality control of probiotics in aquaculture of the Republic of Serbia will become an important issue. The need for the development of adequate technology for the evaluation of the efficiency of microbial agents as probiotics in aquaculture will be also increased.

Acknowledgements

This paper is a result of the research within the project TP 31011 “The influence of the quality of the food components for cyprinid fish species on the quality of meat, losses and the profitability of production”, financed by the Ministry of Science and Technological Development, the Republic of Serbia.

References

1. Aly SM, Abdel-Galil Ahmed Y, Abdel-Aziz Ghareeb A and Mohamed MF2008. Studies on *Bacillus subtilis* and *Lactobacillus acidophilus*, as potential probiotics, on the immune response and resistance of Tilapia nilotica (*Oreochromis niloticus*) to challenge infections. *Fish & Shellfish Immunology* 25, 128-136.
2. Azaza MS, Dhraief MN and Kraiem MM 2008. Effects of water temperature on growth and sex ratio of juvenile Nile tilapia *Oreochromis niloticus*(Linnaeus) reared in geothermal waters in southern Tunisia. *Journal of thermal Biology* 33, 98-105.
3. Balcázar JL, BlasID, Ruiz-Zarzuola I, Cunningham D, Vendrell D and Múzquiz JL 2006. The role of probiotics in aquaculture. *Veterinary microbiology* 114, 173-186.
4. Deane EE and Woo NY 2009. Modulation of fish growth hormone levels by salinity, temperature, pollutants and aquaculture related stress: a review. *Reviews in fish biology and fisheries* 19, 97-120.
5. DeVrese M and Marteau P 2007. Probiotics and Prebiotics: Effects on Diarrhea. *Journal of Nutrition* 137, 803-811.
6. FAO/WHO 2001. Report on Joint FAO/WHO Expert Consultation on Evaluation of Health and Nutritional Properties of Probiotics in Food Including Powder Milk with Live Lactic Acid Bacteria. 1–4 October 2001, Cordoba, Argentina. Available at: ftp://ftp.fao.org/es/esn/food/probio_report_en.pdf.
7. Faramarzi M, Kiaalvandi S and Iranshahi F 2011. The effect of probiotics on growth performance and body composition of common carp (*Cyprinus carpio*). *Journal of Animal and Veterinary Advances* 10, 2408-2413.
8. Fuller R1987. A review, probiotics in man and animals. *Journal of Applied Bacteriology* 66, 365–378.

9. Gatesoupe FJ 1999. The use of probiotics in aquaculture. *Aquaculture* 180, 147–165.
10. Gatesoupe FJ 2007. Live yeasts in the gut: natural occurrence, dietary introduction, and their effects on fish health and development. *Aquaculture* 267, 20–30.
11. Geijtenbeek TB and Gringhuis SI 2009. Signalling through C-type lectin receptors: shaping immune responses. *Nature Reviews Immunology* 9, 465–479.
12. Irianto A and Austin B 2002. Probiotics in aquaculture. *Journal of Fish Diseases* 25, 633–642.
13. Jovanović R, Mišćević M, Ćirković M, Milošević N, Ljubojević D 2011. Effect of probiotic to the production of one month old carp. 7. Međunarodni gospodarsko – znanstveni skup o ribarstvu; Hrvatsko ribarstvo na pragu EU, Riba kao funkcionalna hrana, Croatia, Vukovar, pp 7-11.
14. Klaenhammer TR and Kullen MJ 1999. Selection and design of probiotics. *International journal of food microbiology* 50, 45–57.
15. Kozasa M 1986. Toyocerin (*Bacillus toyoi*) as growth promoter for animal feeding. *Microbiolial Alimentary Nutrition* 4, 121–135.
16. Kumar R, Mukherjee SC, Prasad KP and Pal AK 2006. Evaluation of *Bacillus subtilis* as a probiotic to Indian major carp *Labeo rohita* (Ham.). *Aquaculture Research* 37, 1215–1221.
17. Lall SP and Lewis-McCrea LM 2007. Role of nutrients in skeletal metabolism and pathology in fish—an overview. *Aquaculture* 267, 3–19.
18. Lara-Flores M, Olivera-Castillo L and Olvera-Novoa MA 2010. Effect of the inclusion of a bacterial mix (*Streptococcus faecium* and *Lactobacillus acidophilus*), and the yeast (*Saccharomyces cerevisiae*) on growth, feed utilization and intestinal enzymatic activity of Nile tilapia (*Oreochromis niloticus*). *International Journal of Fisheries and Aquaculture* 2, 93–101.
19. Lilly DM and Stillwell RH 1965. Probiotics: growth promoting factors produced by microorganisms. *Science* 147, 747–748.
20. Ljubojević D, Ćirković M and Mišćević M 2013. Probiotics in carp fish nutrition. *Veterinarski glasnik* 67, 429–439.
21. Mišćević M, Ćirković M, Jovanović R, Ljubojević D, Mašić Z, Novakov N and Marković M 2011. Effect of probiotics to the production of one-year old carp. 22nd International Symposium Food safety production, Bosnia and Hercegovina, Trebinje, pp 50–52.
22. Mišćević M, Ćirković M, Jovanović R, Ljubojević D, Novakov N, Mašić Z and Marković M 2012. Effect of probiotics to the production of one-year old tench and common carp. *Archiva Zootechnica* 15, 29–36.
23. Nayak SK, Swain P and Mukherjee SC 2007. Effect of dietary supplementation of probiotic and vitamin C on the immune response of Indian major carp, *Labeo rohita* (Ham.). *Fish & shellfish immunology* 23, 892–896.
24. Panigrahi A and Azad IS 2007. Microbial intervention for better fish health in aquaculture: the Indian scenario. *Fish physiology and biochemistry* 33, 429–440.
25. Sahandi J, Jafaryan H, Roozbehfar R, Babae S, Dehestani M 2012. The use of two enrichment forms (*Brachionus plicatilis* enrichment and rearing water enrichment) with probiotic bacilli spore on growth and survival of Silver carp (*Hypophthalmichthys molitrix*). *Iranian Journal of Veterinary Research* 13, 289–295.
26. Sahu MK, Swarnakumar NS, Sivakumar K, Thangaradjou T and Kannan L 2008. Probiotics in aquaculture: importance and future perspectives. *Indian journal of microbiology* 48, 299–308.

27. Salinas I, Abelli L, Bertoni F, Picchiatti S, Roque A, Furones D and Esteban MÁ 2008. Monospecies and multispecies probiotic formulations produce different systemic and local immunostimulatory effects in the gilthead seabream (*Sparus aurata*L.). *Fish & shellfish immunology* 25, 114-123.
28. Selvaraj V, Sampath K and Sekar V 2005. Administration of yeast glucan enhances survival and some non-specific and specific immune parameters in carp (*Cyprinus carpio*) infected with *Aeromonas hydrophila*. *Fish & shellfish immunology* 19, 293-306.
29. Suzer C, Çoban D, Kamaci HO, Saka Ş, Firat K, Otgucuoğlu Ö and Küçüksarı H 2008. *Lactobacillus* spp. bacteria as probiotics in gilthead sea bream (*Sparus aurata*, L.) larvae: Effects on growth performance and digestive enzyme activities. *Aquaculture* 280, 140–145.
30. Tewe OO, Losel DM and Abu OA 1999. Solid state fermentation of sweet potato using two monoculture fungi: Changes in protei fatty acids and mineral composition. *Tropical Journal of Animal Science* 5, 219-224.
31. Wang Y and Xu Z 2006. Effect of probiotics for common carp (*Cyprinus carpio*) based on growth performance and digestive enzyme activities. *Animal Feed Science and Technology* 127, 283–92.
32. Wang YB, Li JR and Lin J 2008. Probiotics in aquaculture: challenges and outlook. *Aquaculture* 281, 1-4.
33. Zhang X, Fu L, Deng B, Liang Q, Zheng J, Sun J and Li W 2013. *Bacillus subtilis* SC02 supplementation causes alterations of the microbial diversity in grass carp water. *World Journal of Microbiology and Biotechnology* 29, 1645-1653.