



**The Effectiveness of a Psychoeducation Intervention
Delivered Via WhatsApp for Mothers of Children with Autism
Spectrum Disorder ASD in The Kingdom of Saudi Arabia: A
randomized controlled trial**

Journal:	<i>Child: Care, Health & Development</i>
Manuscript ID	CCH-2016-0428.R4
Manuscript Type:	Research Article
Keywords:	intervention, ASD, Autism Spectrum Disorder, Training, Mother, Well-Being

SCHOLARONE™
Manuscripts

Copy

The effectiveness of ASD parental intervention

The Effectiveness of a Psychoeducation Intervention Delivered Via WhatsApp for Mothers of Children with Autism Spectrum Disorder ASD in The Kingdom of Saudi Arabia: A randomized controlled trial

Abstract

Background: Mothers of children with autism spectrum disorder (ASD) report high levels of stress and lower levels of well-being than parents of typically developing children. Current interventions for ASD typically focus on working with the child rather than delivering strategies to help support parents.

Objective: To evaluate the effectiveness of a psychoeducation intervention developed to support mothers of children with ASD in Saudi Arabia.

Method: Sixty two mothers (23-52 years) of children (26-78 months) were recruited to a multi-site RCTs of the intervention. The intervention consisted of one face-to-face session (60 minutes) and four virtual sessions (30 minutes each) delivered using WhatsApp.

Parenting stress was the primary outcome, with secondary outcomes focusing on maternal depression, anxiety, and happiness, and child behaviour problems and ASD symptoms. Data was collected at baseline T1, immediately post intervention T2 and 8 week follow-up T3.

Results: One way ANCOVA was used at T2 and T3 with T1 scores entered as a covariate. Improvements were found at T2 for stress ($F=234.34$, $P=0.00$, $d=-1.52$), and depression ($F=195.70$, $P=0.00$, $d=-2.14$) but not anxiety and these results were maintained at T3. Changes in child behaviour problems were limited to improvements in hyperactivity at T2

1 The effectiveness of ASD parental intervention
2
3

4 (F=133.66, P=0.00, d= -1.54). Although changes in stress and depression were statistically
5 significant, change to clinically normal levels was limited to depression. None of the
6 participants had recovered after the intervention (PSI-SF stress scores), while 23 mothers
7 (71.87%) in the intervention group had recovered at T2 and 22 (68.75%) at T3 (HADS
8 depression scores).
9
10
11
12
13
14
15
16
17
18
19

20 **Conclusion:** This intervention with WhatsApp support is beneficial but may need to be
21 augmented with other forms of support for mothers of children with ASD including more
22 condensed sessions on stress and interventions targeting anxiety.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Introduction

50
51
52
53
54

55 Autism Spectrum Disorder (ASD) is a neurodevelopmental disorder associated with
56 'persistent impairment in reciprocal social communication and social interaction, and
57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 restricted, repetitive patterns of behaviour, interest, or activities' with a worldwide prevalence
5
6 of around 1% (American Psychiatric Association 2013, p.53); and 0.6% in the Kingdom of
7
8 Saudi Arabia (KSA) (Al Jarallah *et al.* 2005). Research shows that parents' of child with
9
10 Autism experience high levels of stress after the diagnosis of their children (Gupta 2007,
11
12 Meirsschaut *et al.* 2010); higher than stress levels in parents of children with other disabilities
13
14 (Phetrasuwan and Miles 2009, Estes *et al.* 2009). Research also shows that parents of
15
16 children with ASD suffer from higher levels of depression (Benson and Karlof 2009, Carter
17
18 *et al.* 2009). Mothers usually experience higher levels of stress than other members of the
19
20 family (Ingersoll and Dvortcsak 2006a). Most evidence suggests a bi-directional relationship
21
22 between parenting stress and childhood behaviour problems (Osborne and Reed 2009,
23
24 Sofronoff and Farbotko 2002, Jones *et al.* 2014). This suggests that mothers' who experience
25
26 high levels of stress require strategies to manage their own stress levels, the additional stress
27
28 that may arise from behaviour problems in their child, as well as strategies to address
29
30 behaviour problems in the child that may manifest because of their own stress. Evidenced
31
32 also suggests that parent-focused interventions using cognitive behaviour techniques
33
34 elements (e.g., problem solving and cognitive restructuring) were found to be effective in
35
36 reducing maternal stress in parents of children with intellectual disabilities (Hastings and
37
38 Beck 2004)
39
40
41
42
43
44

45 The majority of parenting interventions have been designed to teach parents to apply
46
47 specific strategies focused on supporting the child (Ingersoll and Dvortcsak 2006b, Stiebel
48
49 1999). Only a few studies have evaluated the efficacy of parent training interventions, where
50
51 interventions were designed mainly for the benefit of parents, focusing on parental well-being
52
53 for parents of children with ASD (Bendixen *et al.* 2011, Keen *et al.* 2010, Ferraioli and Harris
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 2013). However, the majority of these studies contain very small sample sizes and are often
5
6 not adequately controlled (Chiang 2014, Samadi *et al.* 2013).
7
8

9
10 In addition, access to support for parents of children with ASD is often limited in
11
12 Western countries due to barriers including insurance cover, parents' time constraints, and
13
14 distance from major cities so self-help intervention might be promising (Karst and Van
15
16 Hecke 2012). However, self-help interventions with minimal therapists' support for parents
17
18 of children with ASD appear to have limited evidence to date due to small sample size, lack
19
20 of adequate control, and poor tailoring of intervention strategies to underlying parental needs
21
22 (Ingersoll *et al.* 2016). Almansour *et al.* (2013) acknowledged the shortage of available and
23
24 affordable treatment options and training services for parents of children with ASD in KSA.
25
26 Mothers of children with ASD in KSA face additional restrictions accessing parenting
27
28 training including for example limited available transportation (e.g., buses and metros).
29
30 Despite increasing access to technology, researchers in mental health are only slowly
31
32 adopting the opportunities it brings to support the delivery of appropriate treatments (Hollis
33
34 *et al.* 2015). The goal of this study was to evaluate the efficacy of a self-help
35
36 Psychoeducation intervention with minimal therapists' support and delivered via WhatsApp.
37
38
39

40 **Hypotheses**

- 41
42
43 • Mothers in the intervention group (IG) will report lower stress than mothers in
44
45 the care as usual (CAU) group.
- 46
47
48 • Mothers in the IG will report lower depression, anxiety, and child behaviour
49
50 problems than mothers in the CAU group.
51
52

53 **Methods**

54 55 56 57 **Trial design**

1 The effectiveness of ASD parental intervention
2
3

4 The design was a pragmatic randomized two-arm controlled trial with two conditions
5 (IG versus CAU) and three time points (pre- intervention (T1), post-intervention (T2), and an
6 eight week follow-up (T3)).
7
8
9

10 11 12 13 **Participants**

14 Mothers of children with ASD were recruited from autism organizations at KSA.
15
16 Inclusion criteria for the study were i) mothers above the age of 18, living with their child
17 with ASD, ii) Mothers capable of reading and writing in Arabic, iii) Mothers who had smart
18 phones with a WhatsApp application
19
20
21
22
23
24

25 26 **Sample size.**

27 Based on the large effects sizes reductions in parenting stress reported by Ferraioli et al
28 (2013), an alpha level of 0.05 and an analysis strategy focused on One Way ANCOVA with
29 T1 scores entered as a covariate, a provisional estimate of power based on Cohen (1992)
30 suggested a minimum of 26 participant per trial arm.
31
32
33
34
35
36
37
38

39 This study received ethics approval from The University of X, School of Medicine
40 Ethics Committee and the committee at X in KSA. Consent forms were signed by all
41 participants.
42
43
44

45 46 **Randomisation/Blinding**

47 Simple randomisation based on the odd and even method yielded blinded envelopes to
48 inform therapist of allocation to IG or CAU. The researchers were blind to allocation
49
50
51
52
53
54

55 56 **Procedure**

57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 Study information packs were distributed by the organisations to all eligible mothers with
5 children with an ASD diagnosis from February 29th 2015 until March 30th 2015. Mothers
6 who consented to participate were randomly allocated to trial arms. Copies of the training
7 manual were distributed to mothers in the IG before the start of the first face-to-face session.
8
9 Sessions 2 – 5 consisted of 30 minutes therapist support via WhatsApp. Mothers in the CAU
10 group received advice about their child’s educational and behavioural problems from the
11 organizations. Three therapists assisted in delivering the intervention. One has a master’s
12 degree in psychology and four years’ experience working with children with ASD and their
13 mothers in an autism organisation. Two had a minimum of three years’ of experience
14 working with children with ASD, one with a graduate certificate in special education and the
15 other a certified clinical psychologist.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **Intervention**

33
34
35 Psychological models of stress and coping, including the transactional model of stress
36 developed by Lazarus and Launier (1978) and the Double ABCX Model developed by
37 McCubbin and Patterson (1983), have contributed to the development of the intervention.
38
39 One session was devoted to stress reduction, with the aim of reducing maternal stress in
40 participants by changing the way mothers appraise stressful situation and their children’s
41 behaviour. The intervention targeted different stages of the Double ABCX Model (Pickard
42 and Ingersoll 2016): devoting session one to inform mothers about the aetiology of ASD (aA
43 component refers to any stressor); session two to target stress in mothers and how can they
44 approach different stressful situations (Component cC refers to appraisal of a stressful
45 situation); session three to child behaviour problems to help the mothers cope with the initial
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 stressor of having a child with ASD; and session five to inform mothers about the available
5
6 resources in KSA (Component bB refers to resources).
7
8
9

10
11 Intervention materials were developed in the format of a booklet divided into five
12 sections (Table 1). The intervention was developed as a guided self-help (GSH) intervention
13 in line with the main principles and recommendations of NHS *Good Practice Guidance on*
14 *the use of self-help materials within Increasing Access to Psychological Therapies IAPT*
15 *services (Baguley et al. 2010)*. WhatsApp is a free mobile messaging application in which
16 people can exchange private or group messages as well as visual and audio media (How it
17 Works 2016).
18
19
20
21
22
23
24
25
26

27 **Measures**

28
29
30 **Parents' demographic questionnaire.** Demographic questions collected information about
31 participants including mothers' age, child with ASD age, number of siblings with ASD and
32 their ages, marital status, mothers' education and occupation, and fathers' occupation.
33
34
35
36

37
38 Arabic standardized versions of the following measures were administered to
39 participants in this study as it is the official and mother language for people in KSA.
40
41
42

43 **Parent Stress Index Short Form (PSI-SF).** The PSI-SF is a self-reported questionnaire that
44 aims to measure the stress associated with parenting (Abidin 1995). It includes 36 items
45 divided into three subscales, each contains 12 items: Parental Distress (PD); Parent–Child
46 Dysfunctional Interaction (PCDI) and Difficult Child (DC). The later measures parents'
47 perceptions of their children's temperament and compliance (Abidin 1995). The score of
48 most items range between one (strongly disagree) and five (strongly agree). Raw total scores
49 above 90 on the total score indicates clinically significant high level of stress.
50
51
52
53
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention
2
3

4 **Hospital Anxiety and Depression scale (HADS).** This measure is used to determine the
5 level of anxiety and depression (Zigmond and Snaith 1983) . It has 14 items, seven to
6 measure anxiety and seven for measuring depression. Each item has a choice of four
7 responses that are scored from 0 to 3 yielding a maximum score of 21 for each subscale.
8 Scores of 11 or more for either subscale of depression or anxiety are considered to indicate a
9 significant “case” of psychiatric co-morbidity
10
11
12
13
14
15
16
17
18
19

20
21 **Strength and Difficulties Questionnaire (SDQ).** This is a self-report questionnaire to
22 identify behavioural problems in children (Goodman 1997) . The SDQ parents’ version of
23 children aged 4-12 was used. It includes 25 items divided into five subscales (five items each)
24 including emotional symptoms, conduct problems, hyperactivity/inattention, peer relationship
25 problems, and prosocial behaviour. Parents are asked to indicate on a 3-point response scale
26 the extent to which their child shows symptoms of behavioural difficulties (1 = not true, 2 =
27 somewhat true, 3 = certainly true), and the total SDQ score is based on the sum of all the
28 subscales scores except for the pro-social one
29
30
31
32
33
34
35
36
37
38
39

40 **The Indian Scale for assessment of Autism (ISAA).** This scale was developed by the
41 National Institute of Mentally Handicapped for measuring the severity of autism. It is based
42 on Childhood Autism Rating Scale CARS by Schopler *et al.* (1980). It has 40 items rated on
43 a five-point scale (rarely, sometimes, frequently, mostly, and always)(National Institute for
44 the Mentally Handicapped 2009). According to the ISSA manual, autism is defined by a
45 score of 70. Total scores of 70 to 106 indicates mild autism, 107-153 moderate autism, and
46 scores of 153 and above indicate severe autism
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention
2
3

4 **The Arabic Scale of Happiness (ASH).** This is a measure developed originally in Arabic
5 language then translated to English (Abdel-Khalek 2013). It was designed to measure
6 happiness in adults. It has 20 items (five items are only fillers that do not contribute to
7 scoring). Each item could be answered on a 5-point intensity scale. The total score ranges
8 from 15 to 75.
9
10
11
12
13

14 **Intervention implementation fidelity, Participants' compliance, and Satisfaction**

15
16
17
18

19 After each session, the therapists sent questions asking participants to report the time
20 spent reading the intervention manual, and to rate the usefulness of the training session on a
21 scale from zero (unhelpful) to ten in (very helpful). They also kept records of attendance and
22 were given *a Therapist's scripts* to remind them to go through the same essential topics.
23
24
25
26
27
28

29 **Analysis strategy**

30
31

32 Baseline equivalence between groups was analysed using a series of one-way
33 ANOVA for continuous data and chi-square tests for discrete variables. Differences between
34 groups at T2 and T3 were assessed using a series of one way ANCOVAS with T1 scores
35 entered as covariates. Clinical significance was investigated using the two criteria: being
36 closer to the mean of the normal population and the Reliable Change Index RCI (Jacobson
37 and Truax 1991). Effect sizes were calculated based on change scores an interpreted based on
38 the traditional conventions of effect size proposed by Cohen (1988).
39
40
41
42
43
44
45
46
47

48 A visual inspection of the data after T1 data collection revealed five cases that were
49 not suitable for inclusion in the study as they scored within the normal range (>70) in ASD
50 symptoms on ISSA. These five cases were withdrawn from the analysis because of the
51 uncertainty of whether or not they met the inclusion criteria.
52
53
54
55
56
57

58 **Results**

59
60

The effectiveness of ASD parental intervention

Participants' characteristics and flow

One hundred and thirty eight mothers were referred into the study from organizations and baseline data were collected from 67 mothers who consented, who then were randomised to either an IG or CAU. It was not possible to explore whether mothers who consented were different from mothers who did not consent. See Figure 1 for the flow of participants through the trial. The demographic characteristics are presented in Table 2. Children with ASD were all at preschool age. Their mothers were predominantly in their thirties. Eight percent of mothers had a second child with autism. The majority of mothers in the sample were highly educated and much more educated than the average Saudi female (The Saudi Central Department of Statistics and Information, 2013). The majority of the mothers were married, with the sample containing less divorced women than might be expected given the Saudi average divorce rate (The Saudi Central Department of Statistics and Information, 2013). A minority of the mothers in the sample were working in professions that are typical occupations for Saudi women. All fathers of children in the study were working. Fathers worked in a range of professions, with a large proportion of soldiers and policeman.

Examination of differences at baseline between trial arms indicated no significant differences.

Parental outcomes

The means and SDs for each parental outcome measure at T1 and T2 are presented in Table 3. There was a significant effect of treatment on PSI-SF. An examination of the corresponding effect sizes calculated using Cohen *d* revealed the differences at T2 to be very large but was reduced at T3, to a large effect (Table 4). All three PSI-SF subscales demonstrated a significant effect of intervention at T2, and also at T3. An examination of the

The effectiveness of ASD parental intervention

1
2
3
4 corresponding effect sizes at T2 (Table 3) exhibited a very large effect size difference for the
5
6 PCDI subscale, a large effect size for the parental distress subscale and a small effect size for
7
8 the difficult child subscale in favour of intervention. At T3 the magnitude of effect sizes in
9
10 favour of intervention were attenuated for all three subscales, although the PCDI subscale
11
12 remain very large ($d = -1.51$) and the difficult child subscale remain small ($d = -0.28$) (but the
13
14 parental distress scale reduced from a large to a small effect size ($d = -0.33$). There was also a
15
16 significant effect of treatment group on HADS depression at T2 and T3 but not HADS
17
18 anxiety (Table 3). An examination of the corresponding effect sizes revealed the differences
19
20 at T2, and T3 to be very large and therefore meaningful (Table3). There was no meaningful
21
22 effect sizes differences at T2 and T3 for HADS anxiety which corroborated the lack of
23
24 statistical significance between the groups. There was also a significant effect of treatment
25
26 group on the Arabic happiness scale at T2 and T3 although the corresponding effect sizes
27
28 were small.
29
30
31
32

33 34 **Child outcomes**

35
36
37 There was a significant effect of treatment group on the Indian Scale for Autism
38
39 Assessment at T2 and T3 although the corresponding effect sizes were small There was a
40
41 significant effect of treatment group on the SDQ total at T2 (and T3 and the hyperactivity
42
43 scale at T2 and T3 (. However, no significant effect was shown for emotions, conduct, and
44
45 prosocial subscales either at T2 or T3. A close examination of the corresponding effect sizes
46
47 for SDQ revealed a large effect size at T2 and T3 for SDQ total and a very large effect size
48
49 for the hyperactivity subscale at T2) and T3. Not surprisingly, there was no effect on SDQ
50
51 emotions, conduct, and prosocial subscales at T2 and T3, which mirrors the lack of statistical
52
53 significance from the ANCOVA.
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 An exploration of the relationship between participants' implementation fidelity and
5
6 change in outcomes between T1 and T3 revealed two marginally significant effects for total
7
8 number of minutes spent reading the manual and HADS depression and happiness).
9

10 11 12 **Clinical Significance**

13
14 Clinical significant change was examined in this study for the significant parental
15
16 outcomes of PSI-SF and HADS depression. The number and percentage of participants who
17
18 experienced clinical significance are displayed in table 5. For a participant to be considered
19
20 as recovered after the course of an intervention, he or / she should achieve clinically
21
22 significant change on both criteria: becoming close to the functional rather than dysfunctional
23
24 population; and experiencing RC in their scores on the same measure (Jacobson, Roberts,
25
26 Berns, & McGlinchey, 1999). A thorough inspection of the data in this study showed that no
27
28 participant in this sample had recovered after the intervention (based on their PSI-SF scores).
29
30 Even though three mothers in the intervention group had experienced clinically significant
31
32 change on their PSI-SF scores by moving to the side of the normal population, these three
33
34 participants did not experience any reliable significant change. Twenty three mothers
35
36 (71.87%) in the intervention group had recovered at T2 and 22 (68.75%) at T3 based on their
37
38 HADS depression scores.
39
40
41
42
43
44
45

46 47 48 **Discussion**

49 This study adds to the evidence based supporting the effectiveness of parent training
50
51 interventions in enhancing parental well-being in mothers of children with ASD. Results of
52
53 this study indicated that the intervention was effective in reducing maternal stress,
54
55 depression, parent reported ASD symptoms, child behaviour problems, and increasing
56
57 maternal happiness in mothers of children with ASD in KSA.
58
59
60

The effectiveness of ASD parental intervention

1
2
3
4 Parental mean total stress score was significantly reduced in mothers in the IG and
5 maintained at follow-up. This is consistent with other studies (Bendixen *et al.* 2011, Chiang
6 2014, Ferraioli and Harris 2013). Mean scores for all PSI-SF subscales were significantly
7 reduced at T2 which supports Bendixen *et al.* (2011).
8
9
10
11
12

13
14 A significant reduction in maternal depression post intervention was maintained at
15 follow-up, a finding consistent with Bristol *et al.* (1993) Surprisingly, there was no reduction
16 in maternal anxiety which contradicts the assertion that depression and anxiety scores are
17 usually correlated (Almansour *et al.* 2013, Lee 2009 b). This might be due to the low internal
18 consistency of the HADS anxiety or that the proposed intervention was not sufficient to
19 reduce anxiety in mothers of children with ASD, as anxiety reduction was not a specific
20 intervention target. Alternatively the positive affect of how mothers enjoy their environment
21 in this particular sample which responded to intervention, might be different to the negative
22 affect of depression which usually correlates with anxiety Watson *et al.* (1988). This
23 assertion would fit with longitudinal data from Barker *et al.* (2011) showing a lack of
24 association between anxiety and depression scores over time on mothers of children with
25 ASD.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 Parental reported ASD symptoms were significantly reduced at T2 and these results
42 were maintained at follow-up. The theoretical explanation for why mothers reported reduced
43 ASD symptoms post intervention is not clear, as strategies to directly target ASD were not
44 included in the intervention. The complex relationship between parental well-being and child
45 behaviour and the putative role of potential mediating variables was beyond the scope and
46 statistical power of the current study. Child behaviour problems were also found to
47 significantly decrease at T2, a finding that is consistent with the results of other studies
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 evaluating interventions for parents of children with ASD (Tellegen and Sanders 2014,
5
6 Whittingham *et al.* 2009).
7

8
9 Although changes in stress and depression were statistically significant, change to
10 clinically normal levels was limited to depression. None of the participants had recovered
11 after the intervention (PSI-SF stress scores), while 23 mothers in the IG had recovered at T2
12 and 22 at T3 (HADS depression scores). However, Jacobson and Truax (1991) noted that the
13 aim for participants to be recovered after a course of an intervention might be too strict for
14 people with pervasive and serious conditions. Therefore, as the participants of this study were
15 mothers of children with ASD, it was not peculiar that they did not fully recover at the end of
16 training.
17
18
19
20
21
22
23
24
25
26
27
28

29 **Strengths**

30
31 This study has many strong points. This study used a RCT design to evaluate the efficacy of
32 the intervention, which is considered to be the golden design for programme evaluation
33 (Akobeng 2005). The CONSORT statement guidelines for reporting RCTs were used in this
34 study, which ensured a clear and comprehensive reporting of RCTs. Moreover, a follow-up
35 data collection were used in this study to test the stability of findings overtime.
36
37
38
39
40
41
42
43

44 **Limitations**

45
46 The lack of objective measures of maternal well-being and child behaviour problems
47 was a limitation in this study, however, the aim of the intervention was to evaluate the effect
48 of the intervention on maternal well-being and mothers remain the best informants about their
49 own well-being. Another limitation of this study is that it did not examine moderation due to
50 the limited sample size. The design of this study did not allow an exploration of the
51 differences in outcomes if the intervention was delivered face to face compared to WhatsApp
52
53
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 (dismantling design). This study did not provide any information about health economics,
5
6 even though the intervention was brief, self-administered, and delivered via WhatsApp,
7
8 which could likely to be cost effective.
9

10
11 The low value of alpha in some of the measures in this study (HADS anxiety $\alpha=0.55$, conduct
12
13 subscale of SDQ $\alpha=0.49$) could be cultural factors and these measures may not be entirely
14
15 suitable for non-western populations. The Peer sub-scale in SDQ was removed from the
16
17 analysis as it showed a very low internal consistency alpha value of 0.19.
18
19

20 21 22 23 **Key messages**

- 24
25
26 • As this was a small-scale study, it would be important in the future to replicate the
27
28 study in a larger sample, especially as we now have information to inform a full
29
30 power calculation.
31
- 32
33 • It will be important in future studies to explore the cost effectiveness of this
34
35 intervention as a greater understanding of costs and benefits of the intervention could
36
37 be used to build a business case for its implementation.
38
- 39
40 • It will be necessary to conduct a study on implementation, to be able to explore
41
42 additional barriers to the use of the intervention in routine clinical practice within
43
44 ASD organisations in KSA. Moreover, is to explore the relationship between mode of
45
46 delivery, intervention intensity, and outcome.
47
- 48
49 • A larger study powered for mediation and moderation would help inform clinical
50
51 servicers about why the intervention leads to change in outcomes and for whom the
52
53 intervention is best suited.
54
55
56
57
58
59
60

The effectiveness of ASD parental intervention

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review Copy

References

The effectiveness of ASD parental intervention

Abdel-Khalek, A. M. (2013) The Arabic Scale of Happiness (ASH): psychometric characteristics. *Comprehensive Psychology*, **2**, Article 5.

Abidin, R. R. (1995) *Parenting Stress Index: Professional Manual*, Psychological Assessment Resources, Odessa, FL.

Akobeng, A. K. (2005) Understanding randomised controlled trials. *Arch Dis Child*, **90**, 840-844.

Al-Shodokhi, Y. & Al-Jabr, A. (2010) *Al Daleel Al Shamel Le Khadmat Al Tawahod Bi Al Mamlakah Alarabia Alsaudia*

[Comprehensvie guide to Autism Spectrum Disorder services in the Kingdom of Saudi Arabia]. <https://slpemad.files.wordpress.com/2015/04/d8a7d984d8afd984d98ad984-d8a7d984d8b4d8a7d985d984-d984d8aed8afd985d8a7d8aa-d8b0d988d98a-d8a7d984d8aad988d8add8af-d8a8d8a7d984d8b3d8b9.pdf>

Al Jarallah, A., Al Waznah, T., Al Ansari, S. & Al Hazmi, M. (2005) Studying autism and related developmental disorders. King Saudi University, Riyadh.

Almansour, M. A., Alateeq, M. A., Alzahrani, M. K., Algeffari, M. A. & Alhomaidan, H. T. (2013) Depression and anxiety among parents and caregivers of autistic spectral disorder children. *Neurosciences Journal*, **18**, 58-63.

American Psychiatric Association (2013) *Diagnostic and Statistical Manual of Mental Disorders: Dsm-5*, Amer Psychiatric Pub Incorporated.

Apa, A. P. A. (2013) *Diagnostic and Statistical Manual of Mental Disorders: Dsm-5*, Amer Psychiatric Pub Incorporated.

Baguley, C., Farrand, P., Hope, R., Leibowitz, J., Lovell, K., Lucock, M., O'neill, C., Paxton, R., Pilling, S., Richards, D., Turpin, G., White, J. & Williams, C. (2010) Good

The effectiveness of ASD parental intervention

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- practice guidance on the use of self-help materials within IAPT services. In:
Improving Access to Psychological Therapy (ed. Nhs). Technical Report
- Barker, E. T., Hartley, S. L., Seltzer, M. M., Floyd, F. J., Greenberg, J. S. & Orsmond, G. I.
(2011) Trajectories of Emotional Well-Being in Mothers of Adolescents and Adults
with Autism. *Developmental Psychology*, **47**, 551-561.
- Bendixen, R. M., Elder, J. H., Donaldson, S., Kairalla, J. A., Valcante, G. & Ferdig, R. E.
(2011) Effects of a Father-Based In-Home Intervention on Perceived Stress and
Family Dynamics in Parents of Children With Autism. *American Journal of
Occupational Therapy*, **65**, 679-687.
- Benson, P. & Karlof, K. (2009) Anger, Stress Proliferation, and Depressed Mood Among
Parents of Children with ASD: A Longitudinal Replication. *Journal of Autism and
Developmental Disorders*, **39**, 350-362.
- Bristol, M. M., Gallagher, J. J. & Holt, K. D. (1993) Maternal depressive symptoms in
autism: Response to psychoeducational intervention. *Rehabilitation Psychology*, **38**,
3-10.
- Carter, A. S., Martinez-Pedraza, F. D. L. & Gray, S. a. O. (2009) Stability and Individual
Change in Depressive Symptoms Among Mothers Raising Young Children With
ASD: Maternal and Child Correlates. *Journal of Clinical Psychology*, **65**, 1270-1280.
- Chiang, H.-M. (2014) A Parent Education Program for Parents of Chinese American
Children With Autism Spectrum Disorders (ASDs) A Pilot Study. *Focus on Autism
and Other Developmental Disabilities*, **29**, 88-94.
- Cohen, J. (1988) *Statistical Power Analysis for the Behavioral Sciences*, L. Erlbaum
Associates.

The effectiveness of ASD parental intervention

Estes, A., Munson, J., Dawson, G., Koehler, E., Zhou, X.-H. & Abbott, R. (2009) Parenting stress and psychological functioning among mothers of preschool children with autism and developmental delay. *Autism*, **13**, 375-387.

Ferraioli, S. J. & Harris, S. L. (2013) Comparative Effects of Mindfulness and Skills-Based Parent Training Programs for Parents of Children with Autism: Feasibility and Preliminary Outcome Data. *Mindfulness*, **4**, 89-101.

Goodman, R. (1997) The Strengths and Difficulties Questionnaire: a research note. *Journal of child psychology and psychiatry, and allied disciplines* **38**, 581-586.

Gupta, V. B. (2007) Comparison of parenting stress in different developmental disabilities. *Journal of Developmental and Physical Disabilities*, **19**, 417-425.

Hastings, R. P. & Beck, A. (2004) Practitioner review: stress intervention for parents of children with intellectual disabilities. *Journal of Child Psychology and Psychiatry*, **45**, 1338-1349.

Hollis, C., Morriss, R., Martin, J., Amani, S., Cotton, R., Denis, M. & Lewis, S. (2015) Technological innovations in mental healthcare: harnessing the digital revolution. *The British journal of psychiatry : the journal of mental science* **206**, 263-265.

How It Works. (2016) In *WhatsApp*. Available at: <https://www.WhatsApp.com/>

Ingersoll, B. & Dvortcsak, A. (2006a) Including parent training in the early childhood special education curriculum for children with autism spectrum disorders. *Journal of Positive Behavior Interventions*, **8**, 79-87.

1 The effectiveness of ASD parental intervention
2
3

4 Ingersoll, B. & Dvortcsak, A. (2006b) Including Parent Training in the Early Childhood
5

6 Special Education Curriculum for Children With Autism Spectrum Disorders. *Topics*
7
8 *in Early Childhood Special Education*, **26**, 179-187.
9

10 Ingersoll, B., Wainer, A. L., Berger, N. I., Pickard, K. E. & Bonter, N. (2016) Comparison of
11

12 a Self-Directed and Therapist-Assisted Telehealth Parent-Mediated Intervention for
13
14 Children with ASD: A Pilot RCT. *Journal of Autism and Developmental Disorders*,
15
16 **46**, 2275-2284.
17

18 Jacobson, N. S. & Truax, P. (1991) Clinical significance: A statistical approach to defining
19

20 meaningful change in psychotherapy research. *J Consult Clin Psychol*, **59**, 12-19.
21

22 Jacobson, N. S., Roberts, L. J., Berns, S. B., & McGlinchey, J. B. (1999). Methods for
23

24 defining and determining the clinical significance of treatment effects: description,
25
26 application, and alternatives. *Journal of consulting and clinical psychology*
27

28
29
30
31
32 , *67*(3), 300-307.
33

34 Jones, L., Hastings, R. P., Totsika, V., Keane, L. & Rhule, N. (2014) Child behavior
35

36 problems and parental well-being in families of children with autism: the mediating
37
38 role of mindfulness and acceptance. *American journal on intellectual and*
39
40 *developmental disabilities*
41
42
43

44 **119**, 171-185.
45

46 Karst, J. S. & Van Hecke, A. V. (2012) Parent and Family Impact of Autism Spectrum
47

48 Disorders: A Review and Proposed Model for Intervention Evaluation. *Clinical Child*
49
50 *and Family Psychology Review*, **15**, 247-277.
51

52 Keen, D., Couzens, D., Muspratt, S. & Rodger, S. (2010) The effects of a parent-focused
53

54 intervention for children with a recent diagnosis of autism spectrum disorder on
55
56 parenting stress and competence. *Research in Autism Spectrum Disorders*, **4**, 229-241.
57
58
59
60

1 The effectiveness of ASD parental intervention
2
3

4 Lazarus, R. S. & Launier, R. (1978) Stress-Related Transactions between Person and
5 Environment. In: *Perspectives in Interactional Psychology* (eds. Pervin, L. A. &
6 Lewis, M.), pp. 287-327. Springer US, City.
7
8

9
10
11 Lee, G. K. (2009 b) Parents of Children with High Functioning Autism: How Well Do They
12 Cope and Adjust? *Journal of Developmental and Physical Disabilities*, **21**, 93-114.
13

14
15 Mccubbin, H. I. & Patterson, J. M. (1983) The Family Stress Process. *Marriage & Family*
16 *Review*, **6**, 7-37.
17

18
19
20 Meirsschaut, M., Roeyers, H. & Warreyn, P. (2010) Parenting in families with a child with
21 autism spectrum disorder and a typically developing child: Mothers' experiences and
22 cognitions. *Research in Autism Spectrum Disorders*, **4**, 661-669.
23

24
25
26 National Institute for the Mentally Handicapped. (2009) Indian Scale for the Assessment of
27 Autism (ISAA). From Government of India: Ministry of Social Justice and
28 Empowerment. Available at: [http://autism-india.com/autism-diagnosis/indian-scale-for-](http://autism-india.com/autism-diagnosis/indian-scale-for-assessment-of-autism-isaa/)
29 [assessment-of-autism-isaa/](http://autism-india.com/autism-diagnosis/indian-scale-for-assessment-of-autism-isaa/)
30
31

32
33
34
35
36 Osborne, L. A. & Reed, P. (2009) The Relationship Between Parenting Stress and Behavior
37 Problems of Children With Autistic Spectrum Disorders. *Exceptional Children*, **76**,
38 54-73.
39

40
41
42 Phetrasuwan, S. & Miles, M. S. (2009) Parenting Stress in Mothers of Children With Autism
43 Spectrum Disorders. *Journal for Specialists in Pediatric Nursing*, **14**, 157-165.
44

45
46
47 Pickard, K. E. & Ingersoll, B. R. (2016) Using the Double ABCX Model to Integrate
48 Services for Families of Children with ASD. *Journal of Child and Family Studies*, 1-
49 14.
50
51
52
53
54
55
56
57
58
59
60

1 The effectiveness of ASD parental intervention

2
3
4 Samadi, S. A., Mcconkey, R. & Kelly, G. (2013) Enhancing parental well-being and coping
5
6 through a family-centred short course for Iranian parents of children with an autism
7
8 spectrum disorder. *Autism*, **17**, 27-43.

9
10
11 Schopler, E., Reichler, R. J., Devellis, R. F. & Daly, K. (1980) Toward objective
12
13 classification of childhood autism: Childhood Autism Rating Scale (CARS). *J Autism*
14
15 *Dev Disord*, **10**, 91-103.

16
17
18 Simpson, R. L. & De Boer-Ott, S. R. (2005) *Autism spectrum disorders: interventions and*
19
20 *treatments for children and youth*, Corwin Press.

21
22 Sofronoff, K. & Farbotko, M. (2002) The Effectiveness of Parent Management Training to
23
24 Increase Self-Efficacy in Parents of Children with Asperger Syndrome. *Autism*, **6**,
25
26 271-286.

27
28
29 Stiebel, D. (1999) Promoting Augmentative Communication During Daily Routines: A
30
31 Parent Problem-Solving Intervention. *Journal of Positive Behavior Interventions*, **1**,
32
33 159-169.

34
35
36 Tellegen, C. L. & Sanders, M. R. (2014) A Randomized Controlled Trial Evaluating a Brief
37
38 Parenting Program With Children With Autism Spectrum Disorders. *J Consult Clin*
39
40 *Psychol*, **82**, 1193-1200.

41
42 The Saudi Central Department of Statistics and Information. (2013). *Latest statistical*
43
44 *proceeding*. Retrieved from <http://www.cdsi.gov.sa/>

45
46
47
48
49 Treacy, L., Tripp, G. & Baird, A. (2005) Parent stress management training for attention-
50
51 deficit/hyperactivity disorder. *Behavior Therapy*, **36**, 223-233.

The effectiveness of ASD parental intervention

Watson, D., Clark, L. A. & Carey, G. (1988) Positive and negative affectivity and their relation to anxiety and depressive disorders. *Journal of Abnormal Psychology*, **97**, 346-353.

Whittingham, K., Sofronoff, K., Sheffield, J. & Sanders, M. R. (2009) Stepping Stones Triple P: An RCT of a Parenting Program with Parents of a Child Diagnosed with an Autism Spectrum Disorder. *Journal of Abnormal Child Psychology*, **37**, 469-480.

Wright, B. & Williams, C. (2007) *Intervention and Support for Parents and Carers of Children and Young People on the Autism Spectrum: A Resource for Trainers*, Jessica Kingsley Publishers.

Zigmond, A. S. & Snaith, R. P. (1983) The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, **67**, 361-370.

Table 1: Summary of the intervention sessions

Session' topic	Summary	Reference
What is Autism?	Aetiology of autism, including its causes, how it affects the lives of people with ASD, and the formal DSM-5 diagnosis criteria.	(APA 2013, Simpson and de Boer-Ott 2005)
Stress	Definition of stress, examples, causes and tips to help reduce stress in parents.	(Treacy <i>et al.</i> 2005)
Managing behaviour	Explaining behaviour, how behaviour occurs and different ways that will help parents to manage their children's behaviour problems.	(Wright and Williams 2007)

The effectiveness of ASD parental intervention

Mood	Explaining mood, how can it affects the way people respond to different situations and how can different ways of thinking aggravate negative mood.	(L. Treacy, personal communication, October 10, 2015)
Resources for families	Listing the available resources to parents of children with ASD in KSA and contact information of ASD organisations and services in their regions.	(Al-Shodokhi and Al-Jabr 2010)

Review Copy

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

The effectiveness of ASD parental intervention

Table 2 : Demographic characteristics of study sample (n=62)

Characteristics	IG	CAU
	N = 32	N = 30
Child age (months), M(SD)	63.18 (13.68)	58.73 (14.07)
Mother's age (years), M(SD)	32.90 (7.26)	34.43 (6.65)
Mothers with another ASD child, n (%)	3 (9.37)	2 (6.66)
Age of another ASD child, M(SD)	52.00 (15.62)	71.00 (52.32)
Education level, n (%), *[% in SA population]		
Less than high school	5 (15.62)	5 (16.66)
High school	9 (28.12)	11 (36.66)
Bachelor's degree	18 (56.25)	14 (46.66)

1
2 The effectiveness of ASD parental intervention
3
4
5

6
7

Marital status, n (%), [% in SA population]

8
9 Married 29 (90.62) 27 (90.00)

10
11
12
13 Divorced 3 (9.37) 3 (10.00)
14
15
16
17
18
19

20
21 Working females, n (%), [% in SA population]
22

7 (21.87) 6 (20.00)

23
24
25
26
27
28
29
30
31
32
33 Teachers 6 (18.75) 4 (13.33)
34
35
36
37

38 Nurses 1 (3.12) 2 (6.66)
39
40

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

The effectiveness of ASD parental intervention

	Soldiers	9 (28.12)	11 (36.66)
	IT	3 (9.37)	3 (10.00)
	Teachers	6 (18.75)	4 (13.33)
Fathers' jobs, n (%)	Administrative work	11 (34.37)	8 (26.66)
	Business	1 (3.12)	2 (6.66)
	Physicians	2 (6.25)	2 (6.66)

*Only available information about Percentages in KSA population is presented (from Saudi central department of Statistics and Information based on 2010 census)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

The effectiveness of ASD parental intervention

HADS (anxiety)	7.71(2.45)	7.59(2.09)	7.63(2.77)	7.83(2.64)	3.76(0.057)	-0.12
Happiness	43.71(9.33)	47.50(8.56)	43.13(10.26)	43.46(9.80)	121.32(<0.001)	0.35
ISAA	100.50(13.60)	91.71(12.76)	101.20(14.16)	95.66(13.82)	21.85(<0.001)	-0.23
SDQ (revised)	14.40(4.44)	10.40(3.60)	14.83(4.37)	14.76(3.89)	100.63(<0.001)	-0.89
Emotional	4.50(2.09)	4.50(2.09)	4.50(2.31)	4.56(2.29)	.69(0.40)	-0.03
Hyperactivity	6.68(2.40)	2.84(1.95)	7.00(2.21)	6.70(2.26)	133.66(<0.001)	-1.54
Conduct	3.21(1.80)	3.21(1.80)	3.33(1.93)	3.43(1.85)	2.20(0.14)	-0.05
Prosocial	3.78(2.35)	3.78(2.35)	4.33(2.39)	4.60(2.12)	7.03(0.10)	-0.11

The effectiveness of ASD parental intervention

Table 4 : Means and SDs for outcomes at T1 and T3

Measure	Intervention		CAU		F (P value)	Cohen's d T1 to T3
	M (SD)		M (SD)			
	T1	T3	T1	T3		
PSI-SF	129.96(13.82)	116.50(12.46)	132.80(13.07)	132.56(12.86)	131.47(<0.001)	-0.98
Parental distress	43.84(6.74)	41.50(6.42)	45.53(6.24)	45.36(6.13)	64.39(<0.001)	-0.33
Parent child dysfunctional interaction	40.75(6.94)	30.96(5.32)	41.5(5.95)	41.50(5.78)	137.92(<0.001)	-1.51
Difficult child(revised)	41.28(4.21)	40.03(4.12)	41.90(4.23)	41.83(4.29)	28.95 (<0.001)	-0.28
HADS (depression)	10.68(3.56)	4.12(2.49)	10.66(3.47)	11.33(3.39)	183.58 (<0.001)	-2.05
HADS (anxiety)	7.71(2.45)	7.65(2.00)	7.63(2.77)	7.86(2.67)	1.69(0.19)	-0.11

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

The effectiveness of ASD parental intervention

Happiness	43.71(9.33)	47.62(8.47)	43.13(10.26)	43.56(9.63)	95.46 (<0.001)	0.35
ISAA	100.50(13.60)	91.68(12.72)	101.20(14.16)	95.66(13.84)	23.23 (<0.001)	-0.23
SDQ (revised)	14.40(4.44)	14.75(4.21)	14.83(4.37)	19.53(4.09)	41.80 (<0.001)	-0.98
Emotional	4.50(2.09)	4.56(2.03)	4.50(2.31)	4.63(2.25)	0.28(0.59)	-0.03
Hyperactivity	6.68(2.40)	3.09(1.85)	7.00(2.21)	6.80(2.34)	115.85 (<0.001)	-1.47
Conduct	3.21(1.80)	3.25(1.77)	3.33(1.93)	3.46(1.81)	1.14(0.28)	-0.05
Prosocial	3.78(2.35)	3.84(2.39)	4.33(2.39)	4.63(2.15)	3.63(0.06)	-0.10

Review Copy

The effectiveness of ASD parental intervention

Table 5: The number and percentage of reliably and clinically improved mothers for each significant measure for the IG and CAU

Measures	Reliably improved		Clinically improved		
	N (%)		N (%)		
	T2	T3	T2	T3	
PSI-SF	IG	12 (37.5)	11 (34.37)	3 (9.37)	3 (9.37)
	CAU	0 (0.00)	0 (0.00)	0 (0.00)	1 (3.33)
HADS-depression	IG	24 (75)	24 (75)	30 (93.75)	29 (90.62)
	CAU	0 (0.00)	0 (0.00)	3 (10)	3 (10)

The effectiveness of ASD parental intervention

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Flowchart diagram showing the flow of participants through each stage of the trial.

Review Copy




Figure 1: Flowchart diagram showing the flow of participants through each stage of the trial.

599x776mm (72 x 72 DPI)