

1 **Nutrient limitation or home field advantage: does microbial community adaptation**
2 **overcome nutrient limitation of litter decomposition in a tropical peatland?**

3

4 Jorge Hoyos-Santillan^{1*}, Barry H. Lomax¹, Benjamin L. Turner² and Sofie Sjögersten¹

5

6 ¹The University of Nottingham, School of Biosciences, Division of Agricultural and
7 Environmental Science, Room C-36 Gateway Building, Sutton Bonington Campus,
8 Loughborough, LE12 5RD, UK

9 ²Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Republic of
10 Panama

11

12 *Corresponding author: Jorge Hoyos-Santillan

13 e-mail: jhoyosantillan@hotmail.com

14 tel: +52 012292695268

15

16 Running headline: Nutrient limitation or home field advantage

17

18 **Summary**

19 **1.** Litter decomposition is an important control on carbon accumulation in tropical peatlands.

20 Stoichiometric theory suggests that decomposition is regulated by elemental ratios in litter while
21 the home field advantage hypothesis predicts that decomposer communities are adapted to local
22 conditions. To date, the relative importance of these contrasting theories for litter decomposition
23 and therefore the carbon balance of tropical peatlands remains poorly understood.

24 **2.** We conducted two *in situ* litter decomposition experiments in a lowland tropical peatland. The
25 first experiment tested the importance of the stoichiometric theory using a factorial nutrient
26 addition experiment at two sites with contrasting vegetation (*Raphia taedigera* and
27 *Camposperma panamensis*) to assess how nutrient addition affected microbial enzyme activity
28 and litter mass loss at the peat surface and at 50 cm depth. The second experiment tested the
29 importance of home field advantage by reciprocal translocation of leaf litter from *R. taedigera*
30 and *C. panamensis* forests, which differed in both litter chemistry and soil nutrient availability,
31 to separate the influence of litter chemistry and soil/site properties on litter mass loss.

32 **3.** The activities of hydrolytic enzymes involved in the decomposition of large plant polymers
33 were stimulated by nitrogen addition only where nitrogen availability was low relative to
34 phosphorus, and were stimulated by phosphorus addition where phosphorus availability was low.

35 **4.** The addition of nitrogen, but not phosphorus, increased leaf litter decomposition under
36 waterlogged conditions at 50 cm depth, but not at the peat surface.

37 **5.** Decomposition was greatest for autochthonous litter irrespective of site nutrient status,
38 indicating that adaptation of the microbial community to low nutrients can partly overcome
39 nutrient limitation, and suggesting that home field advantage can influence litter decomposition
40 rates.

41 **6.** Synthesis. Our study shows that leaf litter decomposition and the activity of microbial
42 enzymes in tropical peatlands are constrained in part by nutrient availability. However, such
43 nutrient limitation of litter decomposition can be overcome by adaptation of the microbial
44 community.

45

46 **Highlights:**

- 47 • Nitrogen and phosphorus stimulated activity of hydrolytic enzymes associated with
48 decomposition in agreement with stoichiometric theory.
- 49 • Nitrogen availability limited leaf litter decomposition under anoxic conditions,
50 suggesting environmental and litter chemistry controls of nutrient limitation.
- 51 • Litter decomposition was greatest at the site where the litter originated, irrespective of
52 site nutrient status, supporting home field advantage theory.

53

54

55

56

57

58

59

60 **Keywords:** carbon turnover, nitrogen, phosphorus, tropical peat, nutrient dynamics

61 **1. Introduction**

62 Decomposition rates of organic matter influence carbon storage and regulate nutrient availability
63 in natural ecosystems. Decomposition is carried out by complex groups of microorganisms and
64 the rate of decomposition is controlled by how the substrate properties, together with the abiotic
65 environment, meet the demands of the microbial communities (Kaiser *et al.* 2014). According to
66 stoichiometric theory, the balance of carbon (C), nitrogen (N) and phosphorus (P) that
67 decomposer organisms must maintain to regulate metabolic function and growth limits
68 decomposition rates when nutrient ratios in the substrate do not match demand by individual
69 microorganisms (Sterner & Elser 2002; Manzoni & Porporato 2009). If this holds true,
70 decomposition rates should not be limited by nutrient availability when the composition (with
71 regards to C, N and P) of the substrate (*e.g.* leaf litter) is similar to that of the decomposer
72 organisms. Indeed, greater nutrient availability enhances decomposition in a wide range of
73 ecosystems (*e.g.*, subarctic heaths, mangroves), supporting the notion of nutrient limitation of
74 decomposition (Quested *et al.* 2005; Cornwell *et al.* 2008; Keuskamp *et al.* 2015b). However, in
75 other instances nutrient addition has had limited effects on decomposition rates (*e.g.*
76 decomposition of low quality litter has been found to be energy rather than nutrient limited;
77 Knorr *et al.* 2005; Keuskamp *et al.* 2013). Furthermore, nutrient limitation of litter
78 decomposition fluctuates over time, reflecting changing nutrient demands of the decomposer
79 organisms as well as changes in litter chemistry as decomposition progresses (Kaiser *et al.*
80 2014).

81

82 Microorganisms can overcome resource limitation by up-regulating the production of
83 extracellular enzymes involved in C, N and P acquisition, depending on which nutrients are
84 limiting their growth (Sinsabaugh & Follstad Shah 2012). For example, low P availability
85 increases the activity of acid phosphatases in a range of soils (Olander & Vitousek 2000; Allison
86 *et al.* 2007; Sjögersten *et al.* 2011), while low nutrient availability can drive tight nutrient cycling
87 within microbial communities (Kaiser *et al.* 2014). Strong interactions between the composition
88 and functioning of the microbial community and the dominant litter inputs are one of the
89 explanations of the so called “home field advantage” (HFA), whereby the decomposer
90 community becomes adapted, or optimized, to degrade the litter at a given site (Austin *et al.*
91 2014). This results in faster litter decomposition rates when litter decomposes adjacent to the
92 plants that produced it (Vivanco & Austin 2008). This pattern is relatively weak, but holds true at
93 the global scale, with an approximately 8% greater mass loss when litter material was
94 decomposing at “home” (Ayres *et al.* 2009; Veen *et al.* 2015). However, the effects of the home
95 field advantage are more pronounced when sites differ considerably in soil nutrient availability
96 and plant species composition, suggesting lower degree of redundancy among decomposer
97 communities across locations with strongly contrasting soil and litter type characteristics (Veen
98 *et al.* 2015).

99 Lowland tropical peatlands have the fastest rates of peat accumulation in the world – up to 10
100 times faster than temperate, subarctic and boreal peatlands (Gorham, Janssens & Glaser 2003;
101 Chimner & Ewel 2005; Dommain, Couwenberg & Joosten 2011) – and contain 40-90 Gt of C
102 (Kurnianto *et al.* 2015). The functioning of tropical peatlands as a C store is currently under
103 threat as land use change, climate change and increasing levels of atmospheric N deposition
104 accelerate decomposition rates (Galloway *et al.* 2004; Bragazza *et al.* 2012; IPCC 2013). If

105 nutrient availability is a key limitation of decomposition in tropical peatlands, as has been found
106 at higher latitudes (Wang *et al.* 2014), then greater nutrient availability might reduce C storage.
107 In addition, nutrient availability shapes the species composition of peat swamp forests (Brady
108 1997; Page *et al.* 1999; Troxler 2007; Sjögersten *et al.* 2011) and hence the quality and the
109 quantity of litter inputs (Wright *et al.* 2013; Hoyos-Santillan *et al.* 2016), with implications for
110 the composition of the decomposer community (Troxler *et al.* 2012) and decomposition rates
111 (Yule & Gomez 2009; Hoyos-Santillan *et al.* 2015).

112 Litter decomposition of tropical peatland tree species varies among species (*e.g.*, between palms
113 and hardwoods) and tissue types (*e.g.*, between roots and leaves) (Yule & Gomez 2009; Hoyos-
114 Santillan *et al.* 2015). Furthermore, the degree of waterlogging and nutrient availability, as well
115 as microbial community composition, pH, and concentrations of dissolved oxygen and phenolic
116 compounds, vary within peat profiles (Freeman, Ostle & Kang 2001; Jackson, Liew & Yule
117 2009; Hoyos-Santillan *et al.* 2015, 2016). Therefore, decomposition rates of the same litter
118 material differs depending on its position within the peat profile (Hoyos-Santillan *et al.* 2015).

119 To test the importance of stoichiometric theory (Sterner & Elser 2002; Manzoni & Porporato
120 2009) and HFA (Austin *et al.* 2014) for litter decomposition, we carried out two experiments in a
121 tropical peatland in Panama. The first experiment was a factorial N and P addition experiment in
122 two contrasting forest types, a nutrient rich palm swamp and a relatively less nutrient rich mixed
123 forest (Sjögersten *et al.* 2011). This experiment tested the hypothesis that nutrient availability
124 controls (i) activities of extra cellular hydrolytic enzymes, which are involved in microbial
125 nutrient and carbon acquisition and (ii) litter decomposition. The experiment involved
126 decomposing different litter tissue types (leaves, roots and stems) at the peat surface and at 50
127 cm depth. We predicted that if the microbial community at a site was nutrient limited, nutrient

128 addition would reduce microbial C:N and C:P ratios and down-regulate enzymes involved in
129 nutrient acquisition, resulting in a subsequent up-regulation on enzymes involved in the
130 breakdown of sugars, hemi-cellulose and cellulose (Sinsabaugh & Follstad Shah 2012).
131 According to stoichiometric theory, we expected nutrient addition to accelerate litter mass loss at
132 the low nutrient mixed forest site, and that litter decomposition would be greatest for both litter
133 types at the more nutrient rich palm swamp site. We also predicted that litter with high C:N and
134 C:P ratios would be more responsive to nutrient addition with respect to mass loss, in agreement
135 with Baumann *et al.* (2009).

136 The second experiment was a reciprocal leaf litter translocation experiment between the two
137 contrasting forest types. This experiment tested the hypothesis that litter is decomposed faster at
138 “home” than “away” irrespective of site nutrient status (Kaiser *et al.* 2014; Austin *et al.* 2014).
139 For this experiment we carried out reciprocal transplants of leaf litter material from two different
140 trees species that were the dominant trees at two peatlands sites with contrasting nutrient status
141 (*i.e.* the same two sites that were used for the nutrient addition experiment). We predicted that
142 according to the HFA theory the “home” palm leaf litter would degrade more rapidly at the palm
143 site while the litter from the low nutrient mixed forest would degrade fastest at the mixed forest
144 site, *i.e.* its home location (Veen *et al.* 2015). This contrasts with our prediction above of greater
145 decomposition at the nutrient rich site following stoichiometric theory (*e.g.* Sterner & Elser
146 2002), allowing us to investigate the respective influences of HFA and stoichiometric theory on
147 C dynamics and, by extension, peat accumulation in tropical peatlands.

148

149 **2. Materials and methods**

150 2.1. Study sites

151 The study was conducted in the north-west Caribbean coast of Panama where several large
152 peatlands are located within the Bocas del Toro province (Phillips, Rouse & Bustin 1997).
153 Rainfall averages 3092 ± 181 mm yr⁻¹, with a mean annual air temperature of 25.9 ± 0.3 °C
154 (2003 to 2011; Smithsonian Tropical Research Institute Physical Monitoring Program). There is
155 no pronounced seasonality (Wright *et al.* 2011), although there are two periods of reduced
156 rainfall from February to April and August to September.

157 Seven phasic communities have been identified in these peatlands (Phillips *et al.* 1997). We
158 studied two of these: palm swamp dominated by *Raphia taedigera* (Mart.), a canopy forming
159 palm in the Arecaceae family (9°25'29.20"N, 82°24'05.60"W), and mixed forest dominated by
160 *Camposperma panamensis* (Standl), an evergreen broadleaved hardwood tree in the
161 Anacardiaceae family (9°25'15.00"N, 82°24'14.64"W). The sites were located within the
162 Changuinola peat deposit in the San San Pond Sak wetland (Ramsar site No. 611; ≈ 164 km²).
163 The distance between the sites was approximately 300 m. Both sites are freshwater (surface
164 water conductivity < 200 μ S cm⁻¹), with the water table predominantly at or just below (10 cm)
165 the peat surface. Maximum recorded water fluctuations were + 15 to – 40 cm relative to the peat
166 surface, with surface water consistently above the peat surface during periods of high rainfall.
167 Dissolved O₂ concentrations in the pore water were up to 3.3 ppm at the surface (Palm swamp:
168 1.35 ± 0.25 ppm; Mixed forest: 2.15 ± 0.34), but as low as 0.2 ppm at 50 cm belowground (Palm
169 swamp: 0.72 ± 0.27 ppm; Mixed forest: 0.68 ± 0.19). Nutrient levels at the two sites differ with
170 respect to total and exchangeable P (higher at the palm swamp), as well as microbial N and P
171 (higher at the palm site) and peat C:N and C:P ratios (higher at the palm swamp site) (Sjögersten

172 *et al.* 2011). Palm sites had large amounts of palm leaf litter at the surface and a dense but
173 shallow (1.1 m depth) fibrous root system (Wright *et al.* 2011). The mixed forest sites had large
174 amounts of *C. panamensis* leaf litter at the surface but leaf litter from other species was also
175 present (for further details on of the forest structure and composition see Sjögersten *et al.*, 2011
176 and Hoyos-Santillan *et al.*, 2016). *C. panamensis* is characterized by woody lignified structural
177 roots reaching at least 1 m depth and abundant surface knee roots (Wright *et al.* 2011).
178 Microtopography within all sites consisted of shallow ponds and raised areas (close to trees
179 associated with root structures).

180

181 2.2. Experimental design and methodology

182

183 2.2.1. Nutrient addition experiment

184 The potential role of nutrient limitation on microbial activity and litter decomposition was
185 explored by a 5 month (October 2011 to March 2012) litterbag experiment. The nutrient
186 treatments were: N, P, N+P and control (Ctrl). The experiment consisted of ten blocks distributed
187 along 150 m transects running from south-east to north-west at both the palm swamp and mixed
188 forest sites (20 blocks in total). Each block was 10 × 10 m with the nutrient enrichment
189 treatments applied at each corner, blocks were 5 m apart (Fig. 1). Adjacent corners had the same
190 nutrient treatment.

191 *R. taedigera* and *C. panamensis* litter for the decomposition study was collected from the palm
192 swamp and mixed forest, respectively. The collected litter consisted of recently senesced leaves,

193 freshly cut leaf stalks (petioles) or stems (~ 5 cm in diameter), and fine lateral roots (2–4 mm
194 diameter) from the top 20 cm of the soil profile. After collection, the litter was cleaned with
195 deionized water (DI) and air dried for five days. To allow comparable masses to be weighed out,
196 the litter material was cut into smaller pieces: leaves were cut into ~ 2 × 2 cm pieces, roots were
197 cut into ~ 2 cm lengths, and stems were cut into ~ 1 cm thick discs to ensure that a cross section
198 of the stem tissue was used. Litter was weighed (leaves: ~ 2 g; whereas stems and roots: ~ 1 g),
199 placed separately into pre-weighed polyester mesh litterbags (10 × 10 cm; 560 µm mesh), and
200 tied with polyamide thread (∅ = 0.8 mm). Litter bags were placed directly on the peat surface
201 avoiding hollows. For the belowground incubation (50 cm depth), a narrow slit was cut into the
202 peat and litterbags were manually pushed to the right depth. To aid recovery, litter bags were
203 tied to a string which was securely attached to the ground surface. One litterbag of each tissue
204 type was placed at each of the incubations locations at the start of the experiment, giving a total
205 of 480 litterbags (2 depths (surface and 50 cm depth) × 3 tissue types (leaves, stems, roots) × 4
206 nutrient treatments (Ctrl, N, P and N+P) × 2 sites (palm swamp and mixed forest) × 10 blocks).

207 Nutrient enrichment was applied once at the beginning of the experiment by filling 25 cm
208 sections of dialysis tubing (Spectra/Por[®] membrane: 40mm diameter, 6000 to 8000 molecular
209 weight cut off) with 0.86 mol of either N (Urea: CO(NH₂)₂) or P (calcium phosphate monobasic
210 monohydrate: Ca(H₂PO₄)₂•H₂O) fertilizer. This allowed a slow release of nutrients through the
211 membrane (Feller 1995). Within each block, fertilizer was applied at both the surface and
212 belowground (50 cm) adjacent to the litterbags (< 10 cm from litterbags). For the belowground
213 treatment the dialysis tubes were inserted in a narrow vertical slit cut into the peat.

214 After five months, soil samples were collected to evaluate the impact of the nutrient treatments
215 on surface peat properties (*i.e.* extractable and microbial nutrients, and hydrolytic enzyme

216 activities). To do this, $10 \times 10 \times 10$ cm samples of peat were carefully cut from the surface peat
217 where the litterbags were incubated. Soil samples were stored in plastic bags at 4 °C for one
218 week prior to nutrient and enzymatic analyses.

219 The increase of available nutrients after the nutrient addition treatment, dissolved organic C
220 (DOC) and dissolved N fractions (TDN = dissolved organic nitrogen (DON) + inorganic fraction
221 (nitrate-nitrite and ammonium)) were extracted from surface peat (10 cm depth) by shaking 40 g
222 (fresh weight) of peat in 75 mL of 0.5 M K_2SO_4 for 1 h (Sjögersten *et al.* 2011). Extracts were
223 centrifuged (8000 g, 15 min) and DOC and TDN in the supernatant were determined after a five-
224 fold dilution by TOC-TN analyzer (Shimadzu, Columbia, MD). Readily-exchangeable P was
225 determined by extraction with anion exchange membranes (AEM) (Myers, Thien & Pierzynski
226 1999; Turner and Romero 2009). For this purpose, surface peat (20 g fresh weight) was shaken
227 for 24 h with 80 mL deionized water and five anion-exchange resin strips (1×40 mm;
228 manufactured by BDH Prolabo). The strips were rinsed in deionized water and the phosphate
229 recovered by shaking for 1 h in 50 mL of 0.25 M H_2SO_4 . Phosphate was determined in the acid
230 solution at 880 nm following online neutralization and automated molybdate colorimetry using a
231 flow injection analyzer (Lachat Quikchem 8500, Hach Ltd, Loveland, CO).

232 To investigate the relationship between the nutrient treatments and microbial activity, we
233 measured, in the peat, microbial biomass C, N and P, and extracellular hydrolytic enzyme
234 activities; these parameters were used as indicators of the functioning of the microbial
235 community at the two experimental sites. Microbial C and N were estimated by $CHCl_3$
236 fumigation and 0.5 M K_2SO_4 extraction using a correction factor of 2.64 to account for the
237 unrecovered biomass C (Vance, Brookes & Jenkinson 1987) and 1.85 to account for unrecovered
238 biomass N (Brookes *et al.* 1985). Microbial P was determined by extraction by hexanol

239 fumigation and anion-exchange membranes as described previously (Myers *et al.* 1999; Turner
240 & Romero 2009). Microbial P was calculated as the difference between phosphate in fumigated
241 and unfumigated samples.

242 Total C and N were measured in initial litter and peat samples collected from the peat surface
243 and 50 cm depth. Litter and peat samples were ball milled prior to analysis on a total element
244 analyzer (Thermo Flash EA 1112, CE Instruments, Wigan, UK). Peat and litter ash from loss on
245 ignition analysis was dissolved in 6 M HNO₃ to estimate P concentration by molybdate
246 colorimetry (Andersen 1976). For detailed methods see Hoyos-Santillan (2014).

247 To assess if nutrient addition altered the activity of enzymes involved in the release of C, N, P
248 and sulfur from organic compounds, the activities of five different extracellular hydrolytic
249 enzymes were measured at the end of the experiment using fresh surface peat collected from
250 three of the nutrient addition experimental blocks at the palm swamp and the mixed forest.

251 Assays were conducted using methylumbelliferone-linked fluorogenic substrates (Turner &
252 Romero 2009; Turner 2010). Specifically, enzymes and substrates were: i)
253 phosphomonoesterase: 4-methylumbelliferyl phosphate (MUP); ii) phosphodiesterase: bis-(4-
254 methylumbelliferyl) phosphate (BisMUP); iii) arylsulfatase: 4-methylumbelliferyl sulfate
255 (MUS); iv) β -glucosidase: 4-methylumbelliferyl β -D-glucopyranoside (MUBG); v) *N*-acetyl- β -
256 glucosaminidase: 4-methylumbelliferyl *N*-acetyl- β -D-glucosaminide (MUNA). For the assays,
257 peat (2 g fresh weight) was added to 200 mL of 1 mM sodium azide (NaN₃) solution and stirred
258 for 10 min. Aliquots (50 μ L) of peat suspension were dispensed into a 96-well microplate
259 containing 100 μ L of 200 μ M substrate and 50 μ L of sodium acetate-acetic acid buffer adjusted
260 to pH 4 (the mean peat pH). Microplates were incubated at 30 °C for 30 min; following
261 incubation 50 μ L of 0.5 M NaOH was added to terminate the reaction, and fluorescence was

262 determined immediately on a FLUOstar Optima spectrofluorometer (BMG Labtech, Offenburg,
263 Germany).

264

265 2.2.2. Litter translocation experiment

266 The reciprocal litter translocation experiment involved incubating *R. taedigera* leaf litter in both
267 a palm swamp and a mixed forest and vice versa for *C. panamensis*. The litter translocation used
268 five of the ten blocks at the palm swamp and mixed forest; litterbags with leaves were installed
269 at surface of the control corners of the odd numbered blocks. Total number of litterbags was 20
270 (*i.e.*, 2 species \times 2 sites \times 5 replicates). The incubation time was five months and litter mass loss
271 was quantified as in 2.2.3.

272

273 2.2.3. Litterbag recovery

274 After collection, the litterbags were carefully rinsed with deionized water. It is possible that fine
275 litter ($<560 \mu\text{m}$) was lost during the cleaning process, resulting in a slight overestimation of the
276 mass loss during the incubation. After rinsing, bags were opened and the litter visually inspected
277 to remove new root growth. Litter was then dried at 70 °C for a minimum of 48 h to constant
278 weight in pre-weighted aluminum trays. The remaining mass of litter were calculated as a
279 proportion of the initial mass remaining at the end of the experiment (Wieder & Lang 1982).

280

281 2.3. Statistical analyses

282 We used linear mixed models to assess the impact of the nutrient treatments on the measured
283 parameters. The models were fitted using Residual Maximum Likelihood (REML). To analyze
284 the effect of the nutrients addition experiment on nutrient concentrations, ratios (C:N, C:P and
285 N:P), hydrolytic enzyme activity, phasic community and nutrient treatment (Ctrl, N, P and N+P)
286 were used as fixed factors and block as random factor. To analyze the effect of the nutrient
287 addition on % litter remaining, the nutrient treatment (Ctrl, N, P and N+P), the different tissues,
288 and the incubation depth were used as fixed factors, and block as random factor. The
289 relationships between nutrient ratios in the extractable and microbial fractions were analyzed
290 using linear regression. For the analysis of the litter translocation experiment (% remaining
291 mass_{dw}), the sites (palm swamp and mixed forest), and the translocation treatment were used as
292 fixed factors, and block was the random factor. Residual plots were checked to ensure the
293 assumption of normality and homogeneity of the residuals were met. We calculated the home
294 field advantage index (HFAI), which quantifies the extent to which decomposition is faster or
295 slower at home. Results throughout the text and figures are presented as mean ± SE. Statistical
296 analyses were performed in GenStat (VSN International 2011).

297 2.3.1. HFAI calculation

298 The HFAI is useful to evaluate the results obtained from the reciprocal experiment in the context
299 of the home field advantage theory. The calculation was done according to Ayres *et al.* (2009).
300 In order to do so, we calculated A_{RMLa} , A_{RMLb} , B_{RMLa} and B_{RMLb} ; which represent the Relative
301 Mass Loss (RML) of leaves from one specie at a certain site. For instance, A_{RMLa} represents the
302 relative mass loss of leaves from specie A at site a:

$$303 \quad A_{RMLa} = \frac{A_a}{A_a + B_a} \times 100$$

304 where A_a and B_a correspond to the percent mass loss of leaf litter of two different species (*i.e.*, A
305 and B) at site a. From these, HFAI was calculated as follows:

$$306 \quad HFAI = \left[\frac{A_{RMLa} + B_{RMLb}}{2} / \frac{A_{RMLb} + B_{RMLa}}{2} \right] \times 100 - 100$$

307 **3. Results**

308 3.1. Differences in site and litter nutrient status

309 The two study sites differed in their nutrient status with greater TDN and readily-exchangeable P
310 concentrations at the palm swamp site, in line with Sjögersten *et al.* (2011). This difference was
311 reflected in the nutrient status of the microbial community, which differed between the two sites:
312 the palm swamp had higher microbial N and P concentrations, lower microbial C:N ratios, and
313 higher microbial C:P ratios.

314 In control plots, C:N ratios were higher in the microbial fraction than in the extractable fraction,
315 while C:P and N:P ratios were lower in the microbial fraction (Table 1). Freshly fallen litter had
316 high C:N ratios but varied considerably among tissues and species with *R. taedigera* stems
317 having the highest C:N ratio, and *R. taedigera* leaves having the lowest C:N ratio. *R. taedigera*
318 litter C:P ratios were comparable to the surface peat, while the peat C:P ratios at 50 cm depth
319 were much higher. The *C. panamensis* litter had a more variable C:P ratio than *R. taedigera* with
320 leaf litter having four times as high ratios as root and stem tissue. The C:P ratios for all litter
321 types for both species was considerably greater than in the peat extractable and microbial
322 fractions. Litter N:P ratios were less than half of those found for surface peat, but higher than the
323 N:P ratios in the microbial and extractable fractions.

324

325 3.2. Nutrient addition experiment

326

327 3.2.1. Extractable and microbial nutrients

328 Five months after the nutrient addition, TDN and readily-exchangeable P were significantly
329 greater in plots where nutrients were applied (N_{add} : $F_{1,28} = 8.71$, $P < 0.01$; P_{add} : $F_{1,56} = 7.67$, $P <$
330 0.01 ; Fig. 2), apart from TDN concentrations at the palm site. Neither DOC nor microbial C
331 varied significantly with nutrient addition (N_{add} : $F_{1,30} = 1.53$; $P > 0.05$; P_{add} : $F_{1,30} = 0.02$; $P >$
332 0.05). Microbial biomass N and P did not increase in response to the fertilization treatment (N_{add} :
333 $F_{3,6} = 0.87$; $P > 0.05$; $F_{3,6} = 1.16$; $P > 0.05$; at the palm and mixed forest, respectively; P_{add} : $F_{2,11}$
334 $= 1.04$; $P > 0.05$; $F_{2,10} = 1.71$; $P > 0.05$; at the palm and mixed forest, respectively; Fig. 2).
335 However, both the DOC/TDN (*i.e.* the extractable fraction) ($\text{Site} \times N_{\text{add}}$: $F_{1,12} = 13.66$; $P < 0.001$)
336 and microbial C:N ($\text{Site} \times N_{\text{add}}$: $F_{1,12} = 5.59$; $P < 0.05$) ratios decreased significantly in response
337 to N addition at the low nutrient mixed forest site (Fig. 3a,b); and there was a positive
338 relationship ($F_{1,23} = 30.09$; $P < 0.001$; $R^2 = 0.56$) between the DOC/TDN and microbial C:N
339 ratios (Fig. 3c).

340

341 3.2.2. Impacts of nutrient addition on extracellular enzymatic activity

342 Phosphomonoesterase activity was higher in the mixed forest site than at the palm swamp site
343 ($F_{1,4} = 58.28$, $P < 0.01$) but was not affected by nutrient addition ($F_{3,12} = 1.95$, $P > 0.05$) (Fig.
344 4a). The activity of phosphodiesterase did not vary between sites ($F_{1,4} = 4.23$, $P > 0.05$) or
345 treatments ($F_{3,12} = 1.9$, $P > 0.05$) (Fig. 4b). Arylsulfatase activity decreased with P addition ($F_{1,12}$

346 = 5.72, $P < 0.05$), while N addition increased arylsulfatase activity at the palm swamp but not at
347 the mixed forest site (Site \times N_{add}: $F_{1,12} = 5.5$, $P < 0.05$) (Fig. 4c). β -glucosidase activity did not
348 vary between sites (Fig. 4d), but was increased by N addition at the palm swamp but not at the
349 mixed forest (Site \times N_{add}: $F_{1,12} = 4.03$, $P < 0.05$). In contrast, P addition increased *N*-acetyl- β -
350 glucosaminidase activity at the mixed forest but not at the palm swamp (Site \times P_{add}: $F_{1,12} =$
351 14.19, $P < 0.01$) (Fig. 4e).

352

353 3.2.3. Impacts of nutrient addition on litter decomposition

354 When decomposed at the surface, roots were the most recalcitrant tissue of *R. taedigera*; whereas
355 stems were the most recalcitrant tissue of *C. panamensis* (Fig. 5c,e). Leaves of *R. taedigera*
356 decomposed slower than *C. panamensis* leaves at the surface and belowground (Fig. 5a,d).
357 Leaves decomposed fastest among *C. panamensis* tissues; whereas stems decomposed fastest
358 among *R. taedigera* tissues (Fig. 5b,d).

359 Nitrogen addition increased the belowground mass loss of both *R. taedigera* and *C. panamensis*
360 leaves by ~ 10% (Fig. 5a,d). However, this effect was not observed when N and P were applied
361 together. Phosphorus addition in isolation slightly reduced mass loss of *R. taedigera* and *C.*
362 *panamensis* leaves belowground (Fig. 5a,d).

363

364 3.3. Translocation experiment

365 Mass loss was consistently greater at the site of litter origin ($F_{2,55} = 101.48$, $P < 0.001$) (Fig. 6).
366 Specifically, mass loss of *R. taedigera* leaves was approximately 6 % higher at the palm swamp
367 site compared to the *R. taedigera* litter translocated to the mixed forest. This pattern was
368 repeated on *C. panamensis* leaves, with mass loss being 9 % higher in the mixed forest site
369 compared to the *C. panamensis* leaves translocated to the palm swamp. The home field
370 advantage index (HFAI) demonstrated a positive effect of 28 %.

371

372 **4. Discussion**

373 *4.1. Nutrient controls of extra cellular hydrolytic enzyme activities and litter decomposition*

374 As expected the mixed forest site had lower nutrient availability than the palm swamp site (Fig.
375 2) and we observed strong effects of the nutrient addition on both extractable (*i.e.* DOC/TDN)
376 and microbial C:N in the low nutrient mixed forest, but not in the nutrient rich palm swamp (Fig.
377 3). In contrast to our prediction that sites with microbial nutrient limitation would respond to
378 nutrient addition by down-regulating enzymes involved in nutrient acquisition, we found no
379 down-regulation of phosphomonoesterase activity at either site. However, in line with our
380 prediction, the activity of enzymes involved in the decomposition of large plant-derived
381 polymers, including β -glucosidase, arylsulfatase and *N*-acetyl- β -glucosaminidase, were enhanced
382 by N addition in surface peat in the palm swamp and by P addition in the mixed forest,
383 respectively (Fig. 4d, e). This reflects differences in the nutrient levels at the two sites: low N
384 relative to P concentrations at the palm swamp and low P concentrations in the mixed forest
385 (Olander & Vitousek 2000; Sjögersten *et al.* 2011) and suggests that the degradation of sugars as
386 well as more complex organic molecules in this peatland are in part limited by variation in forest

387 nutrient status in agreement with findings from higher latitude peatlands (Bubier *et al.* 2003;
388 Wang *et al.* 2014).

389

390 In contrast to our prediction that nutrient addition would accelerate litter mass loss at the low
391 nutrient site, but have little effect, at the high nutrient palm site, N addition increased mass loss
392 of leaf litter deeper in the peat profile by ~ 10 % (Fig. 5a,d). This is important because foliar
393 litter inputs represent a sizable fraction (~ 30%) of the total C inputs from net primary
394 productivity (NPP; 333 g C m⁻² yr⁻¹; Sjögersten *et al.* 2014) and partially decomposed leaf litter
395 contributes to long term C storage in peatlands as it becomes buried and preserved over time due
396 to water logged conditions (Hoyos-Santillan *et al.* 2015). Nitrogen addition affected leaf litter
397 decomposition only at depth, indicating that nutrient limitation is an additional constraint on
398 decomposition under anaerobic conditions, and/or that nutrient limitation is more pronounced in
399 deeper, more degraded peat. Furthermore, shifts in the microbial community composition and a
400 reduction in microbial activity in response to anaerobic conditions are likely to slow nutrient
401 mineralization at depth (Jackson *et al.* 2009).

402 Variation in mass loss responses to N addition among litter types, with leaves decomposing
403 faster with N addition but roots and stems being unaffected, is presumably linked to differences
404 in litter organic chemistry among tissue types (Hobbie & Vitousek 2000). For example, root and
405 stem tissues from the two study species contained greater concentrations of lignin than leaves,
406 making them more recalcitrant to decomposition (Hoyos-Santillan *et al.* 2015). As lignin
407 decomposition is strongly limited by oxygen availability (Zeikus 1981), it is plausible that
408 aeration was a greater limitation of degradation of lignin rich roots and stems than nutrient

409 availability, explaining why only decomposition of labile leaf litter tissue was enhanced by the N
410 addition under the water logged conditions at 50 cm depth.

411 Phosphorus addition reduced litter mass loss, in contrast to our prediction. This might be linked
412 to suppression of phenol oxidase activity, as suggested by findings from mangrove and mineral
413 soil systems (Keuskamp *et al.* 2015a; Qi *et al.* 2016), possibly due to a reduction in fungal
414 activity in response to greater concentrations of mineral P (Tien & Myer 1990; Hobbie 2000). As
415 a reduction in phenoloxidase activity may reduce decomposition of complex C (Freeman *et al.*
416 2004), suppression of phenol oxidase activity by P addition in our study might therefore explain
417 the reduction in decomposition in P treated plots.

418 Taken together, our findings in part support our hypothesis that nutrient availability influences
419 litter decomposition and activities of extra cellular hydrolytic enzymes. Nutrient addition
420 increased the activities of extra cellular enzymes involved in degradation of large plant
421 molecules, and increased leaf litter decomposition under anoxic conditions at depth following N
422 addition. However, high C:N or C:P ratios in the bulk litter tissues, relative to low C:N and C:P
423 ratios in the microbial biomass, which are at the lower range of C:N and C:P ratios for the
424 microbial biomass reported in the literature (Cleveland & Liptzin 2007; Xu, Thornton & Post
425 2013), did not predict which litter types were most affected by nutrient addition. Instead,
426 microbial C:N ratios were clearly related to the C:N ratios in the extractable dissolved fraction,
427 suggesting a decoupling between bulk litter chemistry and microbial stoichiometric ratios in line
428 with Fanin *et al.* (2013), although the slope of the relationship shown in this study is steeper.
429 Furthermore, decomposition of leaf litters, which has the lowest lignin:N ratios of the different
430 tissue types (Hoyos-Santillan *et al.* 2015), were most responsive to N addition. These somewhat
431 contrasting findings suggests that although nutrient availability clearly affects some of the

432 processes controlling litter decomposition in line with stoichiometric theory (Sterner & Elser
433 2002), low nutrient availability does not seem to exert a strong control of litter decomposition in
434 these two peat swamp forest communities. Instead, nutrient limitation appears to be mediated by
435 litter chemistry and position in the peat profile, reflecting peat oxygen levels (Hoyos-Santillan *et*
436 *al.* 2016).

437

438 4.2 Home field advantage in the context of contrasting site nutrient status

439 Our findings of a strong positive HFA effect supported our prediction that palm leaf litter would
440 degrade faster at the palm site, while the litter from the low nutrient mixed forest would degrade
441 fastest at the mixed forest site (Fig. 6). The HFA index (28 %) was at the upper range for HFAI
442 reported in the literature (Ayres *et al.* 2009; Veen *et al.* 2015), which we speculate was driven by
443 the two litter species belonging to contrasting plant functional types (*i.e.* palm vs evergreen
444 broad leaved), which has previously been show to result in strong HFA effects and the
445 contrasting site nutrient levels (Ayres *et al.* 2009; Veen *et al.* 2015).

446 Furthermore, the alternative prediction that a site with higher nutrient status would increase litter
447 decomposition rates was not supported by our findings, because *C. panamensis* leaf litter
448 degraded at a marginally greater rate at the low nutrient mixed forest site than the *R. taedigera*
449 leaf litter at the palm site, while root litter decomposition was comparable when incubated at the
450 peat surface (Fig. 2, 5). Although the slower stem decomposition of *C. panamensis* compared
451 with *R. taedigera* might be linked to low nutrient levels at the mixed forest site, contrasting
452 tissue chemistry (*i.e.* lignified woody vs palm stem tissue structure) between the two species

453 might also influence decomposition rates (Hoyos-Santillan *et al.* 2015), as tissue chemistry
454 strongly affect decomposition rates (*e.g.* Baumann *et al.* 2009).

455 The translocation experiment clearly supported the HFA theory. Despite greater fertility at the
456 palm swamp site (Fig. 2), which we assumed would enhance decomposition rates, decomposition
457 was always greater for autochthonous litter even when litter was decomposing in the lower
458 nutrient environment. This suggests that the microbial community is adapted to decompose site-
459 specific litter and that a well-adapted decomposer community is more important for
460 decomposition than nutrient availability. This notion is supported by the fact that distinct soil
461 microbial communities accompany particular forest communities within the peatland (Troxler *et*
462 *al.* 2012), suggesting that different consortia of microorganisms are responsible for litter
463 decomposition at the two different sites. This is consistent with previous findings in temperate,
464 subtropical and tropical forests (Hunt *et al.* 1988; Gholz *et al.* 2000; Mayor & Henkel 2006;
465 Zhou *et al.* 2008; Austin *et al.* 2014). For example, it is plausible that different microbial
466 communities produce different enzymes (Kaiser *et al.* 2014) suggesting that microbial
467 communities involved in decomposition are specialized rather than being functionally redundant
468 (Schimel & Schaeffer 2012; Keiser *et al.* 2014).

469

470 4.3 Peatland C dynamics in the context of nutrient limitation and HFA

471

472 Our results indicate that nutrient limitation is an important control of decomposition processes in
473 tropical peatlands and could account for the persistence of relatively labile leaf material deeper in

474 the peat profile where nutrient levels tend to be low (Hoyos-Santillan *et al.* 2015). However,
475 given that nutrient addition did not accelerate litter mass loss at the peat surface, which is
476 governed by oxic conditions and generally has a faster decomposition rate (Hoyos-Santillan *et al.*
477 2015), there does not appear to be a “nutrient latch” on C loss from litter decomposition in this
478 peatland. Our results also support HFA theory, indicating that microbial adaptations to the
479 conditions found at a given site can overcome factors often considered to exert strong controls of
480 litter decomposition rates, such as low nutrient availability.

481 In the context of long-term peatland carbon dynamics, our study demonstrates that stoichiometric
482 ecological theory applies to peatland decomposition processes, particularly under conditions
483 where oxygen and nutrient levels are low but the organic material is relatively labile (*i.e.* long
484 term preservation of leaf litter through the water logged parts of the peat profile). Our study also
485 suggests that decomposition rates at the peatland surface may remain high across contrasting
486 plant phasic communities as a result of a specialised decomposer communities adapted to these
487 “home” conditions. Finally, our results show that contrasting tissue chemistry should not be used
488 as a predictor of *in situ* decomposition rates, or different litters contribution to long term peatland
489 C storage without considering the associated decomposer community at a given site.

490

491 **Acknowledgments**

492 J.H.S. thanks The National Council on Science and Technology (CONACyT-Mexico) for his
493 PhD scholarship (211962). The authors also thank the Light Hawk program for its support in the
494 aerial surveys. We thank Erick Brown for field assistance, and Gabriel Jácome, Plinio Góndola,
495 Dayana Agudo, Tania Romero, Luis A. Ramos, Dianne de la Cruz, Vanessa Pardo, John Corrie

496 and Darren Hepworth for logistical and laboratory support. The authors declare that there are no
497 conflicts of interest.

498 **Data accessibility**

499 Data to support this article is publicly available at Dryad Digital Repository (Hoyos-Santillan *et*
500 *al.* 2017; doi:10.5061/dryad.460mc).

501

502 **References**

503 Allison, V.J., Condon, L.M., Peltzer, D.A., Richardson, S.J. & Turner, B.L. (2007) Changes in
504 enzyme activities and soil microbial community composition along carbon and nutrient
505 gradients at the Franz Josef chronosequence, New Zealand. *Soil Biology and Biochemistry*,
506 **39**, 1770–1781.

507 Andersen, J. (1976) An ignition method for determination of total phosphorus in lake sediments.
508 *Water Research*, **10**, 329–331.

509 Austin, A.T., Vivanco, L., González-Arzac, A. & Pérez, L.I. (2014) There's no place like home?
510 An exploration of the mechanisms behind plant litter-decomposer affinity in terrestrial
511 ecosystems. *The New phytologist*, **204**, 307–314.

512 Ayres, E., Steltzer, H., Simmons, B.L., Simpson, R.T., Steinweg, J.M., Wallenstein, M.D., Mellor,
513 N., Parton, W.J., Moore, J.C. & Wall, D.H. (2009) Home-field advantage accelerates leaf
514 litter decomposition in forests. *Soil Biology and Biochemistry*, **41**, 606–610.

515 Baumann, K., Marschner, P., Smernik, R.J. & Baldock, J.A. (2009) Residue chemistry and
516 microbial community structure during decomposition of eucalypt, wheat and vetch residues.

517 *Soil Biology and Biochemistry*, **41**, 1966–1975.

518 Brady, M.A. (1997) *Organic Matter Dynamics of Coastal Peat Deposits in Sumatra, Indonesia*.
519 The University of British Columbia.

520 Bragazza, L., Buttler, A., Habermacher, J., Brancaleoni, L., Gerdol, R., Fritze, H., Hanajík, P.,
521 Laiho, R. & Johnson, D. (2012) High nitrogen deposition alters the decomposition of bog
522 plant litter and reduces carbon accumulation. *Global Change Biology*, **18**, 1163–1172.

523 Brookes, P.P.C., Landman, A., Pruden, G. & Jenkinson, D.D.S. (1985) Chloroform fumigation
524 and the release of soil nitrogen: A rapid direct extraction method to measure microbial
525 biomass nitrogen in soil. *Soil Biology and Biochemistry*, **17**, 837–842.

526 Bubier, J., Crill, P., Mosedale, A., Frohking, S. & Linder, E. (2003) Peatland responses to varying
527 interannual moisture conditions as measured by automatic CO₂ chambers. *Global*
528 *Biogeochemical Cycles*, **17**, 35.1-35.15.

529 Chimner, R.A. & Ewel, K.C. (2005) A tropical freshwater wetland: II. Production, decomposition,
530 and peat formation. *Wetlands Ecology and Management*, **13**, 671–684.

531 Cleveland, C.C. & Liptzin, D. (2007) C:N:P stoichiometry in soil: is there a “Redfield ratio” for
532 the microbial biomass? *Biogeochemistry*, **85**, 235–252.

533 Cornwell, W.K., Cornelissen, J.H.C., Amatangelo, K., Dorrepaal, E., Eviner, V.T., Godoy, O.,
534 Hobbie, S.E., Hoorens, B., Kurokawa, H., Pérez-Harguindeguy, N., Quested, H.M., Santiago,
535 L.S., Wardle, D.A., Wright, I.J., Aerts, R., Allison, S.D., van Bodegom, P., Brovkin, V.,
536 Chatain, A., Callaghan, T. V., Díaz, S., Garnier, E., Gurvich, D.E., Kazakou, E., Klein, J.A.,
537 Read, J., Reich, P.B., Soudzilovskaia, N.A., Vaieretti, M.V. & Westoby, M. (2008) Plant
538 species traits are the predominant control on litter decomposition rates within biomes

539 worldwide. *Ecology Letters*, **11**, 1065–1071.

540 Dommain, R., Couwenberg, J. & Joosten, H. (2011) Development and carbon sequestration of
541 tropical peat domes in south-east Asia: links to post-glacial sea-level changes and Holocene
542 climate variability. *Quaternary Science Reviews*, **30**, 999–1010.

543 Fanin, N., Fromin, N., Buatois, B. & Hättenschwiler, S. (2013) An experimental test of the
544 hypothesis of non-homeostatic consumer stoichiometry in a plant litter-microbe system (ed E
545 Cleland). *Ecology Letters*, **16**, 764–772.

546 Feller, I.C. (1995) Effects of Nutrient Enrichment on Growth and Herbivory of Dwarf Red
547 Mangrove (*Rhizophora Mangle*). *Ecological Monographs*, **65**, 477.

548 Freeman, C., Ostle, N.J., Fenner, N. & Kang, H. (2004) A regulatory role for phenol oxidase during
549 decomposition in peatlands. *Soil Biology and Biochemistry*, **36**, 1663–1667.

550 Freeman, C., Ostle, N. & Kang, H. (2001) An enzymic “latch” on a global carbon store. *Nature*,
551 **409**, 149.

552 Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer, E.W., Howarth, R.W., Seitzinger, S.P.,
553 Asner, G.P., Cleveland, C.C., Green, P.A., Holland, E.A., Karl, D.M., Michaels, A.F., Porter,
554 J.H., Townsend, A.R. & Vöosmarty, C.J. (2004) Nitrogen Cycles: Past, Present, and Future.
555 *Biogeochemistry*, **70**, 153–226.

556 Gholz, H.L., Wedin, D.A., Smitherman, S.M., Harmon, M.E. & Parton, W.J. (2000) Long-term
557 dynamics of pine and hardwood litter in contrasting environments: toward a global model of
558 decomposition. *Global Change Biology*, **6**, 751–765.

559 Gorham, E., Janssens, J.A. & Glaser, P.H. (2003) Rates of peat accumulation during the postglacial
560 period in 32 sites from Alaska to Newfoundland, with special emphasis on northern

561 Minnesota. *Canadian Journal of Botany*, **81**, 429–438.

562 Hobbie, S.E. (2000) Interactions between litter lignin and soil nitrogen availability during leaf
563 litter decomposition in a Hawaiian montane forest. *Ecosystems*, **3**, 484–494.

564 Hobbie, S.E. & Vitousek, P.M. (2000) Nutrient limitation of decomposition in hawaiian forests.
565 *Ecology*, **81**, 1867–1877.

566 Hoyos-Santillan, J. (2014) *Controls of Carbon Turnover in Lowland Tropical Peatlands*. The
567 University of Nottingham.

568 Hoyos-Santillan, J., Lomax, B.H., Large, D., Turner, B.L., Boom, A., Lopez, O.R. & Sjögersten,
569 S. (2015) Getting to the root of the problem: litter decomposition and peat formation in
570 lowland Neotropical peatlands. *Biogeochemistry*, **126**, 115–129.

571 Hoyos-Santillan, J., Lomax, B.H., Large, D., Turner, B.L., Boom, A., Lopez, O.R. & Sjögersten,
572 S. (2016) Quality not quantity: Organic matter composition controls of CO₂ and CH₄ fluxes
573 in neotropical peat profiles. *Soil Biology and Biochemistry*, **103**, 86–96.

574 Hoyos-Santillan, J., Lomax, B.H., Turner, B.L. & Sjögersten, S. (2017) Data from: Nutrient
575 limitation or home field advantage: does microbial community adaptation overcome nutrient
576 limitation of litter decomposition in a tropical peatland?

577 Hunt, H.W., Ingham, E.R., Coleman, D.C., Elliott, E.T. & Reid, C.P.P. (1988) Nitrogen Limitation
578 of Production and Decomposition in Prairie, Mountain Meadow, and Pine Forest. *Ecology*,
579 **69**, 1009–1016.

580 IPCC. (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group*
581 *I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds TF
582 Stocker, D Qin, G-K Plattner, M Tignor, SK Allen, J Boschung, A Nauels, Y Xia, V Bex,

583 and PM Midgley). Cambridge University Press, Cambridge, United Kingdom and New York,
584 NY, USA.

585 Jackson, C.R., Liew, K.C. & Yule, C.M. (2009) Structural and functional changes with depth in
586 microbial communities in a tropical Malaysian peat swamp forest. *Microbial ecology*, **57**,
587 402–12.

588 Kaiser, C., Franklin, O., Dieckmann, U. & Richter, A. (2014) Microbial community dynamics
589 alleviate stoichiometric constraints during litter decay (ed N Johnson). *Ecology Letters*, **17**,
590 680–690.

591 Keuskamp, J.A., Feller, I.C., Laanbroek, H.J., Verhoeven, J.T.A. & Hefting, M.M. (2015a) Short-
592 and long-term effects of nutrient enrichment on microbial exoenzyme activity in mangrove
593 peat. *Soil Biology and Biochemistry*, **81**, 38–47.

594 Keuskamp, J.A., Hefting, M.M., Dingemans, B.J.J., Verhoeven, J.T.A. & Feller, I.C. (2015b)
595 Effects of nutrient enrichment on mangrove leaf litter decomposition. *The Science of the total*
596 *environment*, **508**, 402–10.

597 Keuskamp, J.A., Schmitt, H., Laanbroek, H.J., Verhoeven, J.T.A. & Hefting, M.M. (2013)
598 Nutrient amendment does not increase mineralisation of sequestered carbon during
599 incubation of a nitrogen limited mangrove soil. *Soil Biology and Biochemistry*, **57**, 822–829.

600 Knorr, M., Frey, S.D., Curtis, P.S. & Knorr, A.M. (2005) Nitrogen additions and litter
601 decomposition: A meta-analysis. *Ecology*, **86**, 3252–3257.

602 Kurnianto, S., Warren, M., Talbot, J., Kauffman, B., Murdiyarso, D. & Frohking, S. (2015) Carbon
603 accumulation of tropical peatlands over millennia: A modeling approach. *Global Change*
604 *Biology*, **21**, 431–44.

- 605 Manzoni, S. & Porporato, A. (2009) Soil carbon and nitrogen mineralization: Theory and models
606 across scales. *Soil Biology and Biochemistry*, **41**, 1355–1379.
- 607 Mayor, J.R. & Henkel, T.W. (2006) Do ectomycorrhizas alter leaf-litter decomposition in
608 monodominant tropical forests of Guyana? *The New phytologist*, **169**, 579–88.
- 609 Myers, R.G., Thien, S.J. & Pierzynski, G.M. (1999) Using an Ion Sink to Extract Microbial
610 Phosphorus from Soil. *Soil Science Society of America Journal*, **63**, 1229.
- 611 Olander, L.P. & Vitousek, P.M. (2000) Regulation of soil phosphatase and chitinase activity by N
612 and P availability. *Biogeochemistry*, **49**, 175–191.
- 613 Page, S.E., Rieley, J.O., Shoty, W. & Weiss, D. (1999) Interdependence of peat and vegetation
614 in a tropical peat swamp forest. *Philosophical transactions of the Royal Society of London.*
615 *Series B, Biological sciences*, **354**, 1885–97.
- 616 Phillips, S., Rouse, G.E.G. & Bustin, R.M. (1997) Vegetation zones and diagnostic pollen profiles
617 of a coastal peat swamp, Bocas del Toro, Panamá. *Palaeogeography, Palaeoclimatology,*
618 *Palaeoecology*, **128**, 301–338.
- 619 Qi, R., Li, J., Lin, Z., Li, Z., Li, Y., Yang, X., Zhang, J. & Zhao, B. (2016) Temperature effects on
620 soil organic carbon, soil labile organic carbon fractions, and soil enzyme activities under long-
621 term fertilization regimes. *Applied Soil Ecology*, **102**, 36–45.
- 622 Quested, H.M., Callaghan, T. V., Cornelissen, J.H.C. & Press, M.C. (2005) The impact of
623 hemiparasitic plant litter on decomposition: Direct, seasonal and litter mixing effects. *Journal*
624 *of Ecology*, **93**, 87–98.
- 625 Schimel, J.P. & Schaeffer, S.M. (2012) Microbial control over carbon cycling in soil. *Frontiers in*
626 *Microbiology*, **3**, 1–11.

- 627 Sinsabaugh, R.L. & Follstad Shah, J.J. (2012) Ecoenzymatic Stoichiometry and Ecological
628 Theory. *Annual Review of Ecology, Evolution, and Systematics*, **43**, 313–343.
- 629 Sjögersten, S., Black, C.R., Evers, S., Hoyos-Santillan, J., Wright, E.L. & Turner, B.L. (2014)
630 Tropical wetlands: A missing link in the global carbon cycle? *Global Biogeochemical Cycles*,
631 **28**, 1371–1386.
- 632 Sjögersten, S., Cheesman, A.W., Lopez, O. & Turner, B.L. (2011) Biogeochemical processes
633 along a nutrient gradient in a tropical ombrotrophic peatland. *Biogeochemistry*, **104**, 147–
634 163.
- 635 Sterner, R.W. & Elser, J.J. (2002) *Ecological Stoichiometry: The Biology of Elements from*
636 *Molecules to the Biosphere*.
- 637 Tien, M. & Myer, S.B. (1990) Selection and characterization of mutants of *Phanerochaete*
638 *chrysosporium* exhibiting ligninolytic activity under nutrient-rich conditions. *Applied and*
639 *Environmental Microbiology*, **56**, 2540–2544.
- 640 Troxler, T.G. (2007) Patterns of phosphorus, nitrogen and $\delta^{15}\text{N}$ along a peat development gradient
641 in a coastal mire, Panama. *Journal of Tropical Ecology*, **23**, 683–691.
- 642 Troxler, T.G., Ikenaga, M., Scinto, L., Boyer, J.N., Condit, R., Perez, R., Gann, G.D. & Childers,
643 D.L. (2012) Patterns of Soil Bacteria and Canopy Community Structure Related to Tropical
644 Peatland Development. *Wetlands*, **32**, 769–782.
- 645 Turner, B.L. (2010) Variation in pH Optima of Hydrolytic Enzyme Activities in Tropical Rain
646 Forest Soils. *Applied and Environmental Microbiology*, **76**, 6485–6493.
- 647 Turner, B.L. & Romero, T.E. (2009) Short-Term Changes in Extractable Inorganic Nutrients
648 during Storage of Tropical Rain Forest Soils. *Soil Science Society of America Journal*, **73**,

649 1972.

650 Vance, E.D., Brookes, P.C. & Jenkinson, D.S. (1987) An extraction method for measuring soil
651 microbial biomass C. *Soil Biology and Biochemistry*, **19**, 703–707.

652 Veen, G.F.C., Freschet, G.T., Ordonez, A. & Wardle, D.A. (2015) Litter quality and environmental
653 controls of home-field advantage effects on litter decomposition. *Oikos*, **124**, 187–195.

654 Vivanco, L. & Austin, A.T. (2008) Tree species identity alters forest litter decomposition through
655 long-term plant and soil interactions in Patagonia, Argentina. *Journal of Ecology*, **96**, 727–
656 736.

657 VSN International. (2011) GenStat for Windows 14th Edition.

658 Wang, M., Moore, T.R., Talbot, J. & Richard, P.J.H. (2014) The cascade of C:N:P stoichiometry
659 in an ombrotrophic peatland: from plants to peat. *Environmental Research Letters*, **9**, 24003.

660 Wieder, R.K. & Lang, G.E. (1982) A Critique of the Analytical Methods Used in Examining
661 Decomposition Data Obtained From Litter Bags. *Ecology*, **63**, 1636.

662 Wright, E., Black, C.R., Cheesman, A.W., Drage, T., Large, D., Turner, B.L. & Sjögersten, S.
663 (2011) Contribution of subsurface peat to CO₂ and CH₄ fluxes in a neotropical peatland.
664 *Global Change Biology*, **17**, 2867–2881.

665 Wright, E., Black, C.R., Cheesman, A.W., Turner, B.L. & Sjögersten, S. (2013) Impact of
666 simulated changes in water table depth on ex situ decomposition of leaf litter from a
667 Neotropical peatland. *Wetlands*, **33**, 217–226.

668 Xu, X., Thornton, P. & Post, W. (2013) A global analysis of soil microbial biomass carbon,
669 nitrogen and phosphorus in terrestrial ecosystems. *Global Ecology and Biogeography*, **22**,
670 737–749.

- 671 Yule, C.M. & Gomez, L.N. (2009) Leaf litter decomposition in a tropical peat swamp forest in
672 Peninsular Malaysia. *Wetlands Ecology and Management*, **17**, 231–241.
- 673 Zeikus, J.G. (1981) Lignin metabolism and the carbon cycle. *Advances in microbial ecology* (ed
674 M. Alexander), pp. 211–243. Springer US, New York.
- 675 Zhou, G., Guan, L., Wei, X., Tang, X., Liu, S., Liu, J., Zhang, D. & Yan, J. (2008) Factors
676 influencing leaf litter decomposition: an intersite decomposition experiment across China.
677 *Plant and Soil*, **311**, 61–72.
- 678
- 679

680 **Table captions**

681 **Table 1** Mass-based ratios among C, N and P in different substrate types at the palm swamp and
 682 mixed forest sites.

Table 1. Mass-based ratios among C, N and P in different substrate types at the palm swamp and mixed forest sites.


Ratio	Palm swamp			Mixed forest		
	C:N	C:P	N:P	C:N	C:P	N:P
Microbial	7.43 ± 0.16	5.29 ± 0.47	0.71 ± 0.07	8.25 ± 0.28	3.49 ± 0.47	0.43 ± 0.07
Extractable	3.87 ± 0.38	8.21 ± 1.11	2.13 ± 0.21	3.80 ± 0.44	28.76 ± 6.93	7.80 ± 2.29
Leaf*	37.71 ± na	911.5 ± na	24.17 ± na	127.9 ± na	3984 ± na	31.16 ± na
Root*	55.91 ± na	1155 ± na	20.65 ± na	78.19 ± na	1034 ± na	13.22 ± na
Stem*	140.2 ± na	1082 ± na	7.71 ± na	117.8 ± na	963.0 ± na	8.18 ± na
Peat (surface) ^a	41.53 ± na	1142 ± na	45.24 ± na	35.13 ± na	1274 ± na	98.78 ± na
Peat (-50 cm) ^b	19.79 ± na	5642 ± na	196.9 ± na	40.73 ± na	3001 ± na	76.27 ± na

*Litter are *R. taedigera* and *C. panamensis* for the palm swamp and mixed forest, respectively.

^{a,b} Peat samples were taken before the nutrient treatment was applied (October 2011) from the top 10 cm of the peat profile.


683 **Figure captions**

684 **Figure 1** Schematic diagram outlining the experimental set up for the nutrient addition, (Ctrl)
 685 control, (N) nitrogen and (P) phosphorous. The same set up was used at the palm swamp and the
 686 mixed forest sites. Ten blocks were set up at each site with litterbags placed both at the peat surface
 687 and at 50 cm depth.




688

689 **Figure 2** Comparison of extractable (solid bars) and microbial (hatched bars) Ctrl, N and P in
 690 surface peat at the two study sites, (a,c,e palm swamp; b,d,f mixed forest). Dissolved organic
 691 carbon (a, b), readily-exchangeable P (c, d) and total dissolved N (e, f), after 5 months of the *in*
 692 *situ* nutrient addition. Note the different scales on the ordinate axis when comparing palm swamp
 693 and mixed forest. Statistical analyses are presented in the text.




694

695 **Figure 3** Effects of the nutrient addition treatment on the C:N ratio in: a) the extractable fraction
 696 (*i.e.* DOC/TDN), b) the microbial biomass and c) the relationship between the C:N ratio in the
 697 extractable fraction and in the microbial biomass. Statistical analyses are presented in the text.



698


699 **Figure 4** Hydrolytic enzymes activity (nmol MU g⁻¹ min⁻¹): a) Phosphomonoesterase (MUP), b)
 700 Phosphodiesterase (BisMUP), c) Arylsulfatase (MUS), d) β-glucosidase (MUBG) and e) N-
 701 acetyl-β-glucosaminidase (MUNA). Surface peat samples were taken 5 months after the *in situ*
 702 nutrient addition. Statistical analyses are presented in text.



703

704 **Figure 5** Effect of nutrient addition (Control (Ctrl), N, P and N+P) on the *in situ* % of mass
 705 remaining. *R. taedigera* litter mass remaining of (a) leaves, (b) stems, (c) roots after 5 months.
 706 REML outputs are: Tissue: $F_{2,215} = 121.12$, $P < 0.001$; Surface/Belowground: $F_{1,215} = 38.88$, $P <$
 707 0.001 ; Treatment: $F_{3,215} = 3.14$, $P < 0.05$; Tissue \times Surface/Belowground: $F_{2,215} = 7.33$, $P <$
 708 0.001 ; Tissue \times Treatment: $F_{6,215} = 2.97$, $P < 0.01$; Surface/Belowground \times Treatment: $F_{3,215} =$
 709 0.19 , $P > 0.05$; Tissue \times Surface/Belowground \times Treatment: $F_{6,215} = 0.44$, $P > 0.05$. *C.*
 710 *panamensis* litter mass remaining of (d) leaves, (e) stems, (f) roots after 5 months. REML
 711 outputs are: Tissue: $F_{2,209} = 95.21$, $P < 0.001$; Surface/Belowground: $F_{1,209} = 15.33$, $P < 0.001$;
 712 Treatment: $F_{3,209} = 5.48$, $P < 0.001$; Tissue \times Surface/Belowground: $F_{2,209} = 0.75$, $P > 0.05$;

713 Tissue × Treatment: $F_{6,209} = 2.38$, $P < 0.05$; Surface/Belowground × Treatment: $F_{3,209} = 4.23$, $P <$
 714 0.01 ; Tissue × Surface/Belowground × Treatment: $F_{6,215} = 3.14$, $P < 0.01$




715

716

717 **Figure 6** Mass remaining (%) of *R. taedigera* (palm swamp species) and *C. panamensis* (mixed
 718 forest species) leaf litter after 5 months of decomposition as part of the translocation experiment
 719 between palm swamp and mixed forest sites. Litterbags were placed at the peat surface.
 720 Statistical analyses are presented in the text.

721



722

723