

1 **Title:** Air pressure changes in the creation and bursting of the type-1 big bubble
2 in deep anterior lamellar keratoplasty: an ex-vivo study.

3 **Authors:** Saief L AlTaan, Imran Mohamed, Dalia G Said, Harminder S Dua

4 **Affiliations:** Larry A Donoso laboratory for eye research, Academic Section of
5 Ophthalmology, Division of Clinical Neuroscience, University of Nottingham,
6 Nottingham.

7 **Corresponding author:**

8 Prof. Harminder S Dua

9 Department of Ophthalmology

10 B Floor, Eye ENT Centre

11 Queens Medical Centre, Derby Road

12 Nottingham, NG7 2UH

13 England

14 Email: Harminder.dua@nottingham.ac.uk

15 Tel.: +44 (0)115 924 9924

16 Fax: +44 (0)115 970 9963

17

18 **Conflict of interest:** the authors have no conflict of interest to declare.

19 **Running title:** Air pressure changes in big bubble DALK.

20 **Financial support:** The Royal College of Surgeons, Edinburgh, Scotland and Royal Blind
21 Scotland.

22 Key words: Deep anterior lamellar keratoplasty (DALK), DALK-triple, Dua's layer

23

24 **Abstract**

25 **Aim:** To measure the pressure and volume of air required to create a big bubble
26 (BB) in simulated deep anterior lamellar keratoplasty (DALK) in donor eyes and
27 ascertain the bursting pressure of the BB.

28 **Methods:** Twenty two human sclera-corneal discs were used. Air was injected
29 into the corneal stroma to create a BB and the pressure measured by means of a
30 pressure converter attached to the system via a side port. A special clamp was
31 designed to prevent air leak from the periphery of the discs. The pressure at
32 which air emerged in the corneal tissue; the bursting pressure measured after
33 advancing the needle into the bubble cavity and injecting more air; the volume
34 of air required to create a BB and the volume of the BB were ascertained.

35 **Results:** Type-1 BB were achieved in 19 and type-2 BB in 3 eyes. The maximum
36 pressure reached to create a BB was 96.25 ± 21.61 kpa; the mean type-1
37 intra-bubble pressure was 10.16 ± 3.65 kpa. The mean bursting pressure of a
38 type-1 BB was 66.65 ± 18.65 kpa while that of a type-2 BB was $14.77 \pm$
39 2.44 kpa. The volume of air required to create a type-1 BB was 0.54ml and the
40 volume of a type-1 BB was consistently 0.1ml.

41 **Conclusions:** Dua's layer baring DALK can withstand high intraoperative
42 pressures compared to Descemet's membrane baring DALK. The study suggests
43 that it could be safe to undertake procedures such as DALK-triple with a type-
44 1BB but not with a type-2 BB.

45

46

47 **Introduction:**

48 Deep anterior lamellar keratoplasty (DALK) has replaced penetrating
49 keratoplasty as the procedure of choice in surgical management of eyes with
50 diseases affecting the corneal stroma and affecting sight such as scars,
51 dystrophy or ectasia. The Big Bubble (BB) technique (ref.1) is the most popular
52 technique wherein air is injected in the corneal stroma to separate either the
53 Descemet's membrane (DM) or the DM together with a layer of deep corneal
54 stroma termed the pre-Descemet's layer (Dua's layer-DL). This allows excision
55 of the affected stroma and recipient epithelium and replacement with healthy
56 stroma and epithelium from a cadaver donor.

57 When air is injected in the corneal stroma either cleaves the DL from the deep
58 stroma to create a big bubble termed type-1 or it accesses the plane between
59 DM and DL to create a thin walled bubble termed type-2. The wall of a type-1 BB
60 is made of DL and DM while of a type-2 BB is made of DM alone and is more
61 vulnerable to major tears or bursting during surgery. Often during injection of
62 air, tiny bubbles escape from the peripheral cornea, in the vicinity of the
63 trabecular meshwork in to the anterior chamber and can cause post-operative
64 raised intraocular pressure (ref. 2,3,4).

65 Dua et al have reported that DL is a strong and resilient layer with bursting
66 pressure 1.45 bars (ref. 5). Based on the above information, Zaki AA et al
67 described a combination of DALK with phacoemulsification and lens implant,
68 termed the DALK-Triple procedure. When confronted with patients requiring
69 DALK who also had dense cataracts they were able to perform cataract surgery
70 under the exposed DL of a type-1 BB. They reported that DL could withstand all
71 pressure fluctuations associated with the phacoemulsification procedure and that

72 despite stromal scarring requiring keratoplasty, the DL was remarkably clear in
73 most cases. (ref. 6). In one instance they attempted DALK-Triple under the DM
74 (type-2 BB), which burst promptly during injection of viscoelastic in the anterior
75 chamber.

76 In this study we report the pressure and volume of air required to create the BB,
77 the volume and pressure of air in the type-1 BB and the bursting pressure of the
78 type-1 BB.

79

80 **Materials and Methods:**

81 Tissue samples: Twenty two human sclero-corneal discs from eye bank donor
82 eyes that were not suitable for transplantation were used. The sclera-corneal
83 discs were maintained in organ culture in Eagle's minimum essential medium
84 with 2% foetal bovine serum for four to eight weeks post-mortem. Donor details
85 are given in table 1.

86 Experiment to measure pressure

87 Air injection: The sclero-corneal disc was placed endothelial side up in a petri
88 dish and kept moist with balanced salt solution. In fifteen samples, under an
89 operating microscope, a 30 gauge needle, bent to an angle of 135 degrees,
90 bevel up, attached to a 20 ml syringe was passed from the scleral rim into the
91 corneal stroma and advanced to the centre of the disc. The needle was passed
92 close to the endothelial surface without perforating it. Air was injected with force
93 to overcome the tissue resistance, until a big bubble was formed. The type of
94 the bubble was determined, type-1 or type-2. The position of the needle tip was
95 kept constant in the centre of the sclera-corneal disc in mid stroma.

96 Pressure measurement: An electronic pressure gauge/converter device was used
97 (Keller, K-114, Winterthur, Switzerland). The tube from the device was linked to
98 the side arm of a 3-way cannula attached between the syringe and needle. The
99 injecting needle was attached to the front end of the cannula and a 20 ml
100 syringe to the other end. The device was also connected to a personal computer
101 (PC) via a USB port. The USB link also powered the device. Pressure readings
102 were recorded in real time and transmitted as serial RS485 half-duplex signals to
103 the PC where the pressure was displayed as a continuous trace on the screen by
104 the software associated with the K-114 device. (Figure 1) The pressure recorded
105 was that in the syringe during injection of air. In validation experiments, when
106 the needle was not inserted in tissue and the piston was advanced rapidly, the
107 pressure recorded was between 0 and 1, indicating that the resistance offered by
108 the needle was not relevant to the pressures measured (data not shown).

109 The maximum pressure required to create the bubble was recorded. The plunger
110 was then released and allowed to attain a stable position. The needle tip was
111 advanced to lie inside the BB and the bubble inflated till its wall was taut. The
112 pressure recorded at this point was taken as the base line intra-bubble pressure.
113 With the needle tip in the BB, the piston was pushed further with force until the
114 BB burst. This recorded the bursting pressure of the bubble (Figure 2 a, b).

115 Experiment to measure Volume:

116 As air leaked through multiple points along the circumference of the corneal
117 periphery a clamp was designed to block the holes and stop air leak. In 7
118 samples, the sclero-corneal discs were clamped in a circular clamp of 10mm
119 diameter that prevented air escape from the periphery. A 30 gauge needle
120 attached to a one millilitre syringe (internal diameter 5 mm) filled with air was

121 passed in to the corneal stroma from the scleral rim as described above. During
122 injection the maximum compression of air (position of piston) at the time air just
123 started to appear in the corneal stroma was recorded. The piston was held in
124 place until a type-1 BB was formed. The pressure on the piston was released and
125 piston allowed to reverse to a stable position. The volume of air lost in the
126 cornea was ascertained from the final position of the piston. The BB diameter
127 was measured with a pair of surgical callipers. The needle was then advanced
128 into the BB and all the air aspirated until the BB had completely collapsed. This
129 provided a measure of the volume of air in the big bubble. The pressure (above
130 atmosphere) in the syringe at the point where air started to emerge in the tissue
131 from the needle tip was deduced by the formula $P_1V_1 = P_2V_2$, where P1 is the
132 initial pressure (atmospheric) and V1 the initial volume (1ml) and P2 is the final
133 pressure (unknown) and V2 the final volume (mean 0.54ml, see results).

134 **Results:**

135 The average age of donors was 66 years (range; 52-80 years). There were 15
136 females and 7 males.

137 Pressure measurements: Twelve type-1 and 3 type-2 BB were obtained (table
138 2). The mean pressure attained to create a BB was 96.25+/- 21.61 kilopascal
139 (kpa) (range 90-130kpa). For type-1 BB the mean intra-bubble pressure was
140 10.16 +/- 3.65kpa (range 5.2-18kpa) and the bursting pressure was 66.65 +/-
141 18.65kpa (range 40-110kpa). The median bursting pressure was 68.5kpa (table
142 2). Accurate measurements of type-2 BB could not be obtained as when
143 advancing the needle into the bubble cavity, while the needle was still in the
144 stroma, the type-2 BB burst in one case and the DM disinserted (separated
145 along its peripheral attachment to the stroma) in one sector before the bubble

146 could be inflated enough to make the DM taut. The mean pressure at the time
147 the type-2 BB burst/disinserted was 14.77 +/- 2.44kpa (range 12.0-17.0kpa)
148 (table 2).

149 Volume measurements: In the bubble volume experiment, the maximum
150 compression of air required to create type-1 BB was 0.54 +/- 0.07 ml (range
151 0.5-0.7 ml), the volume of air lost in the cornea was 0.38 +/- 0.06 ml (range
152 0.3-.5 ml) and the average volume of the BB was 0.1 ml. The mean pressure in
153 the syringe at which air started to emerge in the tissue, as calculated from the
154 volume compression, was 131.82 +/- 50.58kpa (range 101.28 – 236.3kpa above
155 atmosphere). The pressures measured directly with the gauge and by this
156 method were not statistically significant (p= 0.25) (Figure 3).

157 Statistical methods: The data was normally distributed as confirmed by Levene's
158 test. Statistical analysis between two groups was performed by the unpaired
159 student t-test using Graphpad prism version 5.0. (Graphpad software, USA).
160 p<0.05 was considered statistically significant."

161

162 **Discussion:**

163 In DALK by the BB technique, when air is injected in the corneal stroma, a type-
164 1 BB forms by air cleaving in the plane of deep stroma and DL, with a posterior
165 displacement of DL and DM. The cleavage and displacement are related to the
166 pressure of air in the corneal stroma and in the BB. As the BB expands
167 posteriorly the intra bubble pressure is countered by the intraocular pressure,
168 which can rise up to 70 mm of mercury (authors' unpublished observations).
169 This counter pressure and the closed space within which the BB expands limits
170 the posterior expansion of the BB in the eye thus rupture of a type-1BB during
171 inflation is unlikely and has not been reported. However, when the type-1BB is

172 deflated and the corneal stroma anterior to it is removed, the DL + DM bulge
173 anteriorly to assume a convex dome shape. Any pressure applied to the DL+DM
174 from within the eye, as during the DALK-triple procedure, would cause the layers
175 to expand outward, into the atmosphere and theoretically reach a bursting point.
176 In this study we set out to ascertain the minimum and mean popping (bursting)
177 pressure of the layers to establish whether it would always be safe to perform
178 cataract surgery under DL+DM after creating a type-1BB.

179

180 The pressure converter K 114 allowed us to measure in real time the pressure at
181 the tip of the needle during the creation of a BB. On initiation of injection, air is
182 compressed in the syringe on account of the tissue resistance offered by the
183 corneal stroma at the site of the tip of the needle. Once this is overcome, air
184 starts to enter the stroma separating the lamellae and the intrastromal pressure
185 builds up as the cornea gets completely aerated. At a critical tissue pressure, the
186 air forces its way to the plane anterior to DL and cleaves this away from deep
187 stroma as a type-1BB. The volume of air required to achieve the critical tissue
188 pressure depends on the escape of air through the trabecular meshwork or
189 through distinct peripheral holes in the stroma, during injection (ref 7, 8, 9).

190 This confounder was eliminated by the use of the clamp, which prevented any
191 escape of air thus giving us an accurate measure of the mean tissue pressure
192 required to create a BB overcoming tissue resistance, which was 96.25 +/-
193 21.61 kpa. It has been recently demonstrated that air injected in the corneal
194 stroma follows a consistent path regardless of the location, direction of bevel
195 and depth of the needle tip in the stroma (ref 9)

196

197 Once a type-1BB was created the intra-bubble pressure was ascertained by
198 advancing the needle into the cavity of the BB. This measured 10.16 +/- 3.65
199 kpa. In the ex-vivo situation of this study, it was possible to expand the type-
200 1BB to its bursting point by continued forceful injection of air with the needle
201 positioned in the cavity of the bubble. This situation would simulate increased
202 intraocular pressure exerted on the layers during phacoemulsification carried out
203 under the layers (DALK-triple). The lowest pressure at which a type-1BB burst
204 was 40 kpa and the highest was 110 kpa. The mean bursting pressure was
205 66.65 +/- 18.65 kpa. Although we reported the bursting pressure in our original
206 paper (ref 5) we refined the measurement by placing the needle tip in the type-1
207 BB while increasing the pressure to bursting point. This approach eliminated any
208 variations induced by the resistance of the stroma to the passage of air. Any
209 effect of variable leakage of air from the periphery of the sclera-corneal was
210 prevented by the use of the clamp. In addition, the accuracy of the
211 measurements was enhanced by using the continuous digital pressure recording
212 device.

213

214 A number of studies have reported the variations in intraocular pressure during
215 phacoemulsification. By direct measurements during surgery Zhao Y et al found
216 that the IOP fluctuated from 13-96 mm Hg (1.8-13.5 kpa) (ref. 10). Khng C et al
217 state that IOP exceeded 60 mm Hg (8.4 kpa) and the highest IOP occurred
218 during hydro-dissection, viscoelastic injection and intraocular lens insertion (ref.
219 11). Vasavada V et al compared the impact of different fluidic parameters on
220 intraoperative IOP and found that the minimum IOP in the low and high
221 parameters groups was 35 mm Hg (4.9 kpa) and 34.5 mm Hg (4.8 kpa)
222 respectively, and the maximum IOP in the low and high parameters groups was

223 69 (9.7 kpa) and 85 (11.9 kpa) mm Hg respectively (ref. 12). In another study
224 Kamae KK et al monitored IOP during IOL implantation and found that the mean
225 and peak IOPs exceeded 60 mm Hg (8.4 kpa) during IOL implantation (ref. 13).
226 In comparison, the data on bursting pressure of the DL+DM generated in this
227 study show that the pressures attained during cataract surgery are several times
228 less than what is required to burst the layers under which phacoemulsification
229 can be carried out in the DALK-triple procedure. Even the lowest bursting
230 pressure had a safety margin of over 25 kpa (177.5 mm Hg) compared to the
231 highest pressure reached during phacoemulsification. This would indicate that
232 DALK-triple is a viable option with regard to the risk of inadvertent rupture of
233 the DL+DM layers intraoperatively.

234

235 When cataract and DALK surgery are required simultaneously; if the cornea is
236 clear, one could consider performing phacoemulsification as the first step and
237 DALK as the second step of the same procedure. However, when the cornea is
238 scarred to an extent that visualisation is poor, a triple-DALK would be the
239 preferred option. With triple-DALK, when air injection fails to produce a type-
240 1BB, manual dissection allows access to the plane between the deep stroma and
241 DL. Once the opaque cornea, related to the aeration of the stroma anterior the
242 DL is removed, the transparent DL allows phacoemulsification to be carried out.

243

244 We were able to create both type-1 and type-2BB as reported by Dua et al 2013
245 however the type-1BB was more consistent occurring in 86.4% of the 22 sclero-
246 corneal discs. The data provided in this study can help us develop an automated
247 system whereby we can produce big bubbles in vivo with improved consistency.

248

249 **ACKNOWLEDGMENTS**

250 The Royal College of Surgeons, Edinburgh, Scotland and Royal Blind Scotland.

251 Elizabeth C King Trust, Pittsburgh, USA.

252 **CONFLICT OF INTERESTS**

253 The authors declare no conflicting interest to this work

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268 **References**

- 269 1. Anwar M, Teichmann KD. Big-bubble technique to bare Descemet's membrane in anterior
270 lamellar keratoplasty. *Journal of cataract and refractive surgery* 2002 Mar; 28: 398-403.
- 271 2. Tan DT, Dart JK, Holland EJ, Kinoshita S. *Corneal transplantation* 2012 May 5; 379: 1749-
272 1761.
- 273 3. Espandar L, Carlson AN. Lamellar keratoplasty: a literature review. *Journal of*
274 *Ophthalmology*. 2013;2013:894319. Doi: 10.1155/2013/894319. Epub 2013 Oct 7
- 275 4. Maurino V, Allan BD, Stevens JD, Tuft SJ. Fixed dilated pupil (Urrets-Zavalía syndrome) after
276 air/gas injection after deep lamellar keratoplasty for keratoconus. *American journal of*
277 *ophthalmology* 2002 Feb; 133: 266-268.
- 278 5. Dua HS, Faraj LA, Said DG, Gray T, Lowe J. Human corneal anatomy redefined: a novel pre-
279 Descemet's layer (Dua's layer). *Ophthalmology* 2013 Sep; 120: 1778-1785.
- 280 6. Zaki AA, Elalfy MS, Said DG, Dua HS. Deep anterior lamellar keratoplasty--triple procedure: a
281 useful clinical application of the pre-Descemet's layer (Dua's layer). *Eye (London, England)* 2015 Mar;
282 29: 323-326.
- 283 7. Dua HS, Faraj LA, Said DG. Dua's layer: its discovery, characteristics and clinical applications.
284 In: Ángeles del Buey Sayas MA, Cristina Peris Martínez CP, eds. *Biomechanica y arquitectura corneal*
285 *Barcelona: Elsevier;2014:35-47.*
- 286 8. Dua HS, Faraj LA, Said DG. Dua's layer: discovery, characteristics, clinical applications,
287 controversy and potential relevance to glaucoma. *Expert Review of Ophthalmology* 2015; 10: 531-47.
- 288 9. Dua HS, Faraj LA, Kenawy MB. Dynamics of big bubble formation in deep anterior lamellar
289 keratoplasty (DALK) by the big bubble technique: In vitro studies. *ACTA Ophthalmologica 2017*; In
290 press.
- 291 10. Zhao Y, Li X, Tao A, Wang J, Lu F. Intraocular pressure and calculated diastolic ocular
292 perfusion pressure during three simulated steps of phacoemulsification in vivo. *Investigative*
293 *ophthalmology & visual science* 2009; 50: 2927-2931.

294 11. Khng C, Packer M, Fine IH, Hoffman RS, Moreira FB. Intraocular pressure during
295 phacoemulsification. *Journal of cataract and refractive surgery*. 2006; 32: 301-308.

296 12. Vasavada V, Raj SM, Praveen MR, Vasavada AR, Henderson BA, Asnani PK. Real-time
297 dynamic intraocular pressure fluctuations during microcoaxial phacoemulsification using different
298 aspiration flow rates and their impact on early postoperative outcomes: a randomized clinical trial.
299 *Journal of refractive surgery*. 2014; 30: 534-540.

300 13. Kamae KK, Werner L, Chang W, Johnson JT, Mamalis N. Intraocular pressure changes during
301 injection of microincision and conventional intraocular lenses through incisions smaller than 3.0 mm.
302 *Journal of cataract and refractive surgery*. 2009; 35: 1430-1436.

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318 **Titles and legends to figures**

319 Figure 1: (a) Pressure converter system K-144, (b) Real pressure record over time (red
320 graph), the temperature of the pressure sensor (blue graph), the maximum pressure
321 (pink graph), the minus pressure (green line).

322 Figure 2: (a) pressure change over time (red line) in T1BB. (b) Pressure change over
323 time (red line) in T2BB.

324 Figure 3: Compares the pressure calculated from the volume compression of the syringe
325 and that measured directly with gauge ($p = 0.25$).

Table 1: Donor details for the sclera-corneal discs used in the experiments

Sample No	Eye Bank No.	Type of big bubl	Sex	Age	Date of death	Cause of death
E1955	M20599B	T1BB	F	67	08/05/2014	Stroke
E2168	M21433B	T1BB	F	60	29/12/2014	Other (unknown)
E2182	M21468A	T1BB	F	58	07/01/2015	Cancer
E2183	M21468B	T1BB	F	58	07/01/2015	Cancer
E2246	M21715A	T1BB	F	69	15/03/2015	Chronic obstructive pulmonary disease
E2187	M21447B	T1BB	F	65	02/01/2015	Pending
E2385	M22280B	T1BB	M	73	29/06/2015	Respiratory failure
E2347	M22237A	T1BB	F	52	17/06/2015	Encephalopathy
E2278	M22072B	T1BB	F	80	07/05/2015	sepsis
E2276	M22016B	T1BB	M	74	01/05/2015	Brain damage hypoxia
E2275	M22016A	T1BB	M	74	01/05/2015	Brain damage hypoxia
E2309	M21828B	T1BB	M	72	02/04/2015	Cronic obstructive pulmonary disease
E2326	M22034A	T2BB	F	75	04/05/2015	Myocardial infarction
E2348	M22237B	T2BB	F	52	17/06/2015	Encephalopathy
E2384	M22333A	T2BB	F	68	14/07/2015	Myocardial infarction
E2677	M22933B	T1BB	F	81	29/12/2015	Myocardial infarction
E2675	M22956B	T1BB	M	53	02/01/2016	Unknown
E2674	M22956A	T1BB	M	53	02/01/2016	Unknown
E2678	M22913A	T1BB	F	44	14/12/2015	Intracranial heamorrhage
E2679	M22913B	T1BB	F	44	14/12/2015	Intracranial heamorrhage
E2829	M23226B	T1BB	M	80	14/03/2016	Cancer
E2836	M23301B	T1BB	F	88	03/04/2016	Old Age

Table 2: Measurements of the big bubble

	Sample No.	Diameter(mm)	Intrabubble pressure(Kpa)	Bursting pressure(kpa)	
T1BB	E1955	nm	nm	45	
	E2168	7	12	60	
	E2182	9	13	80	
	E2183	8.5	14	73	
	E2246	8.5	11.6	66	
	E2187	8.5	18	40	
	E2309	8.5	7.5	110	
	E2275	8.5	7.5	78	
	E2276	8.5	7.5	55	
	E2278	nm	5.2	71	
	E2347	8.5	6.8	76.8	
	E2385	8.5	8.7	45	
	T2BB	E2326	10	nm	17
		E2348	10.5	nm	12
E2384		10.5	nm	12.7	

Abbreviations: nm, not measured; Kpa, Kilopascal.





