

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

## **Abstract**

The Common Sense Model (CSM) is a useful framework for understanding adjustment (mood and treatment adherence) amongst survivors in the acute phase of stroke ( $\leq$ three-months). CSM stroke studies have thus far focused on the single outcomes, mood and medication adherence, neglecting other pertinent aspects of post-stroke recovery (i.e., Health-Related Quality of Life (HRQL) and disability). The purpose of this study was to examine relationships between baseline illness beliefs and three-month post-stroke HRQL, mood and disability. A longitudinal observational design was adopted, involving 50 survivors (mean age=66.9 years, 68% male). The primary outcome, HRQL, was measured using EQ-5D-5L. The secondary outcome, mood was measured using the Patient Health Questionnaire-9; and disability, using the Nottingham Extended Activities of Daily Living Scale. A stroke-specific version of the Illness Perception Questionnaire-Revised measured illness beliefs. Spearman's correlations showed that beliefs about the fluctuating effects of stroke ( $\rho=0.50$ ,  $p<0.001$ ) and perceptions of considerable distress at baseline were significantly associated with worse mood three-months post-stroke ( $\rho=0.41$ ,  $p<0.001$ ). Baseline illness beliefs were not significantly related to three-month post-stroke HRQL or disability. Despite being limited by a modest sample size, the findings reiterated the need for routine clinical assessment of mood immediately after stroke, and indicated that simultaneous measurement of timeline-cyclical beliefs and emotional representations may also be beneficial.

**Keywords:** Common Sense Model; Illness Beliefs; Stroke; Mood; Depression; Recovery; Disability; Health-Related Quality of Life.

22

## **Introduction**

23       The Common Sense Model (CSM) suggests that when individuals suffer illness, they  
24 experience a disequilibrium that they become motivated to resolve, and do so by constructing  
25 beliefs about their illness and treatment that guide how they cope with their condition  
26 (Leventhal, Meyer, & Nerenz, 1980). Illness beliefs have five core domains: ‘identity’ –  
27 beliefs about the label of illness; ‘timeline’ – beliefs about illness duration; ‘consequences’ –  
28 beliefs about illness severity/impact; ‘cure/control’ – beliefs about amenability to cure,  
29 prevention or treatment; and ‘causes’ – beliefs about internal (e.g., genes) and external (e.g.,  
30 germ or virus) causes of illness. These have been extended to include: ‘timeline-cyclical’ –  
31 beliefs of an episodic illness; ‘personal control’ and ‘treatment control’ – beliefs about own  
32 ability and that of treatment to manage the illness; ‘illness coherence’ – understanding of the  
33 illness; and ‘emotional representations’ – illness-related distress (Moss-Morris et al., 2002).

34       Eleven studies have thus far examined relationships between illness beliefs and the  
35 single post-stroke outcomes, mood and medication adherence (Ford, 2007; Johnston et al.,  
36 2007; Johnston, Morrison, Macwalter, & Partridge, 1999; Joice, Bonetti, MacWalter, &  
37 Morrison, 2003; Joice, Johnston, & Bonetti, 2002; Klinedinst, Dunbar, & Clark, 2012;  
38 O’Carroll, Chambers, Dennis, Sudlow, & Johnston, 2013; O’Carroll et al., 2011; Phillips,  
39 Diefenbach, Abrams, & Horowitz, 2015; Sjölander, Eriksson, & Glader, 2013; Twiddy,  
40 House, & Jones, 2012). These identified multiple illness beliefs that are significantly  
41 associated with post-stroke mood and medication non-adherence, including perceptions of a  
42 highly symptomatic condition; serious consequences; chronicity; fluctuating effects of stroke;  
43 inability of treatment to manage effects of stroke; poor disease understanding; and stroke-  
44 related distress.

45       This short report examines relationships between illness beliefs and mood, as well as  
46 other important markers of post-stroke recovery (HRQL and disability) that have been

47 defined by the International Classification of Functioning (ICF) framework for health and  
48 disability (World Health Organization, 2001), but have mostly been neglected in CSM stroke  
49 studies to date.

50

51

### **Methods and Materials**

52 We employed a longitudinal observational design, collecting data at baseline (after  
53 study enrolment) and three-months after stroke. Participants were recruited from acute stroke  
54 and rehabilitation wards and outpatient clinics in one hospital in the United Kingdom (UK).  
55 Inclusion criteria were adults (>18 years) with a confirmed diagnosis of acute stroke (within  
56 8-weeks) and sufficient language and cognitive ability to participate. Ethical approval was  
57 granted by the National Research Ethics Service Committee East Midlands – Leicester  
58 (13/EM/0392).

59

#### **Measures**

60 Our outcomes were defined according to ICF domains (‘impairments’ – problems or  
61 loss in body function; ‘activities’ – performance of a task or action; and ‘participation’ –  
62 involvement in a life situation) (World Health Organization, 2001).

63 HRQL (ICF Participation) was measured using EQ-5D-5L (Brooks, 1996). Patient  
64 Health Questionnaire-9 (PHQ-9) measured mood (ICF Impairments). We measured disability  
65 (ICF Activities) or ‘instrumental activities of daily living’ (such as shopping, cooking etc.)  
66 using the stroke-specific Nottingham Extended Activities of Daily Living Scale (Nouri &  
67 Lincoln, 1987). Illness beliefs were measured using a version of the IPQ-R adapted to stroke  
68 (Stroke IPQ-R) (Aujla, Vedhara, Walker, & Sprigg, 2018).

69 After providing written informed consent, we collected socio-demographic; medical  
70 and family history; clinical and lifestyle data. Participants also completed the EQ-5D-5L,  
71 PHQ-9, Nottingham Extended Activities of Daily Living Scale, and Stroke IPQ-R, which

72 were repeated at three-months post-stroke. Data were mostly collected via self-report, with  
73 exception of clinician-reported data (e.g. stroke severity) which were abstracted from medical  
74 records.

### 75 **Statistical Analysis**

76 The primary outcome was three-month post-stroke HRQL –a now prioritised outcome  
77 in acute stroke studies (Deshpande et al., 2011). The secondary outcomes were mood and  
78 disability. We estimated needing 55 participants to detect a correlation of 0.4 between illness  
79 beliefs and markers of post-stroke recovery (e.g., mood), with 80% power, alpha=0.05 and  
80 20% attrition.

81 Analyses were conducted using STATA 13 (StataCorp LP College Station, TX, USA).  
82 Statistical significance was assessed at the 5% level ( $p < 0.05$ ), and a Bonferroni adjustment  
83 corrected for multiple testing. We examined associations between illness beliefs and post-  
84 stroke HRQL, mood and disability using Spearman's rho ( $\rho$ ).

85

### 86 **Results**

87 Sample characteristics are summarised in Table 1 and elaborated elsewhere (Aujla,  
88 Walker, Sprigg, & Vedhara, 2018). In brief, 88 of 1085 patients assessed for eligibility over a  
89 12-month period were eligible and approached for participation. The main reasons for non-  
90 eligibility were non-stroke diagnosis ( $N = 249$ ) and stroke onset over 8 weeks before ( $N =$   
91  $186$ ). Fifty patients consented, with 16% attrition. Average age was 66.9 years ( $SD=14.5$   
92 years), with 68% males and 98% White-British ethnicity. Around 78% reported a first stroke  
93 and 18% a recurrence. The majority of participants reported few symptoms, but believed their  
94 stroke to be chronic, with fluctuating effects, greatly impacting on their lives, and leading to  
95 considerable distress, and despite having an unsatisfactory understanding (particularly of the  
96 causes) of their stroke, perceived that it was controllable.

97 Our analysis used complete cases. Following Bonferroni adjustment, Spearman's  
98 correlations showed that participants who perceived the effects of their stroke to be episodic  
99 ( $\rho=0.50$ ,  $p<0.001$ ) and causing considerable distress ( $\rho=0.41$ ,  $p<0.001$ ) at baseline also  
100 reported worse mood three-months after stroke. No significant correlations emerged between  
101 baseline illness beliefs and three-month post-stroke HRQL and disability (see Table 2).

102

103

### **Discussion**

104 We have shown that mood during the acute phase of recovery after stroke is affected by  
105 maladaptive beliefs about the episodic nature of stroke and stroke-related distress. These  
106 findings were consistent with prior CSM stroke studies, including Ford (2007), Joice et al.  
107 (2003), Klinedinst et al. (2012) and Twiddy et al. (2012). We also uniquely examined  
108 relationships between illness beliefs, HRQL and disability within the first three-months of  
109 stroke. It was surprising that significant associations did not emerge given findings from the  
110 wider CSM literature on physical illnesses (e.g., Damman, Liu, Kaptein, Rosendaal, and  
111 Kloppenburg (2014); Dalbeth et al. (2011); and Spain, Tubridy, Kilpatrick, Adams, and  
112 Holmes (2007)). We suspect that this is likely to relate to our sample. In addition to being  
113 modest in size and inevitably resulting in inadequate statistical power and inflated risk of type  
114 2 error, it also comprised highly functioning survivors of a less severe stroke. An important  
115 limitation of ours and prior CSM stroke studies.

116 CSM theory argues that illness beliefs form when people experience illness (Leventhal  
117 et al., 1980). This implies that if people do not experience symptoms (i.e., are functioning  
118 well post-illness), the health threat may not be considered enough of a problem for  
119 (mal)adaptive illness beliefs to manifest. In order to gain a more thorough picture of how  
120 illness beliefs relate to these specific aspects of post-stroke recovery, it may instead be better  
121 to examine patients most affected by stroke (i.e., survivors of more severe strokes).

122           However, this is a hard group to reach in acute stroke research (Newington & Metcalfe,  
123 2014). The post-stroke impairments that commonly affect these patients (e.g., paralysis,  
124 perceptual difficulties, and impaired cognition) undoubtedly limit their ability to engage with  
125 and provide informed consent for complex studies such as ours. Therefore, it is necessary for  
126 future research to consider ways other than questionnaires to elicit illness beliefs in stroke  
127 survivors with complex needs. One possibility is the ‘Talking Mats’ framework, which  
128 supports people with communication problems (including stroke survivors with aphasia) to  
129 express their views (Murphy, 2000; Murphy, Gray, van Achterberg, Wyke, & Cox, 2010).

130           In view of these limitations, our findings should be considered exploratory.  
131 Nonetheless, we have shown that the CSM may be a useful framework for understanding  
132 psychological adjustment during the acute phase of stroke, and in particular, that early post-  
133 stroke mood may be affected by maladaptive timeline-cyclical beliefs and emotional  
134 representations. These relationships were found even in survivors of a less severe stroke with  
135 little residual disability and mild depressive symptomatology. Therefore, our findings further  
136 emphasise an already recognised need to identify patients with low mood early after stroke  
137 and tie in with the most recent UK stroke clinical guidelines (Intercollegiate Stroke Working  
138 Party, 2016).

139           **Acknowledgements:** This research was undertaken as part of a PhD Studentship  
140 awarded to **FIRST AUTHOR** by the **NAME OF UNIVERSITY**. We would like to thank our  
141 participants for giving up their time to be involved in this research.

142           **Disclosures:** None of the authors have any conflicts of interest to declare.

143

144

### **References**

145 Aujla, N., Vedhara, K., Walker, M., & Sprigg, N. (2018). Evaluating a stroke-specific version  
146 of the Illness Perception Questionnaire-Revised, using the Think-Aloud method *J*  
147 *Health Psychol.* doi:10.1177/1359105318781942

- 148 Aujla, N., Walker, M., Sprigg, N., & Vedhara, K. (2018). Do individual versus illness belief  
 149 schema differ in the prediction of post-stroke recovery? *J Health Psychol.*  
 150 doi:10.1177/1359105318785446
- 151 Brooks, R. (1996). EuroQol: the current state of play. *Health Policy*, 37(1), 53-72.  
 152 doi:10.1016/0168-8510(96)00822-6
- 153 Dalbeth, N., Petrie, K. J., House, M., Chong, J., Leung, W., Chegudi, R., . . . Taylor, W. J.  
 154 (2011). Illness perceptions in patients with gout and the relationship with progression  
 155 of musculoskeletal disability. *Arthritis Care and Research*, 63(11), 1605-1612.  
 156 doi:10.1002/acr.20570
- 157 Damman, W., Liu, R., Kaptein, A. A., Rosendaal, F. R., & Kloppenburg, M. (2014).  
 158 Negative illness perceptions are associated with short-term disability in patients with  
 159 hand osteoarthritis. *Osteoarthritis and Cartilage*, 22, S200-S201.  
 160 doi:10.1016/j.joca.2014.02.383
- 161 Ford, C. (2007). *The relationship between beliefs about stroke and post-stroke depression*  
 162 (Doctor of Clinical Psychology), University of East Anglia,
- 163 Intercollegiate Stroke Working Party. (2016). *National clinical guideline for stroke. Fifth*  
 164 *edition*. London, United Kingdom: Royal College of Physicians Retrieved from  
 165 [https://www.strokeaudit.org/SupportFiles/Documents/Guidelines/2016-National-](https://www.strokeaudit.org/SupportFiles/Documents/Guidelines/2016-National-Clinical-Guideline-for-Stroke-5t-(1).aspx)  
 166 [Clinical-Guideline-for-Stroke-5t-\(1\).aspx](https://www.strokeaudit.org/SupportFiles/Documents/Guidelines/2016-National-Clinical-Guideline-for-Stroke-5t-(1).aspx)
- 167 Johnston, M., Bonetti, D., Joice, S., Pollard, B., Morrison, V., Francis, J. J., & MacWalter, R.  
 168 (2007). Recovery from disability after stroke as a target for a behavioural  
 169 intervention: results of a randomized controlled trial. *Disability & Rehabilitation*,  
 170 29(14), 1117-1127. doi:10.1080/03323310600950411
- 171 Johnston, M., Morrison, V., Macwalter, R., & Partridge, C. (1999). Perceived control, coping  
 172 and recovery from disability following stroke. *Psychology & Health*, 14(2), 181-192.  
 173 doi:10.1080/08870449908407322
- 174 Joice, S., Bonetti, D., MacWalter, R., & Morrison, V. (2003). *Illness representations and*  
 175 *distress in stroke patients: an analysis using the SRM*. Paper presented at the British  
 176 Psychological Society, Division of Health Psychology Conference.
- 177 Joice, S., Johnston, M., & Bonetti, D. (2002). *Using Leventhal's self-regulation model to*  
 178 *explore non-adherence to a workbook intervention for stroke patients*. Paper  
 179 presented at the British Psychological Society, Division of Health Psychology  
 180 Conference.
- 181 Klinedinst, N. J., Dunbar, S. B., & Clark, P. C. (2012). Stroke survivor and informal  
 182 caregiver perceptions of post-stroke depressive symptoms. *The Journal of*  
 183 *Neuroscience Nursing*, 44(2), 72-81. doi:10.1097/JNN.0b013e3182477944
- 184 Leventhal, H., Meyer, D., & Nerenz, D. R. (1980). The common sense representation of  
 185 illness danger. In S. Rachman (Ed.), *Contributions to medical psychology* (Vol. 2, pp.  
 186 7-30). New York, United States of America: Pergamon.
- 187 Moss-Morris, R., Weinman, J., Petrie, K. J., Horne, R., Cameron, L. D., & Buick, D. (2002).  
 188 The revised illness perception questionnaire (IPQ-R). *Psychology & Health*, 17(1), 1-  
 189 16. doi:10.1080/08870440290001494
- 190 Murphy, J. (2000). Enabling people with aphasia to discuss quality of life. *British Journal of*  
 191 *Therapy and Rehabilitation*, 7(11), 454-458. doi:10.12968/bjtr.2000.7.11.13835
- 192 Murphy, J., Gray, C. M., van Achterberg, T., Wyke, S., & Cox, S. (2010). The effectiveness  
 193 of the Talking Mats framework in helping people with dementia to express their  
 194 views on well-being. *Dementia*, 9(4), 454-472. doi:10.1177/1471301210381776
- 195 Newington, L., & Metcalfe, A. (2014). Factors influencing recruitment to research:  
 196 qualitative study of the experiences and perceptions of research teams. *BMC Medical*  
 197 *Research Methodology*, 14(1), 10. doi:10.1186/1471-2288-14-10

- 198 Nouri, F. M., & Lincoln, N. B. (1987). An extended activities of daily living scale for stroke  
199 patients. *Clinical Rehabilitation*, *1*(4), 301-305. doi:10.1177/026921558700100409
- 200 O'Carroll, R. E., Chambers, J. A., Dennis, M., Sudlow, C., & Johnston, M. (2013). Improving  
201 adherence to medication in stroke survivors: a pilot randomised controlled trial.  
202 *Annals of Behavioural Medicine*, *46*(3), 358-368. doi:10.1007/s12160-013-9515-5
- 203 O'Carroll, R. E., Whittaker, J., B. H., Johnston, M., Sudlow, C., & Dennis, M. (2011).  
204 Predictors of adherence to secondary preventive medication in stroke patients. *Annals*  
205 *of Behavioral Medicine*, *41*, 383-390. doi:10.1007/s12160-010-9257-6
- 206 Phillips, L. A., Diefenbach, M. A., Abrams, J., & Horowitz, C. R. (2015). Stroke and TIA  
207 survivors' cognitive beliefs and affective responses regarding treatment and future  
208 stroke risk differentially predict medication adherence and categorised stroke risk.  
209 *Psychology & Health*, *30*(2), 218-232. doi:10.1080/08870446.2014.964237
- 210 Sjölander, M., Eriksson, M., & Glader, E.-L. (2013). The association between patients'  
211 beliefs about medicines and adherence to drug treatment after stroke: a cross-sectional  
212 questionnaire survey. *BMJ Open*, *3*(9). doi:10.1136/bmjopen-2013-003551
- 213 Spain, L. A., Tubridy, N., Kilpatrick, T. J., Adams, S. J., & Holmes, A. C. N. (2007). Illness  
214 perception and health-related quality of life in multiple sclerosis. *Acta Neurologica*  
215 *Scandinavica*, *116*(5), 293-299. doi:10.1111/j.1600-0404.2007.00895.x
- 216 Twiddy, M., House, A., & Jones, F. (2012). The association between discrepancy in illness  
217 representations on distress in stroke patients and carers. *Journal of Psychosomatic*  
218 *Research*, *72*, 220-225. doi:10.1016/j.jpsychores.2011.12.004
- 219 World Health Organization. (2001). *International classification of functioning, disability and*  
220 *health*. Geneva, Switzerland: World Health Organization Retrieved from  
221 <http://www.who.int/classifications/icf/en/>  
222



**Table 1: Baseline characteristics of the study sample**

	N Mean (SD)/Frequency (%), unless otherwise stated
<b>Socio-Demographics</b>	
Age	N=50 66.9 (14.5)
Sex-Male	N=50 34 (68.0%)
Ethnic group-White	N=50 49 (98.0%)
University or higher education	N=44 9 (20.5%)
Employment status	N=46
Unemployed	6 (13.0%)
Employed full-time	7 (15.2%)
Employed part-time	3 (6.5%)
Self-employed	5 (10.9%)
Retired	25 (54.4%)
IMD rank*	N=44 Median=20706.5 (IQR=17158.0)
IMD decile*	N=44 Median=7 (IQR=6)
<b>Medical history</b>	
National Institute of Health Stroke Scale score¥	N=44 Median=2 (IQR=4.5)
Pre-morbid Modified Rankin Scale score¥¥	N=48 0 (0)
Previous stroke	N=46 36 (78.3)
Previous TIA	N=46 13 (28.3)
History of heart attack	N=46 6 (13.0%)
History of hypertension	N=46 31 (67.4%)
History of high cholesterol	N=46 24 (52.2%)
History of atrial fibrillation	N=46 10 (21.7%)
History of blood clots	N=46 5 (10.9%)
History of angina	N=46 6 (13.0%)
History of diabetes	N=46 11 (23.9%)
History of depression	N=46 11 (23.9%)
History of anxiety	N=46 9 (19.6%)
Co-morbidities	N=49 34 (69.4%)
<b>Family history-first degree relative (mother, father, sibling)</b>	
History of stroke	N=45 15 (33.0%)
History of TIA	N=46 4 (8.7%)
<b>Clinical data</b>	
Systolic blood pressure (mm/HG)	N=48

	147.6 (33.7)
Diastolic blood pressure (mm/HG)	N=48 78.6 (20.0)
Blood glucose (mmol/L)	N=37 Median = 6.6 (IQR=2.8)
Total cholesterol (mmol/L)	N=44 4.74 (1.30)
HDL cholesterol (mmol/L)	N=40 Median = 1.3 (IQR=0.6)
LDL cholesterol (mmol/L)	N=38 Median = 2.7 (IQR=1.9)
BMI (kg/m <sup>2</sup> )	N=40 Median = 28 (IQR=9.7)
<b>Lifestyle</b>	
Current smoking status	N=41
Non/never smoked	18 (39.1%)
Ex-smoker	24 (52.2%)
Current smoker	4 (8.7%)
Number smoked daily	N=22 10 (13)
Units of beer	N=41 0 (7)
Units of wine	N=41 0 (2)
Units of spirits	N=41 0 (0)
30-minutes of exercise x4 times a week	N=41 36 (78.3%)
Low-fat diet	N=41 24 (52.2%)
Low-sugar diet	N=41 29 (63.0%)
Low-salt diet	N=41 29 (64.4%)

Symbols and abbreviations: \*: Computed using postcode data collected from participants; ¥: High NIHSS scores indicate a more severe stroke; ¥¥: High Modified Rankin Scale scores indicate greater disability; ¥¥¥: High Barthel Index scores indicate greater independence; ¥¥¥¥: BMI: Body mass index; HDL; High Density Lipoprotein; IMD: Index of Multiple Deprivation; IQR: Interquartile range; LDL; Low Density Lipoprotein; SD: Standard deviation; TIA: Transient Ischaemic Attack

**Table 2. Correlation matrix for baseline illness belief domains and follow-up markers of recovery (N=41)**

	Identity	Timeline acute-chronic	Timeline-cyclical	Consequences	Personal control	Treatment control	Illness coherence	Emotional representations	EQ-5D-5L Descriptive System - Index score	EQ-5D-5L 'Your health today' VAS score	Mood	Nottingham Extended ADL
<b>Identity</b>												
<b>Timeline acute-chronic</b>	0.47 p<.01											
<b>Timeline-cyclical</b>	<u>0.62</u> p<.001 *	0.26 p=0.10										
<b>Consequences</b>	<u>0.66</u> p<.001 *	0.45 p<.01	0.35 p<.05									
<b>Personal control</b>	0.19 p=0.23	-0.19 p=0.22	0.12 p=0.45	0.06 p=0.73								
<b>Treatment control</b>	-0.04 p=0.78	-0.17 p=0.30	-0.12 p=0.47	-0.00 p=0.99	0.30 p=0.05							
<b>Illness coherence</b>	0.17 p=0.28	0.00 p=0.99	-0.09 p=0.57	0.11 p=0.49	0.30 p=0.06	0.11 p=0.48						
<b>Emotional representations</b>	<u>0.56</u> p<.001 *	0.27 p=0.09	<u>0.51</u> p<.001*	<u>0.63</u> p<.001*	-0.07 p=0.65	-0.16 p=0.30	-0.00 p=0.98					
<b>EQ-5D-5L Descriptive System - Index score</b>	-0.27 p=0.09	-0.41 p<.01	-0.34 p<.05	-0.19 p=0.24	0.17 p=0.29	-0.11 p=0.49	0.26 p=0.10	-0.26 p=0.10				
<b>EQ-5D-5L 'Your health today' VAS score</b>	-0.11 p=0.49	-0.22 p=0.18	-0.28 p=0.08	-0.10 p=0.53	0.09 p=0.58	0.06 p=0.69	0.17 p=0.30	-0.27 p=0.09	<u>0.51</u> p<.001*			
<b>Mood</b>	0.26 p=0.10	0.04 p=0.80	<u>0.50</u> p<.001*	0.28 p=0.07	-0.06 p=0.71	-0.06 p=0.69	-0.13 p=0.43	<u>0.41</u> p<.001*	-0.21 p=0.18	-0.20 p=0.22		
<b>Nottingham Extended ADL</b>	-0.18 p=0.26	-0.27 p=0.09	-0.02 p=0.90	-0.04 p=0.77	0.02 p=0.88	-0.17 p=0.29	-0.27 p=0.09	-0.03 p=0.84	0.49 p<.01	0.32 p<.05	0.05 p=0.78	

Symbols and abbreviations: \*: P-value significant at the Bonferroni-adjusted significance level ( $p < 0.002$ ); ADL: Activities of Daily Living; VAS: Visual Analogue Scale