

Measurement of color flow in $t\bar{t}$ events from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

V. M. Abazov,³⁵ B. Abbott,⁷² B. S. Acharya,²⁹ M. Adams,⁴⁸ T. Adams,⁴⁶ G. D. Alexeev,³⁵ G. Alkhazov,³⁹ A. Alton,^{60,*} G. Alverson,⁵⁹ G. A. Alves,² L. S. Ancu,³⁴ M. Aoki,⁴⁷ M. Arov,⁵⁷ A. Askew,⁴⁶ B. Åsman,⁴⁰ O. Atramentov,⁶⁴ C. Avila,⁸ J. BackusMayer,⁷⁹ F. Badaud,¹³ L. Bagby,⁴⁷ B. Baldin,⁴⁷ D. V. Bandurin,⁴⁶ S. Banerjee,²⁹ E. Barberis,⁵⁹ P. Baringer,⁵⁵ J. Barreto,³ J. F. Bartlett,⁴⁷ U. Bassler,¹⁸ V. Bazterra,⁴⁸ S. Beale,⁶ A. Bean,⁵⁵ M. Begalli,³ M. Begel,⁷⁰ C. Belanger-Champagne,⁴⁰ L. Bellantoni,⁴⁷ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²² I. Bertram,⁴¹ M. Besançon,¹⁸ R. Beuselinck,⁴² V. A. Bezzubov,³⁸ P. C. Bhat,⁴⁷ V. Bhatnagar,²⁷ G. Blazey,⁴⁹ S. Blessing,⁴⁶ K. Bloom,⁶³ A. Boehnlein,⁴⁷ D. Boline,⁶⁹ T. A. Bolton,⁵⁶ E. E. Boos,³⁷ G. Borissov,⁴¹ T. Bose,⁵⁸ A. Brandt,⁷⁵ O. Brandt,²³ R. Brock,⁶¹ G. Brooijmans,⁶⁷ A. Bross,⁴⁷ D. Brown,¹⁷ J. Brown,¹⁷ X. B. Bu,⁴⁷ M. Buehler,⁷⁸ V. Buescher,²⁴ V. Bunichev,³⁷ S. Burdin,^{41,†} T. H. Burnett,⁷⁹ C. P. Buszello,⁴⁰ B. Calpas,¹⁵ E. Camacho-Pérez,³² M. A. Carrasco-Lizarraga,⁵⁵ B. C. K. Casey,⁴⁷ H. Castilla-Valdez,³² S. Chakrabarti,⁶⁹ D. Chakraborty,⁴⁹ K. M. Chan,⁵³ A. Chandra,⁷⁷ G. Chen,⁵⁵ S. Chevalier-Théry,¹⁸ D. K. Cho,⁷⁴ S. W. Cho,³¹ S. Choi,³¹ B. Choudhary,²⁸ T. Christoudias,⁴² S. Cihangir,⁴⁷ D. Claes,⁶³ J. Clutter,⁵⁵ M. Cooke,⁴⁷ W. E. Cooper,⁴⁷ M. Corcoran,⁷⁷ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ A. Croc,¹⁸ D. Cutts,⁷⁴ A. Das,⁴⁴ G. Davies,⁴² K. De,⁷⁵ S. J. de Jong,³⁴ E. De La Cruz-Burelo,³² F. Déliot,¹⁸ M. Demarteau,⁴⁷ R. Demina,⁶⁸ D. Denisov,⁴⁷ S. P. Denisov,³⁸ S. Desai,⁴⁷ K. DeVaughan,⁶³ H. T. Diehl,⁴⁷ M. Diesburg,⁴⁷ A. Dominguez,⁶³ T. Dorland,⁷⁹ A. Dubey,²⁸ L. V. Dudko,³⁷ D. Duggan,⁶⁴ A. Duperrin,¹⁵ S. Dutt,²⁷ A. Dyshkant,⁴⁹ M. Eads,⁶³ D. Edmunds,⁶¹ J. Ellison,⁴⁵ V. D. Elvira,⁴⁷ Y. Enari,¹⁷ H. Evans,⁵¹ A. Evdokimov,⁷⁰ V. N. Evdokimov,³⁸ G. Facini,⁵⁹ T. Ferbel,⁶⁸ F. Fiedler,²⁴ F. Filthaut,³⁴ W. Fisher,⁶¹ H. E. Fisk,⁴⁷ M. Fortner,⁴⁹ H. Fox,⁴¹ S. Fuess,⁴⁷ T. Gadfort,⁷⁰ A. Garcia-Bellido,⁶⁸ V. Gavrilov,³⁶ P. Gay,¹³ W. Geist,¹⁹ W. Geng,^{15,61} D. Gerbaudo,⁶⁵ C. E. Gerber,⁴⁸ Y. Gershtein,⁶⁴ G. Ginther,^{47,68} G. Golovanov,³⁵ A. Goussiou,⁷⁹ P. D. Grannis,⁶⁹ S. Greder,¹⁹ H. Greenlee,⁴⁷ Z. D. Greenwood,⁵⁷ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,¹⁸ S. Grünendahl,⁴⁷ M. W. Grünewald,³⁰ F. Guo,⁶⁹ G. Gutierrez,⁴⁷ P. Gutierrez,⁷² A. Haas,^{67,‡} S. Hagopian,⁴⁶ J. Haley,⁵⁹ L. Han,⁷ K. Harder,⁴³ A. Harel,⁶⁸ J. M. Hauptman,⁵⁴ J. Hays,⁴² T. Head,⁴³ T. Hebbeker,²¹ D. Hedin,⁴⁹ H. Hegab,⁷³ A. P. Heinson,⁴⁵ U. Heintz,⁷⁴ C. Hensel,²³ I. Heredia-De La Cruz,³² K. Herner,⁶⁰ M. D. Hildreth,⁵³ R. Hirosky,⁷⁸ T. Hoang,⁴⁶ J. D. Hobbs,⁶⁹ B. Hoeneisen,¹² M. Hohlfield,²⁴ S. Hossain,⁷² Z. Hubacek,^{10,18} N. Huske,¹⁷ V. Hynek,¹⁰ I. Iashvili,⁶⁶ R. Illingworth,⁴⁷ A. S. Ito,⁴⁷ S. Jabeen,⁷⁴ M. Jaffré,¹⁶ S. Jain,⁶⁶ D. Jamin,¹⁵ R. Jesik,⁴² K. Johns,⁴⁴ M. Johnson,⁴⁷ D. Johnston,⁶³ A. Jonckheere,⁴⁷ P. Jonsson,⁴² J. Joshi,²⁷ A. Juste,^{47,§} K. Kaadze,⁵⁶ E. Kajfasz,¹⁵ D. Karmanov,³⁷ P. A. Kasper,⁴⁷ I. Katsanos,⁶³ R. Kehoe,⁷⁶ S. Kermiche,¹⁵ N. Khalatyan,⁴⁷ A. Khanov,⁷³ A. Kharchilava,⁶⁶ Y. N. Kharzheev,³⁵ D. Khatidze,⁷⁴ M. H. Kirby,⁵⁰ J. M. Kohli,²⁷ A. V. Kozelov,³⁸ J. Kraus,⁶¹ A. Kumar,⁶⁶ A. Kupco,¹¹ T. Kurča,²⁰ V. A. Kuzmin,³⁷ J. Kvita,⁹ S. Lammers,⁵¹ G. Landsberg,⁷⁴ P. Lebrun,²⁰ H. S. Lee,³¹ S. W. Lee,⁵⁴ W. M. Lee,⁴⁷ J. Lellouch,¹⁷ L. Li,⁴⁵ Q. Z. Li,⁴⁷ S. M. Lietti,⁵ J. K. Lim,³¹ D. Lincoln,⁴⁷ J. Linnemann,⁶¹ V. V. Lipaev,³⁸ R. Lipton,⁴⁷ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,³⁹ M. Lokajicek,¹¹ P. Love,⁴¹ H. J. Lubatti,⁷⁹ R. Luna-Garcia,^{32,||} A. L. Lyon,⁴⁷ A. K. A. Maciel,² D. Mackin,⁷⁷ R. Madar,¹⁸ R. Magaña-Villalba,³² S. Malik,⁶³ V. L. Malyshev,³⁵ Y. Maravin,⁵⁶ J. Martínez-Ortega,³² R. McCarthy,⁶⁹ C. L. McGivern,⁵⁵ M. M. Meijer,³⁴ A. Melnitchouk,⁶² D. Menezes,⁴⁹ P. G. Mercadante,⁴ M. Merkin,³⁷ A. Meyer,²¹ J. Meyer,²³ F. Miconi,¹⁹ N. K. Mondal,²⁹ G. S. Muanza,¹⁵ M. Mulhearn,⁷⁸ E. Nagy,¹⁵ M. Naimuddin,²⁸ M. Narain,⁷⁴ R. Nayyar,²⁸ H. A. Neal,⁶⁰ J. P. Negret,⁸ P. Neustroev,³⁹ S. F. Novaes,⁵ T. Nunnemann,²⁵ G. Obrant,³⁹ J. Orduna,³² N. Osman,⁴² J. Osta,⁵³ G. J. Otero y Garzón,¹ M. Owen,⁴³ M. Padilla,⁴⁵ M. Pangilinan,⁷⁴ N. Parashar,⁵² V. Parihar,⁷⁴ S. K. Park,³¹ J. Parsons,⁶⁷ R. Partridge,^{74,‡} N. Parua,⁵¹ A. Patwa,⁷⁰ B. Penning,⁴⁷ M. Perfilov,³⁷ K. Peters,⁴³ Y. Peters,⁴³ G. Petrillo,⁶⁸ P. Pétroff,¹⁶ R. Piegaia,¹ J. Piper,⁶¹ M.-A. Pleier,⁷⁰ P. L. M. Podesta-Lerma,^{32,¶} V. M. Podstavkov,⁴⁷ M.-E. Pol,² P. Polozov,³⁶ A. V. Popov,³⁸ M. Prewitt,⁷⁷ D. Price,⁵¹ S. Protopopescu,⁷⁰ J. Qian,⁶⁰ A. Quadt,²³ B. Quinn,⁶² M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴¹ I. Razumov,³⁸ P. Renkel,⁷⁶ M. Rijssenbeek,⁶⁹ I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷³ M. Rominsky,⁴⁷ C. Royon,¹⁸ P. Rubinov,⁴⁷ R. Ruchti,⁵³ G. Safronov,³⁶ G. Sajot,¹⁴ A. Sánchez-Hernández,³² M. P. Sanders,²⁵ B. Sanghi,⁴⁷ A. S. Santos,⁵ G. Savage,⁴⁷ L. Sawyer,⁵⁷ T. Scanlon,⁴² R. D. Schamberger,⁶⁹ Y. Scheglov,³⁹ H. Schellman,⁵⁰ T. Schliephake,²⁶ S. Schlobohm,⁷⁹ C. Schwanenberger,⁴³ R. Schwienhorst,⁶¹ J. Sekaric,⁵⁵ H. Severini,⁷² E. Shabalina,²³ V. Shary,¹⁸ A. A. Shchukin,³⁸ R. K. Shivpuri,²⁸ V. Simak,¹⁰ V. Sirotenko,⁴⁷ P. Skubic,⁷² P. Slattery,⁶⁸ D. Smirnov,⁵³ K. J. Smith,⁶⁶ G. R. Snow,⁶³ J. Snow,⁷¹ S. Snyder,⁷⁰ S. Söldner-Rembold,⁴³ L. Sonnenschein,²¹ A. Sopczak,⁴¹ M. Sosebee,⁷⁵ K. Soustruznik,⁹ B. Spurlock,⁷⁵ J. Stark,¹⁴ V. Stolin,³⁶ D. A. Stoyanova,³⁸ M. Strauss,⁷² D. Strom,⁴⁸ L. Stutte,⁴⁷ L. Suter,⁴³ P. Svoisky,⁷² M. Takahashi,⁴³ A. Tanasijczuk,¹ W. Taylor,⁶ M. Titov,¹⁸ V. V. Tokmenin,³⁵ Y.-T. Tsai,⁶⁸ D. Tsybychev,⁶⁹ B. Tuchming,¹⁸ C. Tully,⁶⁵ P. M. Tuts,⁶⁷ L. Uvarov,³⁹ S. Uvarov,³⁹ S. Uzunyan,⁴⁹ R. Van Kooten,⁵¹ W. M. van Leeuwen,³³ N. Varelas,⁴⁸ E. W. Varnes,⁴⁴ I. A. Vasilyev,³⁸ P. Verdier,²⁰ L. S. Vertogradov,³⁵ M. Verzocchi,⁴⁷ M. Vesterinen,⁴³ D. Vilanova,¹⁸

P. Vint,⁴² P. Vokac,¹⁰ H. D. Wahl,⁴⁶ M. H. L. S. Wang,⁶⁸ J. Warchol,⁵³ G. Watts,⁷⁹ M. Wayne,⁵³ M. Weber,^{47,**}
 L. Welty-Rieger,⁵⁰ A. White,⁷⁵ D. Wicke,²⁶ M. R. J. Williams,⁴¹ G. W. Wilson,⁵⁵ S. J. Wimpenny,⁴⁵ M. Wobisch,⁵⁷
 D. R. Wood,⁵⁹ T. R. Wyatt,⁴³ Y. Xie,⁴⁷ C. Xu,⁶⁰ S. Yacoob,⁵⁰ R. Yamada,⁴⁷ W.-C. Yang,⁴³ T. Yasuda,⁴⁷
 Y. A. Yatsunenko,³⁵ Z. Ye,⁴⁷ H. Yin,⁴⁷ K. Yip,⁷⁰ S. W. Youn,⁴⁷ J. Yu,⁷⁵ S. Zelitch,⁷⁸ T. Zhao,⁷⁹
 B. Zhou,⁶⁰ J. Zhu,⁶⁰ M. Zielinski,⁶⁸ D. Zieminska,⁵¹ and L. Zivkovic⁷⁴

(D0 Collaboration)

- ¹Universidad de Buenos Aires, Buenos Aires, Argentina
²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil
³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil
⁴Universidade Federal do ABC, Santo André, Brazil
⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil
⁶Simon Fraser University, Vancouver, British Columbia, and York University, Toronto, Ontario, Canada
⁷University of Science and Technology of China, Hefei, People's Republic of China
⁸Universidad de los Andes, Bogotá, Colombia
⁹Charles University, Faculty of Mathematics and Physics, Center for Particle Physics, Prague, Czech Republic
¹⁰Czech Technical University in Prague, Prague, Czech Republic
¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic
¹²Universidad San Francisco de Quito, Quito, Ecuador
¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France
¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France
¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France
¹⁶LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France
¹⁷LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France
¹⁸CEA, Irfu, SPP, Saclay, France
¹⁹IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France
²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France
²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany
²²Physikalisches Institut, Universität Freiburg, Freiburg, Germany
²³II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany
²⁴Institut für Physik, Universität Mainz, Mainz, Germany
²⁵Ludwig-Maximilians-Universität München, München, Germany
²⁶Fachbereich Physik, Bergische Universität Wuppertal, Wuppertal, Germany
²⁷Panjab University, Chandigarh, India
²⁸Delhi University, Delhi, India
²⁹Tata Institute of Fundamental Research, Mumbai, India
³⁰University College Dublin, Dublin, Ireland
³¹Korea Detector Laboratory, Korea University, Seoul, Korea
³²CINVESTAV, Mexico City, Mexico
³³FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
³⁴Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁵Joint Institute for Nuclear Research, Dubna, Russia
³⁶Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁷Moscow State University, Moscow, Russia
³⁸Institute for High Energy Physics, Protvino, Russia
³⁹Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴⁰Stockholm University, Stockholm and Uppsala University, Uppsala, Sweden
⁴¹Lancaster University, Lancaster LA1 4YB, United Kingdom
⁴²Imperial College London, London SW7 2AZ, United Kingdom
⁴³The University of Manchester, Manchester M13 9PL, United Kingdom
⁴⁴University of Arizona, Tucson, Arizona 85721, USA
⁴⁵University of California Riverside, Riverside, California 92521, USA
⁴⁶Florida State University, Tallahassee, Florida 32306, USA
⁴⁷Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁴⁸University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁴⁹Northern Illinois University, DeKalb, Illinois 60115, USA
⁵⁰Northwestern University, Evanston, Illinois 60208, USA
⁵¹Indiana University, Bloomington, Indiana 47405, USA

⁵²Purdue University Calumet, Hammond, Indiana 46323, USA⁵³University of Notre Dame, Notre Dame, Indiana 46556, USA⁵⁴Iowa State University, Ames, Iowa 50011, USA⁵⁵University of Kansas, Lawrence, Kansas 66045, USA⁵⁶Kansas State University, Manhattan, Kansas 66506, USA⁵⁷Louisiana Tech University, Ruston, Louisiana 71272, USA⁵⁸Boston University, Boston, Massachusetts 02215, USA⁵⁹Northeastern University, Boston, Massachusetts 02115, USA⁶⁰University of Michigan, Ann Arbor, Michigan 48109, USA⁶¹Michigan State University, East Lansing, Michigan 48824, USA⁶²University of Mississippi, University, Mississippi 38677, USA⁶³University of Nebraska, Lincoln, Nebraska 68588, USA⁶⁴Rutgers University, Piscataway, New Jersey 08855, USA⁶⁵Princeton University, Princeton, New Jersey 08544, USA⁶⁶State University of New York, Buffalo, New York 14260, USA⁶⁷Columbia University, New York, New York 10027, USA⁶⁸University of Rochester, Rochester, New York 14627, USA⁶⁹State University of New York, Stony Brook, New York 11794, USA⁷⁰Brookhaven National Laboratory, Upton, New York 11973, USA⁷¹Langston University, Langston, Oklahoma 73050, USA⁷²University of Oklahoma, Norman, Oklahoma 73019, USA⁷³Oklahoma State University, Stillwater, Oklahoma 74078, USA⁷⁴Brown University, Providence, Rhode Island 02912, USA⁷⁵University of Texas, Arlington, Texas 76019, USA⁷⁶Southern Methodist University, Dallas, Texas 75275, USA⁷⁷Rice University, Houston, Texas 77005, USA⁷⁸University of Virginia, Charlottesville, Virginia 22901, USA⁷⁹University of Washington, Seattle, Washington 98195, USA

(Received 4 January 2011; published 10 May 2011)

We present the first measurement of the color representation of the hadronically decaying W boson in $t\bar{t}$ events, from 5.3 fb^{-1} of integrated luminosity collected with the D0 experiment. A novel calorimeter-based vectorial variable, “jet pull,” is used, sensitive to the color-flow structure of the final state. We find that the fraction of uncolored W bosons is $0.56 \pm 0.42(\text{stat} + \text{syst})$, in agreement with the standard model.

DOI: 10.1103/PhysRevD.83.092002

PACS numbers: 12.38.Qk, 12.38.Aw, 14.65.Ha

Color charge is conserved in quantum chromodynamics, the theory that describes strong interactions [1]. At leading order in the strong coupling constant α_s , color can be traced from initial partons to final-state partons in high-energy hadron collisions. Two final-state partons on the same color-flow line are “color connected” and attracted by the strong force. As these colored states shower, the potential energy of the strong force between them is released in the form of hadrons. Thus, knowledge of the color connections between jets can serve as a powerful tool for separating processes that otherwise appear similar. For example, in the decay of a Higgs (H) boson to a pair of bottom (b) quarks, the two b quarks are color connected to

each other, since the H is uncolored (color singlet), whereas in $g \rightarrow b\bar{b}$ background events, they are color connected to beam remnants because the gluon carries a color and an anticolor (color octet). We follow a recent suggestion [2] for reconstructing these color connections experimentally, using observables that can be modeled reliably by available leading-log parton-shower simulations. The technique involves measuring a vectorial quantity called “jet pull,” related to the jet energy pattern in the η - ϕ plane [3], using the measured energy in the calorimeter cells (see Fig. 1). Jets tend to have their pull pointing towards their color-connected partner. For instance, in $H \rightarrow b\bar{b}$ events, the pulls of the two b jets tend to point towards each other, whereas in $g \rightarrow b\bar{b}$ events, they point in opposite directions along the collision axis.

Verification of color-flow simulation and jet pull reconstruction for both color-singlet and color-octet configurations is interesting in its own right [4] and is needed before jet pull can be used in, e.g., $H \rightarrow b\bar{b}$ searches. Color-octet patterns can be studied in many processes, such as W/Z boson production in association with jets. A pure sample of color-singlet hadronic decays is difficult to obtain at a

*Visitor from Augustana College, Sioux Falls, SD, USA.

†Visitor from The University of Liverpool, Liverpool, UK.

‡Visitor from SLAC, Menlo Park, CA, USA.

§Visitor from ICREA/IFAE, Barcelona, Spain.

||Visitor from Centro de Investigacion en Computacion-IPN, Mexico City, Mexico.

¶Visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.

**Visitor from Universität Bern, Bern, Switzerland.




FIG. 1 (color online). Diagram showing two jets in the η - ϕ plane, and the reconstruction of the jet pull vectors (\vec{i}), jet pull angles (θ^{pull}), and relative jet pull angles ($\theta^{\text{pull}}_{\text{rel}}$).

hadron collider, but $t\bar{t}$ events with an $\ell +$ jets final state are good candidates since they have a characteristic signature and contain two jets from the decay of a W boson, which is a color singlet. Each of the two b jets coming from the top quark decays is color connected to one of the beam remnants in a color-octet pattern.

In this paper, we use data collected with the D0 detector [5] at the Fermilab Tevatron $p\bar{p}$ collider, corresponding to 5.3 fb^{-1} of integrated luminosity, to present the first experimental results on the study of jet pull, using $t\bar{t}$ events decaying to $\ell +$ jets ($t\bar{t} \rightarrow WbW\bar{b} \rightarrow \ell\nu bj\bar{j}\bar{b}$, where $\ell = e, \mu$). The object identification, event selection, and simulated Monte Carlo (MC) events are the same as those used in the $t\bar{t}$ cross section analysis [6], except that looser b -tagging criteria [7] are used to increase the statistics of double b -tagged events. We obtain a $\approx 90\%$ pure $t\bar{t}$ sample by requiring an isolated lepton with $p_T > 20 \text{ GeV}$, missing transverse energy $E_T > 20 \text{ GeV}$ ($> 25 \text{ GeV}$ for the $\mu +$ jets channel), and at least four jets, reconstructed with a midpoint cone algorithm [8] of radius 0.5, with $p_T > 20 \text{ GeV}$. At least one jet must have $p_T > 40 \text{ GeV}$, and at least two jets must be identified as b jets. Table I shows the event yields for these selection criteria.

To extract the fraction of color-singlet hadronic W boson decays, the data are compared to both standard model $t\bar{t}$ MC (with a color-singlet W boson) and an alternative model of $t\bar{t}$ with a hypothetical color-octet “ W ” boson decaying hadronically with identical properties except for its color representation. The latter is simulated using the MADGRAPH (MG) [9] event generator interfaced to PYTHIA [10] for showering and hadronization. Simulated events are processed with a GEANT3-based [11] detector simulation, overlaid with random data to account for backgrounds, and reconstructed as data.

D0 uses three liquid-argon/uranium calorimeters to measure the energies of particles: a central section covering $|\eta|$ up to ≈ 1.1 and two end calorimeters that extend

TABLE I. Yields of events passing selections with exactly four or ≥ 5 jets. At least two b -tagged jets are required in the analysis, but the numbers of events with zero or one b -tagged jet are also given. The number of events with zero or one b -tagged jet are also given. The number of $t\bar{t}$ events is calculated using the cross section determined with this data sample, $\sigma_{t\bar{t}} = 8.50 \text{ pb}$. Uncertainties include statistical and systematic contributions. The total uncertainties are smaller than the sum of individual uncertainties due to negative correlations between samples. Detailed selections are described in Ref. [6].

Channel	Sample	Zero b tags	One b tag	≥ 2 b tags
$\ell + 4$ jets	$W +$ jets	576 ± 75	229 ± 32	49 ± 8
	Multijet	115 ± 16	46 ± 7	7 ± 2
	$Z +$ jets	42 ± 6	16 ± 3	4 ± 1
	Other	31 ± 4	19 ± 2	9 ± 1
	$t\bar{t}$	160 ± 22	417 ± 38	519 ± 51
	Total	923 ± 62	727 ± 24	589 ± 48
	Observed	923	743	572
$\ell + \geq 5$ jets	$W +$ jets	60 ± 22	26 ± 11	7 ± 3
	Multijet	17 ± 3	12 ± 2	3 ± 1
	$Z +$ jets	4 ± 1	2 ± 1	1 ± 1
	Other	3 ± 1	3 ± 1	2 ± 1
	$t\bar{t}$	34 ± 6	90 ± 13	132 ± 17
	Total	118 ± 19	132 ± 7	145 ± 15
	Observed	112	127	156

coverage to $|\eta| \approx 4.2$ [3], housed in separate cryostats [12]. In addition, scintillators between the central section and end calorimeter cryostats provide sampling of developing showers for $1.1 < |\eta| < 1.4$. There are approximately ten layers in the radial direction (depending on η), generally composed of cells spanning 0.1×0.1 in $\eta \times \phi$. The energy resolution is about $15\%/\sqrt{E} \oplus 0.3\%$ (in GeV) for electrons and $50\%/\sqrt{E} \oplus 5\%$ for hadrons. Pileup energy from overlapping $p\bar{p}$ interactions results in about 0.5% of cells having energy above the noise-limited energy threshold (≈ 50 – 500 MeV , depending on layer and η). This energy is roughly exponentially distributed, with a mean of $\approx 350 \text{ MeV}$.

The pull is determined for each jet of a pair of reconstructed jets, using the measured energies of the calorimeter cells. Each cell within $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} < 0.7$ of the E_T -weighted center of one of the jets of the pair ($\eta_d^{\text{jet}}, \phi^{\text{jet}}$) is assigned to the jet nearer in ΔR . The contribution of each selected cell to the jet pull is $\vec{i}_{\text{cell}} = E_T^{\text{cell}} |\vec{r}_{\text{cell}}| \vec{r}_{\text{cell}}$, where $\vec{r}_{\text{cell}} = (\eta_d^{\text{cell}} - \eta_d^{\text{jet}}, \phi^{\text{cell}} - \phi^{\text{jet}})$, and E_T^{cell} is the cell’s transverse energy with respect to the nominal center of the detector. The jet pull is $\vec{i} = \sum_{\text{cells}, i} \vec{i}_i / E_T^{\text{jet}}$. The polar angle of the jet pull, θ^{pull} , is defined to be zero when pointing in the positive η direction along the beamline. A small correction to the jet pull is made to account for the energy response and noise in the calorimeters as a function of η_d , particularly in regions between the central and forward cryostats. The angle of the jet pull direction relative to the line defined by the centers of the jet pair

($\theta_{\text{rel}}^{\text{pull}}$) is of primary interest, as we expect color-connected jets to have pulls pointing towards each other. The $\theta_{\text{rel}}^{\text{pull}}$ quantity is calculated for each jet in the pair of highest- p_T b -tagged jets (b pair) and the pair with highest p_T which are not amongst the two highest p_T b -tagged jets (w pair).

To select events with a higher purity of properly identified jet pairs from hadronic W boson decays, we split the sample into events where the invariant mass of the w -pair jets is consistent with the W boson mass, $|m_{jj} - M_W| < 30$ GeV, and events where it is not. For the former, these two jets are found to match the partons from the W boson decay within $\Delta R < 0.5$ in 66% of $t\bar{t}$ MC events with four jets and 46% of events with five or more jets. In the latter case, where the mass of the w -pair jets is outside the W boson mass window, additional gluon radiation in the initial or final state leads to possible additional color configurations, diluting the measurement.

Since the w -pair jets in $t\bar{t}$ events are often from the W boson decay, we expect them to be color connected; thus the jet pulls should generally point towards each other. We expect b -pair jets to have one of the b jets color connected to the proton beam and the other to the antiproton beam; thus the jet pulls should be generally pointing away from each other. This tendency is seen in data, as shown in Fig. 2, with smaller $\theta_{\text{rel}}^{\text{pull}}$ in the w pair than in the b pair. However, the jets in w and b pairs have different kinematics, separation in the detector, and flavor. A direct interpretation of the effects from color flow is therefore not possible from this comparison. Furthermore, there are detector and reconstruction effects on jet pulls from overlapping jet pull cones, calorimeter noise and pileup, and calorimeter response inhomogeneity. For instance, there would be fewer cone overlaps if the jet pull was defined using only calorimeter cells within $\Delta R < 0.5$, producing on average smaller values for $\theta_{\text{rel}}^{\text{pull}}$. With this alternative definition the shape in Fig. 2(a) would peak more towards zero and that in Fig. 2(b) would be flatter. These effects are found to be well modeled by the simulation, and the jet pull definition based on the $\Delta R < 0.7$ cone gives a slightly improved singlet-octet separation. The relative jet pulls




FIG. 2 (color online). The average of the two jet $\theta_{\text{rel}}^{\text{pull}}$ distributions for jets in pairing (a) w and (b) b , in events with exactly four jets, at least two b tags, and the M_W requirement on the w -pair jets. The χ^2/ndf compares the data to the total MC distribution.

$\theta_{\text{rel}}^{\text{pull}}$ in data are also found to be well modeled by simulation for other jet pairings, such as a random w -pair jet and a random b -pair jet. In control samples consisting of events with a leptonic W boson decay, and two, three, or four jets, none identified as b jets, various jet pairings also have jet pulls that agree with simulations. Figure 3 shows the $\theta_{\text{rel}}^{\text{pull}}$ distributions for jets in a control sample with a leptonic W boson decay and two not- b -tagged jets.

To quantify the method's sensitivity to the color-flow structure (color singlet versus color octet) for the hadronic W boson decay, we fit the data to two hypotheses: (i) standard model $t\bar{t}$ with a color-singlet hadronically decaying W boson (singlet MC) and (ii) $t\bar{t}$ with a hypothetical color-octet W boson (octet MC). We determine the fraction of events coming from color-singlet W boson decay (f_{Singlet}) using the fitting procedure from the D0 combined $t\bar{t}$ cross section analysis [6]. We simultaneously measure the $t\bar{t}$ cross section to avoid any possible influence of the $t\bar{t}$ signal normalization on the f_{Singlet} measurement. The discriminating variable used for the fit is derived from the $\theta_{\text{rel}}^{\text{pull}}$ angles of the w -pair jets and depends on the ΔR between the two jets and their η_d . We define the following subdivisions for the data sample, which were optimized by studying the $t\bar{t}$ singlet and octet MC. For events failing the W mass requirement, we do not split the regions further; for other events we split the data sample according to the η_d of the jets and ΔR between the jets. For events where the two jets are highly separated ($\Delta R > 2$), we use the $\theta_{\text{rel}}^{\text{pull}}$ of the leading- p_T jet. Little discrimination is possible for these events, since the additional color radiation is distributed over a large area of the calorimeter. When the two jets are close ($\Delta R < 2$) and $|\eta_d| < 1.0$ for both jets, we use the minimum $\theta_{\text{rel}}^{\text{pull}}$ of the two jets. This is the most sensitive region, and the jet pull is accurately reconstructed in the central calorimeter due to less pileup energy and uniformity of response. Otherwise, if $|\eta_d|$ of the leading- p_T jet is < 1.0 (> 1.0), the $\theta_{\text{rel}}^{\text{pull}}$ of the leading- p_T (second-leading p_T) jet is used.

Table II lists the contribution of each non-negligible source of systematic uncertainty on f_{Singlet} . For all but




FIG. 3 (color online). (a) Leading- p_T and (b) second-leading- p_T jet $\theta_{\text{rel}}^{\text{pull}}$ distributions for w -pair jets, in events with two jets and no b -tagged jets. The χ^2/ndf compares the data to the total MC distribution.

TABLE II. The 1 standard deviation (σ) variation of f_{Singlet} from main systematic uncertainties. The total systematic uncertainty includes all uncertainties, summed in quadrature.

Source	+1 σ	-1 σ
Singlet/octet MC shapes	0.188	-0.188
Jet pull reconstruction	0.100	-0.093
Jet energy resolution	0.033	-0.013
Vertex confirmation	0.028	-0.029
PYTHIA tunes	0.023	-0.025
Jet energy scale	0.024	-0.009
Jet reconstruction and identification	0.017	-0.017
$t\bar{t}$ modeling	0.014	-0.033
Event statistics for matrix method	0.009	-0.010
Other Monte Carlo statistics	0.009	-0.007
Multijet background	0.006	-0.007
Total systematic	0.222	-0.218

the theoretical cross sections, MC statistics, and normalization of the W + heavy flavor jets background uncertainties, we apply the systematic uncertainties just to the $t\bar{t}$ signal sample and ignore the effect on background, as the purity of the $t\bar{t}$ sample is high. To estimate the possible systematic shift of the $\theta_{\text{rel}}^{\text{pull}}$ distribution due to the different energy scale and noise of the calorimeter cells between data and MC as a function of η_d , we apply $\pm 50\%$ of the jet pull η correction and take the resulting difference in shape as the systematic uncertainty for jet pull reconstruction. This covers the differences in the average θ^{pull} when comparing data and MC control samples. We also study systematic uncertainties as in [6], the main ones being from the jet energy scale, jet energy resolution, b -tagging efficiency, and lepton misidentification. Additional systematic uncertainties on $\theta_{\text{rel}}^{\text{pull}}$ are assessed to account for possible differences between MC and data related to the modeling of underlying event, hadronization, and jet showering. To estimate the variation due to these possible mismodelings, we compare $\theta_{\text{rel}}^{\text{pull}}$ distributions in events simulated with PYTHIA to those with ALPGEN [13] or MC@NLO [14], and showering with HERWIG [15]. We also do the comparisons for various PYTHIA parameters for underlying event and color reconnection [16], such as tunes APro and NOCR [17]. When deriving f_{Singlet} from the fit, we use the maximal variation obtained with the different $\theta_{\text{rel}}^{\text{pull}}$ distributions as an estimate of the systematic uncertainty.

Since the results are statistically limited and the analysis does not yet provide sufficient sensitivity for a definitive observation of color flow, we set limits on f_{Singlet} using the likelihood ratio ordering scheme of Feldman and Cousins [18]. We follow the same approach used for the simultaneous extraction of the ratio of branching fractions and the $t\bar{t}$ cross section [19] and generate ensembles of pseudoexperiments for different values of f_{Singlet} between 0 and 1, with the $t\bar{t}$ cross section fixed to the measured value.

We then vary the systematic uncertainties using Gaussian distributions and perform the fit as for the measurement on data. Statistical uncertainties are incorporated by smearing the measured value for each pseudoexperiment with the uncertainty determined in data. We use the nuisance parameters method where the expectation is fit to the data, for a variation of the initial prediction within the systematic uncertainties, allowing also the central result to change [6]. Other methods give compatible results.

We measure $f_{\text{Singlet}} = 0.56 \pm 0.42[\pm 0.36(\text{stat}) \pm 0.22(\text{syst})]$ and $\sigma_{t\bar{t}} = 8.50^{+0.87}_{-0.76}$ pb, consistent with our dedicated cross section measurement [6]. Figure 4 shows




FIG. 4 (color online). The discriminating color-flow variable, the minimum $\theta_{\text{rel}}^{\text{pull}}$ for the w -pair jets, for events passing the M_W requirement, with $\Delta R < 2$, and $\eta_d < 1.0$ for both jets. The $t\bar{t}$ MC shape is obtained using the measured value of f_{Singlet} .




FIG. 5 (color online). Expected C.L. bands for f_{Singlet} . The measured value is shown on the horizontal axis, and the input value on the vertical axis. The wide-dashed line shows the expected value, and the black-white fine-dashed line indicates the measured value of f_{Singlet} .

the distribution for one of the regions of the discriminating color-flow variable, using the measured $t\bar{t}$ cross section and measured f_{Singlet} . The expected 99% C.L. and 95% C.L. limits are $f_{\text{Singlet}} > 0.011$ and $f_{\text{Singlet}} > 0.277$, respectively, corresponding to an expected sensitivity to exclude $f_{\text{Singlet}} = 0$ of about 3 standard deviations, based on pseudoexperiments. The 68% C.L. allowed region from data is $0.179 < f_{\text{Singlet}} < 0.879$. Figure 5 shows the expected 68%, 95%, and 99% C.L. bands for f_{Singlet} .

In summary, we have presented the first study of color flow in $t\bar{t}$ events, with the method of jet pull, using 5.3 fb^{-1} of D0 integrated luminosity. The standard model MC predictions are found to be in good agreement with data, for both the jets from the hadronically decaying W boson, which should be in a color-singlet configuration, and the b -tagged jets from the top quark decays, which should be in a color-octet configuration. To quantify our ability to separate singlet from octet color flow, we measured the color representation of the hadronically decaying W boson

and found $f_{\text{Singlet}} = 0.56 \pm 0.42(\text{stat} + \text{syst})$, while the expected 95% C.L. limit was $f_{\text{Singlet}} > 0.277$. The ability to use color-flow information experimentally will benefit a wide range of measurements and searches for new physics.

We thank Jason Gallicchio, Matthew Schwartz, Steve Mrenna, Peter Skands, and Jay Wacker for discussions and guidance. We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom, and RFBR (Russia); CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC and the Royal Society (United Kingdom); MSMT and GACR (Czech Republic); CRC Program and NSERC (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); and CAS and CNSF (China).

-
- [1] D. Gross and F. Wilczek, *Phys. Rev. D* **8**, 3633 (1973); N. Brambilla and A. Vairo, [arXiv:hep-ph/9904330](https://arxiv.org/abs/hep-ph/9904330).
 - [2] J. Gallicchio and M.D. Schwartz, *Phys. Rev. Lett.* **105**, 022001 (2010).
 - [3] D0 uses a right-handed coordinate system, with the z axis pointing in the direction of the proton beam and the y axis pointing upwards. The azimuthal angle ϕ is defined in the xy plane and is measured from the x axis. The pseudorapidity is defined as $\eta = -\ln[\tan(\theta/2)]$, where θ is the polar angle. Detector η (η_d) is the η of an object measured from the nominal detector center.
 - [4] I. Sung, *Phys. Rev. D* **80**, 094020 (2009).
 - [5] V.M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006); **584**, 75 (2008); **622**, 298 (2010).
 - [6] V.M. Abazov *et al.* (D0 Collaboration), [arXiv:1101.0124](https://arxiv.org/abs/1101.0124).
 - [7] V.M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **620**, 490 (2010).
 - [8] G.C. Blazey *et al.*, in *Proceedings of the Workshop on QCD and Weak Boson Physics in Run II*, edited by U. Baur, R.K. Ellis, and D. Zeppenfeld (Fermilab, Batavia, IL, 2000).
 - [9] J. Alwall *et al.*, *J. High Energy Phys.* **09** (2007) 028.
 - [10] T. Sjöstrand, S. Mrenna, and P.Z. Skands, *J. High Energy Phys.* **05** (2006) 026.
 - [11] R. Brun and F. Carminati, CERN Program Library Long Writeup W5013, 1993 (unpublished).
 - [12] S. Abachi *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **338**, 185 (1994).
 - [13] M.L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A.D. Polosa, *J. High Energy Phys.* **07** (2003) 001.
 - [14] S. Frixione and B.R. Webber, *J. High Energy Phys.* **06** (2002) 029.
 - [15] G. Corcella *et al.*, [arXiv:hep-ph/0210213](https://arxiv.org/abs/hep-ph/0210213).
 - [16] T. Sjöstrand and P.Z. Skands, *J. High Energy Phys.* **03** (2004) 053.
 - [17] P.Z. Skands, *Phys. Rev. D* **82**, 074018 (2010).
 - [18] G. Feldman and R. Cousins, *Phys. Rev. D* **57**, 3873 (1998).
 - [19] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **100**, 192003 (2008).