

**Erratum: Infrared spectra of one- and two-dimensional fullerene polymer structures:  
RbC<sub>60</sub> and rhombohedral C<sub>60</sub>  
[Phys. Rev. B 55, 10 999 (1997)]**

K. Kamarás, Y. Iwasa, and L. Forró

[S0163-1829(98)05509-X]

Table I of the above article contains some mistakes and should be replaced by the table below. The corrections mainly concern the expected Raman lines of rh-C<sub>60</sub> for which we did not present any experimental data and therefore do not influence the conclusions of our measurements. Regarding the infrared lines, the correct statement is that *all* modes are expected to show a twofold splitting in  $D_{3d}$  symmetry, which is borne out by the experimental findings.

We thank Z. A. Nemeth for drawing our attention to the error.

TABLE I. Correlation table of symmetry groups  $I_h$ ,  $D_{2h}$ , and  $D_{3d}$ . Under “splitting,” we denote the change in the number of expected infrared (IR) and Raman ( $R$ ) active modes when the symmetry of the C<sub>60</sub> ball changes from  $I_h$  to the respective point group. The total number and the symmetry species of the allowed modes are summarized in the last two rows.

$I_h$	$D_{2h}$	Splitting	$D_{3d}$	Splitting
$2A_g$	$2A_g$	$R\ 1 \rightarrow 1$	$2A_{1g}$	$R\ 1 \rightarrow 1$
$3F_{1g}$	$3B_{1g} + 3B_{2g} + 3B_{3g}$	$R\ 0 \rightarrow 3$	$3A_{2g} + 3E_g$	$R\ 0 \rightarrow 1$
$4F_{2g}$	$4B_{1g} + 4B_{2g} + 4B_{3g}$	$R\ 0 \rightarrow 3$	$4A_{2g} + 4E_g$	$R\ 0 \rightarrow 1$
$6G_g$	$6A_g + 6B_{1g} + 6B_{2g} + 6B_{3g}$	$R\ 0 \rightarrow 4$	$6A_{1g} + 6A_{2g} + 6E_g$	$R\ 0 \rightarrow 2$
$8H_g$	$16A_g + 8B_{1g} + 8B_{2g} + 8B_{3g}$	$R\ 1 \rightarrow 5$	$8A_{1g} + 16E_g$	$R\ 1 \rightarrow 3$
$1A_u$	$1A_u$	IR 0→0	$1A_{1u}$	IR 0→0
$4F_{1u}$	$4B_{1u} + 4B_{2u} + 4B_{3u}$	IR 1→3	$4A_{2u} + 4E_u$	IR 1→2
$5F_{2u}$	$5B_{1u} + 5B_{2u} + 5B_{3u}$	IR 0→3	$5A_{2u} + 5E_u$	IR 0→2
$6G_u$	$6A_u + 6B_{1u} + 6B_{2u} + 6B_{3u}$	IR 0→3	$6A_{1u} + 6A_{2u} + 6E_u$	IR 0→2
$7H_u$	$14A_u + 7B_{1u} + 7B_{2u}$	IR 0→3	$7A_{2u} + 14E_u$	IR 0→2
$R\ 10(A_g, H_g)$	$R\ 87(A_g, B_{1g}, B_{2g}, B_{3g})$		$R\ 45(A_{1g}, E_g)$	
$IR\ 4(F_{1u})$	$IR\ 66(B_{1u}, B_{2u}, B_{3u})$		$IR\ 44(A_{2u}, E_u)$	