

ALLELOPATHIC EFFECT OF *Xanthium strumarium* L. AND *Abutilon theophrasti* Med. EXTRACTS ON GERMINATION OF MAIZE AND SOYBEAN SEED**Nataša Samardžić^{1*}, Bojan Konstantinović¹, Milena Popov¹, Milan Blagojević¹, Branimir Pavlič², Jelena Vladić²**¹*University of Novi Sad, Faculty of Agriculture, Department for Environmental and Plant Protection, Trg Dositeja Obradovića 8, 21000 Novi Sad, Serbia*²*University of Novi Sad, Faculty of Technology, Bulevar Cara Lazara 1, 21000 Novi Sad, Serbia**e-mail: natasam@polj.uns.ac.rs***Abstract**

During 2014 allelopathic effects of *Xanthium strumarium* L. and *Abutilon theophrasti* Med. extracts to germination and initial development of maize (*Zea mays* L.), and soybean (*Glycine max* L.) were studied in laboratory conditions. In addition to the Water extracts out of dry mass of the tested weed species, extracts made by use of hexane, ethyl acetate and methanol in different concentrations were also used. The applied concentrations were 10, 20, 30 and 40 g/l of dry matter made out of weed species in the 3-4 leaf stage of development. Inhibiting effect of water extract from dry matter of *X. strumarium* and methanol extract from which methanol part was evaporated to maize seed epicotyls and hypocotyls length was established. In comparison to the control, the maximum concentration of 40 g / l of the extract made from Water solution of *A. theophrasti* showed inhibitory effect on soybean seed epicotyls and hypocotyls length. The applied extracts made out of dry matter of the both of the studied weed species *X. strumarium* and *A. theophrasti* reduced maize seed germination for 14.8-26.83% and soybean seed germination for 18.5-35.82%, in comparison to the control in which it was 95% and 92%, respectively. After germination in a climate chamber, epicotyls' and hypocotyls' length of maize and soybean seeds was measured three, six and ten days following spraying by extracts.

Introduction

Weeds effect harmfully to crops by releasing phytotoxines from seeds, by decomposition of remainings, leaching and exudates [1]. Plants release harmful chemicals into the environment, reducing growth and establishment of other plants near them: the process known as allelopathy. Allelopathy is natural ecological phenomenon of relationship between organisms which can be applied for control of weeds, pests and diseases in field crops [2]. High allelopathic potential of *A. theophrasti* is shown due to inhibition of germination and growth of competitive plants, by which it reaches superior position. Although allelopathic interactions of *A. theophrasti* with other crops have been known for several decades, weak attention is paid to the biochemical interactions of this weed species, [3], that indicates negative allelopathic effect of *A. theophrasti* to soybean, maize and tomato crops [4]. Shajie and Saffari (2007) [5] established that extract made of leaves and petioles of *X. strumarium* significantly reduced germination and growth of *Zea mays* L, *Brassica napus* L, *Sesamum indicum* L, *Lens culinaris* Medic. and *Cicer arietinum* L. seedlings. Some researchers studied the effects of Water solutions made from different parts of plants, as well as their impact on other plant species [6] [7]. Different allelopathic activity of different parts of the same weed species also differs in its capabilities of harmful effects to germination and beginning growth of cultivated plants [8] [9]. Plant can

show inhibiting, but also stimulating effect to germination and growth of the nearby plants. In addition to examples of biochemical competition between weed and crop plants, there are examples of allelopathic interference between cultivated plants. This is the best illustrated by long time applied rotation system *Zea mays* (maize), *Glycine max* (soybean). It is observed that the rotation of these crops provides up to 20% higher yields [10].

Experimental

In 2012, at locality Kać near Novi Sad plant parts of *Abutilon theophrasti* Med. and *Xanthium strumarium* L. (stem and leaf) were collected. Water extract was prepared in the following manner: in 0.5 l of water, 150 g of chopped green mass of leaves and stems of *Abutilon theophrasti* Med. and *Xanthium strumarium* L. were immersed. Plant material was left in water for 96 hours at room temperature, after which it was removed, and the extract was filtered by vacuum filter. Filter paper in Petri dishes (150 mm x 25 mm) with germinated seed of the assayed soybean and maize crops was saturated by 8 ml of the extract. Control was moistened by distilled water. The assay was performed according to the method of Šćepanović *et al.*, 2007 [11]. In addition to the applied aqueous extracts, extracts made by use of methanol, hexane and ethyl acetate, from which by evaporation methanol, hexane and ethyl acetate were thrown, were also applied. All four extracts made of the above ground parts of *Abutilon theophrasti* Med. and *Xanthium strumarium* L. were made in concentrations of 10, 20, 30 and 40 g/l. By extracts treated soybean and maize seed were germinated in climatic chamber [12] and seed surface was sterilized according to Elemaru and Filhou (2005) [13].

The assays were set up according to the randomized block design with 4 replications. Each Petri dish contained 25 soybean and maize seeds, i.e. 100 seeds per treatment. All measurements were conducted third, sixth and tenth day after moistening of the studied crops seed. The existence of allelopathic activity of these two studied weed species to the soybean and maize crops were established by measurement of the crops seed epicotyls (mm) and hypocotyls (mm) length and germination (%) [11].

Results and discussion

Experimental data confirm results of other authors [14] [2], in which allelopathic effects were reflected in inhibition of germination, that was even more pronounced for the growth of seedlings. Germination of the studied maize seed was 95% and soybean seed 92%. (Statistica 10) The significant difference of average values of hypocotyls length was tested by statistical data analysis after treatment with extracts made of the above ground parts of *A. theophrasti* and *X. strumarium*. In the study, allelopathic effect of weed species *A. theophrasti* and *X. strumarium* to the beginning growth stages and development of soybean and maize hypocotyls was confirmed, while epicotyls growth was not statistically significantly different from control values for soybean. Water extracts of the above ground (leaves and stem) parts of *A. theophrasti* in concentrations of 20, 30 and 40 g/l showed inhibiting effects to the length of soybean hypocotyls in values (3.88mm; 4.01mm; 4.3mm), lower in comparison to the control value of 20.98mm. All concentrations of methanol extract, hexane and ethyl acetate showed statistically significant difference in comparison to the control. The highest difference in comparison to the control showed hexane extracts made of *A. theophrasti* in concentration of 40 g/l, with measured hypocotyls length of 13.46mm, and control value of 20.98mm (Figure 1).




Figure 1 Measured values of soybean seed hypocotyls after treatment with *Abutilon theophrasti* Med. extracts.

All concentrations of extractions of hexane and ethyl acetate made of *X. strumarium* had inhibiting effect of the growth of maize hypocotyls. Hexane extraction of 40 g/l reduced hypocotyls growth for 22.57% in comparison to the control value, while ethyl acetate extract in concentration of 40g/l reduced maize seed hypocotyls for 24% in relation to the control. Water extracts of *X. strumarium* had statistically significant effect to the hypocotyls growth in higher concentrations of 30 and 40 g/l, reducing hypocotyls for 14.8% and 16.29%, respectively. Methanol extraction of *X. strumarium* did not show statistically significant effect in concentration of 10g/l, but in remaining three concentrations it inhibited maize seed hypocotyls length. Methanol extraction of *X. strumarium* in concentration 40g/l showed the best effect, reducing maize seed hypocotyls growth for 26.83% in comparison to the control (Figure 2).




Figure 2 Measured values of maize seed hypocotyls, after treatment with *Xanthium strumarium* L. extracts.

The increase of maize seed hypocotyls statistically differed in comparison to the control value only in extracts with hexane and methanol in the highest concentrations of 40g/l. Epicotyls length of maize seed treated by hexane extract made of *X. strumarium* was 6.17mm and control value was 8.43mm. Epicotyls length of maize seed treated by methanol extract made of *X. strumarium* was 5.83mm, and control value was 8.43mm.

Conclusions

Based upon conducted studies, data were obtained on the effect of effect of *Abutilon theophrasti* Med. and *Xanthium strumarium* L. extracts on hypocotyls and epicotyls length of soybean and maize seed. Effect of *Abutilon theophrasti* Med. extract in tested concentrations of 10, 20, 30 and 40 g/l to the growth of soybean seed hypocotyls was between 18.5% in water extract, up to 35.82% in hexane extract, which was lower in comparison to the control. Effect of *Xanthium strumarium* L. extract in tested concentrations of 10, 20, 30 and 40 g/l to the growth of maize seed hypocotyls was between 14.8% in water extract, up to 26.83% in methanol extract, which was lower in comparison to the control. The tested concentrations did not show significant deviations in epicotyls length except for the two *Xanthium strumarium* L. extracts in the highest concentration of 40g/l.

References

- [1] S. S. Narwal, Scienlific Publishers, Jodhapur, India, 2004.
- [2] Z. Ashrafi, H. Mashhadi, S. Sadeghi, Pakistan Journal of Weed Science Research, 13(1-2) (2007) 99 -112.
- [3] J. B. Gressel, L.G. Holm, Weed Research 4 (1) (1964) 44-53.
- [4] J.R. Qasem, C.L. Foy, Allelopathy in Agroecosystems, (2001) 43-119.
- [5] E. Shajie, M. Saffari, Allelopathy Journal. 19 (2007) 501-506.
- [6] G. Kazinczi, I. Beres, J. Mikulas, E. Nadasy, Journal of Plant Diseases and Protection, 19 (2004) 301-308.
- [7] B. Konstantinović, N. Samardžić, M. Blagojević, Bo. Konstantinović, The 7th International symposium, Temisoara, (2013), pp. 36-37.
- [8] A. Aziza, A., Tanveer, M., Ali, B., Yasin, H., Babar, M.A. Nadeem, Allelopathy Journal, 22 (2008) 25-34.
- [9] B. Konstantinović, M. Meseldžija, M. Blagojević, N. Samardžić, Bo. Konstantinović, 9 Kongres o korovima, Zlatibor, (2012) 132-133.
- [10] S. J.H. Rizvi, V. Rizvi, Chapman & Hall. London (1992).
- [11] M. Šćepanović, N. Novak, K. Barić, Z. Ostojić, N. Galzina, M. Goršić, Agronomski glasnik 6/2007, Izvorni znanstveni članak (2007):.
- [12] S.U. Chon, Y-M. Kim, J-C. Lee, European Weed Research Society Weed Research, 43,(2003), 444–450.
- [13] V. Elemar, V. Filho, Brazilian Society on Weed Science. Congress Xs 24, Svaio Pedro, BRESIL, 40(1) (2005) 217.
- [14] M. Turk, A. Tawaha, Pakistan Journal Agronomy, 1(2002), 28-30.