



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:

Journal of Sport Psychology in Action

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa40891>

Paper:

Gross, M., Bringer, J., Kilduff, L., Cook, C., Hall, R. & Shearer, D. (2018). A multi-modal biofeedback protocol to demonstrate physiological manifestations of psychological stress and introduce heart rate variability biofeedback stress management. *Journal of Sport Psychology in Action*, 1-11.

<http://dx.doi.org/10.1080/21520704.2018.1496210>

12 month embargo

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>



A Multi-Modal Biofeedback Protocol to Demonstrate Physiological Manifestations of Psychological Stress, and Introduce Heart Rate Variability Biofeedback Stress Management.

Journal:	<i>Journal of Sport Psychology in Action</i>
Manuscript ID	USPA-2017-0045.R1
Manuscript Type:	Special Issue Article
Keywords:	Biofeedback, Heart Rate Variability, Psychophysiology, Stress, Sport

SCHOLARONE™
Manuscripts

Running head: MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY
STRESS MANAGEMENT.

1
2
3 **Abstract**

4
5 2 Physiological monitoring solutions and biofeedback technologies allow sport psychology
6
7 3 practitioners to demonstrate physiological changes due to psychological stress, and provide their
8
9 4 clients with an enhanced awareness of their stress response. These technologies can also provide
10
11 5 evidence for the efficacy of stress management techniques, whether cognitive or physiological in
12
13 6 nature. In this article, we present a protocol that uses multiple physiological signals to demonstrate
14
15 7 how psychological stress can manifest in the body, and then evidence heart rate variability
16
17 8 biofeedback as a stress management technique. Using insights from applied consultations, we focus
18
19 9 on the delivery of the protocol through phases of baseline resting measures, stressor tasks, and post-
20
21 10 stressor recovery. This article provides an accessible use of physiological monitoring and biofeedback
22
23 11 technologies in sport psychology practice, so that practitioners may adopt and adapt this protocol for
24
25 12 their purposes and available equipment.

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

Introduction

Biofeedback technology, where subjects view their live-feed physiological signals, is used to facilitate greater interoceptive awareness and learn individualised stress management techniques via operant conditioning. A growing body of evidence attests the usefulness of biofeedback technologies for directly and indirectly enhancing performance (e.g., Gevirtz, 2013; Jiménez Morgan & Molina Mora, 2017). Indeed, sophisticated biofeedback equipment has been used by elite and professional sports teams such as the World Cup winning Italian football team (Wilson, Peper, & Moss, 2014). However, there are equipment options to suit varied budgets, and the growth of peripheral and inbuilt sensors with mobile devices may be the future of biofeedback interventions (Heathers, 2013). Some detailed example interventions that provide technical details using these products have been published (e.g., Strack, Linden, & Wilson, 2011). However, research focussed literature is often too technical for practitioners who need quick-reference examples of how to deliver biofeedback interventions in sport settings.

Several physiological measures can be used in biofeedback interventions and heart rate variability (HRV) is popular because HRV statistics can index parasympathetic nervous system activation (Laborde, Mosley, & Thayer, 2017), and assess relative states of health and stress (Thayer, Ahs, Fredrikson, Sollers III, & Wager, 2012). HRV is the change in intervals between neighbouring heart beats due to interdependent regulatory processes such as the autonomic nervous system and heart-brain interactions (Shaffer, McCraty, & Zerr, 2014). Broadly, increased HRV indicates effective sport-related decision making under pressure (Laborde, Raab, & Kinrade, 2014), and both acute and chronic adaptive responses to training (Buchheit, 2014). HRV biofeedback is non-invasive and therapeutic benefits have been reported for various clinical conditions (Lehrer & Gevirtz, 2014; Shaffer et al) and sporting performances (Jiménez Morgan & Molina Mora, 2017). Coupled with the ease of learning and scant contraindications, the therapeutic effects of HRV biofeedback make it an attractive stress management intervention in elite sport settings.

The Lehrer, Vaschillo, and Vaschillo (2000) resonance frequency model of HRV biofeedback explains how bespoke abdominal breathing paces, near six breaths per minute, can be identified and

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

1 trained for therapeutic effects on-demand. Resonance frequency breathing utilises respiratory sinus
2
3 arrhythmia (RSA), an innate effect of heart rate increases with inhalation, and decreases with
4
5 exhalation, to create maximal HRV and blood pressure variation. This blood pressure variation
6
7 stimulates the body's control of vascular stiffness known as the baroreflex (Shaffer, McCraty, & Zerr,
8
9 2014), which buffers against extreme increases and decreases in blood pressure and takes a few
10
11 seconds to occur. It is these delays which produce a resonance frequency of approximately 10
12
13 seconds, corresponding to 0.1Hz and six breaths per minute (4 seconds inhale, 6 seconds exhale;
14
15 Lehrer et al). Baroreflex signals provide the brain stem with a visceral context from which autonomic
16
17 emotional responses are formed (Thayer et al, 2012) and stress-related blood pressure increases can
18
19 be inhibited (Saku et al, 2014). In summary, HRV biofeedback works by using RSA (via abdominal
20
21 breathing) to increase HRV and blood pressure variability, which invokes baroreceptor reflexes to
22
23 maintain context-appropriate blood pressure and provide a relaxed yet adaptable internal context for
24
25 emotional responding. When coupled with breathing measures, HRV biofeedback enables dose-
26
27 response observations which can provide participants with insight into physiological and
28
29 psychological stress management.
30
31

32
33 In addition to HRV and breathing, Galvanic Skin Response (GSR) data can reliably reflect
34
35 stress response (sympathetic) activation with a physiological delay of ~3-5 seconds (Critchley, Elliot,
36
37 Mathias, & Dolan, 2000). GSR is the change in skin electro-conductivity from micro secretions of
38
39 sweat due to sympathetic pathways, and is often used for biofeedback interventions (e.g., Toomin &
40
41 Toomin, 1975) because it is an intuitive indicator of stress. Concurrent biofeedback indices can
42
43 complement each other in protocols with baseline, stressor, and recovery phases to provide clients
44
45 with increased awareness of the physiological manifestations of stress and effective management
46
47 techniques thereafter. Indeed, this article describes how biofeedback technology can bring
48
49 participants' data 'to life' to demonstrate the physiological manifestations of psychological stress and
50
51 introduce HRV biofeedback as a stress management intervention. We provide a detailed description
52
53 of the protocol delivery used in our practice, where GSR and breathing data provide reliable and
54
55
56
57
58
59
60

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

1 intuitive indicators of momentary stress levels, then HRV data are explained to evidence the stress
2 management efficacy of HRV biofeedback interventions.

Biofeedback Protocol

4 We deliver this protocol in group or individual sessions with athletes, coaches, and support
5 staff prior to further HRV biofeedback protocols to enhance emotional regulation and performance
6 under pressure (e.g., Gross et al., 2016, 2017). Following sensor fitting and data fidelity checks, the
7 protocol includes a baseline phase and two bouts of stressor and post-stressor recovery phases. In
8 depth discussions of the physiological data occur during post-session reviews, where GSR and
9 breathing data indicate fluctuating stress levels (i.e., comparing stressor impact and recovery
10 effectiveness), and HRV and blood pressure variability are discussed in the context of managing
11 stress for athletic performance.

12 **Sensor Check & Familiarisation.** We inform participants that the protocol involves
13 cognitive tasks designed to be mildly stressful. After fitting sensors to the subject, the practitioner
14 briefly explains the how the sensors work and uses live graphical data displays to support these
15 explanations, confirm sensor accuracy (adjusting if necessary), and explain the physiological
16 concepts. For example, we use finger photoplethysmography (PPG) to record HRV and demonstrate
17 HRV via the variable gaps displayed between peaks in participants' live PPG signal. This helps
18 participants understand that to be adaptable, their heart must rapidly change its pace in response to
19 internal and external stimuli, such as digesting meals or evading threats. Breathing data can be
20 demonstrated by on-demand deep breaths, which cause upward and downward trends in live breathing
21 data with inhalation and exhalation (respectively).

22 GSR can be described as an equivalent of the alert response in cats, where tails raise and hairs
23 stand upright. We also often mention that GSR is used in the 'polygraph test' or 'lie detector', which
24 many are familiar with. The GSR sensors should be placed on participants' dominant hand to control
25 for potential asymmetry in responsiveness (Picard et al., 2015). Light-hearted comments confirming
26 that GSR sensors only *measure* rather than *send* electricity can build rapport and comfort anxious
27 participants. Conversely, confident participants may like to see an instant demonstration of their GSR

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

1 response which can be honoured with an unexpected hand-clap close to their face, reliably followed
2 by an upward trend in GSR values.

3 **Baseline.**

4 Stable measures of HRV, breathing, and GSR at rest are required as a comparison to the
5 stressor and recovery periods, and this should be articulated to participants. Recent advice suggests
6 that 2-minutes are adequate to capture a resting state HRV snapshot (Laborde et al, 2017) and little
7 rationale exists to extend resting measures for breathing or GSR. Practitioners can leave the room
8 during baseline measures to remove the presence of 'other' effects (Kleck et al.,1976).

9 **Stressor 1 & Recovery 1.**

10 Following Baseline measures, the practitioner can provide instructions for the 10-trial simple
11 Stroop test. A simple Stroop test is an interference reaction time test in which names of colours (e.g.,
12 blue) are presented on a screen printed in a different colour font (e.g., red) to the named word. For
13 each trial, participants are required to call out the colour of the font seen (i.e., not read the word)
14 within three seconds. We automate this test in our biofeedback software, however presentation
15 software or free online tests can be used provided test start and end times can be identified in
16 physiological data. Prior to the test, the practitioner can manipulate participants' difficulty
17 expectations to increase the impact on physiological data.

18 A score can be provided after the test and practitioners should move swiftly into Recovery 1
19 which asks participants to use any relaxation techniques they know to reduce GSR values as far as
20 possible in 2-minutes. Almost exclusively, maximal GSR values are reached during the Stroop test
21 and a timely transition into Recovery 1 instructions ensures similar starting values. Maximal and live
22 GSR data are shown on-screen in numerical and graphic representations because they are immediate
23 and intuitive, and participants are instructed to monitor live values as they try to relax. If they have
24 never used relaxation techniques, participants are instructed to recall a time that they were relaxed. If
25 available technology does not allow computed or custom physiological values, practitioners can
26 record peak values in session notes and refer to these in discussions. We have found that commentary
27 on the change in GSR values during Recovery 1 can increase difficulty for the participant. Comments

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

1 such as “*that’s the wrong direction, come on, relax!*”, announcing the remaining time, or providing a
2 target GSR value can achieve this simulation.

3 **Stressor 2 & Recovery 2.**

4 Stressor 2 is an arithmetic task where participants serially subtract 13 from 1084 and vocalise
5 as many answers as possible in 60 seconds, returning to the start if errors are made. The practitioner
6 should call out ‘correct’ for successful calculations and provide a score (number of successful
7 calculations) upon completion. More so than Stressor 1, this task reliably invokes physiological stress,
8 which can be enhanced by distractions such as contaminating participants’ working memory with
9 other numbers, or prior queries into their mental arithmetic capabilities.

10 After the task, practitioners should move swiftly into Recovery 2 which introduces paced
11 abdominal breathing at six breaths per minute (4 seconds inhale, 6 seconds exhale; Lehrer et al.,
12 2000). Breathing is guided by a simple on-screen bar graph in our software, however free websites or
13 mobile device applications can be used. Practitioners should ensure participants’ abdominal breathing,
14 advise them not to completely fill and empty their lungs, and to breathe through their nose or mouth
15 as comfort dictates. A common solution to participants who struggle with abdominal breathing is to
16 place their hands on their chest and navel, then observe as they attempt to move the hand on the navel
17 rather than the chest. We use a dual line graph plotting heart rate and breathing so that the practitioner
18 can monitor breathing against the pacing stimulus and provide coaching tips. Similarly, on-screen
19 numerical displays of breathing rate and both live and new maximal GSR (reliably achieved during
20 Stressor 2) values allow the practitioner to monitor the relaxation achieved.

21 In our experience, six breaths per minute reliably relaxes participants as demonstrated in
22 HRV, GSR, and subjective reports. RSA and increased HRV can usually be seen after two breath
23 cycles, however conscious relaxation commonly requires 4 or more breaths, and we conduct
24 Recovery 2 for 2-minutes for comparable data with both Recovery 1 and Baseline. Physiological
25 variables such as total blood volume impact effectiveness of breathing paces for relaxation (Lehrer et
26 al 2000), so slower paces with the same 2:3 ratio (e.g., five breaths per minute; 4.8 seconds inhale, 7.2
27 seconds exhale) can be used for taller and broader individuals. As per Recovery 1, the practitioner can

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

comment on the change in GSR data and in our experience, these comments do not have the same aversive effects as in Recovery 1. This disparity strengthens post-session discussions on the efficacy of slow abdominal breathing for relaxation.

Review & Discussion

We begin post-session reviews and discussion by ensuring that data are saved, removing physiological sensors, and ensuring participant comfort before showing a session overview (see Figure 1) and explain each data series. At differing points throughout the discussion, we focus on HRV, GSR, PPG, and breathing data, along with computer calculated metrics of heart and breathing rates. Starting with Baseline, we use an on-screen zoom function to view specific phases of the protocol and revisit individualized discussion points.

In our experience, 2-minute Baseline data show rhythmic breath cycles at approximately 12 breaths per minute, variable heart rate reflecting RSA, and decreasing GSR from beginning to end. Participants often request to know their resting heart rate which provides a good opportunity to revisit HRV as reflection of adaptability to internal and external demands. We emphasise that when they are relaxed, breathing patterns are deep and rhythmic and heart rate is variable. However, we rarely mention HRV statistics at this stage because they are not necessary to convey this information and have complicated this insight for some participants. Nevertheless, providing an average heart rate from the Baseline period provides another metric to measure stressor effects and we often set visible threshold lines in our software to enable visual comparisons between phases.

As Figure 1 illustrates, an increase in GSR from Baseline often occurs during Stroop instructions which can be used to highlight arousal and anticipatory stress. GSR commonly peaks during the first 3-5 Stroop trials after participants have realised that the test is within their capabilities. This peak and subsequent 'recovery' can be analogous to settling into a sporting performance. During the Stroop, breathing and heart rates increase while breath depth decreases. We have found that linking these changes to a stress response primes participants for discussions on the effectiveness of regulated breathing for relaxation and recovery from stress.

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

1 To begin Recovery 1 discussions, we ask participants what relaxation technique(s) was used
2 (e.g., fixation on a visual cue, relaxing muscles). We generally find that these techniques moderately
3 decrease GSR, however the trend can reverse if aversive comments are used. Independent of the
4 technique used and as shown in Figure 1, participants show slower, deeper, and rhythmic breathing
5 patterns compared to Stressor 1. Again, we highlight these breathing patterns and the concomitant
6 changes in heart rate to prime Recovery 2 discussions, suggesting that they already know to use
7 breathing to relax, just not at an optimally slow pace.

8 By design, participants find Stressor 2 more stressful than Stressor 1, and this is reflected in
9 physiological data. We frequently see an increase in GSR values during Stressor 2 instructions, and
10 this is a key talking point in our review and discussions. Akin to Stressor 1, GSR may peak early and
11 plateau or decrease towards the end, and shallower breaths with increases in both heart and breathing
12 rates can be seen. Participant calculation errors or practitioner intrusive comments may be reflected in
13 GSR as upward inflexions and further demonstrates the physiological impact of psychological stress.
14 We then discuss how a hyperactive stress response may not support optimal athletic performances.
15 These implications are intuitive for participants from self-paced and closed skill sports, and for others,
16 we focus discussions on the implications for optimal preparation and recovery. The key message we
17 convey here is not that stress is inherently negative for performance, rather that performers should be
18 aware of the effects and learn to manage it to their advantage.

19 We ask participants for their feedback on their experience of the paced breathing activity in
20 Recovery 2, and nearly all report ease in following the pacing stimulus and most report relaxation
21 effects. We initially focus discussions on GSR data as indicators of stress and because they almost
22 unanimously display a decrease towards Baseline values from the session maximum reached during
23 Stressor 2 (see Figure 1). We then mention that there was a larger recovery burden for Recovery 2,
24 and provide a dose-response insight in to how resonance frequency breathing reduces physiological
25 stress by showing the reduction in breathing rate and increase in both RSA and HRV. We then use the
26 zoom function to show two breath cycles of PPG, breathing, and heart rate data to demonstrate how
27 this increased HRV exercises the body's blood pressure regulation system (i.e., the baroreflex;

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

1 Shaffer, McCraty, & Zerr, 2014). As shown in Figure 2, gaps between heart beats show heart rate
2 increasing with inhalation (smaller gaps) and decreasing with exhalation (larger gaps), and varying
3 peak amplitudes indicate changes in artery (and capillary) stiffness in response to changing heart rate
4 (Shaffer et al). For psychological stress, we explain how stimulating the baroreflex inhibits blood
5 pressure increases due to stress (Saku et al., 2014) and forms a relaxed internal context from which
6 autonomic emotional responses are formed (Thayer et al, 2012), among other therapeutic effects (e.g.,
7 mindfulness related focus on breathing; Lehrer & Gevirtz, 2014).

8 Participants can instantly recognise that blood pressure is elevated under stress, but may not
9 realise that HRV and blood pressure variability are decreased and so too is their social, emotional, and
10 environmental adaptability. The best analogy we have found is to discuss the colloquial term ‘hangry’
11 as a state of physiological stress, where hunger increases the propensity for negative emotional states
12 or shortness of patience. We often say “*When hangry, your body is telling your brain to focus on*
13 *acquiring food and that you’re not able take on new tasks or deal with more stress until you eat.*
14 *Blood pressure and heart rate are similar in that elevated levels, but reduced variability, send similar*
15 *messages to the emotional centres of your brain. Breathing at slow paces to create those changes in*
16 *heart rate and stimulate blood pressure regulation inverts those messages into ‘relaxed’ and*
17 *‘adaptable’.*” So as not to over-complicate these insights we refrain from discussing HRV statistics
18 unless participants are particularly interested or have relevant knowledge. To conclude the
19 presentation, we discuss when participants need to manage stress, when they could apply resonance
20 frequency breathing in life and sport, and previous instances when it may have helped.

21 Following discussions of previous stressful experiences, we begin further HRV biofeedback
22 protocols by identifying the participant’s optimal breathing pace as per Gross et al (2016, 2017). We
23 also provide options for free breath pacing applications on mobile devices, and encourage participants
24 to regularly practice in diverse situations in order to learn the breath pace and negate the need for a
25 pacing stimulus. Following this practice, our participants have incorporated their optimal (resonance
26 frequency) breathing paces into daily routines, performance preparations, skill-specific routines, and
27 recovery processes. Consistent with the growing research interest into HRV biofeedback, the reported

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS MANAGEMENT.

1 benefits include competitive anxiety management, relaxation between performance bouts, improved
2 sleep quality, pre-emptive and reactive stress reduction, and minimised pain during exercise recovery.
3 We have also found that these experiences promote further sport psychology support and interest in
4 associated interventions such as mindfulness, likely due to the profound dose-response experience.
5 Indeed, we think that objective demonstrations of psychological skills and their therapeutic
6 mechanisms is a strong rationale for practitioners to regularly use biofeedback technologies.

References

- 7
8 Buchheit, M. (2014). Monitoring training status with HR measurements: do all roads lead to Rome?
9 *Frontiers in Physiology*, 5, 1-19. doi:10.3389/fphys.2014.00073
10
11 Critchley, H. D., Elliott, R., Mathias, C. J., & Dolan, R. J. (2000). Neural Activity Relating to
12 Generation and Representation of Galvanic Skin Conductance Responses: A Functional
13 Magnetic Resonance Imaging Study. *Journal of Neuroscience*, 20(8).
14 <http://www.jneurosci.org/content/20/8/3033.short>
15
16 Gevirtz, R. (2013). The promise of heart rate variability biofeedback: Evidence-based applications.
17 *Biofeedback*, 41, 110-120. doi:10.5298/1081-5937-41. 3.01
18
19 Gross, M. J., Shearer, D. A., Bringer, J. D., Hall, R., Cook, C. J., & Kilduff, L. P. (2016). Abbreviated
20 resonant frequency training to augment heart rate variability and enhance on-demand emotional
21 regulation in elite sport support staff. *Applied Psychophysiology and Biofeedback*, 263-274.
22 doi:10.1007/s10484-015-9330-9
23
24 Gross, M. J., Hall, R., Bringer, J. D., Cook, C. J., Kilduff, L. P., & Shearer, D. A. (2017). Resonant
25 frequency training in elite sport: A case study example. *Journal of Sport Psychology in Action*,
8, 173-183.
26
27 Heathers, J. A. J. (2013). Smartphone-enabled pulse rate variability: An alternative methodology for
28 the collection of heart rate variability in psychophysiological research. *International Journal of*
29 *Psychophysiology*, 89(3), 297–304. <https://doi.org/10.1016/j.ijpsycho.2013.05.017>

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS
MANAGEMENT.

- 1 Jimémez Morgan, S., & Molina Mora, J. A. (2017). Effect of Heart Rate Variability Biofeedback on
2 Sport Performance, a Systematic Review. *Applied Psychophysiology and Biofeedback*, *42*(3),
3 235–245. doi.org/10.1007/s10484-017-9364-2
- 4 Kleck, R. E., Vaughan, R. C., Cartwright-Smith, J., Vaughan, K. B., Colby, C. Z., & Lanzetta, J. T.
5 (1976). Effects of being observed on expressive, subjective and physiological responses to
6 painful stimuli. *Journal of Personality and Social Psychology*, *34*, 1211-1218.
7 doi:10.1037/0022-3514.34.6.1211
- 8 Laborde, S., Mosley, E., & Thayer, J. F. (2017). Heart Rate Variability and Cardiac Vagal Tone in
9 Psychophysiological Research - Recommendations for Experiment Planning, Data Analysis, and
10 Data Reporting. *Frontiers in Psychology*, *8*, 213. https://doi.org/10.3389/fpsyg.2017.00213
- 11 Laborde, S., Raab, M., & Kinrade, N. (2014). Is the ability to keep your mind sharp under pressure
12 reflected in your heart? Evidence for the neurophysiological bases of decision reinvestment.
13 *Biological Psychology*, *100*, 34–42. doi:10.1016/j.biopsycho.2014.05.003.
- 14 Lehrer, P. M., & Gevirtz, R. (2014). Heart rate variability biofeedback: How and why does it work?
15 *Frontiers in Psychology*, *5*(JUL), 1–9. https://doi.org/10.3389/fpsyg.2014.00756
- 16 Lehrer, P. M., Vaschillo, E., & Vaschillo, B. (2000). Resonant frequency biofeedback training to
17 increase cardiac variability: Rationale and manual for training. *Applied Psychophysiology and*
18 *Biofeedback*, *25*, 177–189.
- 19 Picard, R. W., Fedor, S., & Ayzenberg, Y. (2016). Multiple arousal theory and daily-life
20 electrodermal activity asymmetry. *Emotion Review*, *8*(1), 62-75.
- 21 Saku, K., Kishi, T., Sakamoto, K., Hosokawa, K., Sakamoto, T., Murayama, Y., . . . Sunagawa, K.
22 (2014). Afferent vagal nerve stimulation resets baroreflex neural arc and inhibits sympathetic
23 nerve activity. *Physiological Reports*, *2*, 1-12. doi:10.14814/phy2.12136
- 24 Shaffer, F., McCraty, R., & Zerr, C. L. (2014). A healthy heart is not a metronome: an integrative
25 review of the heart's anatomy and heart rate variability. *Frontiers in Psychology*, *5*(September),
26 1–19. doi.org/10.3389/fpsyg.2014.01040

MULTI-MODAL BIOFEEDBACK AND HEART RATE VARIABILITY STRESS
MANAGEMENT.


- 1 Strack, B., Linden, M., & Wilson, V. S. (Eds.). (2011). Biofeedback & neurofeedback applications in
2 sport psychology. *Association for Applied Psychophysiology and Biofeedback*.
- 3 Thayer, J. F., Ahs, F., Fredrikson, M., Sollers, J. J., I. I. I., & Wager, T. D. (2012). A meta-analysis of
4 heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker
5 of stress and health. *Neuroscience and Biobehavioral Reviews*, 36, 747–756.
6 doi:10.1016/j.neubiorev.2011.11.009.
- 7 Toomin, M. K., & Toomin, H. (1975). GSR biofeedback in psychotherapy: Some clinical
8 observations. *Psychotherapy: Theory, Research & Practice*, 12(1), 33-38.
- 9 Wilson, V. E., Peper, E., & Moss, D. (2006). “The Mind Room” in Italian Soccer Training: The Use of
10 Biofeedback and Neurofeedback for Optimum Performance. *Biofeedback*, 34(3), 79-81.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Biofeedback protocol session overview.


Figure 2. Final two breath cycles in Recovery 2.

For Peer Review Only



Note. Data collection begins with resting Baseline measures (left side), and includes two bouts of Stressor Instructions, Stressor, Recovery. Physiological data series in descending order: Galvanic Skin Response (GSR), Photoplethysmograph (PPG), Breathing, Recovery.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Note. Physiological data series in descending order: Photoplethysmograph (PPG), Breathing, Heart Rate.

View Only