

1 **Intake of Animal Protein Blend Plus Carbohydrate Improves Body Composition with**
2 **no Impact on Performance in Endurance Athletes**

3 Fernando Naclerio¹

4 Eneko Larumbe-Zabala²

5 Mar Larrosa³

6 Aitor Centeno⁴

7 Jonathan Esteve-Lanao³

8 Diego Moreno-Pérez⁵

9

10 ¹ Department of Life and Sport Science, Faculty of Engineering and Science, University of
11 Greenwich, United Kingdom

12 ² Clinical Research Institute, Texas Tech University Health Sciences Center, Lubbock, TX,
13 USA

14 ³ Department of Pharmacy, Biotechnology, Nutrition, Optics and Optometry. Faculty of
15 Biomedical and Health Sciences. Universidad Europea de Madrid.

16 ⁴ Sports Training Laboratory, Universidad Europea, Madrid, Spain.

17 ⁵ Department of Education, Research and Evaluation Methods, Comillas Pontifical
18 University, Madrid, Spain

19

20 Corresponding author: Fernando Naclerio. E-mail: f.j.naclerio@gre.ac.uk

21 Department of Life and Sport Science, University of Greenwich, Sparrows Farm. Sparrows
22 Lane. Avery Hill, New Eltham SE9 2BT.

23 Tel: +44(0) 20831 8441

24

25 Accepted for publication on 18th December 2018 at the International Journal of Sport
26 Nutrition & Exercise Metabolism

27 **ABSTRACT**

28 The impact of animal blend protein supplements in endurance athletes is scarcely researched.
29 We investigated the effect of ingesting an admixture providing orange juice and protein from
30 beef and whey versus carbohydrate alone on body composition and performance over a 10-
31 week training period in male endurance athletes. Participants were randomly assigned to a
32 protein (CHO+PRO, n=15) or a non-protein isoenergetic carbohydrate (CHO, n=15) group.
33 Twenty grams of supplement mixed with orange juice was ingested post-workout or before
34 breakfast on non-training days. Measurements were performed pre- and post-intervention on
35 body composition (by dual-energy X-ray absorptiometry), peak oxygen consumption
36 ($\dot{V}O_{2peak}$), and maximal aerobic speed (MAS). Twenty-five participants (CHO+PRO, n=12;
37 CHO, n=13) completed the study. Only the CHO+PRO group significantly ($p<0.05$) reduced
38 whole body fat (mean \pm SD) (-1.02 ± 0.6 kg), total trunk fat (-0.81 ± 0.9 kg) and increased
39 total lower body lean mass ($+0.52 \pm 0.7$ kg), showing close to statistically significant
40 increases of whole-body lean mass ($+0.57 \pm 0.8$ kg, $p=0.055$). Both groups reduced ($p<0.05$)
41 visceral fat (CHO+PRO, -0.03 ± 0.1 kg; CHO, -0.03 ± 0.5 kg) and improved the speed at
42 MAS (CHO+PRO, $+0.56 \pm 0.5$ km·h⁻¹; CHO, $+0.35 \pm 0.5$ km·h⁻¹). Although consuming
43 animal blend protein mixed with orange juice over 10 weeks helped to reduce fat mass and
44 to increase lean mass, no additional performance benefits in endurance runners were
45 observed.

46

47 **Keywords:** Whey; beef; lean mass; trunk fat; visceral adipose tissue; aerobic, runners.

48 **Introduction**

49 The current daily protein recommendation for regular endurance exercisers is
50 between 1.2 to 1.6 (Thomas et al., 2016) or up to 1.8 g·kg⁻¹·body mass for trained endurance
51 athletes (Jager et al., 2017). Accordingly, Kato et al. (2016), using the amino acid oxidation
52 method, suggested an average daily consumption of 1.65 or up to 1.83 g·kg⁻¹·body mass to
53 satisfy protein requirements in endurance trained males. Such an amount of daily protein
54 intake should be administered evenly spaced throughout the day. Moreover, the consumption
55 of protein during the post-workout time has been proposed as a pragmatic and sensible
56 strategy (Kerksick et al., 2017) for supporting recovery and the adaptational processes
57 (Doering et al., 2016). While no ergogenic outcomes may be evident, research has reported
58 that the post-workout ingestion of protein and carbohydrate admixtures are effective to
59 attenuate markers of muscle damage, decrease muscular soreness (Kerksick et al., 2017), and
60 maintain or increase muscle mass in endurance athletes compared to the ingestion of only
61 carbohydrate (D'Lugos et al., 2016). Consequently, the post-workout ingestion of protein-
62 carbohydrate admixtures may attenuate muscle disruption and optimize changes in body
63 composition but this practice may not have a meaningful effect on performance compared to
64 the ingestion of carbohydrate alone (McLellan et al., 2014).

65 Both whey and beef are high-quality protein sources with a very similar amino acid
66 composition to that found in skeletal muscle (Cruzat et al., 2014). Although whey contains
67 higher concentrations of leucine, which seems to be an important essential amino acid for
68 starting muscle protein synthesis (Naclerio and Larumbe-Zabala, 2016), beef is a source of
69 heme-iron, zinc, vitamin B12, and essential fatty acids that are relevant nutrients in
70 supporting muscle remodeling (Phillips, 2012). Indeed, the ingestion of a post-workout
71 hydrolyzed beef protein was effective to protect muscle mass in male endurance athletes
72 (Naclerio et al., 2017). On the other hand, whey is composed of several bioactive fractions
73 (glycomacropeptide, β -lactoglobulin, α -lactalbumin and lactoferrin), with multiple health
74 (Zapata et al., 2017) and weight control benefits (Miller et al., 2014). Although the positive

75 effects of protein supplementation to support lean mass in endurance athletes is well
76 documented (Doering et al., 2016), its effects to reduce total and abdominal fat have been
77 mainly observed in overweight and obese adults (Arciero et al., Ormsbee, 2014). The aim of
78 the current study, therefore, was to compare the effects of combining a 10-week endurance
79 training program with one of the following commercially available products: (i) Beef and
80 Whey protein blend (Crown® Sport Nutrition, Spain) providing hydrolyzed 100% All Beef
81 and whey isolate (Optipep, Carbery) mixed with orange juice; and (ii) non-protein,
82 carbohydrate-only (maltodextrin and orange juice), on body composition and performance in
83 well-trained male endurance runners. The primary outcomes measures were whole body fat
84 mass, whole body lean mass, total trunk fat mass, trunk lean mass, visceral fat mass, total
85 (right and left) upper and lower body limb lean and fat mass. Secondary outcomes measures
86 included peak oxygen consumption, and maximal aerobic speed. Based on the available
87 literature, we hypothesized that compared to an isoenergetic-only carbohydrate supplement,
88 the post-workout ingestion of a carbohydrate-protein admixture would protect muscle mass,
89 and promote fat reduction with no additional performance benefit in well-trained endurance
90 athletes.

91 **Methods**


92 **Participants**

93 After a pre-screening of the individuals characteristics and training background, thirty
94 endurance athletes met the inclusion criteria: (a) >18–45 years of age; (b) only those who
95 consistently trained between 6 to 10 hours per week (four to seven workout per week) for the
96 last five years were considered for the study; (c) free from musculoskeletal limitations.
97 Exclusion criteria were: (a) history of metabolic conditions and/or diseases; (b) consuming
98 any medication including those with androgenic and/or anabolic effects, nutritional
99 supplements affecting performance and body composition (e.g. creatine, essential amino
100 acids, proteins, dehydroepiandrosterone, etc.) during the previous 8 weeks prior to the start

101 of the study; (c) current use of tobacco products; (d) the presence of any soft tissue or
102 orthopedic limitations.

103 Compliance was confirmed verbally and prior to providing written consent. The study
104 was approved by the Institution Ethics Committee for Clinical Research (ID: 2016 RM/05).
105 All experimental procedures were conducted in accordance with the Declaration of Helsinki
106 and registered as Clinical Trial at ClinicalTrials.gov, U.S. National Institutes of Health
107 (Identifier: NCT02954367).

108 Twenty-five of the 30 recruited participants completed all aspects of the study (Figure.
109 1).



110 **Figure 1:** Flow diagram of participants throughout the course of the study.

111 The study was designed as a double-blind, two parallel group, randomized control
112 trial for between-participant comparisons. After assessing for eligibility, the participants
113 were randomly allocated into two equal-size treatment groups: protein (CHO+PRO), n=15;
114 or carbohydrate only (CHO), n=15. Following a pre-assessment of body composition and

115 performance, the participants were matched by their fat, fat-free and $\dot{V}O_{2\text{peak}}$ values. In a
116 double-blind fashion, the assignment of participants to two treatments was performed by
117 block randomization using a block size of two. Initial groups characteristics (mean \pm SD)
118 were not significantly different at baseline: CHO+PRO: age 30.3 ± 8.8 years, 1.74 ± 0.59 m
119 height, 68.9 ± 4.4 kg body mass, 60.5 ± 7.3 ml/kg/min⁻¹ $\dot{V}O_{2\text{peak}}$; CHO: 34.1 ± 7.8 years, 1.76
120 ± 0.51 m height, 66.2 ± 4.0 kg body mass, 61.49 ± 6.8 ml/kg/min⁻¹, $\dot{V}O_{2\text{peak}}$.

121 Sample size estimations were calculated assuming a two group by two repeated
122 measures model, where the α -error probability was set at 0.05 and the statistical power was
123 established at 0.80 (1- β). Based upon an effect size of $\eta^2=0.035$ for the primary outcome
124 variable, fat mass (kg), and an interaction effect between groups conducted upon an interim
125 analysis of the first 12 participants, a sample size estimation of n=24 was determined as
126 appropriate. Nonetheless, assuming an anticipated attrition rate of 20%, we enrolled 15
127 participants per group.

128 **Assessments**

129 Before and after a 10-week intervention period, measurements of body composition
130 followed by an endurance test were determined. Prior to the assessments, participants were
131 instructed to refrain from any vigorous activity and avoid caffeine ingestion for at least 48-
132 h. All tests were performed at the same time of the day for the same participant.

133 Body mass, whole body fat mass, whole body lean mass, total trunk fat mass,
134 estimated visceral fat mass, and fat and lean mass for upper and lower limbs (right and left)
135 were measured using dual-energy X ray absorptiometry (General Electric Healthcare,
136 Madison, WI). These measurements were performed in standardized conditions, in the
137 morning and in a fasted state.

138 A progressive to volitional exhaustion running test was used to determine peak
139 oxygen consumption ($\dot{V}O_{2\text{peak}}$) and maximal aerobic speed (MAS). After a general warm-up,
140 starting at 10 km·h⁻¹, running speed was increased by 0.3 km·h⁻¹ every 30s until volitional
141 exhaustion. Gas exchange data were collected continuously using an automated breath-by-

142 breath system (Ultima™ Series, MGC Diagnostic Corporation, St. Paul, Minnesota, USA
143 Vmax 29C); which was calibrated according to the manufacturer's instructions. The volume
144 calibration was performed at different flow rates with a 3-L calibration syringe allowing an
145 error <3%. The calibration of gas analyzers was performed automatically using reference
146 values of environmental gases and cylinders (16% O₂, 4% CO₂). $\dot{V}O_{2peak}$ was recorded as the
147 highest $\dot{V}O_2$ value obtained for any continuous 30s period. The maximal aerobic speed
148 (MAS) was associated with the last completed 30s stage before exhaustion (Esteve-Lanao,
149 Foster, Seiler, & Lucia, 2007).

150 **Control of training**

151 All participants were trained by the same coach. All of them committed to follow a
152 10-week training program using a polarized intensity distribution (Esteve-Lanao et al., 2007).
153 Participants trained 5 to 6 sessions per week controlling the duration, distance and quantified
154 intensity by continuous heart rate registration. All the participants trained during the
155 afternoon (12 to 6:00 pm).

156 **Dietary Monitoring**

157 Each participant's baseline diet (3 days, 2 weekdays, and 1 weekend day) was
158 analyzed using Dietplan 6 software (Microsoft Forestfield Software Ltd. 14). Participants
159 were instructed to maintain their normal diet. To evaluate differences caused by treatments,
160 diet was analyzed again during the last week of the intervention.

161 **Supplementation and Control of the Intervention Compliance**

162 The two supplements were presented as 24 g sachets of vanilla-flavored powder diluted
163 in ~250 mL of orange juice. The mixed drinks were similar in appearance, texture and taste,
164 and were isoenergetic. The nutritional composition of each product is presented in Table 1.
165 On training days, supplements were ingested within 20 min after training, whereas on non-
166 training days supplement was administered before breakfast. To avoid missing doses, on non-
167 training days, automatic text messages were sent to all the participants. Additionally,
168 participants were allowed to drink water at libitum but not to consume any food during the

169 training sessions.

170 After completing the first assessment, each participant was given a batch of one of
171 the two products, assigned according to randomization.

172 Tolerance collected from any adverse events and compliance with supplement intake
173 (determined by an individual follow-up) was evaluated continuously. Only participants who
174 completed the 70 days of treatment with a minimum of 4 sessions per week (40 workouts in
175 total) were analyzed. The diary training report was used to determine participant compliance.

176

177

178 **Table 1.** Nutritional composition of drinks per intake (24 g of powder plus 250 ml of
 179 orange juice)

Nutrient	CHO+PRO	CHO
Energy value (kcal)	204	204
Carbohydrates (g)	27.70	50.10
Lipids (g)	1.05	0
Proteins (g)	19.84	0.40
Alanine (g)	1.14	-
Arginine (g)	0.82	-
Aspartic acid (g)	1.94	-
Cysteine (g)	0.33	-
Glutamic acid (g)	3.33	-
Glycine (g)	0.79	-
Histidine (g)	0.48	-
Isoleucine (g)	1.16	-
Leucine (g)	1.76	-
Lysine (g)	1.82	-
Methionine (g)	0.45	-
L-Ornithine	0.02	-
Phenylalanine (g)	0.67	-
Proline (g)	1.08	-
Serine (g)	0.88	-
L-Taurine	0.02	-
Threonine (g)	1.13	-
Tryptophan (g)	0.28	-
Tyrosine (g)	0.58	-
Valine (g)	1.13	-
Total EAA (g)	10.64	-
Heme Iron (mg)	1.93	-
Zinc (mg)	2.26	-
Potassium (mg)	2012.16	-
Magnesium (mg)	15.90	-
Selenium (µg)	2.88	-
Calcium (mg)	59.25	-
Folic Acid (µg)	10.04	-
Niacin (mg)	13.04	-
Vitamin B 6 (mg)	0.04	-
Vitamin B 12 (µg)	0.39	-

180 **Notes:** EAA: essential amino acids; CHO+PRO: supplement admixture including orange juice
 181 mixing with a beef and whey protein blend, CHO: supplement admixture including orange juice mixing with
 182 maltodextrin.

183
 184 **Statistical Analysis**

185 A descriptive analysis was performed and subsequently the Kolmogorov-Smirnov

186 and Shapiro-Francia test were applied to assess normality. Sample characteristics at baseline
187 were compared between conditions (CHO+PRO vs. CHO) using two-tailed independent
188 samples t test. Changes from pre to post treatment in body composition, and performance
189 were assessed using a 2 (treatments) \times 2 (times) repeated measures ANOVA. As suggested
190 by Castañeda et al. (1993), changes over time were analyzed using a priori Bonferroni-
191 adjusted pairwise comparisons. Generalized eta squared (η_G^2) and Cohen's *d* values were
192 reported to provide an estimate of standardized effect size (small $d=0.2$, $\eta_G^2=0.01$; moderate
193 $d=0.5$, $\eta_G^2=0.06$; and large $d=0.8$, $\eta_G^2=0.14$). Significance level was set to 0.05 but *p* values
194 between >0.05 and 0.1 were considered indicative of a trend. Results are reported as mean \pm
195 SD unless stated otherwise. Data analyses were performed with Stata 13.1 (StataCorp,
196 College Station, TX).

197 **Results**

198 Due to non-intervention related reasons, five participants (3 from CHO+PRO and 2
199 from CHO) dropped out of the study. At baseline, all the analyzed variables were not
200 significantly different between groups. Table 2 shows the dietary monitoring results,
201 determined before and after the intervention.

202 At baseline, no between-group differences were observed. However, as a result of the
203 intervention, CHO+PRO group significantly increased both the protein and carbohydrate
204 intakes while CHO group increased the consumption of carbohydrates. Despite no changes
205 observed in the overall caloric intake, both groups increased the energy contribution from
206 carbohydrates and decreased the proportion from fat. However, only CHO+PRO increased
207 the proportion of energy from proteins. Despite the observed changes, no between-treatment
208 differences were observed at post-intervention. No complaints about any negative symptoms
209 (i.e. hypoglycemic reaction) or gastric discomfort due to the ingestion of supplement were
210 reported. Table 3 summarizes the pre and post values of the analyzed variables.

211

212 **Table 2.** Descriptive analysis of the participants diet composition

Treatment	CHO+PRO (n=12)		CHO (n=13)	
	Pre	Post	Pre	Post
Protein				
g·d ⁻¹	122.5 ± 23.4	143.2 ± 29.5*	125.1 ± 28.6	125.4 ± 26.3
g·kg ⁻¹ ·d ⁻¹	1.7 ± 0.3	2.1 ± 0.4*	1.9 ± 0.4	1.9 ± 0.4
% of total energy	21 ± 0.4	23 ± 0.3*	22 ± 0.4	21 ± 0.3
Carbohydrate				
g·d ⁻¹	255.6 ± 102.9	304.5 ± 108.0*	238.82 ± 73.9	281.9 ± 59.3*
g·kg ⁻¹ ·d ⁻¹	3.6 ± 1.4	4.3 ± 1.5*	3.6 ± 1.1	4.2 ± 0.9*
% of total energy	41 ± 0.6	47 ± 0.5*	41 ± 0.6	48 ± 0.5*
Fat				
g·d ⁻¹	97.6 ± 27.8	103.98 ± 31.01	96.07 ± 29.6	93.5 ± 21.1
g·kg ⁻¹ ·d ⁻¹	1.4 ± 0.4	1.48 ± 0.40	1.42 ± 0.4	1.4 ± 0.3
% of total energy	38 ± 0.5	30 ± 0.3*	38 ± 0.5	31 ± 0.4*
Energy				
Total daily energy	2433.5 ± 726.7	2561.0 ± 797.7	2339.8 ± 600.9	2373.9 ± 471.5
Kcal·kg ⁻¹ ·d ⁻¹	34.8 ± 10.5	36.4 ± 10.5	34.7 ± 8.3	35.2 ± 6.4

213 Notes: Pre and post intervention values are presented as mean ± standard deviation

214 *P<0.05; **P<0.001 and ^Tp<0.10 from pre to post-intervention (last week of intervention).215 CHO+PRO = participants ingesting orange juice mixed with beef and whey protein, CHO
216 participants ingesting orange juice mixing with maltodextrin.

217

218 Main time effects were observed for body mass [F(1,23)=7.86, p=0.010, $\eta_G^2=0.26$],
219 whole body fat [F(1,23)=15.83, p=0.001, $\eta_G^2=0.41$], whole body lean mass [F(1,23)=4.75,
220 p=0.040, $\eta_G^2=0.17$], total trunk fat mass [F(1,23)=12.04, p=0.002, $\eta_G^2=0.34$], visceral fat mass
221 [F(1,23)=14.83, p=0.001, $\eta_G^2=0.39$], total lower body limb fat mass [F(1,23)=6.07, p=0.022,
222 $\eta_G^2=0.21$] and total lower body limb lean mass [F(1,23)=5.06, p=0.034, $\eta_G^2=0.18$]. No
223 interaction or between-groups effects were identified. Pairwise comparisons revealed that
224 only CHO+PRO significantly reduced body mass (p=0.039). Both groups reduced whole
225 body fat mass (CHO+PRO, p=0.004; CHO, p=0.024), but neither group increased trunk or
226 upper body lean mass. No change in arm fat was observed. Furthermore, only CHO+PRO
227 produced a significant increase in the total lower body limb lean mass (p=0.016) along with
228 a very close to significant increase (p=0.055) in the whole-body lean mass. Additionally, both
229 groups showed close to significant decreases in total lower body limb fat mass (CHO+PRO
230 p=0.098; CHO p=0.075).

231 **Table 3.** Descriptive analysis of the body composition and performance variables

Variables	CHO+PRO (n=12)				CHO (n=13)			
	Pre	Post	Change	ES	Pre	Post	Change	ES
Body mass (kg)	69.6 ± 4	68.8 ± 4*	-0.87 ± 0.9	0.63	67.2 ± 3.6	66.5 ± 4.3 [†]	-0.67 ± 1.6	0.49
Whole body fat mass (kg)	14.5 ± 3.4	13.4 ± 2.8**	-1.02 ± 0.6	0.92	14.1 ± 2.8	13.4 ± 2.3*	-0.74 ± 1.3	0.67
Whole body lean mass (kg)	53.1 ± 3.3	53.6 ± 3.4 [†]	+0.57 ± 0.8	0.58	51.6 ± 3.8	51.9 ± 3.7	+0.28 ± 1.0	0.29
Total trunk fat mass (kg)	6.8 ± 2.1	6.0 ± 1.5**	-0.81 ± 0.9	0.94	6.3 ± 1.5	5.9 ± 1.4	-0.39 ± 0.8	0.45
Trunk lean mass (kg)	24.2 ± 1.8	24.0 ± 1.7	-0.19 ± 0.9	0.20	23.3 ± 1.6	23.4 ± 1.4	+0.13 ± 0.8	0.15
Visceral fat mass (kg)	0.34 ± 0.1	0.31 ± 0.1**	-0.03 ± 0.1	0.82	0.32 ± 0.1	0.28 ± 0.1*	-0.03 ± 0.5	0.72
Total lower body limb fat mass (kg)	4.9 ± 1.0	4.7 ± 1.0 [†]	-0.22 ± 0.2	0.47	5.1 ± 1.0	4.9 ± 1.0 [†]	-0.24 ± 0.58	0.54
Total lower body limb lean mass (kg)	18.9 ± 1.4	19.4 ± 1.6*	+0.52 ± 0.7	0.75	18.4 ± 1.6	18.5 ± 1.7	+0.10 ± 0.6	0.16
Total upper body limb fat mass (kg)	1.7 ± 0.5	1.7 ± 0.6	+0.01 ± 0.1	0.01	1.7 ± 0.5	1.5 ± 0.4	-0.11	0.30
Total upper body limb lean mass (kg)	6.4 ± 0.6	6.6 ± 0.7	+0.22 ± 0.6	0.38	6.1 ± 0.8	6.2 ± 0.8	+0.02 ± 0.5	0.04
$\dot{V}O_{2peak}$ (ml·kgm ⁻¹ ·min ⁻¹)	61.0 ± 5.6	61.2 ± 4.0	+0.24 ± 2.8	0.07	60.1 ± 6.9	60.8 ± 5.0	0.15 ± 3.7	0.04
Maximal aerobic speed (km·h ⁻¹)	17.8 ± 1.3	18.4 ± 1.0**	+0.56 ± 0.5	1.01	17.7 ± 1.0	18.1 ± 0.9*	+0.35 ± 0.5	0.64

232

233


234

235

236

Note: Values determined at pre, post and the corresponding calculated change (post – pre) are presented as mean ± standard deviation. Pairwise comparison *p<0.05; **p<0.01 respect to pre-intervention values. [†]p >0.05 and <0.1. ES= Cohen's d, effects size for two dependent means. CHO+PRO = participants ingesting orange juice mixing with beef and whey protein, CHO participants ingesting orange juice mixing with maltodextrin.

237 Only CHO+PRO significantly decreased total trunk fat ($p=0.004$, Figure 2A).
 238 However, both treatments decreased visceral fat (CHO+PRO, $p=0.009$; CHO, $p=0.016$,
 239 Figure 2B). No statistically significant differences between groups were observed after
 240 intervention in any of the body composition variables.



241

242 **Figure 2.** Observed changes in the total trunk fat (A) and estimated visceral fat (B).
 243 CHO+PRO = participants ingesting orange juice mixed with beef and whey protein,
 244 CHO = participants ingesting orange juice mixed with maltodextrin.
 245 Data are presented as mean (95% CI). ** $p<0.01$, * $p<0.05$; respect to pre-intervention values.
 246

247 Training time distribution was as follows: 75–80% in Zone 1, ~5% in Zone 2, and
 248 15–20% in Zone 3. The resulted training load using the ECOs methods described by Esteve-
 249 Lanao et al., (2017) was ~43%-7%-50% for Zone 1; Zone 2 and Zone 3 respectively.

250 No time, group or time x group interaction effects were determined for $\dot{V}O_{2peak}$,
 251 however, main time ($F(1,23)=17.11$, $p=0.001$, $\eta_G^2=0.43$) but no group or interaction effects
 252 were determined for MAS. Pairwise comparisons revealed that both groups significantly
 253 increased the speed at MAS (CHO+PRO $p=0.001$; CHO $p=0.03$).

254 Discussion

255 The present study shows that ingesting a 20 g post workout protein blend (beef and

256 whey) mixed with orange juice over 10 weeks promoted positive changes in body
257 composition, reduced body mass, total trunk fat and increased lean mass in endurance-trained
258 runners. Despite the observed modification in the CHO+PRO group and the improved MAS
259 determined in both groups (CHO+PRO and CHO), no significant differences between
260 conditions were noticed at post-intervention.

261 Compared to CHO, the decrease in body mass in CHO+PRO was associated with a
262 higher amount of fat mass loss (CHO+PRO: -1.02 ± 0.6 vs. CHO: -0.74 ± 1.3) alongside a
263 superior increase of the whole-body lean mass (CHO+PRO: $+0.57 \pm 0.8$ vs. CHO: $+0.28 \pm$
264 1.0). Indeed, only the CHO+PRO group showed higher effect sizes to increase lower body
265 limb lean and whole-body lean mass respectively (Table 3).

266 The observed results emphasize the positive effects of ingesting a protein supplement
267 to preserve or promote muscle mass in endurance athletes (Doering et al., 2016; Naclerio et
268 al., 2017). Maintaining appropriate levels of lower body limb lean mass in long distance
269 runners has been associated with more efficient recovery, reduced overload related injuries
270 and generally better training outcomes (Doering et al., 2016). Moreover, the ingestion of a
271 post-workout admixture providing carbohydrates and 0.25 to 0.4 g·kg·body mass⁻¹ of high-
272 quality protein has been shown to favor body net protein balance and support recovery after
273 endurance exercises (Jager et al., 2017). Participants allocated to CHO+PRO were ingesting
274 between 0.26 to 0.31 g·kg·body mass⁻¹ immediately post-workout or before breakfast during
275 non-training days. The administered amount falls within the recommended protein intake to
276 maximize muscle protein synthesis at rest (Areta et al., 2013) or to significantly improve
277 muscle repair after exercise (Morton et al., 2015).

278 There was no apparent effect due to energy or macronutrient difference as an effect
279 of the intervention. Thus, the only main difference between conditions was the composition
280 of the post-workout supplement. According to the diet records, the amount of carbohydrates
281 consumed by the two groups (CHO+PRO: 4.33 ± 1.47 ; CHO: 4.20 ± 0.87 g·kg⁻¹·d⁻¹, Table 2)
282 was below the recommended dose of 5 to 7 g·kg⁻¹·d⁻¹ for endurance athletes (Thomas et al.,

283 2016). The limited carbohydrate intake could have negatively influenced performance or
284 induced loss of lean body mass. However, no negative effects on body composition or
285 performance were observed for both treatments. When carbohydrates are provided below the
286 required amount, a higher daily protein intake toward $2 \text{ g}\cdot\text{kg}^{-1}$ would be necessary to support
287 metabolic adaptation including optimal glycogen replenishment and muscle remodeling
288 (Thomas et al., 2016). Participants in both groups were consuming a relatively high amount
289 of daily protein. Furthermore, no participant was ingesting less than $1.4 \text{ g}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$ which is
290 well above than the minimum daily amount of protein ($1.2 \text{ g}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$) recommended for
291 endurance exercisers (Thomas et al., 2016). Additionally, only one participant in CHO+PRO
292 and three in CHO ingested more than $1.65 \text{ g}\cdot\text{kg}^{-1}$ of protein which is the suggested average
293 intake to satisfy the metabolic demands of endurance training (Kato et al., 2016). Our results
294 seem to support the recommendation of ingesting high-quality protein-carbohydrate
295 admixtures immediately after training for maintaining lean mass and reducing trunk fat
296 (Kerksick et al., 2017). Although both CHO+PRO and CHO decreased whole body fat, only
297 the CHO+PRO group significantly reduced total trunk fat (Table 3 and Figure 2A) and
298 increased lower body lean mass (Table 3). The beneficial effect of ingesting high-quality
299 protein supplements on body composition has been extensively reported in active or
300 sedentary (Miller et al., 2014) overweight/obese (Arciero et al., 2014), as well as in physically
301 active (Monteyne et al., 2018) or trained individuals (Morton et al., 2018; Taylor et al., 2016).
302 Nonetheless, as visceral fat decreased in both conditions, it seems that regular exercise
303 represents the main stimulus for mobilizing internal fat in normal weight trained athletes.
304 The ingestion of animal protein, particularly whey, rather than vegetable protein has been
305 associated with suppressed appetite, increased satiety (Miller et al., 2014), and favors protein
306 synthesis which in turn would increase thermogenesis after ingesting high-protein meals
307 (Acheson et al., 2011). Therefore, a hypothetically higher use of fat as the predominant fuel
308 to support muscle-remodeling during the early recovery phase after ingesting a post-workout
309 protein-carbohydrate admixture could be the cause of the more favorable changes in body

310 composition observed in CHO+PRO compared to CHO. Moreover, recent evidences in
311 rodents suggest that some components of whey protein such as Lactalbumin and Lactoferrin
312 may increase postprandial lipolysis markers (Mobley et al., 2015), improve energy balance
313 and decrease adiposity (Zapata et al., 2017) .

314 The present study is not without limitations; the diet was not strictly controlled but
315 only recorded over 3 days before and after intervention. Although this approach has been
316 extensively used, providing a pre-packed daily-meal scheme to participants would offer an
317 ideal scenario to standardize and control their diet (Jeacocke and Burke, 2010). Although the
318 observed trend to increase in lean mass for the CHO+PRO group could be explained by a
319 gain in musculature, it is possible that non-muscle lean tissue in the trunk region made
320 substantial contribution (Mitchell et al., 2017). Magnetic Resonance Imaging techniques
321 would have been required to identify the contribution of skeletal muscle, viscera, and gut to
322 the observed changes in lean mass indistinguishable with the use of DEXA as in the current
323 study.

324 Considering the research design, the current findings support that the ingestion of a
325 post-workout admixture providing protein from beef and whey mixed with orange juice
326 represents a suitable alternative to improve body composition (trunk fat mass loss, increase
327 whole and lower body limb lean mass) compared with the ingestion of carbohydrates alone.
328 Nonetheless, no impact on performance has been observed.

329 **Acknowledgements**

330 FN, DM and ELZ designed the study. DM and AC collected data. ELZ and FN analyzed the
331 data. Interpretation and manuscript preparation was conducted by FN and supported by all
332 authors. All authors edited and approved the final version of the paper.

333 The authors want to thank Kelsey Jendrzey for grammar review and editing of the
334 manuscript.

335 **Declaration of funding sources and conflict of interest**

336 Research funding Real Madrid-European University of Madrid 2016/RM05. M. L. holds a
337 Ramón y Cajal Fellowship 2012 11910. However, these do not affect this original research
338 content and purpose.

339 The authors declare that they have no conflicts of interest relevant to the content of this
340 manuscript.

341

342

343 **REFERENCES**

- 344 Acheson, K. J., Blondel-Lubrano, A., Oguey-Araymon, S., Beaumont, M., Emady-Azar, S.,
 345 Ammon-Zufferey, C., ... Bovetto, L. (2011). Protein choices targeting thermogenesis and
 346 metabolism. *The American Journal of Clinical Nutrition*, *93*(3), 525–534.
 347 <https://doi.org/10.3945/ajcn.110.005850>
- 348 Arciero, P. J., Baur, D., Connelly, S., & Ormsbee, M. J. (2014). Timed-daily ingestion of
 349 whey protein and exercise training reduces visceral adipose tissue mass and improves insulin
 350 resistance: the PRISE study. *Journal of Applied Physiology* (1985), *117*(1), 1–10.
 351 <https://doi.org/10.1152/jappphysiol.00152.2014>
- 352 Areta, J., Burke, L. M., Ross, M. L., Camera, D. M., West, D. D., Broad, E. M., ... Coffey,
 353 V. G. (2013). Timing and distribution of protein ingestion during prolonged recovery from
 354 resistance exercise alters myofibrillar protein synthesis. *The Journal of Physiology*, *591*(9),
 355 2319–2331.
- 356 Castañeda, M. B., Levin, J. R., & Dunham, R. B. (1993). Using planned comparisons in
 357 management research: A case for the Bonferroni procedure. *Journal of Management*, *19*(3),
 358 707–724. <https://doi.org/10.1177/014920639301900311>
- 359 Chernoff, R. (2004). Protein and older adults. *Journal of the American College of Nutrition*,
 360 *23*(6 Suppl), 627S–630S. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15640517>
- 361 Cruzat, V. F., Krause, M., & Newsholme, P. (2014). Amino acid supplementation and impact
 362 on immune function in the context of exercise. *Journal of the International Society of Sports*
 363 *Nutrition*, *11*(1), 61. <https://doi.org/10.1186/s12970-014-0061-8>
- 364 D'Lugos, A. C., Luden, N. D., Faller, J. M., Akers, J. D., McKenzie, A. I., & Saunders, M.
 365 J. (2016). Supplemental Protein during Heavy Cycling Training and Recovery Impacts
 366 Skeletal Muscle and Heart Rate Responses but Not Performance. *Nutrients*, *8*(9).
 367 <https://doi.org/10.3390/nu8090550>
- 368 Doering, T. M., Reaburn, P. R., Phillips, S. M., & Jenkins, D. G. (2016). Postexercise Dietary
 369 Protein Strategies to Maximize Skeletal Muscle Repair and Remodeling in Masters
 370 Endurance Athletes: A Review. *International Journal of Sport Nutrition and Exercise*
 371 *Metabolism*, *26*(2), 168–178. <https://doi.org/10.1123/ijsnem.2015-0102>
- 372 Esteve-Lanao, J., Foster, C., Seiler, S., & Lucia, A. (2007). Impact of training intensity
 373 distribution on performance in endurance athletes. *Journal of Strength and Conditioning*
 374 *Research*, *21*(3), 943–949. <https://doi.org/10.1519/R-19725.1>
- 375 Esteve-Lanao, J., Moreno-Perez, D., Cardona, C. A., Larumbe-Zabala, E., Munoz, I., Selles,
 376 S., & Cejuela, R. (2017). Is Marathon Training Harder than the Ironman Training? An ECO-
 377 method Comparison. *Frontiers in Physiology*, *8*, 298.
 378 <https://doi.org/10.3389/fphys.2017.00298>
- 379 Jager, R., Kerksick, C. M., Campbell, B. I., Cribb, P. J., Wells, S. D., Skwiat, T. M., ...
 380 Antonio, J. (2017). International Society of Sports Nutrition Position Stand: protein and
 381 exercise. *Journal of the International Society of Sports Nutrition*, *14*, 20.
 382 <https://doi.org/10.1186/s12970-017-0177-8>
- 383 Jeacocke, N. A., & Burke, L. M. (2010). Methods to standardize dietary intake before
 384 performance testing. *International Journal of Sport Nutrition and Exercise Metabolism*,

- 385 20(2), 87–103. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20479482>
- 386 Kato, H., Suzuki, K., Bannai, M., & Moore, D. R. (2016). Protein Requirements Are Elevated
387 in Endurance Athletes after Exercise as Determined by the Indicator Amino Acid Oxidation
388 Method. *PLoS One*, 11(6), e0157406. <https://doi.org/10.1371/journal.pone.0157406>
- 389 Kerksick, C. M., Arent, S., Schoenfeld, B. J., Stout, J. R., Campbell, B., Wilborn, C. D., ...
390 Antonio, J. (2017). International society of sports nutrition position stand: nutrient timing.
391 *Journal of the International Society of Sports Nutrition*, 14, 33.
392 <https://doi.org/10.1186/s12970-017-0189-4>
- 393 McLellan, T. M., Pasiakos, S. M., & Lieberman, H. R. (2014). Effects of protein in
394 combination with carbohydrate supplements on acute or repeat endurance exercise
395 performance: a systematic review. *Sports Medicine*, 44(4), 535–550.
396 <https://doi.org/10.1007/s40279-013-0133-y>
- 397 Miller, P. E., Alexander, D. D., & Perez, V. (2014). Effects of whey protein and resistance
398 exercise on body composition: a meta-analysis of randomized controlled trials. *Journal of*
399 *the American College of Nutrition*, 33(2), 163–175.
400 <https://doi.org/10.1080/07315724.2013.875365>
- 401 Mitchell, C. J., Milan, A. M., Mitchell, S. M., Zeng, N., Ramzan, F., Sharma, P., ... Cameron-
402 Smith, D. (2017). The effects of dietary protein intake on appendicular lean mass and muscle
403 function in elderly men: a 10-wk randomized controlled trial. *The American Journal of*
404 *Clinical Nutrition*, 106(6), 1375–1383. <https://doi.org/10.3945/ajcn.117.160325>
- 405 Mobley, C. B., Fox, C. D., Ferguson, B. S., Pascoe, C. A., Healy, J. C., McAdam, J. S., ...
406 Roberts, M. D. (2015). Effects of protein type and composition on postprandial markers of
407 skeletal muscle anabolism, adipose tissue lipolysis, and hypothalamic gene expression.
408 *Journal of the International Society of Sports Nutrition*, 12, 14.
409 <https://doi.org/10.1186/s12970-015-0076-9>
- 410 Monteyne, A., Martin, A., Jackson, L., Corrigan, N., Stringer, E., Newey, J., ... James, L. J.
411 (2018). Whey protein consumption after resistance exercise reduces energy intake at a post-
412 exercise meal. *European Journal of Nutrition*, 57(2), 585–592.
413 <https://doi.org/10.1007/s00394-016-1344-4>
- 414 Morton, R. W., McGlory, C., & Phillips, S. M. (2015). Nutritional interventions to augment
415 resistance training-induced skeletal muscle hypertrophy. *Frontiers in Physiology*, 6, 245.
416 <https://doi.org/10.3389/fphys.2015.00245>
- 417 Morton, R. W., Murphy, K. T., McKellar, S. R., Schoenfeld, B. J., Henselmans, M., Helms,
418 E., ... Phillips, S. M. (2018). A systematic review, meta-analysis and meta-regression of the
419 effect of protein supplementation on resistance training-induced gains in muscle mass and
420 strength in healthy adults. *British Journal of Sports Medicine*, 52(6), 376–384.
421 <https://doi.org/10.1136/bjsports-2017-097608>
- 422 Naclerio, F., & Larumbe-Zabala, E. (2016). Effects of Whey Protein Alone or as Part of a
423 Multi-ingredient Formulation on Strength, Fat-Free Mass, or Lean Body Mass in Resistance-
424 Trained Individuals: A Meta-analysis. *Sports Medicine*, 46(1), 125–137.
425 <https://doi.org/10.1007/s40279-015-0403-y>
- 426 Naclerio, F., Seijo, M., Larumbe-Zabala, E., Ashrafi, N., Christides, T., Karsten, B., &
427 Nielsen, B. V. (2017). Effects of Supplementation with Beef or Whey Protein Versus
428 Carbohydrate in Master Triathletes. *Journal of the American College of Nutrition*, 38(8), 1–

- 429 9. <https://doi.org/10.1080/07315724.2017.1335248>
- 430 Phillips, S. M. (2012). Nutrient-rich meat proteins in offsetting age-related muscle loss. *Meat*
431 *Science*, *92*(3), 174–178. <https://doi.org/10.1016/j.meatsci.2012.04.027>
- 432 Taylor, L. W., Wilborn, C., Roberts, M. D., White, A., & Dugan, K. (2016). Eight weeks of
433 pre- and postexercise whey protein supplementation increases lean body mass and improves
434 performance in Division III collegiate female basketball players. *Applied Physiology*
435 *Nutrition and Metabolism*, *41*(3), 249–254. <https://doi.org/10.1139/apnm-2015-0463>
- 436 Thomas, T., Erdman, K. A., & Burke, L. M. (2016). American College of Sports Medicine
437 Joint Position Statement: Nutrition and athletic performance. *Medicine and Science in Sports*
438 *& Exercise*, *48*(3), 543–568. <https://doi.org/10.1001/jama.1956.02970290016006>
- 439 Zapata, R. C., Singh, A., Pezeshki, A., Nibber, T., & Chelikani, P. K. (2017). Whey Protein
440 Components - Lactalbumin and Lactoferrin - Improve Energy Balance and Metabolism.
441 *Scientific Reports*, *7*(1), 9917. <https://doi.org/10.1038/s41598-017-09781-2>
- 442