

***Homo sapiens* in Arabia by 85,000 years ago**

Huw S. Groucutt^{1,2*}, Rainer Grün^{3,4}, Iyad S.A. Zalmout⁵, Nick A. Drake⁶, Simon J. Armitage^{7,8}, Ian Candy⁷, Richard Clark-Wilson⁷, Julien Louys³, Paul S. Breeze⁶, Mathieu Duval^{3,9}, Laura T. Buck^{10,11}, Tracy L. Kivell^{12,13}, Emma Pomeroy^{10,14}, Nicholas B. Stephens¹³, Jay T. Stock^{10,15}, Mathew Stewart¹⁶, Gilbert J. Price¹⁷, Leslie Kinsley⁴, Wing Wai Sung¹⁸, Abdullah Alsharekh¹⁹, Abdulaziz Al-Omari²⁰, Muhammad Zahir²¹, Abdullah M. Memesh⁵, Ammar J. Abdulshakoor⁵, Abdu M. Al-Masari⁵, Ahmed A. Bahameem⁵, Khaled S.M. Al Murayyi²⁰, Badr Zahrani²⁰, Eleanor M.L. Scerri^{1,2} & Michael D. Petraglia^{2,22*}

¹ School of Archaeology, Research Laboratory for Archaeology and the History of Art, Hayes House, 75 George Street, University of Oxford, Oxford, OX1 2BQ, UK.

² Department of Archaeology, Max Planck Institute for the Science of Human History, Kahlaische Stasse 10, D-07743, Jena, Germany.

³ Australian Research Centre for Human Evolution (ARCHE), Environmental Futures Research Institute, Griffith University, Nathan, QLD 4111, Australia.

⁴ Research School of Earth Sciences, The Australian National University, Canberra ACT, 0200, Australia.

⁵ Saudi Geological Survey, Sedimentary Rocks and Palaeontology Department, Jeddah 21514, Saudi Arabia.

⁶ Department of Geography, King's College London, Strand, London WC2R 2LS, UK

⁷ Department of Geography, Royal Holloway, University of London, TW20 0EX UK

⁸ SFF Centre for Early Sapiens Behaviour (SapienCE), University of Bergen, Post Box 7805, 5020, Bergen, Norway.

⁹ Geochronology, Centro Nacional de Investigación sobre la Evolución (CENIEH), Paseo Sierra de Atapuerca, 3, 09002 Burgos, Spain

27 ¹⁰ PAVE Research Group, Dept. of Archaeology, University of Cambridge, Cambridge, CB2 3QG,
28 UK.

29 ¹¹ Earth Sciences Department, Natural History Museum, Cromwell Road, London, SW7 5BD, UK.

30 ¹² Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent,
31 Canterbury, CT2 7NR, UK.

32 ¹³ Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher
33 Platz 6, 04103, Leipzig, Germany.

34 ¹⁴ School of Natural Sciences and Psychology, Liverpool John Moores University, Liverpool, L3 3AF,
35 UK.

36 ¹⁵ Department of Anthropology, University of Western Ontario, London, Ontario, N6A 3K7, Canada.

37 ¹⁶ Palaeontology, Geobiology and Earth Archives Research Centre, School of Biological, Earth and
38 Environmental Science, University of New South Wales, Sydney 2052, Australia.

39 ¹⁷ School of Earth and Environmental Sciences, The University of Queensland, St Lucia 4072,
40 Queensland, Australia.

41 ¹⁸ Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD,
42 UK.

43 ¹⁹ Department of Archaeology, King Saud University, Riyadh, 12372 Saudi Arabia.

44 ²⁰ Saudi Commission for Tourism and National Heritage, Riyadh 11586, Saudi Arabia.

45 ²¹ Department of Archaeology, Hazara University, Manshra, 21300, Pakistan.

46 ²² Human Origins Program, National Museum of Natural History, Smithsonian Institution,
47 Washington, DC, 20560, USA.

48

49 Corresponding authors: HSG (huw.groucutt@rlaha.ox.ac.uk) and MDP (petraglia@shh.mpg.de)

50

51

52

53 Understanding the timing and character of *Homo sapiens* expansion out of Africa is critical
54 for inferring the colonisation and admixture processes that underpin global population
55 history. It has been argued that dispersal out of Africa had an early phase, particularly ~130-
56 90 thousand years ago (ka), that only reached the East Mediterranean Levant, and a later
57 phase, ~60-50 ka, that extended across the diverse environments of Eurasia to Sahul.
58 However, recent findings from East Asia and Sahul challenge this model. Here we show that
59 *H. sapiens* was in the Arabian Peninsula before 85 ka. We describe the Al Wusta-1 (AW-1)
60 intermediate phalanx from the site of Al Wusta in the Nefud Desert, Saudi Arabia. AW-1 is
61 the oldest directly dated fossil of our species outside Africa and the Levant. The
62 palaeoenvironmental context of Al Wusta demonstrates that *H. sapiens* using Middle
63 Palaeolithic stone tools dispersed into Arabia during a phase of increased precipitation driven
64 by orbital forcing, in association with a primarily African fauna. A Bayesian model
65 incorporating independent chronometric age estimates indicates a chronology for Al Wusta of
66 ~95-86 ka, which we correlate with a humid episode in the later part of Marine Isotope Stage
67 5 known from various regional records. Al Wusta shows that early dispersals were more
68 spatially and temporally extensive than previously thought. Early *H. sapiens* dispersals out of
69 Africa were not limited to winter rainfall-fed Levantine Mediterranean woodlands
70 immediately adjacent to Africa, but extended deep into the semi-arid grasslands of Arabia,
71 facilitated by periods of enhanced monsoonal rainfall.

72

73 **Background**

74

75 *Homo sapiens* evolved in Africa in the late Middle Pleistocene¹. Early dispersals out of
76 Africa are evidenced at the Levantine site of Misliya at ~194-177 ka², followed by Skhul and
77 Qafzeh, where *H. sapiens* fossils have been dated to ~130-100 and ~100-90 ka respectively³.

78 While the Levantine fossil evidence has been viewed as the onset of a much broader dispersal
79 into Asia^{4,6}, it has generally been seen as representing short-lived incursions into the
80 woodlands of the Levant immediately adjacent to Africa, where relatively high precipitation
81 is produced by winter storms tracking across the Mediterranean^{7,8}. While the Levantine
82 record indicates the subsequent local replacement of early *H. sapiens* by Neanderthals, the
83 failure of early dispersals to extend beyond the Levant is largely inferred from interpretations
84 of genetic data⁹. Genetic studies have suggested that recent non-African populations stem
85 largely¹⁰, if not entirely⁹, from an expansion ~60-50 ka, but this model remains debated. The
86 absence of low latitude Pleistocene human DNA and uncertainties regarding ancient
87 population structure undermine conclusions drawn from genetic studies alone. The paucity of
88 securely dated archaeological, palaeontological and ancient DNA data - particularly across
89 southern Asia - has made testing dispersal hypotheses challenging^{4,7,11}.

90

91 Recent fossil discoveries in East Asia indicate that the early (particularly Marine Isotope
92 Stage 5) dispersals of *Homo sapiens* extended across much of southern Asia. At Tam Pa Ling
93 in Laos, *Homo sapiens* fossils date to between 70 and 46 ka¹². Teeth assigned to *Homo*
94 *sapiens* from Lida Ajer cave, Sumatra, were recovered from a breccia dating to 68 ± 5 ka,
95 with fauna from the site dating to 75 ± 5 ka¹³. Several sites in China have produced fossil
96 material claimed to represent early *Homo sapiens*¹⁴. These include teeth from Fuyan Cave
97 argued to be older than 80 ka based on the dating of an overlying speleothem a few metres
98 from the fossils¹⁵, and teeth from Luna Cave that were found in a layer dating to between
99 129.9 ± 1.5 ka and 70.2 ± 1.4 ka¹⁶. Teeth and a mandible from Zhiren Cave, China, date to at
100 least 100 ka and have been argued to represent *Homo sapiens*, but other species attributions
101 are possible¹⁷. The recent documentation of a human presence in Australia from ~65 ka is
102 consistent with these findings¹⁸. Likewise, some interpretations of genetic data are consistent

103 with an early spread of *Homo sapiens* across southern Asia¹⁰. These discoveries are leading
104 to a radical revision of our understanding of the dispersal of *Homo sapiens*, yet there remain
105 stratigraphic and taxonomic uncertainties for many of the east Asian fossils^{14,19}, and
106 thousands of kilometers separate these findings from Africa.

107

108 The Arabian Peninsula is a vast landmass at the crossroads of Africa and Eurasia. Growing
109 archaeological evidence demonstrates repeated hominin occupations of Arabia^{20,21} each
110 associated with a strengthened summer monsoon which led to the re-activation of lakes and
111 rivers²²⁻²⁴, as it did in North Africa²⁵. Here we report the discovery of the first pre-Holocene
112 human fossil in Arabia, Al Wusta-1 (AW-1), as well as the age, stratigraphy, vertebrate
113 fossils and stone tools at the Al Wusta site (Fig. 1, see also Supplementary Information).

114

115 ***Figure 1 hereabouts***

116

117 **Results**

118 AW-1 is an intermediate manual phalanx, most likely from the 3rd ray (Fig. 2a,
119 Supplementary Information 1: see below for detail on siding and species identification). It is
120 generally well-preserved, although there is some erosion of the cortical/subchondral bone,
121 and minor pathological bone formation (likely an enthesophyte) affecting part of the
122 diaphysis (Supplementary Information 1). The phalanx measures 32.3 mm in proximo-distal
123 length, and 8.7 mm and 8.5mm in radio-ulnar breadth of the proximal base and midshaft,
124 respectively (Supplementary Table 1).

125

126 AW-1 is more gracile than the robust intermediate phalanges of Neanderthals²⁶⁻²⁸, which are
127 broader radio-ulnarly relative to their length and have a more 'flared' base. AW-1's proximal

128 radio-ulnar maximum breadth is 14.98 mm, which provides an intermediate phalanx breadth-
129 length index (proximal radio-ulnar maximum breadth relative to articular length) of 49.6.
130 This is very similar to the mean (\pm SD) for the Skhul and Qafzeh *H. sapiens* of 49.7 (\pm 4.1)
131 and 49.1 (\pm 4.0) for Upper Palaeolithic Europeans, but 1.89 standard deviations below the
132 Neanderthal mean of 58.3 (\pm 4.6)²⁹.

133

134 ***Figure 2 hereabouts***

135

136 To provide a broad interpretive context for the Al Wusta phalanx, we conducted linear and
137 geometric morphometric (GMM) landmark analyses (Supplementary Information 1) on
138 phalanges from non-human primates, fossil hominins and geographically widespread recent
139 *H. sapiens*. Comparative linear analyses (Supplementary Information 1, Supplementary
140 Tables 2 and 3, Supplementary Figure 1) reveal that there is substantial overlap across most
141 taxa for all shape ratios, so AW-1 falls within the range of variation of *H. sapiens*, cercopiths,
142 *Gorilla*, *Australopithecus afarensis*, *A. sediba* and Neanderthals. However, AW-1 is most
143 similar to the median value or falls within the range of variation of recent and early *H.*
144 *sapiens* for all shape ratios.

145

146 Geometric morphometric (GMM) analyses of AW-1 and various primate groups including
147 hominins (see Supplementary Table 4 and Supplementary Figure 2 for landmarks, and
148 Supplementary Table 5 for sample) are illustrated in Figure 3 and Supplementary Figure 3.
149 PC1 and PC2 together account for 61% of group variance in shape. AW-1 is separated on
150 these two shape vectors from the non-human primates and most of the Neanderthals. AW-1
151 falls closest to the recent and early *H. sapiens* and is clearly differentiated from all non-

152 human primates. This is also shown by the Procrustes distances from AW-1 to the mean
153 shapes of each taxonomic group (Supplementary Table 6).

154

155 ***Figure 3 hereabouts***

156

157 Three of the Neanderthal phalanges (from Kebara 2 and Tabun C1) are quite disparate from
158 the main Neanderthal cluster and fall closer to the *H. sapiens* and Al Wusta cluster on PC1
159 and 2 (Figure 3 and Supplementary Figure 3). Having established the hominin affinity of
160 AW-1, shape was analysed in more detail using a smaller hominin sample for which ray
161 number and side were known, which included Kebara 2 and Tabun C1. The broader primate
162 sample used in the first GMM analysis was not used for the more detailed shape analysis, as
163 the initial comparisons show clearly that AW-1 is not a non-human primate and including
164 this level of variation could potentially mask more subtle shape differences between
165 hominins. The side and ray are also not known for most of the Neanderthal and non-human
166 primate samples, meaning it would be impossible to evaluate the effect of these factors using
167 this sample.

168

169 The more in-depth shape comparison and modelling using the hominin sample of phalanges
170 of known ray and side (Supplementary Table 7) demonstrates that the long and slender
171 morphology of AW-1 falls just outside the range of variation of comparative Middle
172 Palaeolithic modern humans, but that its affinity is clearly with *H. sapiens* rather than
173 Neanderthals (Fig. 4, Supplementary Table 8). Although both Pleistocene *H. sapiens* and
174 Neanderthal landmark configurations fall almost completely inside the scatter for the
175 Holocene *H. sapiens* sample in the principal components analysis (Figure 4), AW-1 is closest
176 to Holocene *H. sapiens* 3rd intermediate phalanges. AW-1 overlaps with the Holocene *H.*

177 *sapiens* sample, but is separated from the Pleistocene *H. sapiens* specimens by a higher score
178 on PC2 and from the Neanderthal group by a simultaneously higher score on PC1 and PC2.
179 The Procrustes distances (Supplementary Table 8), also show that AW-1 is most distinct
180 from the Neanderthal phalanges, which fall towards the lower ends of both PCs and are
181 characterised by shorter and broader dimensions. PC1 and PC2 in this analysis show that
182 AW-1 is taller and narrower (in all directions: dorso-palmarly, proximo-distally and radio-
183 ulnarly) than almost all the phalanges in the comparative sample and is particularly distinct
184 from most of the Neanderthal phalanges. In this analysis AW-1 is closest in shape to 3rd
185 phalanges of individuals from (in descending order of proximity) Egyptian Nubia, and
186 Medieval Canterbury (UK), and Maiden Castle (Iron Age Dorset, UK) (Supplementary Table
187 9), although there is not a great difference in its distance to any of these specimens. These
188 analyses suggest that the AW-1 phalanx is likely to be a 3rd intermediate phalanx from a *H.*
189 *sapiens* individual.

190

191 ***Figure 4 hereabouts***

192

193 The third ray is the most symmetrical ray in the hand and is therefore difficult to side,
194 particularly when not all of the phalanges of a particular individual are present. Comparing
195 AW-1 separately to right and to left phalanges (Supplementary Information 1.4) gives results
196 which are very similar to the pooled sample, such that AW-1 is closest to Holocene *H.*
197 *sapiens* 3rd rays for both right and left hands (Supplementary Figure 4, Supplementary Table
198 10). There is little difference in morphological closeness between AW-1 and its nearest
199 neighbour in the samples of right and left bones (Supplementary Table 11), reflecting the lack
200 of difference in morphology between the sides. It is therefore not possible to suggest whether
201 AW-1 comes from a right or a left hand using these analyses.

202

203 AW-1 is unusual in its more circular midshaft cross-sectional shape (Fig. 2B), which is
204 confirmed by cross-sectional geometric analyses (Supplementary Information 1.5). This may
205 reflect the pronounced palmar median bar that makes the palmar surface slightly convex at
206 the midshaft rather than flat, the latter being typical of most later *Homo* intermediate
207 phalanges. However, more circular shafts may reflect greater loading of the bone in multiple
208 directions and enthesophytes are a common response to stress from high levels of physical
209 activity³⁰. This morphology may reflect high and varied loading of the fingers during intense
210 manual activity.

211

212 To determine the age of AW-1, and associated sediments and fossils, we used a combination
213 of uranium series (U-series), electron spin resonance (ESR) and optically stimulated
214 luminescence (OSL) dating (Methods, Supplementary Information 2 and 3). U-series ages
215 were produced for AW-1 itself (87.6 ± 2.5 ka) and hippopotamus dental tissues (WU1601),
216 which yielded ages of 83.5 ± 8.1 ka (enamel) and 65.0 ± 2.1 ka (dentine). They should be
217 regarded as minimum estimates for the age of the fossils. In addition, a combined U-series-
218 ESR age calculation for WU1601 yielded an age of $103 +10/-9$ ka. AW-1 was found on an
219 exposure of Unit 3b, and WU1601 excavated from Unit 3a, one metre away (Fig 1b). Unit 1
220 yielded OSL ages of 85.3 ± 5.6 ka (PD17), 92.2 ± 6.8 ka (PD41) and 92.0 ± 6.3 ka (PD15),
221 while Unit 3a yielded an OSL age of 98.6 ± 7.0 ka (PD40). The OSL age estimates agree
222 within error with the US-ESR age obtained for WU-1601 and the minimum age of ~ 88 ka
223 obtained for AW-1. These data were incorporated into a Bayesian sequential phase model³¹
224 which indicates that deposition of Unit 1 ceased 93.1 ± 2.6 ka (Phase 1: PD15, 17, 41) and
225 that Units 2 and 3 and all associated fossils were deposited between 92.2 ± 2.6 ka and $90.4 \pm$
226 3.9 ka (Phase 2: all other ages) (Supplementary Information 4, Supplementary Figure 11).

227

228 This ~95-86 ka timeframe is slightly earlier than most other records of increased humidity in
229 the region in late MIS 5^{32,33}, which correlate with a strengthened summer monsoon
230 associated with an insolation peak at 84 ka (Fig. 6). The underlying (Unit 3) aeolian sand
231 layer at Al Wusta correlates with an insolation minimum at the end of MIS 5c. The
232 chronometric age estimates for the site suggest that lake formation and the associated fauna
233 and human occupation occurred shortly after this in time. Regional indications of increased
234 humidity around the 84 ka insolation peak include speleothem formation at ~88 ka in the
235 Negev³⁴, and the formation of sapropel S3 beginning ~86 ka³⁵. In both the Levant and
236 Arabia, records are consistent with this switch from aridity to humidity around this time³²⁻⁴⁰.
237 Precisely reconstructing regional palaeoclimate at this time and relating it to human
238 demographic and behavioural change has proved challenging. This reflects both rapid
239 changes in climate, as well as the complexities involved in dating relevant deposits⁴¹. In
240 summary, combining chronological data (Supplementary sections 2-4), interpretation of the
241 sedimentary sequence (described below), and the regional setting of Al Wusta, we conclude
242 that lake formation and associated finds such as the AW-1 phalanx relate to the late MIS 5
243 humid period associated with the 84 ka insolation peak.

244

245 The sedimentary sequence at Al Wusta consists of a basin-like deposit of exposed carbonate-
246 rich sediments (Unit 2, 0.4-0.8 m thick), underlain by wind-blown sand (Unit 1) and overlain
247 by water-lain sands (Unit 3). The carbonate rich sediments of Unit 2 are interpreted as
248 lacustrine marl deposits on the basis of their sedimentology, geochemistry, and diatom
249 palaeoecology (Figure 1c, Methods, Supplementary Information 5). At both the macro- and
250 micro-scale, these beds are relatively massive and comprise fine-grained calcite, typical of
251 material precipitating and accumulating in a still-water lacustrine environment⁴². At the

252 micro-scale there is no evidence for the desiccation or fluctuation of water levels typical of
253 palustrine/wetland environments⁴², implying that the lake body was perennial. The diatom
254 flora support this, containing species such as *Aulacoseira italica* and *Aulacoseira granulata*
255 throughout the sequences, indicating an alkaline lake a few metres deep. The water was fresh,
256 not saline or brackish, since saline tolerant species and evaporitic minerals are absent
257 throughout. While $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of continental carbonates are controlled by a wide-
258 range of variables, the values derived from the Al Wusta marl beds are compatible with the
259 suggestion of marls precipitated in a perennial lake basin. The Al Wusta carbonate beds
260 therefore indicate a perennial lake body a few metres in depth. The existence of a marl
261 precipitating lake basin implies that this system was groundwater fed (to allow for sufficient
262 dissolved mineral material to be present in the lake waters). Although the Al Wusta sequence
263 represents a single lake basin, the development of such a feature over highly permeable
264 aeolian sands in a region where no lake systems exist at the present day implies a local
265 increase in water table that would require an increase in mean annual rainfall. Consequently,
266 the Al Wusta sequence represents the occurrence of a humid interval at this time. The Unit 2
267 marl is overlain by a medium-coarse sand (Unit 3) with crude horizontal laminations,
268 occasional clasts, fragments of ripped up marl and shells of *Melanooides tuberculata* and
269 *Planorbis* sp. While some vertebrate fossils and lithics were found in the upper part of Unit 2,
270 most were found in or on the surface of Unit 3. Unit 3a sands are waterlain and represent the
271 encroachment of fluvial sediment as the lake environment shallowed and contracted. Unit 3b
272 represents a winnowed lag formed by aeolian deflation of 3a. The sequence is capped by a
273 dense network of calcitic rhizoliths marking the onset of fully terrestrial conditions.
274
275 A total of 860 vertebrate fossils were excavated from Unit 3 and the top of Unit 2 (n=371)
276 and systematically surface collected (n=489). These include specimens attributed to Reptilia,

277 Aves, and Mammalia (Supplementary Table 19, Methods, Supplementary Information 6).
278 Notable taxa now extinct in Arabia are predominately grazers and include *Hippopotamus*,
279 *Pelorovis*, and *Kobus*. The faunal community demonstrates a clear preference for temperate
280 to semi-arid grasslands, and the presence of *Hippopotamus* and *Kobus* indicate permanent
281 muddy, fluvial, or lacustrine conditions⁴³ not currently found in the Nefud Desert, but
282 consistent with the geological evidence from the site. The faunal assemblages show a strong
283 affinity to African fauna, particularly *Hippopotamus*, *Pelorovis*, and *Kobus*⁴⁴. Many large
284 tooth pits on fossils indicate that large carnivores played a role in the accumulation of the
285 deposit. Long bone circumference, completeness and numbers of green fractures suggests
286 modification of bones by bone-breaking agents such as large carnivores or hominins
287 (Supplementary Information 6). However, no evidence of cut-marks or hammerstone damage
288 to the bones was observed.

289

290 An assemblage of 380 lithic artefacts (stone tools) was recovered from the excavation of
291 upper Unit 2 and Unit 3 and systematic surface collection (Methods, Figure 5, Supplementary
292 Information 7). They are of Middle Palaeolithic character and most are chert and quartzite.
293 The assemblage demonstrates a focus on centripetal Levallois reduction, and is similar to
294 other late Marine Isotope Stage 5 assemblages in the west and north of Arabia⁴⁵, and
295 contemporaneous assemblages in east (e.g. Aduma, BNS at Omo Kibish) and northeast
296 Africa (e.g. Bir Tarfawi), as well as those from the Levant (e.g. Qafzeh)¹¹ (Fig. 5).

297

298 ***Figure 5 hereabouts***

299 ***Figure 6 hereabouts***

300

301

302 **Discussion**

303

304 Al Wusta-1 is the oldest directly dated *H. sapiens* fossil outside Africa and the Levant. It
305 joins a small but growing corpus of evidence that the early dispersal of *H. sapiens* into
306 Eurasia was much more widespread than previously thought. The site of Al Wusta is located
307 in the Nefud desert more than 650 km southeast of Skhul and Qafzeh (Fig. 1A). This site
308 establishes that *H. sapiens* were in Arabia in late MIS 5, rather than being restricted to Africa
309 and the Levant as suggested by traditional models (Fig. 6). With Skhul dating to ~130-100
310 ka, Qafzeh to ~100-90 ka^{3,46} and Al Wusta to ~95-85 ka it is currently unclear if the
311 southwest Asian record reflects multiple early dispersals out of Africa or a long occupation
312 during MIS 5. The association of the Al Wusta site with a late MIS 5 humid phase (Fig. 6),
313 suggests that significant aspects of this dispersal process were facilitated by enhanced
314 monsoonal rainfall. While changes in behaviour and demography are crucial to understanding
315 the dispersal process, climatic windows of opportunity were also key in allowing *H. sapiens*
316 to cross the Saharo-Arabian arid belt, which often constituted a formidable barrier^{24,25}.

317

318 **Conclusion**

319

320 Al Wusta shows that the early, Marine Isotope Stage 5, dispersals of *H. sapiens* out of Africa
321 were not limited to the Levantine woodlands sustained by winter rainfall, but extended deep
322 into the Arabian interior where enhanced summer rainfall created semi-arid grasslands
323 containing abundant fauna and perennial lakes. After long being isolated in Africa^{1,47,48}, the
324 Late Pleistocene saw the expansion of our species out of Africa and into the diverse ecologies
325 of Eurasia. Within a few thousand years of spreading into Eurasia our species was occupying
326 rainforest environments and making long sea crossings to remote islands^{13,18}. Adapting to the

327 semi-arid conditions of the Saharo-Arabian arid belt represented a crucial step on this
328 pathway to global success and the Al Wusta *Homo sapiens* fossil demonstrates this early
329 ability to occupy diverse ecologies which led to us becoming a cosmopolitan species.

330

331 **References**

332

- 333 1. Stringer, C. The origin and evolution of *Homo sapiens*. *Philos. Trans. R. Soc. B. Biol.*
334 *Sci.* **371**, 20150237-20150237 (2016).
- 335 2. Hershovikitz, I., *et al.* The earliest modern humans outside Africa. *Science* 359, 456-
336 459 (2018).
- 337 3. Grün, R. *et al.* U-series and ESR analyses of bones and teeth relating to the human
338 burials from Skhul. *J. Hum Evol.* **49**, 316-334 (2005).
- 339 4. Groucutt, H. S. *et al.* Rethinking the dispersal of *Homo sapiens* out of Africa. *Evol.*
340 *Anthropol.* **24**, 149-164 (2015).
- 341 5. Petraglia, M. D. *et al.* Middle Paleolithic assemblages from the Indian subcontinent
342 before and after the Toba super-eruption. *Science* **317**, 114-116 (2007).
- 343 6. Bae, C.J., Douka, K., Petraglia, M.D. On the origin of modern humans: Asian
344 perspectives. *Science* **358**, DOI: 10.1126/science.aai9067 (2017).
- 345 7. Mellars, P., Gori, K. C., Carr, M., Soares, P. A. & Richards, M. B. Genetic and
346 archaeological perspectives on the initial modern human colonization of southern
347 Asia. *Proc. Natl Acad. Sci. USA* **110**, 10699–10704 (2013).
- 348 8. Shea, J. J. Transitions or turnovers? Climatically-forced extinctions of *Homo sapiens*
349 and Neanderthals in the east Mediterranean Levant. *Quatern. Sci. Rev.* **27**, 2253-2270
350 (2008).

- 351 9. Mallick, S. *et al.* The Simons Genome Diversity Project: 300 genomes from 142
352 diverse populations. *Nature* **538**, 201-206 (2016).
- 353 10. Pagani, L. *et al.* Genomic analyses inform on migration events during the peopling of
354 Eurasia. *Nature* **538**, 238-242 (2016).
- 355 11. Groucutt, H. S. *et al.* Stone tool assemblages and models for the dispersal of *Homo*
356 *sapiens* out of Africa. *Quatern. Int.* **382**, 8-30 (2015).
- 357 12. Demeter, F., *et al.* Early Modern Humans from Tam Pà Ling, Laos. Fossil Review
358 and Perspectives. *Curr. Anthropol.* **57**, S17, DOI: 10.1086/694192 (2017)
- 359 13. Westaway, K.E., *et al.* An early modern human presence in Sumatra 73,000-63,000
360 years ago. *Nature* **548**, 322-325 (2017).
- 361 14. Michel, V., *et al.* The earliest modern *Homo sapiens* in China? *J. Hum. Evol.* **101**,
362 101-104 (2016).
- 363 15. Liu, W., *et al.* The early unequivocally modern humans in southern China? *Nature*
364 **526**, 696-699 (2015).
- 365 16. Bae, C., *et al.* Modern human teeth from Late Pleistocene Luna Cave (Guangxi,
366 China). *Quatern. Int.* **354**, 169-183 (2015).
- 367 17. Liu, W., *et al.* Human remains from Zhiredong, South China, and modern human
368 emergence in East Asia. *Proc. Natl. Acad. Sci. USA* **107**, 19201-19206 (2010).
- 369 18. Clarkson, C., *et al.* Human occupation of northern Australia by 65,000 years ago.
370 *Nature* **547**, 306-310 (2017).
- 371 19. Martínón-Torres, M., Wu, X., de Castro, J.M.B., Xing, S., Liu, W. *Homo sapiens* in
372 the Eastern Asian Late Pleistocene. *Curr. Anthropol.* **58**, S17, DOI: 10.1086/694449.
- 373 20. Groucutt, H. S. & Petraglia, M. D. The prehistory of Arabia: Deserts, dispersals and
374 demography. *Evol. Anthropol.* **21**, 113-125 (2012).

- 375 21. Petraglia, M. D., Groucutt, H. S., Parton, A. & Alsharekh, A., Green Arabia: Human
376 prehistory at the Cross-Roads of continents. *Quatern. Int.* **382**, 1-7 (2015).
- 377 22. Jennings, R. P. The greening of Arabia: Multiple opportunities for human occupation
378 in the Arabian Peninsula during the Late Pleistocene inferred from an ensemble of
379 climate model simulations. *Quatern. Int.* **205**, 181-199 (2015).
- 380 23. Rosenberg, T. M. *et al.* Middle and Late Pleistocene humid periods recorded in
381 palaeolake deposits in the Nafud desert, Saudi Arabia. *Quatern. Sci. Rev.* **70**, 109-123
382 (2013).
- 383 24. Breeze, P. S. *et al.* Palaeohydrological corridors for hominin dispersals in the Middle
384 East ~250-70,000 years ago. *Quatern. Sci. Rev.* **11**, 155-185 (2016).
- 385 25. Scerri, E. M. L., Drake, N. A., Jennings, R., Groucutt, H.S. Earliest evidence for the
386 structure of *Homo sapiens* populations in Africa. *Quatern. Sci. Rev.* **101**, 207-216
387 (2014).
- 388 26. Trinkaus, E. *The Shanidar Neandertals*. (Academic Press, New York, 1981).
- 389 27. McCown T. D. & Keith, A. *The Stone Age of Mount Carmel Volume 2: The fossil*
390 *human remains from the Levalloiso-Mousterian* (Clarendon Press, Oxford, 1939).
- 391 28. Vandermeersch, B. *Les hommes fossiles de Qafzeh (Israel)* (CNRS, Paris, 1981).
- 392 29. Walker, M. J., Ortega, J., López, M. V., Parmová, K. & Trinkaus, E. Neanderthal
393 postcranial remains from the Sima de las Palomas del Cabezo Gordo, Murcia,
394 Southeastern Spain. *Am. J. Phys. Anthropol.* **144**, 505-515 (2011).
- 395 30. Benjamin, M. *et al.* Where tendons and ligaments meet bone: attachment sites
396 ('entheses') in relation to exercise and/or mechanical load. *J. Anat.* **208**, 471-490.
- 397 31. Bronk Ramsey, C. Bayesian analysis of radiocarbon dates. *Radiocarbon* **51**, 337-360
398 (2009).

- 399 32. Drake, N. A., Breeze, P., Parker, A., 2013. Palaeoclimate in the Saharan and Arabian
400 Deserts during the Middle Palaeolithic and the potential for hominin dispersals.
401 *Quatern. Int.* **300**, 48-61 (2013).
- 402 33. Parton, A., et al. Orbital-scale climate variability in Arabia as a potential motor for
403 human dispersals. *Quatern. Int.* **382**, 82-97 (2015).
- 404 34. Vaks, A., Bar-Matthews, M., Matthews, A., Ayalon, A., Frumkin, A. Middle-Late
405 Quaternary paleoclimate of northern margins of the Saharan-Arabian Desert:
406 reconstruction from speleothems of Negev Desert, Israel. *Quatern. Sci. Rev.* **29**, 2647-
407 2662 (2010).
- 408 35. Grant, K.M., et al. The timing of Mediterranean sapropel deposition relative to
409 insolation, sea-level and African monsoon changes. *Quatern. Sci. Rev.* **140**, 125-141
410 (2016).
- 411 36. Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A. & Hawkesworth, C. J.
412 Sea-land oxygen isotopic relationships from planktonic foraminifera and speleothems
413 in the Eastern Mediterranean region and their implication for paleorainfall during
414 interglacial intervals. *Geochim. Cosmochim. Acta* **67**, 3181–3199 (2003).
- 415 37. Lisiecki, L. E. & Raymo, M. E. A Pliocene-Pleistocene stack of 57 globally
416 distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography* **20**, 1-17 (2005).
- 417 38. Berger, A. & Loutre, M. F. Insolation values for the climate of the last 10 million
418 years. *Quatern. Sci. Rev.* **10**, 297-317 (1991).
- 419 39. Fleitmann, D., Burns, S.J., Neff, U., Mangini, A., Matter, A. Changing moisture
420 sources over the last 333,000 years in Northern Oman from fluid-inclusion evidence
421 in speleothems. *Quatern. Res.* **60**, 223-232 (2003).
- 422 40. Rosenberg, T.M., et al. Humid periods in southern Arabia: Windows of opportunity
423 for modern human dispersal. *Geology* **39**, 1115-1118 (2011).

- 424 41. Clark-Balzan, L., Parton., A., Breeze, P.S., Groucutt, H.S., Petraglia, M.D. Resolving
425 problematic luminescence chronologies for carbonate- and evaporite-rich sediments
426 spanning multiple humid periods in the Jubbah Basin, Saudi Arabia. *Quatern.*
427 *Geochron.* <https://doi.org/10.1016/j.quageo.2017.06.002> (2018).
- 428 42. Alonso-Zarza, A. M. Palaeoenvironmental significant of palustrine carbonates and
429 calcretes in the geological record. *Earth Sci. Rev.* **60**, 261-298 (2003).
- 430 43. Estes, R. D. *The Behaviour Guide to African Mammals* (University of California
431 Press, Berkeley, 1991).
- 432 44. O'Regan, H. J., Turner, A., Bishop, L. C., Elton, S., Lamb, A. L. Hominins without
433 fellow travellers? First appearances and inferred dispersals of Afro-Eurasian large-
434 mammals in the Plio-Pleistocene. *Quatern. Sci. Rev.* **30**, 1343-1352 (2011).
- 435 45. Groucutt, H.S. et al. Human occupation of the Arabian Empty Quarter during MIS 5:
436 evidence from Mundafan al-Buhayrah. *Quatern. Sci. Rev.* **119**, 116-135 (2015).
- 437 46. Millard, A. R. A critique of the chronometric evidence for hominid fossils: I. Africa
438 and the Near East 500-50 ka. *J. Hum. Evol.* **54**, 848-874 (2008).
- 439 47. Hublin, J.J., et al. New fossils from Jebel Irhoud, Morocco and the pan-African origin
440 of *Homo sapiens*. *Nature* **546**, 289-292 (2017).
- 441 48. Richter, D., et al. The age of the hominin fossils from Jebel Irhoud, Morocco, and the
442 origins of the Middle Stone Age. *Nature* **546**, 293-296 (2017).
- 443
444
445
446
447
448

449 **Supplementary Information** is available in the online version of the paper.

450

451 **Acknowledgements.** We thank HRH Prince Sultan bin Salman bin Abdulaziz Al-Saud,
452 President of the Saudi Commission for Tourism and National Heritage (SCTH), and Prof. Ali
453 Ghabban, Vice President of the SCTH for permission to carry out this study. Dr Zohair
454 Nawab, President of the Saudi Geological Survey, provided research support and logistics.
455 Fieldwork and analyses were funded by the European Research Council (no. 295719, to MDP
456 and 617627, to JTS), the SCTH, the British Academy (HSG and EMLS), The Leverhulme
457 Trust, the Australian Research Council (DP110101415 to RG, FT150100215 and
458 TF15010025 to MD, and FT160100450 to JL), and the Research Council of Norway (SFF
459 Centre for Early Sapiens Behaviour, 262618). We thank Patrick Cuthbertson, Klint Janulis,
460 Marco Bernal, Salih Al-Soubhi, Mohamad Haptari, Adel Matari, and Yahya Al-Mufarreh for
461 assistance in the field. We thank Ian Cartwright (Institute of Archaeology, University of
462 Oxford) for the photographs of AW-1 (Fig. 2a), Ian Matthews (RHUL) for producing the
463 Bayesian age model, and Michelle O'Reilly (MPI-SHH) for assistance with the preparation
464 of figures. We acknowledge the Max Planck Society for supporting us with comparative
465 fossil data, and we thank curators for access to comparative extant and fossil material in their
466 care (Supplementary Tables 5 and 7).

467

468 **Author Contributions** H.S.G. and M.D.P. designed, coordinated and supervised the study.
469 H.S.G., I.S.Z., N.D, S.A., I.C., R.C-W., J.L., P.S.B., M.S., G.J.P., A.A., A.A.-O., A.M. B.A.,
470 E.M.L.S. and M.D.P. conducted excavation, survey and multidisciplinary sampling at Al
471 Wusta. L.T.B., T.L.K., E.P., N.B.S and J.T.S. conducted the morphological analysis and
472 comparative study of the AW-1 phalanx. R.G., M.D. and L.K. carried out the U-series and
473 ESR analyses. S.J.A. and R.C.W carried out the OSL dating. I.C. and R.C.W conducted the

474 stratigraphic and sedimentological analysis of the site, with input from N.D., J.L. and G.J.P.
475 W.W.S. analysed the diatoms. M.S. and J.L. analysed the vertebrate fossils, with input from
476 G.J.P. Lithic analysis was conducted by H.S.G. and E.M.L.S. Spatial analyses were
477 conducted by P.S.B. All authors helped to write the paper.

478

479 **Author Information** The authors declare no competing financial interests.

480 Readers are welcome to comment on the online version of the paper.

481 Correspondence and requests for materials should be addressed to

482 H.S.G. (huw.groucutt@rlaha.ox.ac.uk) or M.D.P. (petraglia@shh.mpg.de).

483

484 **Data availability statement.** Authors can confirm that all relevant data are included in the
485 paper and/ or its supplementary information files.

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500 **Figure captions**

501

502 **Figure 1. Al Wusta location, map of site and stratigraphy.** A: The location of Al Wusta
503 and other key MIS 5 sites in the region¹¹; B: Al Wusta digital elevation model showing
504 location of AW-1 phalanx, marl beds, lithics and vertebrate fossils, and the locations of the
505 trenches and sections. The inset shows a satellite image of the site; C: Stratigraphic log of Al
506 Wusta showing the sedimentology of the exposed carbonate beds, isotopic values, OSL ages
507 for sand beds and U-series and ESR ages for AW-1 and WU-1601. Sands are shown in
508 yellow: lower massive sands are aeolian (Unit 1), upper laminated sands are waterlain (Unit
509 3a) and have been locally winnowed to generate a coarse desert pavement (Unit 3b),
510 lacustrine marls are shown (Unit 2) in grey (for full key and description see Supplementary
511 Figures 13 and 14 and Supplementary Information 5). Section PD40 is shown as it contains
512 the thickest sequence and is most representative of Al Wusta, chronometric age estimates
513 (marked *) from the site are depicted in their relative stratigraphic position, see
514 Supplementary Figure 14 for their absolute positions.

515

516 **Figure 2. Photographs and micro-CT scans of Al Wusta-1 *Homo sapiens* phalanx.** A:
517 photographs in (left column, top to bottom) distal, palmar and proximal views, and (middle
518 row, left to right) lateral 1, dorsal and lateral 2 views. Micro-CT cross-sections (illustrated at
519 2x magnification) include B (54% from proximal end) and C (illustrating abnormal bone).

520

521 **Figure 3. Scatterplot of the first two principal components (PC) scores of the geometric**
522 **morphometric analysis of the Al Wusta-1 phalanx compared with a sample of primates,**
523 **including hominins.** Non-human hominoids: lilac; *Gorilla*: circles, *Pan*: triangles.

524 Cercopithecoids: red; *Colobus*: triangles, *Mandrillus*: squares, *Papio*: circles. Neanderthals:
525 blue diamonds. *H. sapiens*: green; early *H. sapiens*: circles, Holocene *H. sapiens*: squares. Al
526 Wusta-1: black star, circled in red.

527

528 **Figure 4: Scatterplot of the first two principal component (PC) scores from the**
529 **geometric morphometric analyses of AW-1 and sample of comparative hominin 2nd, 3rd,**
530 **and 4th intermediate phalanges.** Wireframes show mean configuration warped to extremes
531 of PC axes in dorsal (left), proximal (middle) and lateral (right) views. Convex hulls added
532 post-hoc to aid visualisation.

533

534 **Figure 5. Selected Al Wusta lithic artefacts.** A: argillaceous quartzite flake; B: quartz
535 hammerstone; C: ferruginous quartzite Levallois flake; D: chert Levallois flake; E: Quartz
536 recurrent centripetal Levallois core; F: quartzite preferential Levallois core with centripetal
537 preparation and pointed preferential removal.

538

539 **Figure 6. The chronological and climatic context of Al Wusta.** The Al Wusta lake phase
540 falls chronologically at the end of the time-range of MIS 5 sites from the Mediterranean
541 woodland of the Levant (~130-90 ka) and earlier than the late dispersal(s) (~60-50 ka) as
542 posited in particular by genetic studies. The chronology of these dispersals and occupations
543 correspond with periods of orbitally modulated humid phases in the eastern Mediterranean³⁶
544 that are important intervals for human dispersals into Eurasia, and are also proposed to
545 correspond with episodes of monsoon driven humidity in the Negev and Arabian desert³⁴.
546 Environmental amelioration of the Saharo-Arabian belt, therefore, appears to be crucial for
547 allowing occupation at key sites that document dispersal out of Africa. A: East Mediterranean
548 speleothem $\delta^{18}\text{O}$ record from Soreq and Pequin Caves³⁶; B: global $\delta^{18}\text{O}$ record³⁷; C:

549 Insolation at 30 degrees north³⁸, showing the temporal position of key sites relating to
550 dispersal out of Africa^{2,3,11,48}. The chronology for Al Wusta shows the phases defined by the
551 Bayesian model at 2σ .

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575 **Methods**

576

577 **Site identification, survey and excavation.** The site of Al Wusta (field code WNEF16_30)
578 was discovered in 2014 as part of a programme of joint survey fieldwork of the Palaeodeserts
579 Project, the Saudi Commission for Tourism and National Heritage, and the Saudi Geological
580 Survey. It is located in the western Nefud desert, a few kilometres from the Middle
581 Pleistocene fossil locality of Ti's al Ghaddah⁴⁹. The locations of all materials of interest
582 (fossils, stone tools, geomorphological features, excavations and sample points) were
583 recorded using a high-precision Trimble XRS Pro Differential GPS system and a total station,
584 and entered into a GIS (Fig. 1). Elevation data (masl) were recorded as a series of transects
585 across the site, and a digital elevation model (DEM) and contours interpolated (Spline) from
586 all data with precisions of better than 10 cm in all (x,y,z) dimensions (22,047 points). This
587 allowed visualisation and recording of the spatial relationships between materials in three
588 dimensions (Fig. 1). Eight trenches were excavated into the fossil and artefact bearing
589 deposits. These trenches revealed vertebrate remains and lithics, but no further human fossils
590 were recovered.

591

592 **Morphological analysis of Al Wusta-1 phalanx.** The phalanx was scanned using micro-
593 computed tomography (micro-CT) on the Nikon Metrology XT H 225 ST High Resolution
594 scanner and X-Tek software (Nikon Metrology, Tring, UK) housed in the Cambridge
595 Biotomography Centre, University of Cambridge, UK. Scan parameters were: a tungsten
596 target; 0.5 mm copper filter; 150 kV; 210 mA; 1080 projections with 1000 ms exposure, and
597 resulted in a voxel size of 0.02 mm³. The micro-CT data were reconstructed using CT-PRO
598 3D software (Nikon Metrology) and exported as an image (.tif) stack. Other CT data were

599 obtained from the institutions cited in Supplementary Table 5 with permissions following the
600 memoranda of understanding with each institution.

601

602 3D landmarks and semilandmarks were chosen to best describe the overall shape of the
603 morphology of the AW-1 phalanx (Supplementary Table 4, Supplementary Figure 2), and
604 were digitised on virtual reconstructions of phalanges created from micro-CT data in AVIZO
605 8 and 9.1 (FEI Software, Burlington, Mass.). Landmark coordinates were exported for use in
606 Morphologika⁵⁰. In Morphologika, generalized Procrustes analyses were performed to
607 superimpose landmark coordinate data, and principal components analyses (PCA) were run
608 to investigate similarities in shape between specimens. Shape differences along principal
609 componentss were visualised and wireframes were produced in Morphologika, PC scores
610 were exported to create graphs in R⁵¹. Procrustes distances between specimens were
611 calculated using MorphoJ⁵². To avoid representing the same phalanges from different sides of
612 a single individual as independent data points and to maximise sample sizes in pooled
613 analyses, right phalanges were used in cases where the phalanges from both sides were
614 present. Where only the left was present, this was used and ‘reflected’ (i.e. mirrored) in
615 Morphologika to generate landmark configurations consistent with right phalanges.

616

617 **U-series and combined US-ESR dating of fossil bone and teeth.** The AW-1 phalanx (lab
618 number 3675) and a hippopotamus tooth fragment (lab number WU1601) were collected
619 from Trench 1 (Fig.1) for U-series and combined US-ESR dating, respectively. The external
620 dose rate utilised the data of OSL sample PD40, which was collected in an equivalent
621 position within unit 3a.

622

623 *U-series analysis.* U-series analyses were conducted at the Research School of Earth
624 Sciences, The Australian National University, Canberra. The experimental setup for the U-
625 series analysis of the phalanx was described in detail by Grün and colleagues⁵³
626 (Supplementary Figures 2 and 3, Supplementary Information 2). Laser ablation (LA) was
627 used to drill a number of holes into AW-1 following the approach of Benson and
628 colleagues⁵⁴. After a cleaning run with the laser set at a diameter of 460 µm, seven holes were
629 drilled for 1000 s with the laser set at 330 µm. The isotopic data streams were converted into
630 $^{230}\text{Th}/^{234}\text{U}$ and $^{234}\text{U}/^{238}\text{U}$ activity ratios and apparent Th/U age estimates and subsequently
631 binned into 30 successive sections (each containing 33 cycles) for the calculation of average
632 isotopic ratios and ages. A similar experimental setup and methodology were employed for
633 the LA U-series analysis of tooth sample WU1601. The whole closed system U-series
634 analytical datasets of the enamel and dentine sections were integrated to provide the data
635 input for the ESR age calculations.

636

637 *Combined US-ESR dating of the fossil tooth: ESR dose evaluation.* The ESR dose evaluation
638 of the hippo tooth was carried out at CENIEH, Burgos, Spain, following a similar procedure
639 to that described in Stimpson and colleagues⁴⁹. Enamel was collected from WU1601 and
640 powdered <200 µm. The sample was then divided into 11 aliquots and gamma irradiated with
641 a Gammacell-1000 Cs-137 source to increasing doses until 3.4 kGy. ESR measurements were
642 carried out at room temperature with an EMXmicro 6/1 Bruker ESR spectrometer coupled to
643 a standard rectangular ER 4102ST cavity. ESR intensities were extracted from T1-B2 peak-
644 to-peak amplitudes of the ESR signal of enamel. Fitting procedures were carried out with a
645 single saturating exponential (SSE) function through the pooled ESR experimental data
646 derived from the repeated measurements, with data weighting by the inverse of the squared
647 ESR intensity ($1/I^2$) and following the recommendations by Duval and Grün⁵⁵. Full details

648 about the experimental conditions and analytical procedure may be found in Supplementary
649 Information 2.

650

651 *Combined US-ESR dating of the fossil tooth: Dose rate evaluation and age calculations.* The
652 combined US-ESR age of WU1601 was calculated with the DATA programme⁵⁶ using the
653 US model defined by Grün and colleagues⁵⁷. The following parameters were used for the
654 dose rate evaluation: an alpha efficiency of 0.13 ± 0.02 ⁵⁸, Monte-Carlo beta attenuation
655 factors from Marsh⁵⁹, dose-rate conversion factors from Guerin and colleagues⁶⁰, external
656 sediment (beta and gamma) dose rate from the OSL sample PD40, a depth of 25 ± 10 cm,
657 resulting in an age of $103 \pm 10/-9$ ka.

658

659 **Optically Stimulated Luminescence Dating.** Three samples (PD15, PD17 and PD41) were
660 collected from the aeolian sands (Unit 1) underlying the southern marl outcrop (Unit 2, Fig
661 1B). A fourth sample (PD40) was taken from the main fossil bearing bed (Unit 3). Individual
662 quartz grains were measured on a Risø TL/OSL-DA-15 instrument using the single-aliquot
663 regenerative-dose (SAR) method⁶¹. The burial dose for each sample (D_b) was calculated
664 using the central age model (CAM)⁶².

665

666 Environmental dose rates were determined using a Risø GM-25-5 low-level beta counting
667 system⁶³ (beta dose rate), field gamma spectrometry (gamma dose rate), and an estimate of
668 the cosmic dose rate derived using site location and present day sediment burial depths⁶⁴. Full
669 optically stimulated luminescence dating methods and results are presented in Supplementary
670 Information Section 3. All analyses were carried out in the Royal Holloway Luminescence
671 Laboratory by SA and R C-W.

672

673 **Age modelling.** Chronometric ages for samples from the Al Wusta site were incorporated
674 into a Bayesian sequential phase model implemented in OxCal v4.2³¹ (Supplementary
675 Information 4; Supplementary Figure 11. The model consists of two discrete phases separated
676 by a hiatus. Phase 1 was defined by the three OSL ages (PD15, 17 and 41) for samples from
677 the aeolian sands (Unit 1) underlying the lacustrine marls (Unit 2). Phase 2 was defined by
678 the ages for the sand (PD40) and fossils (AW-1 and WU1601) from the waterlain sediments
679 (Unit 3) overlying Unit 2. U-series ages for WU1601 and AW-1 were treated as minimum
680 age estimates, whereas PD40 and the combined U-series-ESR age on WU1601 were treated
681 as finite age estimates. Since the Al Wusta sequence accumulated over a short period of time,
682 and contains only five finite ages (and three minimum ages), the General Outlier Model³¹ was
683 unable to function, and instead a simpler model using agreement indices was employed. This
684 analysis yielded A_{model} (76) and A_{overall} (79) values well in excess of the generally
685 accepted threshold (60³¹), with only one age yielding an individual agreement index below
686 this threshold (PD17, 51). These data indicate that no ages should be excluded from the
687 model, and that the age model itself is robust. The Bayesian sequential model yielded an age
688 for the end of Phase 1 of 93.1 ± 2.6 ka (1 σ uncertainties), while Phase 2 yielded start and end
689 dates of 92.2 ± 2.6 ka and 90.4 ± 3.9 ka respectively. The end date for phase 2 should be
690 treated as a maximum value since no overlying material is present, precluding the possibility
691 of further constraining the end of this phase.

692

693 **Stratigraphy and sedimentology.**

694 *Sediment analysis.* Bulk samples (in the form of coherent blocks) were taken at 10 cm
695 intervals through each of the marl beds in four sections (Fig. 1C and Supplementary Figures
696 13 and 14). Each block was air-dried and subsamples (ca 0.5 g) were removed, powdered and
697 analysed for percentage carbonate content using Bascomb calcimetry, which measures the

698 volume of carbon dioxide liberated from a known sample mass during reaction with 10%
699 HCl⁶⁵. Thin sections were prepared from fresh sediment blocks. The sediments did not
700 require acetone treatment as they were already dry and, due to their permeability, were
701 impregnated with a bonding resin. Standard thin section preparation was then carried out
702 using techniques developed in the Centre for Micromorphology at Royal Holloway,
703 University of London⁶⁶. Thin sections were analysed using an Olympus BX-50 microscope
704 with magnifications from 20x to 200x and photomicrographs were captured with a Pixera
705 Penguin 600es camera. A point-count approach was used to produce semi-quantified data
706 from the thin sections, based on counting micro-features at 3 mm intervals along linear
707 transects 1 cm apart. Kemp⁶⁷, Stoops⁶⁸ and Alonso-Zarza⁴² were referred to when identifying
708 features. X-ray diffraction analysis (XRD) was carried out in the Department of Earth
709 Sciences (Royal Holloway, University of London). Powdered samples were analysed on a
710 Philips PW1830/3020 spectrometer with copper K α X-rays. Mineral peaks were identified
711 manually from the ICDD Powder Diffraction File (PDF) database. The methods and results
712 are described further in Supplementary Information 5.

713

714 *Diatoms.*

715 *Sample preparation.* Samples were analysed using the standard method of Renberg⁶⁹
716 (Supplementary Information 5). Thus, all samples were treated with 30% H₂O₂ and 5% HCl
717 to digest organic material and remove calcium carbonate. Distilled water was added to dilute
718 the samples after heating, which were then stored in the refrigerator for four days to minimise
719 further chemical reactions. The samples were rinsed daily and allowed to settle overnight. A
720 known volume of microspheres was added to the supernatant after the last rinse to enable
721 calculation of the diatom concentration⁷⁰. The slides were air-dried at room temperature in a
722 dust free environment before mounting with Naphrax diatom mountant. Diatom taxonomy

723 followed Krammer and Lange-Bertalot⁷¹⁻⁷³ and taxonomic revisions^{74,75} with at least 300
724 valves enumerated for a representative sample at x1000 magnification.
725
726 *Numerical analysis.* Prevalent trends in the diatom assemblage were explored using
727 ordination analyses using CANOCO 4.5 of ter Braak and Šmilauer⁷⁶. Detrended
728 Correspondence Analysis (DCA⁷⁷) with detrending by segments and down-weighting of rare
729 species was used to investigate taxonomic variations within each site and to determine
730 whether linear or unimodal models should be used for further analyses. If the gradient length
731 of the first axis is <1.5 SD units, linear methods (Principle Component Analysis, PCA)
732 should be used; however, if the gradient length is >1.5 SD units, unimodal methods
733 (Correspondence Analysis) should be used⁷⁸. Detrended Canonical Correspondence Analysis
734 (DCCA⁷⁹) was also used to show changes in compositional turnover scaled in SD units.
735 Therefore, variations in the down-core DCCA first axis sample scores show an estimate of
736 the compositional change between samples along an environmental or temporal gradient.
737 Depth was used as the sole constraint as the samples in each site are in a known temporal
738 order⁸⁰. The dataset was square-root transformed to normalise the distribution prior to
739 analyses. Optimal sum-of-squares partitioning⁸¹ with the program ZONE⁸² and comparison of
740 the zones with the Broken-stick model using the program BSTICK⁸³ were used to determine
741 significant zones. The planktonic: benthic ratio, habitat summary, concentration and the F
742 index (a dissolution index⁸⁴) were calculated for all the samples.

743

744 *Stable isotopes*

745 It is common practice, when analysing the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of lacustrine/palustrine
746 carbonates to either: 1) sieve the sediment and analyse the <63 μm fraction, or 2) use the
747 microstructure of the sample, as identified under thin section, to identify pure, unaltered

748 fabrics, which can then be drilled out and analysed⁸⁵. The former procedure ensures that the
749 analysed fraction comprises pure authigenic marl (rather than a mixture of ostracod,
750 mollusc, chara and marl components that will contain different isotopic values). The latter is
751 done to ensure that any carbonate that has been affected by diagenesis is sampled. Neither of
752 these approaches were carried out here as; 1) microfabric analysis showed no evidence for
753 diagenesis (although some of the samples are cemented the cement makes a negligible
754 component of sample mass), and 2) some of the samples have incipient cementation, which
755 means that they cannot be sieved. Bulk carbonate powders were consequently analysed for
756 $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$. To show that the analysis of bulk samples had no impact on the derived
757 isotopic data, samples that were friable enough to be sieved were treated with sodium
758 hexametaphosphate to disaggregate them and then homogenised and separated into two
759 subsamples for isotopic analysis; (1) a sieved $<63\mu\text{m}$ fraction and (2) a homogenised bulk
760 sample. The resulting isotopic data showed no difference between the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values
761 of the sieved and bulk samples (Supplementary Figure 13b), highlighting that the
762 homogenous and unaltered nature of the material results in bulk carbonate isotopic analysis
763 generating valid data. Two samples were taken from different locations of each sampled
764 block to generate a larger dataset of independent samples. The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of each
765 samples were determined by analysing CO_2 liberated from the reaction of the sample with
766 phosphoric acid at 90°C using a VG PRISM series 2 mass spectrometer in the Earth Sciences
767 Department at Royal Holloway. Internal (RHBNC) and external (NBS19, LSVEC) standards
768 were run every 4 and 18 samples respectively. 1σ uncertainties are 0.04‰ ($\delta^{18}\text{O}$) and 0.02‰
769 ($\delta^{13}\text{C}$). All isotope data presented in this study are quoted against the Vienna Pee Dee
770 Belemnite (VPDB) standard.

771

772 **Vertebrate fossil analyses.** Each fossil specimen was identified to lowest taxonomic and
773 anatomical level possible (Supplementary Figure 20, Supplementary Table 19 and
774 Supplementary Information 6). Taxonomic identification and skeletal element portions were
775 determined based on anatomical landmarks, and facilitated by comparisons with the
776 Australian National University Archaeology and Natural History reference collection
777 (Canberra), unregistered biological collections held at the University of New South Wales
778 (Sydney), and the large mammal collections of the Zoologische Staatssammlung München
779 (Munich). Each specimen was assigned a size category (small, medium, and large) following
780 Dominguez-Rodrigo and colleagues⁸⁶, and corresponding to the five size classes described in
781 Bunn⁸⁷, where small, medium and large denote size classes 1-2, 3A-3B and 4-6, respectively.
782 Element abundance is reported as Number of Identified Specimens (NISP).
783
784 Each specimen was examined for modification by eye and hand-lens (10x) under both natural
785 and high-incidence light, and examined at different angles to assist identification of fine-scale
786 surface modifications. Where required, further examination and photography was carried out
787 using a digital microscope (Model: Dino-lite, AM7013MZ). Morphometric data (length,
788 breadth and width) was measured using digital callipers (Model: Mitutoyo Corp, CD-
789 8"PMX), and specimen weights using a digital scale. Bone surface modifications were
790 identified and recorded following standard methodologies: butchery and tooth marks⁸⁸⁻⁹⁴,
791 burning⁹⁵⁻⁹⁶, rodent gnawing^{97,98}, weathering⁹⁹ and trampling¹⁰⁰. Carnivore damage was
792 categorized as pit, score, furrow or puncture, and the location noted⁹⁴. Tooth mark
793 morphometric data – short and long axes – was also recorded. Any additional modifications,
794 i.e. polish, manganese staining, and root etching, were also reported and described. Bone
795 breakage was recorded as green, dry, or both, following Villa and Mahieu¹⁰¹. Long bone

796 circumference completeness was recorded using the three categories described by Bunn¹⁰²:
797 type 1 (<1/2), type 2 (>1/2 but < complete) and type 3 (complete).

798

799 **Lithic analysis.** Lithics were systematically collected during pedestrian transects and
800 excavations of Al Wusta. This produced a total studied assemblage of 380 lithics
801 (Supplementary Information 7). Further lithics extended for a considerable distance to the
802 north, seeming to track the outlines of the palaeolake, but we only conducted detailed
803 analysis on lithics from the southern part of the site, close to AW-1 and the sedimentary ridge
804 on which it was found (i.e. south of the Holocene playa). These were analysed using the
805 methodology described in Scerri and colleagues^{25,103,104} and Groucutt and colleagues^{45,105}. As
806 well as qualitative analysis of technological features indicating particular techniques and
807 methods of reduction, a variety of quantitative features such as dimensions, the number of
808 scars and % of cortex were recorded. Informative examples were selected for photography
809 and illustration. This approach allows both a characterisation and description of the
810 assemblage and broad comparison with other assemblages from surrounding regions.

811

812

813

814

815

816

817

818

819

820

821 **Methods References**

822

823 49. Stimpson, C. et al. Middle Pleistocene vertebrate fossils from the Nefud Desert, Saudi
824 Arabia: Implications for biogeography and palaeoecology. *Quatern. Sci. Rev.* **143**, 13-
825 36 (2016).

826 50. O'Higgins, P., Jones, N. Facial growth in *Cercocebus torquatus*: an application of
827 three-dimensional geometric morphometric techniques to the study of morphological
828 variation. *J. Anat.* **193**, 251-72 (1998).

829 51. R Core Team. *R: A Language and Environment for Statistical Computing.* (Vienna,
830 Austria: R Foundation for Statistical Computing, <http://www.R-project.org>, 2015).

831 52. Klingenberg C, P. MorphoJ: an integrated software package for geometric
832 morphometrics. *Mol. Ecol. Resour.* **11**, 353-7 (2011)

833 53. Grün, R., Eggins, S., Kinsley, L., Mosely, H., Sambridge, M. Laser ablation U-series
834 analysis of fossil bones and teeth. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **416**,
835 150-167 (2014).

836 54. Benson, A., Kinsley, L., Defleur, A., Kokkonen, H., Mussi, M., Grün, R. Laser
837 ablation depth profiling of U-series and Sr isotopes in human fossils. *J. Arch. Sci.* **40**,
838 2991-3000 (2013).

839 55. Duval, M., Grün, R. Are published ESR dose assessments on fossil tooth enamel
840 reliable? *Quat. Geochron.* **31**, 19-27 (2016).

841 56. Grün, R. The DATA program for the calculation of ESR age estimates on tooth
842 enamel. *Quatern. Geochron.* **4**, 231-232 (2009).

843 57. Grün, R., Schwarcz, H.P., Chadam, J. ESR dating of tooth enamel: Coupled
844 correction for U-uptake and U-series disequilibrium. *Int. J. Radiat. Appl. Instrum.*
845 *Nucl. Tracks. Radiat. Meas.* **14**, 237-241 (1988).





- 846 58. Grün, R., Katzenberger-Apel, O. An alpha irradiator for ESR dating. *Ancient TL* **12**,
847 35-38 (1994).
- 848 59. Marsh, R. E. Beta-gradient Isochrons Using Electron Paramagnetic Resonance:
849 Towards a New Dating Method in Archaeology. (MSc thesis, McMaster University,
850 Hamilton, 1999).
- 851 60. Guérin, G., N. Mercier and G. Adamiec. Dose-rate conversion factors: update.
852 *Ancient TL* 29(1): 5-8 (2011).
- 853 61. Murray, A. S., Wintle, A. G. Luminescence dating of quartz using an improved
854 single-aliquot regenerative-dose protocol. *Radiat. Meas.* **32**, 57-73 (2000).
- 855 62. Galbraith, R. F., Roberts, R. G., Laslett, G. M., Yoshida, H., Olley, J. M. Optical
856 dating of single and multiple grains of quartz from Jinmium rock shelter, northern
857 Australia: Part I, experimental design and statistical models. *Archaeometry* **41**, 339-
858 364 (1999).
- 859 63. Bøtter-Jensen, L., Mejdahl, V. Assessment of beta dose-rate using a GM multicounter
860 system. *Int. J. Rad. Appl. Instrum. B.* **14**, 187-191 (1988).
- 861 64. Prescott, J. R., Hutton, J. T. Cosmic ray and gamma ray dosimetry for TL and ESR.
862 *Int. J. Rad. Appl. Instrum.* **14**, 223-227 (1988).
- 863 65. Gale, S., Hoare, P. *Quaternary Sediments: Petrographic Methods for the Study of*
864 *Unlithified Rocks* (Belhaven and Halsted Press, 1991).
- 865 66. Palmer, A. P., Lee, J. A., Kemp, R. A., Carr, S. J. *Revised laboratory procedures for*
866 *the preparation of thin sections from unconsolidated sediments.* (Centre for
867 micromorphology publication, Royal Holloway, University of London, 2008).
- 868 67. Kemp, R. A. *Soil micromorphology and the Quaternary.* (Quaternary Research
869 Association, 1985).



- 870 68. Stoops, G. *Interpretation of micromorphological features of soils and regoliths*
871 (Elsevier, 2010).
- 872 69. Rengberg, I. A procedure for preparing large sets of diatom sets from sediment. *J.*
873 *Palaeolimnol* **4**, 87-90 (1990).
- 874 70. Battarbee, R. W., Knen, M. J. The use of electronically counter microspheres in
875 absolute diatom analysis. *Limnol. Oceanogr.* **27**, 184-188 (1982).
- 876 71. Krammer, K., Lange-Bertalot, H. *Bacillariophyceae 2. Teil Epithemiaceae,*
877 *Suirellaceae* (Gustav-Fisher Verlag, 1988).
- 878 72. Krammer, K., Lange-Bertalot, H. *Bacillariophyceae 3. Teil Centrales,*
879 *Fragicariaceae, Eunotiaceae* (Gustav-Fisher Verlag, 1991a).
- 880 73. Krammer, K., Lange-Bertalot, H. *Bacillariophyceae 4. Teil Achnantheaceae, Kritshe*
881 *Ergänzungen zu Navicula (Lineolate) und Gomphonema* (Gustav-Fisher Verlag,
882 1991b).
- 883 74. Crawford, R. M., Likhoshway, Y. V., Jahn, R. Morphology and identity of
884 *Aulacoseira italic* and typification of *Aulacoseira* (Bacillariophyta). *Diatom*
885 *Research* **18**, 1-19 (2003).
- 886 75. Navok, T., Guillory, W. X., Julius, M. L., Theriort, E. C., Alverson, A. J. Towards a
887 phylogenetic classification of species belonging to the diatom genus *Cyclotella*
888 (Bacillariophyceae): Transfer of species formerly placed in *Puncticulata,*
889 *Handmannia, Pliocaenicus* and *Cyclotella* to the genus *Lindavia*. *Phytotaxa* **217**, 249-
890 264 (2015).
- 891 76. Ter Braak, C. J. F., Šmilauer, P. *CANOCO reference manual and CanoDraw for*
892 *Windows user's guide: software for canonical community ordination (version 4.5).*
893 (Microcomputer Power, 2002).


- 894 77. Hill, M. O., Gauch, H. G. Detrended correspondence analysis: An improved
895 ordination technique. *Plant. Ecol.* **42**, 47-58 (1980).
- 896 78. Ter Braak, C. J. F., Prentice, I. C. A theory of gradient analysis. *Adv. Ecol. Res.* **18**,
897 271-317 (1988).
- 898 79. Ter Braak, C. J. F. Canonical Correspondence Analysis: A new eigenvector technique
899 for multivariate direct gradient analysis. *Ecology* **67**, 1167-1179 (1986).
- 900 80. Smol, J. P. et al. Climate-driven regime shifts in the biological communities of arctic
901 lakes. *Proc. Natl. Acad. Sci. USA* **102**, 4397-4402 (2005).
- 902 81. Birks, H. J. B., Gordon, A. D. *Numerical methods in Quaternary Pollen Analysis*.
903 (Academic Press, 1985).
- 904 82. Juggins, S. *ZONE software, version 1.2*. (University of Newcastle, 1985).
- 905 83. Bennett, K. D. Determination of the number of zones in a biostratigraphical sequence.
906 *New Phytologist* **132**, 155-170 (1996).
- 907 84. Ryves, D. B. Juggins S., Fritz, S. C., Battarbee, R. W. Experimental diatom
908 dissolution and the quantification of microfossil preservation in sediments.
909 *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **172**, 99-113 (2001).
- 910 85. Candy, I. et al. The evolution of Palaeolake Flixton and the environmental context of
911 Starr Carr: an oxygen and carbon isotopic record of environmental change for the
912 early Holocene. *Proc. Geol. Assoc.* **126**, 60-71 (2015).
- 913 86. Domínguez-Rodrigo, M., Barba, R., De la Torre, I., Mora, R. in *Deconstructing*
914 *Olduvai: A Taphonomic Study of the Bed I Sites* (eds. Domínguez-Rodrigo, M.,
915 Barba, R., Egeland, C. P.) 101-125 (New York, Springer, 2007).
- 916 87. Bunn, H. T. *Meat-eating and human evolution: Studies on the diet and subsistence*
917 *patterns of Plio-Pleistocene hominids in East Africa* (University of Wisconsin,
918 Madison, Unpublished PhD thesis, 1982).






- 919 88. Bunn, H. T., Kroll, E. M. Systematic butchery by Pilo/ Pleistocene Hominids at
920 Olduvai Gorge , Tanzania. *Curr. Anthropol.* **27**, 431–452 (1986).
- 921 89. Binford, L. R. *Faunal remains from Klasies River Mouth.* (Academic Press, 1984).
- 922 90. Andrews, P. & Cook, J. Natural modifications to bones in a temperature setting. *Man*
923 **20**, 675–691 (1985).
- 924 91. Blumenshine, R. J., Selvaggio, M. M. Percussion marks on bone surfaces as a new
925 diagnostic of hominid behaviour. *Nature* **333**, 763–765 (1988).
- 926 92. Fisher, J. W. Bone Surface modifications in zooarchaeology. *J. Archaeol. Method*
927 *Theory* **2**, 7–68 (1995).
- 928 93. Noe-Nygaard, N. Man-made trace fossils on bones. *J. Hum. Evol.* **4**, 461–461 (1989).
- 929 94. Binford, L. R. *Bones: Ancient Men and Modern Myths.* (Academic Press, 1981).
- 930 95. Stiner, M., Kuhn, S., Weiner, S., Bar-Yosef, O. Differential burning, recrystallization,
931 and fragmentation of archaeological bone. *J. Archaeol. Sci.* **22**, 223–237 (1995).
- 932 96. Shipman, P., Foster, G., Schoeninger, M. Bunt bones and teeth: an experimental study
933 of color, morphology, crystal structure and shrinkage. *J. Archaeol. Sci.* **11**, 307–325
934 (1984).
- 935 97. Tong, H. W., Zhang, S., Chen, F., Li, Q. Rongements sélectifs des os par les porcs-
936 épics et autres rongeurs : cas de la grotte Tianyuan, un site avec des restes humains
937 fossiles récemment découvert près de Zhoukoudian (Choukoutien). *Anthropologie*
938 **112**, 353–369 (2008).
- 939 98. Dart, R. A. Bone tools and Porcupine gnawing. *Am. Anthropol.* **60**, 715–724 (1958).
- 940 99. Behrensmeyer, A. K. Taphonomic and ecological information from bone weathering.
941 *Paleobiology* **4**, 150–162 (1978).
- 942 100. Behrensmeyer, A. K., Gordon, K., Yanagi, G. T. Trampling as a cause of bone
943 surface damage and psuedo-cutmarks. *Nature* **319**, 402–403 (1986).

- 944 101. Villa, P., Mahieu, E. Breakage pattern of human long bones. *J. Hum. Evol.* **21**,
945 27–48 (1991).
- 946 102. Bunn, H. T. in *Animals and Archaeology, Volume 1* (eds. Clutton-Brock, J. &
947 Grigson, C.) **1977**, 143–148 (Oxford, BAR International Series 163, 1983).
- 948 103. Scerri, E. M. L., Groucutt, H. S., Jennings, R. P., Petraglia, M. D. Unexpected
949 technological heterogeneity in northern Arabia indicates complex Late Pleistocene
950 demography at the gateway to Asia. *J. Hum. Evol.* **75**, 125-142 (2014).
- 951 104. Scerri, E. M. L., Gravina, B., Blinkhorn, J., Delagnes, A. Can lithic attribute
952 analyses identify discrete reduction trajectories? A quantitative study using refitted
953 lithic sets. *J. Arch. Method Theory* **23**, 669-691 (2016).
- 954 **105.** Groucutt, H.S., et al. Late Pleistocene lakeshore settlement in northern Arabia:
955 Middle Palaeolithic technology from Jebel Katefeh, Jubbah. *Quatern. Int.* **382**, 215-
956 236 (2016).









Group

- Holocene *H. sapiens*
- Neanderthals
- Pleistocene *H. sapiens*
- ▲ Al Wusta

Ray

- ▲ Al Wusta
- 2nd
- 3rd
- ◆ 4th

a



b



c



d




e



f



Marine Isotope Stages (Lisiecki & Raymo, 2004)



SI 1. Description and comparison of the of Al Wusta-1 phalanx.

1.1 Pathology

The Al Wusta-1 (AW-1) phalanx shows evidence of pathological changes to the bone surface. Additional pathological bone formation affects the proximal half of the shaft, covering approximately one quarter of the dorsal surface, measuring 11.9 mm proximo-distally and 5.9 mm radio-ulnarly, and projecting approximately 2.5 mm from inferred ‘normal’ bone surface. Micro-CT scanning confirms that the additional bone is continuous with the cortical bone of the shaft, but there is no evidence of a fracture or other trauma (Fig. 2B, C). Its irregular, angular morphology suggests that this additional bone may be due to the ossification of the central slip of the extensor digitorum muscle (i.e., a “bony spur” or enthesophyte), which attaches to the intermediate phalanx in this region. The unusual, relatively circular cross-sectional shape of AW-1 may also reflect these pathological changes.

1.2 Linear metric analysis of the Al Wusta-1 intermediate phalanx

Linear measurements of AW-1 are presented in Supplementary Table 1. We conducted an analysis of nine linear measurements of intermediate phalanx shape across a sample of extant primates and fossil hominins (Supplementary Table S2). For extant non-human primates, intermediate phalanges (IPs) from all rays (2-5) of one side (either left or right) were included as it is possible that all non-human primate IPs may show similar morphology to human IPs^{1,2}. However, human and fossil hominin IPs from the fifth ray (IP5) show a distinctive, asymmetrical shape that is not present in AW-1 and thus all *H. sapiens* IP5 specimens and potential IP5 fossil hominin specimens were excluded from the analysis. Although data from

multiple IPs from a single individual are not independent, without knowing the exact ray to which AW-1 belongs, nor the exact ray or number of individuals associated with several of the comparative fossil hominin intermediate phalanges, it is more conservative to include the range of morphological variation across multiple rays.

Linear measurements included the maximum proximo-distal length of the phalanx (i.e. total length), maximum dorso-palmar height of the proximal base, the dorso-palmar height and radio-ulnar breadth of the proximal articular facet, radio-ulnar breadth of the proximal shaft, and dorso-palmar height and radio-ulnar breadth of the midshaft and distal shaft, all of which could be confidently measured on AW-1.

All metrics were assessed and compared as a ratio of the total length of the phalanx.

Comparisons across extant taxa, Neanderthals and *H. sapiens* (i.e. all taxonomic groups with large enough sample sizes) were evaluated using Mann-Whitney U pairwise comparisons with a Bonferroni correction for multiple comparisons (Supplementary Table 3). Relative comparisons of AW-1 and other fossil specimens were visually assessed via box-and-whisker plots (Supplementary Figure 1).

Comparative analyses reveal that there is substantial overlap across most taxa in all shape ratios. For any given shape ratio, AW-1 falls within the range of variation of cercopiths, *Gorilla*, *A. afarensis*, *A. sediba*, Neanderthals and *H. sapiens*. However, AW-1 is most similar to the median value or falls within the range of variation of recent and early *H. sapiens* for all shape ratios (Supplementary Figure 1), confirming its affiliation with *H. sapiens* revealed by the 3D geometric morphometric analyses (see main text and below). More specifically, AW-1 is very similar to the *H. sapiens* median value in the relative

radioulnar breadth of the proximal base and the proximal shaft, and the dorso-palmar height at midshaft. AW-1 falls within the lower range of variation of *H. sapiens*, and outside or at the extreme of the Neanderthal range of variation, in its dorso-palmar height and radioulnar breadth proximal facet, and its radioulnar breadth at midshaft and the distal shaft.

Note that published values for the controversial *H. sapiens* specimen Cueva Victoria CV-0 specimen are included in the proximal base breadth and midshaft breadth and height shape ratios (Supplementary Figure 1). This specimen is always the most extreme outlier in the box-and-whisker plots, and falls in the direction of the cercopithecoid median value, suggesting that this specimen is indeed that of *Theropithecus*, and not *H. sapiens*, supporting Martínez-Navarro and colleagues^{1,2}.

1.3 Geometric morphometric comparison of non-human primate, fossil hominin and AW-1 Wusta-1 phalanges

To provide a broader interpretive context for AW-1, we provide a principal components analysis of geometric morphometric landmark data (Supplementary Table 4, Supplementary Figure 2) on a sample of phalanges from a range of primates including fossil hominins (Supplementary Table 5). In Figure 3 (main text) and Supplementary Figure 3, PC1 and PC2 together account for 61% of group variance in shape. AW-1 is separated on these two shape vectors from the non-human primates and most of the Neanderthals by a shorter, wider diaphysis and palmarly flatter proximal base. It shares a proximal head that is higher to the right (dorsal view) with *H. sapiens*, although this may be a function of the proportion of left and right sides in each sample. AW-1 falls closest to the Holocene and early *H. sapiens* and is well differentiated from all non-human primates. This is shown by the Procrustes distances

from AW-1 to the mean shapes of each taxonomic group (Figure 3, Supplementary Figure 3 and Supplementary Table 6).

1.4 Geometric morphometric analysis restricted to AW-1 and hominin phalanges of known side and digit numbers

Details of the sample are given in Supplementary Table 7. Methods and Results for pooled left and right hands are given in the main text (see Figure 4 and also Supplementary Tables 8-9.)

1.4.1 Left and right 2nd, 3rd and 4th intermediate phalanges separated.

The results showing AW-1 compared separately to right and to left phalanges (Supplementary Figure 4, Supplementary Tables 10-11) are very similar to the pooled sample (see main text, Figure 4 and Supplementary Tables 8-9), such that AW-1 is closest to Holocene *H. sapiens* 3rd rays for both right and left hand, although Pleistocene *H. sapiens* configurations fall almost completely inside the scatter for the Holocene *H. sapiens* sample. AW-1 is most distinct from the Neanderthal phalanges of both the left and right hands. The greatest separation between AW-1 and other groups is described by PC2 for both the right and left phalanges. These vectors describe the shape difference between shorter and stockier vs. longer and narrower configurations. AW-1 is taller and narrower (in all directions: dorso-palmarly, proximo-distally and radio-ulnarly) than shapes towards the other end of the PC2s, which describe most of the Neanderthal phalanges. Again, these analyses suggest that AW-1 is likely to be a 3rd intermediate phalanx from a *H. sapiens* individual.

1.5 Cross sectional geometry analyses

1.5.1 Materials and Methods

Cross-sectional geometry (CSG) of bones examines the amount and distribution of cortical bone in the cross section, which reflects primarily the impacts of body size, body shape, and activity on the skeleton³⁻⁶. CSG of AW-1 and the comparative 2nd-4th phalanges (Supplementary Table S7) were calculated in ImageJ⁷ using the BoneJ plugin⁸ and using the same microCT data as for the GMM analyses. Slices at 54% of total AW-1 phalanx length (measured from the proximal end) were analysed to avoid the influence on cross-sectional properties of the pathological bone formation on the shaft. Total area (TA) of the cross section was calculated by filling the medullary cavity with the 'fill holes' function of ImageJ and rerunning the slice through BoneJ. Percent cortical area (%CA) reflecting cortical bone thickness was calculated as $100 \times \text{cortical area} / \text{TA}$. J , a measure of torsional and twice average bending rigidity, was calculated as the sum of maximum and minimum bending rigidities (I_{max} and I_{min} respectively)⁹.

GMM analyses suggest that AW-1 is a 3rd intermediate phalanx, but plots were generated for each of manual rays 2-4 in case these analyses suggested otherwise. Where left and right sides were present for the same ray of the same individual, the mean was used.

As body size and activity are both important determinants of bone cross-sectional properties (see above), CA and J were plotted against bone length to examine whether the cross-sectional properties relative to body size could differentiate Neanderthal, Pleistocene *H. sapiens* and Holocene *H. sapiens* and thus be informative regarding the taxonomic affiliation

of AW-1. However, it must be noted that CSG of the phalanges, unlike the limb long bones^{4,8}, is not well documented in the literature and the relative importance of body size, activity and taxonomy remain to be investigated in detail. The relationship between I_{max} and I_{min} , which reflects the circularity of bone distribution was also examined by plotting I_{max} against I_{min} . Plots were generated using IBM SPSS Statistics v. 23.

1.5.2 Results

In general, AW-1 lies outside of the range of CSG for intermediate phalanges from ray 2, well within the range for ray 3, and at the upper end of the range for ray 4 (Supplementary Figure 5), supporting the interpretation that AW-1 is most likely to be a 3rd intermediate phalanx. For all cross-sectional properties, Holocene *H. sapiens* show a large range of variation and the small sample of Neanderthals and Pleistocene *H. sapiens* do not appear well differentiated from the Holocene specimens. While generally within the range of the comparative specimens, AW-1 falls just outside the range of the sample for I_{max} relative to I_{min} , with a low ratio indicating an unusually circular cross-section. In the long bones of the lower limb, more circular shafts indicate similar loading in multiple directions¹⁰⁻¹¹, but its precise interpretation for manual phalanges remains to be explored.

Further work to document the range of variation in phalanx CSG and its relationship to ancestry and behaviour patterns would be required to further interpret the cross-sectional circularity of the AW-1. A relationship between this high level of circularity and the pathological bone formation on the dorsal surface of AW-1's shaft cannot be excluded, since the shaft could be expanded in a dorso-palmar direction even where external appearance is

normal, which would serve to lower the I_{max}/I_{min} ratio. Alternatively, a generally high level of loading might account for both the enthesophyte and more circular cross-section of the shaft.

SI 2. U-series and combined US-ESR dating of fossil bone and teeth from Al Wusta.

2.1 Materials and Methods

2.1.1 Material

The human phalanx (AW-1, lab code for U-series = 3675) and a hippopotamus tooth fragment (lab code WU1601) were collected from Trench 1. The external dose rate calculations are based on the data from OSL sample PD40 (Supplementary Table 16), which was collected at the equivalent position within unit 3a.

2.1.2 U-series analysis

U-series analysis of bones can be used to reconstruct U-uptake phases. Modern bones are virtually U free. All the uranium that is measured in fossil samples migrated into the skeletal tissues after these were buried. However, it is difficult to establish whether this U-uptake was a single stage process that occurred a short time after burial, or whether the U-accumulation was a complex, multistage process that may have commenced a significant time after the original burial¹². In any case, as long as there is no indication for uranium leaching, the calculated U-series age results have to be regarded as minimum age estimates with respect to the age of the fossil.

The experimental setup for the U-series analysis of the AW-1 phalanx was previously described in Grün and colleagues¹². Laser ablation (LA) was used to drill a number of holes the finger bone following the approach of Benson et al.¹³. After a cleaning run with the laser

set at a diameter of 460 μm , seven holes were drilled for 1000 s (Supplementary Figure 6A) with the laser set at 330 μm . The isotopic data streams (Supplementary Figure 6B) were converted into $^{230}\text{Th}/^{234}\text{U}$ and $^{234}\text{U}/^{238}\text{U}$ activity ratios and apparent Th/U age estimates (Supplementary Figure 6C) and subsequently binned into 30 successive sections (each containing 33 cycles) for the calculation of average isotopic ratios and ages (Supplementary Figure 6D; Supplementary Table 12).

A similar experimental setup and methodology were employed for the LA U-series analysis of tooth sample WU1601 (Supplementary Figure 8). Individual closed system U-series age estimates were calculated for each ablation spot and the whole analytical data of the enamel and dentine sections were integrated to provide the data input for the ESR age calculations (Supplementary Table 13).

2.1.3 ESR dose evaluation

Enamel was collected from tooth WU1601 and powdered $<200\ \mu\text{m}$. The sample was then divided into 11 aliquots and gamma irradiated with a Gammacell-1000 Cs-137 source to the following doses: 0, 49, 97, 146, 243, 340, 486, 873, 1457, 2430 and 3397 Gy. ESR measurements were carried out at room temperature with an EMXmicro 6/1 Bruker ESR spectrometer coupled to a standard rectangular ER 4102ST cavity. The following procedure was used to minimise the analytical uncertainties: (i) all aliquots of a given sample were carefully weighed into their corresponding tubes and a variation of $<1\ \text{mg}$ was tolerated from one aliquot to another; (ii) ESR measurements were performed using a Teflon sample tube holder inserted from the bottom of the cavity to ensure that the vertical position of the tubes

remains exactly the same for all aliquots. The following acquisition parameters were used: 3-30 scans (depending on the sample and aliquot measured), 1 mW microwave power, 1024 points resolution, 15 mT sweep width, 100 kHz modulation frequency, 0.1 mT modulation amplitude, 20 ms conversion time and 5 ms time constant.

ESR intensities were extracted from T1-B2 peak-to-peak amplitudes of the ESR signal of enamel¹⁴, and then corrected by the corresponding number of scans and aliquot mass. Fitting procedures were carried out with the Microcal OriginPro 9.5 software using a Levenberg-Marquardt algorithm by chi-square minimization. D_E values were obtained by fitting a single saturating exponential (SSE) function through the experimental data, with data weighting by the inverse of the squared ESR intensity ($1/I^2$)¹⁵.

The sample was measured three times on different days in order to evaluate the precision in measurement and D_E . Only small variation of between 1.5% and 2.7% ($1-\sigma$) were found over the three days, respectively. The final D_E value was calculated by pooling all the ESR intensities derived from the three repeated measurements in a single dose response curve (DRC)¹⁶. In order to avoid D_E overestimation caused by the use of the SSE function, an appropriate maximum irradiation dose (D_{max}) was selected in accordance with the recommendations by Duval and Grün¹⁷: given the magnitude of the D_E value (~ 100 Gy), D_{max}/D_E should be between 5 and 10, and final dose evaluation were thus done with $D_{max}=873$ Gy ($D_{max}/D_E=9$). The final DRC is shown in Supplementary Figure 9.

2.1.4 Dose rate evaluation and US-ESR age calculation

For the dose rate calculations, the following parameters were used: an alpha efficiency of 0.13 ± 0.02 ¹⁸, Monte-Carlo beta attenuation factors¹⁹, recently published dose-rate conversion factors²⁰. For the calculation of the external sediment dose rate, the data of OSL sample PD40 (Supplementary Table 16) were used. The total external dose rate of 438 ± 27 $\mu\text{Gy/a}$ consists of 254 ± 25 $\mu\text{Gy/a}$ cosmic dose rate (for the actual depth of 25 cm below surface), 180 ± 10 $\mu\text{Gy/a}$ external gamma dose rate, plus the external beta dose rate was corrected for the tooth configuration, resulting in 4 $\mu\text{Gy/a}$.

Combined US-ESR ages were calculated with DATA²¹ using the US model defined by Grün et al²². Further details about this dating method applied to fossil teeth may be found in Duval²³. The other details of the age result of $103^{+10/-9}$ ka are shown in Supplementary Table 14.

2.2 Results and discussion

2.2.1 U-series dating of the Al Wusta-1 phalanx

Some scans show elevated ²³²Th levels (see Supplementary Figure 6B, around cycle 250), which is the result of probing some pores that may be filled with detrital material. However, all ²³⁸U/²³²Th elemental ratios are well above 50. Supplementary Figure 6E shows that the calculated ages of the cycles with lower U/Th ratios are not affected by detrital ²³⁰Th, which would systematically increase the results.

All age calculations are based on closed system assumptions, they are thus apparent age estimates. U/Th ages were calculated using the Isoplot 3.75 Excel add-on²⁴. Note that this program does not calculate asymmetric errors. All reported errors are 2- σ . We did not use the diffusion-adsorption decay (DAD) model of Sambridge and colleagues²⁵ as there is evidence of at least two discrete U-uptake phases.

All data steams have in common that the outer domains yield younger lower apparent ages than those further inside the bone (Supplementary Figure 7A). This is clearly associated with higher U concentrations (Supplementary Figure 7B and C). This indicates a second discrete phase of U-accumulation, which overprinted the original isotopic signature. The more U was taken up at that later phase, the younger becomes the apparent U-series age. For each of the holes, the apparent age plateaus after an initial increase. These age plateaux systematically increase from Hole 1 to Hole 7 (see Supplementary Figure 7A). Interestingly, there is a reverse trend with older plateau ages associated with slightly higher U-concentrations (Supplementary Figure 7D). This is could be related to a process where the domains, which had accumulated more U during an initial uptake phase, were relatively less affected by any subsequent U-migration processes.

To conclude, the age plateaux of hole #7 of 87.6 ± 2.5 ka represents the minimum age of the finger bone. The true age of the bone may be older because (i), the age plateaux of holes #1 to 6 are affected by later U-uptake phases, and it is not possible to ascertain whether the age plateau of hole #7 was not affected; (ii), the observed, complex U-migration processes may have commenced a considerable time after the initial burial.

2.2.2 Combined US-ESR dating of the fossil tooth

Apparent U-series age results obtained for the dental tissues are close to those obtained in the human phalanx: 83.5 ± 8.1 ka in the enamel and 65.0 ± 2.1 ka in dentine. These results suggest that the tooth sample is at least 83 ka old.

The combined US-ESR age calculation yields a result of 103^{+10}_{-96} ka. This ESR result agrees with OSL sample PD40 (98.6 ± 7 ka) within error.

SI 3: Optically stimulated luminescence (OSL) dating of Al Wusta.

3.1 Sample collection and preparation

Trenches were dug through the marl beds into the underlying aeolian sand (Unit 1) at Al Wusta. Three samples (PD15, PD17 and PD41, Supplementary Figure 13) were collected from Unit 1 sands underlying the southern marl outcrop, upon which the AW-1 phalanx and the majority of the animal fossils were found. A fourth sample (PD40, Supplementary Figure 13) was taken from Unit 3. PD40 and PD41 were taken from the same trench. OSL samples were collected by hammering opaque tubes into cleaned section faces. In the laboratory, quartz was extracted from the portion of each sample which had not been exposed to sunlight. Samples were initially wet-sieved to isolate the 212-180 μm size fraction, and carbonates and organic matter were subsequently removed using 1M HCl and H_2O_2 respectively. The samples were then re-sieved at 180 μm and quartz was extracted from the >180 μm fraction using density separations at 2.62 and 2.70 g/cm^3 and a subsequent HF acid etch (23M HF for 60 min followed by a 10M HCl rinse). Since Nefud quartz samples are prone to feldspar contamination, an additional one week H_2SiF_6 treatment followed by an HCl rinse was also performed. Etched samples were re-sieved at 150 μm , to remove partially dissolved grains, and stored in opaque containers prior to measurement.

3.2 Luminescence measurements

3.2.1 Equipment

All OSL measurements presented here were carried out using a Risø TL/OSL-DA-15 automated dating system²⁶, fitted with a single-grain OSL attachment^{27,28}. Optical stimulation

of single-grains used a 10 mW Nd: YVO₄ solid-state diode-pumped green laser (532 nm) focussed to yield a nominal power density of 50 W/cm², following²⁶. All infra-red (IR) stimulation was carried out using an IR (870 nm) laser diode array yielding a power density of 132 mW/cm². OSL passed through 7.5 mm of Hoya U-340 filter and was detected using an Electron Tubes Ltd 9235QB15 photomultiplier tube. Irradiation was carried out using a 40 mCi ⁹⁰Sr/⁹⁰Y beta source giving ~6 Gy/min. This source is calibrated relative to the National Physical Laboratory, Teddington ⁶⁰Co γ -source (Hotspot 800²⁹). Due to the spatial inhomogeneity of beta emitters across the active face of our ⁹⁰Sr/⁹⁰Y beta source, it was necessary to calibrate the dose rate to each individual grain position on a single-grain disc³⁰ using the method of Armitage and colleagues³¹.

3.2.2 Single-grain measurement and analysis

Measurements were made on 2800-3600 individual quartz grains per sample (Supplementary Table 15), using the single-aliquot regenerative-dose (SAR) method³², to estimate the equivalent dose (D_e). Since optimum measurement conditions vary between samples^{32,33}, single-grain dose recovery tests^{34,35} were performed on two of the four samples (PD15 and PD17) using a known dose of ~50 Gy. A preheating regime of 260 °C held for 10 seconds prior to measurement of the natural/regenerated dose, and 220 °C held for 10 seconds prior to measurement of the test dose, yielded dose recovery ratios (measured dose/known dose) of 1.01 ± 0.03 for both samples and was adopted for subsequent D_e measurements. Optical stimulation was carried out at 125 °C for 2 s using the green laser. The OSL signal was that recorded during the first 0.3 s of stimulation, with a background signal from the last 0.3 s of stimulation subtracted^{36,37}. Dose response curves were fitted with a saturating exponential or a saturating-exponential-plus-linear function. The standard error associated with each

individual D_e determination was estimated by Monte Carlo simulation. Curve fitting, D_e determination and Monte Carlo simulation were performed using version 4.31.9 of the Luminescence Analyst software³⁸.

It has been observed widely that the majority of quartz grains from unheated sedimentary deposits do not yield a measureable OSL signal³⁹⁻⁴¹ or display luminescence characteristics which indicate that they are unsuitable for age determination. Consequently, single-grain dating studies must adopt criteria for rejecting uninformative grains⁴⁰. In this study, grains were rejected where one or more of the following conditions are met: (1) the OSL signal from the grain is too low to distinguish it from the variability in the background signal, determined using the “Tn signal more than 3 sigma above BG” rejection criterion in Luminescence Analyst³⁸; (2) the recycling ratio⁴² differs from unity by greater than two standard errors; (3) the IR-depletion ratio¹⁷ is greater than two standard errors below unity; (4) recuperation exceeds 5% of the natural signal and (5) the sensitivity-corrected natural luminescence intensity is greater than the saturation intensity of the measured growth curve (termed “oversaturation” in Supplementary Table 15). In addition, grains were rejected where their measured D_e was within measurement uncertainty of 0 Gy at 2σ . This last criterion excludes intruded modern grains, and has been found necessary for the analysis of samples from similar contexts elsewhere in the Nefud Desert⁴³. These rejection criteria were applied in order, and only the first cause of rejection is recorded in Supplementary Table 15. Of the 17,900 grains measured, only 265 displayed acceptable luminescence properties, a yield of 1.5%. The majority of grains (94%) were rejected due to low OSL signal intensity. Despite rigorous attempts to remove feldspar from the quartz fraction, fewer grains were accepted than failed the IR-depletion ratio criterion. This phenomenon has been observed in other single-grain studies on Nefud “quartz” e.g Petraglia and colleagues⁴³.

To determine the age of a sample from a single-grain dataset it is necessary to calculate a single value for the burial dose (D_b). Since the OSL samples presented here were taken from aeolian sand where complete resetting of the OSL signal prior to deposition may be assumed, and the small number of intruded grains were rejected prior to analysis, D_b was calculated for all samples using the central age model (CAM)⁴⁴. Overdispersion (OD) values for the Al Wusta samples ranged from $16 \pm 3\%$ to $26 \pm 3\%$ (PD17 and PD40 respectively), which is consistent with values reported from well-bleached undisturbed sediments^{33,45,46}, supporting the use of the CAM to derive estimates of D_b . Equivalent dose distributions for each sample are presented in Supplementary Figure 10.

3.3 Environmental dose rate calculations

For HF acid etched sand-sized quartz grains, the environmental dose rate consists of external beta, gamma and cosmic ray components. Beta dose rates were measured using a Risø GM-25-5 low-level beta counting system⁴⁷ using MgO and Volkagem loess⁴⁸ standards, while gamma dose rates were measured using an EG&G Ortec digiDart-LF gamma-spectrometer using the “threshold” method. Dose rates were corrected for the effects of HF etching⁴⁹, grain size⁵⁰ and a water content of $5 \pm 2.5\%$. The 2σ uncertainty on water content encompasses completely dry conditions (0%) and saturation for 25% of the burial period (10%), representing the full range of reasonable mean water content scenarios of a freely draining aeolian sand. Cosmic ray dose rates were calculated using site location (27.4°N , 39.4°E , 925 m elevation) and present day sediment burial depths⁵¹. For samples other than PD40, the overburden was assumed to consist of 40 cm of sand (the target sampling depth below the base of the marl) containing 5% water by mass (assumed bulk density 1.74 g/cm^3), with the remainder of the overburden depth being carbonates also containing 5% water by mass

(measured bulk density 1.36 g/cm^3). For PD40, the entire overburden was assumed to be sand with an assumed bulk density of 1.74 g/cm^3 . Using these assumptions, cosmic rays contribute between 33 % (PD15) and 44% (PD40) of the total calculated dose rate, meaning that accurate estimation of the cosmic ray dose rate is more important for these samples than is normally the case. Therefore, the assumptions regarding burial depth require detailed consideration. Firstly, it was assumed that Unit 2 was deposited instantaneously above PD15-17 and PD41, while the full depth of Unit 3a was deposited instantaneously above PD40. This assumption allows the cosmic dose rate to be regarded as constant throughout the sample's burial period. For Unit 2, this assumption must approximate reality since the OSL ages for samples below Unit 2 are indistinguishable from those above. For unit 3 geologically instantaneous deposition cannot be demonstrated, but seems likely based on the interpretation that this deposit represents the encroachment of fluvial sedimentation during the final drying of Al Wusta lake (SI Section 5). Uncertainties on the cosmic ray dose rate were set at $\pm 10\%$. Assuming that the overlying sediments were deposited rapidly, mean overburden during a sample's burial period is unlikely to have been lower than that in the present day. Consequently, cosmic ray dose rates calculated as described above are either accurate or overestimates of mean burial dose rates. Cosmic ray dose rate overestimation will occur where appreciable reduction in the depth of overlying sediments has occurred since burial. In the case of samples overlain by carbonate beds (all except PD40), the cohesive nature of these sediments suggests that limited removal of overlying carbonates has occurred since burial. Conversely, the coarse surface lag (Unit 3b) above Unit 3a (Sample PD40) suggests that some overburden loss due to deflation has occurred (SI Section 5). It is difficult to estimate the quantity of material lost due to this process, but the sensitivity of the true cosmic ray dose rate to deflation may be estimated. In the present case, the lower (-10%) boundary of the estimated cosmic ray dose rate would be achieved by adding $\sim 40 \text{ cm}$ of sand (bulk

density 1.74 g/cm^3) to the burial depth of the shallow samples (PD17 and 40), or ~ 70 cm to the burial depth of the deeper samples (PD15, 16, 41). However, this calculation represents the worst-case scenario, since it implies greater overburden throughout the burial period, followed by instantaneous removal of $\sim 40/\sim 70$ cm of sand immediately prior to sampling. If a more realistic model, assuming continuous deflation throughout the burial period is used, then the lower boundary of the estimated cosmic ray dose rate is achieved after removal of 80 cm of sand overlying the shallow samples, and 140 cm of sand overlying deeper samples. These considerations suggest that the $\pm 10\%$ (1σ) uncertainties assumed for our cosmic ray dose rates encompass reasonable variations in overburden density over time.

Dose rates and sample ages are presented in Supplementary tables 16 and 17 respectively. All analyses were carried out in the Royal Holloway Luminescence Laboratory by SA and R C-W.

SI 4 Site chronology and Bayesian model.

The combination of different numerical dating methods enables us to provide ages for AW-1 and associated sediment and fossils. The chronostratigraphic evidence available may be summarized as follows (Supplementary Figures 11 and 12):

- The human phalanx was found on the surface of Trench 1, on an exposure of Unit 3b. Direct U-series dating of AW-1 itself provides an age of 87.6 ± 2.5 ka (2σ confidence level). This result should be regarded as a minimum age for the fossil.
- A hippopotamus tooth (WU1601) was collected from Trench 1 within Unit 3a, one metre away from AW-1 (Supplementary Figure 14). U-series dating of the dental tissues provides apparent ages of 83.5 ± 8.1 ka (enamel) and 65.0 ± 2.1 ka (dentine) (2σ). These minimum age results are consistent with that obtained for AW-1, suggesting that both fossils are coeval.
- Combined US-ESR dating of WU1601 yields an age of $103 +10/-9$ ka (1σ), indicating a relatively early uranium uptake in dental tissues ($p=-0.83$ and -0.53 for enamel and dentine, respectively). This age estimate is compatible with the minimum age results derived from U-series dating of AW-1 and WU1601.
- The OSL sample collected from PD40 section within the same Unit 3a provides an age of 98.6 ± 7.0 ka (PD40). This estimate is consistent with the US-ESR age obtained for WU-1601 and the minimum age of ~ 88 ka obtained for AW-1.
- Three OSL samples collected from Unit 1 provides ages of 85.3 ± 5.6 ka (PD17), 92.2 ± 6.8 ka (PD41) and 92.0 ± 6.3 ka (PD15). These ages are internally consistent and a weighted mean age of 89.3 ± 3.6 ka may be derived. Because Unit 1 is stratigraphically located below Unit 3a, these results provide a maximum possible age for Unit 3 and the associated fossils.

The combination of these data suggests a 2σ time interval of 85.1 to 96.5 ka for AW-1 (86.5-92.9 ka at 1σ) based on the minimum and maximum age constraints derived from the direct U-series age ($87.6-2.5 = 85.1$ ka) and the weighted mean OSL age of Unit 1 ($89.3+7.2=96.5$ ka). This age range agrees well with the US-ESR age result obtained for WU1601 collected close to AW-1 (Supplementary Figure 14).

In order to further constrain the chronology of the deposits, all these data were incorporated into a Bayesian sequential phase model with phase 1 (underlying aeolian sand) corresponding to Unit 1 and Phase 2 (lake phase) to both Units 2 (no samples dated) and 3. We have not excluded shared systematic uncertainties (e.g. uncertainties shared between OSL ages) from the uncertainty term on individual ages. The exclusion of shared systematic uncertainties may be appropriate where: 1) a number of age estimates for a single event are being combined (e.g. 10 OSL ages for a single stratum) or 2) where the principal aim is to determine the duration of time represented in a phase of activity/deposition. However, we use the Bayesian sequential phase model to determine the timing of deposition of the AW1 specimen. Here the uncertainties on individual ages (whether shared or not) do represent the uncertainty on the timing (but not, in the case of shared uncertainties, the duration) of deposition. With this type of model removing systematic uncertainties would give a false sense of precision to the depositional age for AW1. Modelled ages indicate that deposition of Unit 1 ceased 93.1 ± 2.6 ka (Phase 1, PD15, 17, 41) and that Units 2 and 3 and all associated fossils were deposited between 92.2 ± 2.6 ka and 90.4 ± 3.9 ka (Phase 2, all other ages). The end date for phase 2 should be treated as a maximum value since no overlying material is present, precluding the possibility of further constraining the end of this phase. The timing of the lake phase is earlier than that indicated by other humidity records from the region. Given that the three samples from sands underlying the lake (arid phase indicators) are internally consistent, and coincide with the MIS 5c

insolation minimum, we propose that the subsequent lake phase correlates with the strengthened summer monsoon associated with the insolation peak at 84 ka (Fig. 6).

The code used to produce the Bayesian sequence model is as follows:

```
Plot()
{
Sequence()
{
Boundary("Start 1");
Phase("1")
{
Date("PD41",N(calBP(92200),6800))
{
};
Date("PD17",N(calBP(85300),5600))
{
};
Date("PD15",N(calBP(92000),6300))
{
};
};
Boundary("End of sand");
Boundary("Start of lake");
Phase("2")
{
Date("PD40",N(calBP(98600),7000))
{
};
Before("i")
{
Date("AW-1",N(calBP(87600),1250))
{
};
};
Before("ii")
{
Date("Hippo enamel",N(calBP(83500),4050))
{
};
};
Before("iii")
{
Date("Hippo dentine",N(calBP(65000),1050))
{
};
};
Date("US-ESR",N(calBP(103000),10000))
{
};
};
Boundary("End of lake");
};
};
};
```


SI 5. Stratigraphy, sedimentology and palaeoecology.

5.1 Results

5.1.1 Sedimentology

Unit 1 consists of loose, cross-bedded medium to coarse sands that contain evidence for bioturbation and iron-staining, this deposit underlies the sequence across the whole Al Wusta site (Supplementary Figures 13 and 14). The carbonate beds (Unit 2) exposed at Al Wusta are structureless and homogeneous, with some evidence for weak, horizontal laminations in PD16 sediments at depths between 70-40 cm. Carbonate content is high throughout all three sequences (typically >50%, Supplementary Figure 13), but the sediments have a low density typical of diatomite when dry. Shell fragments are rare but are present in the base of PD15 and PD16 and towards the top of PD16. These fragments are obvious in thin section but not in hand specimen and any picked remains could not be identified to specific taxa. During sieving of samples for isotopic analysis the coarse residue, >63 μm , was examined and ostracod fossils were not observable. XRD whole rock analysis (that characterises the mineralogy of crystalline mineral composition, and consequently will not identify organics and non-crystalline biominerals, i.e. diatoms), indicate that the samples are dominated by calcite (>90%) with minor amounts of quartz (Supplementary Table 18 and Supplementary Figure 15). Peaks that are attributable to evaporitic minerals such as gypsum and halite are present but the peak heights are so small that they are indistinguishable from background noise. Unit 3a consists of medium sands that are weakly horizontally laminated and contain fragments of eroded marls, occasional clasts and shells (*Melanoides tuberculata* and *Planorbis* sp). The upper part of Unit 3 (Unit 3b) is effectively desert pavement and has been

formed by the aeolian deflation of Unit 3a, concentrating clasts, shells, lithic and fossils at the land surface. The upper parts of Unit 3a contain calcite-cemented rhizolith that, as a result of progressive sediment deflation, now occur at the land surface as a cemented cap.

5.1.2 Micromorphology

All samples are dominated by homogeneous microsparite with some local zones of spar and micrite (Supplementary Figure 16a). Most sediments are massive with no clear sedimentary structures except for the sediments in PD16 (40-70 cm) which show evidence for finely laminated calcite, with laminations reflecting alternations between fine-grained micrite, coarser microspar and organics (Supplementary Figure 16b). Towards the top of PD17 laminations of allogenic quartz grains occur but these are infrequent (30-40 cm). Where quartz grains occur, their surfaces show signs of etching and replacement with calcite.

Organic remains are present in either an amorphous form or showing clear signs of cellular preservation. Diatoms and sponge spicules are visible in many slides (Supplementary Figure 16c) and the latter are typically most abundant at the sample levels with the lowest % carbonate values. Iron staining occurs towards the top of PD15 (40-50 cm) and towards the base of PD16 (0-10 cm). There is no evidence for neomorphism and negligible evidence for pore-infilling of secondary cement.

5.1.3 $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ analysis

The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of the sub-samples from the four sampling locations (PD15, 16, 17 and 40) are shown in Supplementary Figure 17 Data from all four sections show very similar

$\delta^{18}\text{O}$ values), although there is a significant scatter in each dataset. Greater variation occurs within the mean $\delta^{13}\text{C}$ values, although the standard deviations of each dataset are again large. Although the dataset from each sampling locality is small PD15 (n = 8), PD 16 (n = 15), PD17 (n = 7) and PD40 (n = 12) it is sufficient to show the following patterns: 1) there is no co-variance in $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values at any site or within the whole dataset level (r^2 ca 0.2) and 2) there is no consistent pattern of variation upwards through the profiles. With regard to the second point, while there is a clear trend at site PD40 of increasingly positive values upwards through the profile the reverse is true of PD16. Interpretation of these conflicting signals is problematic because it is unclear whether the deposits in each sequence are absolutely contemporaneous. One sample of calcite rhizolith was analysed from the calcrete capping PD40. This sample has the highest $\delta^{18}\text{O}$ value of the dataset, but it is the $\delta^{13}\text{C}$ value that is most significant as the high value (-1.12‰) implies that the vegetation that became established after the lake basin infilled/desiccated was a C4 grassland.

5.1.4 Diatoms

The diatom flora at PD15 and PD16 are shown in Supplementary Figures 18 and 19. Both sites show very similar species assemblages. The diagrams are divided into statistically significant zones based on a comparison with the broken-stick model using the program BSTICK version 1⁵². The species diagrams of each site can be divided into three statistically significant zones, the main characteristics of which are outlined below, along with the main environmental implications of these assemblages:

PD15

Zone I

This zone comprises planktonic species (*Lindavia rossi*, *Cyclotella krammeri* and *Lindavia ocellata*) which is illustrated by the high planktonic: benthic ratio. The CA and DCCA axis 1 scores decrease slightly in this zone. The log concentration, however, increases in this section meanwhile there is a minor decrease in the F index.

Zone II

This section reflects a change in species from mainly planktonic taxa to an increase in benthic taxa (*Cymbella affinis* and *Nitzschia angustata*) and periphytic (*Staurosirella lapponica* and *Ulnaria ulna* (agg.)). The increase in benthic and periphytic taxa is notable in the habitat composition diagram. The CA and DCCA axis 1 sample scores both decline considerably. There is also a decrease in the log concentration and the F index remains low.

Zone III

Aulacoseira italica and *Aulacoseira granulata* are the dominant taxa of this zone. The sample at 40 cm does not contain any benthic taxa therefore the planktonic: benthic ratio cannot be calculated for this level which is reflected in the habitat composition diagram. There is an increase in the CA axis 1 sample scores meanwhile a decrease in the DCCA axis 1 sample scores and log concentration occurs. The F index, which is a dissolution index, shows that the level of preservation is high for this sample in comparison to the rest of the profile.

pH

The assemblage reflects increasing pH as *Lindavia ocellata* usually occurs in mesotrophic lakes⁵³, whereas *Staurosirella lapponica*, *Stephanodiscus hantzschii*, *Aulacoseira granulata* and *Aulacoseira italica* are indicative of eutrophic lakes^{54,55}. *Staurosirella lapponica* and

Stephanodiscus hantzschii are good indicators of high alkalinity lakes meanwhile *Aulacoseira granulata* and *Aulacoseira italica* are characteristic of carbonate rich lakes.

Lake depth

The planktonic: benthic ratio and the habitat summary show that zone I is the deepest as planktonic (e.g. *Lindavia rossi* and *Cyclotella krammeri*) and tychoplanktonic (e.g. *Lindavia ocellata*) taxa increase to reach a peak at 0cm^{56,57}. The proportion of periphytic taxa increases throughout the zone II meanwhile benthic (e.g. *Cymbella affinis* and *Nitzschia angustata*) taxa reach a peak at 20 cm suggesting a decline in water depth which remains low for the remainder of the sequence. *Aulacoseira italica* and *Aulacoseira granulata*, which are periphytic and planktonic taxa respectively, dominate in zone III. *Aulacoseira granulata* suggests that lake levels were quite shallow⁵⁸ up to a few metres and quite turbulent^{55,59}. *Aulacoseira italica* accounts for 65% of the assemblage at 40 cm also suggesting shallower conditions as the habitat summary shows that this level comprises of periphytic and planktonic taxa with a small proportion of taxa with unknown ecology.

Salinity

There are very few saline tolerant species except for *Epithemia argus* which can be found in harsh conditions from saline to high alkaline, nutrient poor conditions. Due to its low abundance and the lack of other saline species salinity is not a major driver of this assemblage.

Phosphorous

The taxa (*Lindavia rossi*, *Cyclotella krammeri* and *Cyclotella ocellata*) show increasing nutrient enrichment from a mesotrophic lake, which suggests moderate phosphorous concentration to eutrophic conditions with high phosphorous concentrations characterised by *Stephanodiscus hantzschii*, *Aulacoseira granulata* and *Aulacoseira italica*⁵⁶⁻⁵⁹.

PD16

Zone I

Stephanodiscus and *Aulacoseira* are the dominant taxa in this zone with small increases in *Fragilaria delicatissima*, *Staurosirella lapponica* and *Cymbella affinis*. There is an increase in the planktonic: benthic ratio at the 10 cm which remains stable until zone II. The PCA and DCCA axis 1 scores diverge at 0 cm; however, both show little change meanwhile the log concentration and the F Index both increase until 10 cm before decreasing.

Zone II

There is a large decline in *Aulacoseira italica* in this section, while a decrease in the periphytic taxa occurs with a contemporaneous increase in benthic taxa. The PCA axis 1 scores remain stable; meanwhile there is an increase in the DCCA axis 1 sample scores, fluctuations in the log concentration and an increase in preservation in this zone.

Zone III

Lindavia rossi, *Cyclotella krammeri* and *Lindavia ocellata* are the predominant taxa of this zone; however, *Aulacoseira italica* and *Lindavia ocellata* are the main taxa at 70 cm. The reduced abundance of benthic taxa is reflected by the increased planktonic: benthic ratio. The PCA and DCCA axis 1 sample scores also illustrate the change in composition. The log concentration and F Index both follow the same trend and decrease overall in this section.

pH

The assemblage shows generally declining pH from *Stephanodiscus hantzschii*, *Stephanodiscus parvus*, *Aulacoseira angustata*, *Staurosirella lapponica*, *Aulacoseira italica*, *Cymbella affinis*, *Lindavia ocellata*, *Cyclotella krammeri* and *Lindavia rossi*. The interaction of other variables, for examples, depth, light and phosphorous can also affect the abundance of different taxa^{56,61}. The taxa present in zones I and II are associated with higher pH consistent with lower water levels which can concentrates the pH. In zone III the taxa changes and reflects lower pH conditions coherent with higher water levels which dilutes the pH level and other nutrients^{62,63}.

Lake depth

The planktonic: benthic ratio and habitat summary shows that zone I and II consists of mainly benthic taxa and periphytic taxa suggesting lower water levels. Zone III comprises of mostly planktonic and tychoplanktonic species, indicating higher water levels. The PCA and DCCA axis 1 sample scores both show a big change in the composition and, over time, of the assemblage, reflecting the change in water level.

Salinity

Epithemia argus is the only saline species present in the diagram which indicates that assemblage is not significantly influenced by salinity.

Phosphorous

The response to nutrient enrichment, which is usually linked to phosphorous, follows the same trend as pH of decreasing nutrient status as reflected by the declining nutrient tolerant

taxa. Nutrient cycling within the lake is also affected by other variables (e.g. depth, light and pH) thus in zones I and II where there is lower water depth the phosphorous concentration increases enabling *Stephanodiscus hantzschii* and *Aulacoseira granulata* to thrive. Conversely in Zone III the assemblage consists of species with a low threshold for phosphorous which suggests a higher water level as the phosphorous becomes more dilute^{62,63}.

The overall change in the diatom assemblages can be assessed by the changes in the species composition of the sample (PCA or CA) and over time between the samples (DCCA). CA was used in PD15 which shows that there are substantial changes from 20 cm to the top of the core as the exploratory DCA axis 1 gradient length was ≥ 1.5 SD units. The DCCA also shows a corresponding high species turnover which suggests a large change in species composition between samples which the decrease in the planktonic:benthic ratio reflects. PCA was used in PD16 (gradient length of exploratory DCA was ≤ 1.5 SD units) that shows there are smaller species compositional changes within this site than PD15. The largest change occurs between 40-50 cm at PD15 which is simultaneous with a change in taxa reflected by the sudden increase in the DCCA axis 1 sample scores and the planktonic:benthic ratio. Although the changes appear larger in PD15 there are caveats to consider; the small sample size of 5 samples at this site may not be as representative of the environment as a larger dataset so the results may change if more samples were analysed at more frequent intervals. The planktonic:benthic ratio of PD15 is smaller than that of PD16 suggesting less fluctuation in the water level.

5.2 Interpretation

The proxy data outlined above allows the environment and hydrology of the Al Wusta lake beds to be reconstructed. This has implications for both the environment of the site itself but also for the regional palaeoclimate of the time.

5.2.1 Environment and hydrology of the Al Wusta lake beds

The loose and cross-bedded nature of the sands of Unit 1 are typical of aeolian dune sands and indicate that, prior to carbonate accumulation, the Al Wusta site was affected by active dune migration. Unit 2, characterised by massive carbonate beds and their microfabrics, is typical of the accumulation of authigenic carbonate marl on the bed of a lacustrine environment^{64,65}. The precipitation of carbonate in the water column and its deposition out of suspension leads to the development of fine-grain marl beds. The structureless character of the beds is indicative of shallow water resulting in the sediment being oxidised and consequently bioturbated, or exposed to water column turbulence by surface winds. The preservation of laminations within PD16 most probably relates to short-term episodes of deeper water or more rapid rates of sedimentation. Despite being shallow there is no sedimentary evidence to indicate that the lacustrine environment underwent desiccation, as there is a complete absence of surface exposure features, and the character of the sediments are consistent throughout the entirety of the bed⁶⁴. The shallow nature of the water is supported by the diatom assemblage which contains significant proportions of *Aulacoseira italica* and *Aulacoseira granulata* which indicates a few metres of water depth, with the latter being indicative of turbulent water conditions. While changes in the planktonic/benthic ratio of the diatom assemblage suggest progressive changes in water level there is no evidence to indicate that the water level dropped to sub-aerially expose the lake beds. Consequently, the beds record the existence of a perennial water body.

The diatom flora is dominated by freshwater species with negligible evidence for salinity tolerant species. The lake was, therefore, a freshwater environment; an interpretation supported by the dominance of calcite in the lake XRD mineralogy traces and an absence of evaporitic minerals such as halite and gypsum.

A lack of covariance in $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values is often used to indicate whether a lake basin was part of an open system (i.e. a system with an active inflow and outflow to the basin) or not. Generally, the absence of co-variance is indicative of open-system lake waters that are regularly recharged; this has the effect of limiting any in-basin modification of the isotopic signal by processes such as evaporation⁶⁶. Such a scenario could be true of the Al Wusta lake beds as the persistence of freshwater conditions (as evidenced by the diatom assemblages) and the lack of evaporative minerals would indicate that this system never underwent sufficient evaporation to produce brackish or saline conditions. However, two points are important to consider here. Firstly, a number of studies exist that report closed lake systems that do not show co-variance in their isotopic signals⁶⁷. The exact cause of this is not always clear, however, it highlights the fact that the absence of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ co-variance at Al Wusta should not be used as definitive evidence of this site recording an open lake system. Secondly, it is important to acknowledge that, even when there is no isotopic covariance, the $\delta^{18}\text{O}$ of the water in open system lakes can still become enriched as a result of the evaporation of surface waters. However, as such systems have regular recharge this does not result in the waters becoming brackish or saline⁶⁸. This is an important point as it is impossible to say whether minor or major evaporative enrichment in lake $\delta^{18}\text{O}$ water values occurred at Al Wusta, but the diatom and mineralogical evidence shows that if such evaporation did occur, it was not sufficient to cause the waters to become brackish.

In summary, the proxy evidence indicates that Unit 2 represents the deposition of marl in a perennial water body that was entirely fresh. At this time the Al Wusta lake basin was an accessible and permanent freshwater resource for any early humans in the region. The pattern in the $\delta^{18}\text{O}$ values seen at site PD40 (an increase in $\delta^{18}\text{O}$ values upward through the sequence) could reflect a trend to greater aridity during the accumulation of this sequence (with increasing values reflecting greater evaporation and hence greater aridity). However, as the reverse is true for PD16 it is difficult to know how robust this interpretation is.

The mean $\delta^{18}\text{O}$ values of this sequence (-2.4 to -1.9‰) reflect both the $\delta^{18}\text{O}$ value of the lake water and the temperature at which carbonate precipitation occurred⁶⁶. Both of these are unknown, however, the nearest local isotopic dataset indicates that modern rainfall has a $\delta^{18}\text{O}$ value of -4.0 to -4.5‰⁶⁹. If such values occurred at the time of marl development at Al Wusta then it is likely that either the temperature of mineralisation was surprisingly low (i.e. a carbonate with a mean $\delta^{18}\text{O}$ value of -2.2‰ precipitating from water with a mean $\delta^{18}\text{O}$ value of -4.25‰ would form in isotopic equilibrium at a temperature of ca 11°C) or that significant isotopic enrichment, through evaporation, of the lake water had occurred. It is also important to consider, however, that the modern climate system that generates rainfall with a value of -4.0 to -4.5‰ is a function of a westerly dominated system. As most humid phases in the Arabian Peninsula are suggested to relate to increased monsoon strength⁷⁰, it is likely that the $\delta^{18}\text{O}$ value of the rainfall during the formation of the lake system at Al Wusta was significantly different, and most likely heavier, than modern day values. A strong south to north gradient in the $\delta^{18}\text{O}$ value of modern rainfall appears to exist in Arabia in the present day, with values in the southern part of this region being >0‰ in contrast with those in the north that are around 4‰⁷¹. Only a small northward shift in the air masses bringing rainfall

from a southerly source would, therefore, produce a significant increase in the $\delta^{18}\text{O}$ value of rainfall at Al Wusta.

Consequently, it is currently unclear whether the marl $\delta^{18}\text{O}$ value of ca -2.2‰ is a function of low water temperatures, isotopic enrichment, different air mass trajectory or a combination of all of these factors. Although many of the artefacts and fossils are found within Unit 3 it is important to stress that *in situ* artefacts are found with Unit 2 implying that humans were present in the landscape during the existence of the perennial freshwater lake body.

The deposits of Unit 3a are interpreted as waterlain sands based on the presence of freshwater molluscs, the horizontal laminations and the incorporation of ripped up fragments of the underlying marl. The sands are interpreted as reflecting the encroachment of fluvial sedimentation during the final phase of basin infilling, either as a result of desiccation or simply due to the progressive reduction in available accommodation space. It is considered that the former is more likely as the inclusion of ripped up marl fragments would imply that the upper surface of Unit 2 had dried out and fragmented prior to the deposition of Unit 3a. The waterlain sands are capped by a horizon of calcified rhizoliths that represent the final transition from sub-aqueous to terrestrial conditions at the site. The coarse surface lag (Unit 3b) reflects the winnowing of Unit 3a by aeolian processes and the formation of a desert pavement. The relatively resistant nature of the rhizolith horizon makes this layer resistant to aeolian erosion resulting in the occurrence of these features as a cemented cap at the top of the sequence.

The majority of the excavated Al Wusta artefacts and fossils are found in Unit 3a. As the sediments of this unit contain evidence of eroded and reworked marl from Unit 2 it is

possible that the archaeological and faunal material has been derived from Unit 2 and reworked into the fluvial sands. If this is the case then it supports the idea that human occupation at the site is associated with the existence of a perennial, freshwater lake. If the artefacts and fossils are *in situ* then it implies that occupation of the site must have persisted after the contraction of the lake basin and the desiccation of the water body.

5.2.2. Implications of the Al Wusta sequence for regional palaeoclimate

The proxy record of the Al Wusta sequence records the existence of a perennial, freshwater lake system. Regarding human migration into this region, it is important to establish whether the existence of such a system can be explained by local hydrogeology or by increased regional moisture (i.e. the occurrence of a humid phase). It is argued here, despite evidence from only a single site being presented, that an increase in mean annual precipitation is required to explain the development of the Al Wusta lake basin. This suggestion is based upon three main observations. Firstly, the fact that Unit 2 comprises well-developed marl beds means that the lake waters must have been fed by groundwater recharge. The presence of sufficient dissolved Ca^{2+} in the lake waters to produce extensive marl precipitation requires migration of the waters through an aquifer. While much of the underlying geology is quartz rich sands, these are interbedded with units of carbonate rich marls that reflect phases of lake activity that pre-date the formation of the Al Wusta sequence. It is likely that these beds are the source of the dissolved minerals that fed into the Al Wusta basin. Secondly, no groundwater fed lake systems currently exist in the Nefud as the water table is too low. Surface water bodies are, therefore, restricted to highly ephemeral recharge playas. Thirdly, the underlying sediments and bedrock in this region are highly permeable aeolian sands; no local impermeable strata exist that could generate a locally perched water table that would

explain the existence of a lake system at this site. It is, therefore, proposed that the genesis of a groundwater fed lake system at this locality requires a regional increase in mean annual rainfall and the persistence of a humid phase in the Nefud Desert.

5.2.3 Summary

The Al Wusta sequence records the evolution of the site across an entire humid phase from aeolian deposition (Unit 1) to the development of lacustrine conditions (Unit 2) to the progressive drying out of the lake basin and the reversion to terrestrial conditions (Units 3a and b). In particular, the diverse range of proxies discussed here indicate that the Unit 2 carbonate deposits at Al Wusta were deposited in a perennial shallow (a few metres of water depth) alkaline lake environment. This lake water body formed as a result of a humid phase and was fresh throughout its existence, making it a major resource for humans and fauna during this interval.

SI 6. Al Wusta vertebrate palaeontology, biogeography and taphonomy.

The Al Wusta vertebrate fossil assemblage (Supplementary Table 19; Supplementary Figure 19) is very fragmented, with few complete bones recovered. Fossil weathering is extensive and much of the cortical surfaces of the bones are missing, making skeletal element and taxonomic identification difficult. Of the 860 specimens, 305 are identifiable to skeletal element, with long bone shafts most common (Supplementary Table 20). Of the 12 distinct taxa identified, medium-sized bovids are most abundant, followed by small-sized bovids and *Hippopotamus*. *Hippopotamus* is represented solely by incisor and tusk fragments recovered *in situ*. *Kobus* sp. is represented by partial right M₂ and M₃ teeth (Supplementary Figure 20). The M₃ has a distally positioned lingual accessory cusp, simple and flattened U-shaped infundibulum, and a rounded buccally projecting hypoconulid. The occlusal length is within the range of *K. ellipsiprymnus*; nevertheless, the specimen is too fragmented to allow positive species identification. Rodentia is represented by two specimens, a maxilla fragment missing all molars, and an almost complete cranium including part of the mandible. Despite the completeness of this specimen, it is distorted, fragile and partially covered in matrix, making a more precise taxonomic identification difficult. Two isolated reptile teeth belonging to a species of *Varanus* were identified. The presence of *Struthio* is confirmed by numerous egg shell fragments recovered *in situ*. Additionally, two smaller-bodied species of birds are represented by incomplete long bone fragments.

While *Hippopotamus* is restricted to Africa today, during the Pleistocene it was common throughout parts of Europe and Asia^{72,73}. Dispersals out of Africa led to localised speciation events in Europe and the Levant⁷⁴; however, there is some debate surrounding the precise number of *Hippopotamus* species^{75,76}. While it seems likely that the Al Wusta specimens

represent an out of Africa dispersal, it should be noted that *H. amphibius* has been identified at sites in Britain and Central Europe dated near the Middle to Late Pleistocene transition^{72,77,78}. *Pelorovis* is common in East Africa throughout the Pleistocene and is also found at a few sites in North Africa but is absent from Europe and the Levant after the Early Pleistocene⁷⁹⁻⁸¹. Both *Pelorovis* and *Hippopotamus* have been identified at other sites in Saudi Arabia^{82,83} suggesting repeated dispersal events into the Arabian Peninsula during periods of climate amelioration. *Kobus* typically inhabit flood-plains and grasslands bordering water⁸⁴. During the Middle and Late Pleistocene *Kobus* was restricted to Africa and mostly south of the Maghreb. Notable northern occurrences were reported from Algeria and Egypt, associated with the Last Interglacial^{85,86}, but is unknown from the Levant during this period. This may support a tendency toward longitudinal dispersals⁸⁷, in this case eastward dispersal out of Africa and into the Arabian Peninsula via the Southern Levant. *Struthio* is common in Africa, but has also been identified by eggshell fragments on the Indian subcontinent from as early as the Middle Miocene and as recently as 18 ka^{87,88}. *Struthio* has been found at several sites in the Levant, and was extant in Southwest Asia until the 20th century.

Weathering of many specimens, particularly those collected from the surface at the western edge of the southern ridge at Al Wusta, suggests that they were likely exposed for a significant period of time post-fossilisation, with abrasion and polishing attributable to aeolian processes. Surface fossils included several refits recovered in close proximity, suggesting limited post-exposure transport. It is likely that such specimens became concentrated on the surface as a result of winnowing of the main fossiliferous layer. For these specimens, post-mortem weathering (i.e. prior to fossilisation) was impossible to determine due to the extensive fossil weathering of the cortical surfaces (i.e. following fossilization).

Specimens collected *in situ* were typically better preserved and lacked the abrasion and polishing common to the specimens from the surface. Pre-depositional weathering of these remains, on the other hand, was difficult to determine due to the fragmentary nature of the fossils and the fact that many specimens remain encrusted in carbonate. Despite the absence of large-bodied mammalian carnivores in the fossil assemblage, tooth pits potentially attributable to hyenas, large canids, or large felids are evident on some bones (Supplementary Figure 20H; Supplementary Table 21). Manganese staining is also present on numerous specimens. Two small bone fragments may have been burnt: one is blackened and the other one is dark brown in colour. However, their preservation is too poor to confidently rule out diagenetic discolouration. No additional bone surface modifications were observed, although this was not surprising considering that fine scale modifications are likely to have been removed during weathering.

Analysis of long bone circumference revealed a majority of type 1 shafts (75%), with type 2 and type 3 occurring much less frequently (12% and 13%, respectively). The type 3 and type 2 to type 1 index is 0.31, falling within the typical range identified for assemblages accumulated by carnivores and hominins⁸⁹. While extensive fossil weathering may also have contributed to the production of type 1 shafts; nevertheless, there is evidence for large carnivores and hominins at Al Wusta and it is likely that they contributed to the production of type 1 shafts during prey processing and consumption. Furthermore, the under-representation of epiphyses (N = 11) at Al Wusta supports carnivore ravaging of long bone ends⁹⁰ although this may also be an artefact of other non-biological preservation biases. Additional support that bone breaking agents were, at least in part, responsible for the accumulation of bones is the presence of green breaks (N = 12). Unfortunately, the assemblage is too poorly preserved to determine a primary accumulator.

SI 7. Al Wusta lithic technology.

7.1 Introduction

The Al Wusta lithic (stone tool) assemblage consists of 380 artefacts, systematically collected in transects and piece plotted from the southern end of the site in close proximity to the marl beds and from excavations in 2016 and 2017 (Figure 1). Lithics continued to the north, but were not included in the analysis here, which is limited to material south of the Holocene playa (Figure 1). The assemblage is Middle Palaeolithic in its characteristics, with a focus on centripetal Levallois reduction, particularly of chert and quartzite. Retouched tools are predominantly side-retouched flakes. The lithics, all from the same spatially restricted area, also display similar raw materials, weathering and technology and are therefore treated here as a single assemblage. The Al Wusta lithic assemblage is a valuable reference point for late MIS 5 Arabia, and displays similar technological features to other contemporary Arabian assemblages^{43,91-93}.

The Al Wusta lithic assemblage can be divided into the following categories (% in brackets shows % of total assemblage): 229 flakes (60.20%) [of which 36 are Levallois flakes, 9.47%], 55 chips and chunks (14.47%), 21 retouched flakes (5.52%), one hammerstone (0.26%), and 74 cores (19.47%). Our aim here is to describe the basic technological characteristics of this lithic assemblage, which we analysed using the methodology and terminology described by Scerri and colleagues⁹⁴⁻⁹⁶ and Groucutt and colleagues^{93,97}.

The surface lithic assemblage was found closely associated with the marl deposits. Eleven lithics were excavated at the site, in the upper part of the marl and in the overlying *in situ*

sand layer. The excavated material demonstrates similar characteristics to the larger sample from the surface, and includes similar raw material (focus on local lacustrine chert), three Levallois cores, flakes with faceted platforms and a debordant flake. The density of lithics on the surface, together with the considerable variation in their size suggests that they are not extensively redeposited, and we suggest that they were made by hominins beside the late MIS 5 lakeshore and then deposited into the lake by gravity and low energy movement.

7.2 Raw materials

In terms of raw material composition, the most common material consists of a local lacustrine chert (65.09%), followed by quartzite (17.59%), ferruginous quartzite (11.02%), quartz (5.25%), argillaceous sandstone (0.52%), and other sandstone (0.52%). This raw material structure is unusual for the area. Al Wusta is the only identified Middle Palaeolithic assemblage in the region not dominated by quartzite, generally highly ferruginized^{91,92,97-99}. It appears that suitable quartzite is rare in the environs of Al Wusta, with our survey of the area showing an absence of large beds of ferruginous quartzite (Figure 5c). Non-ferruginous forms of quartzite (e.g. Figure 5f) are relatively common in the assemblage, and appear to occur as rounded pebbles of fluvial or conglomerate origin. Aside from chert, other materials occur in low frequencies, such as quartz (Figure 5e), which occurs as small pebbles in the bedrock of the area, and a rare type of rock which can be described as argillaceous sandstone⁹⁹ (Figure 5A).

The dominant raw material consists of chert, which occurs with varying colours and textures. The most common is brown in colour (e.g. Figure 5d). Our surveys of the area show that this material has formed in lakebeds in the area, which form a raw material source when exposed

by subsequent erosion. We have conducted knapping experiments with this raw material, which shows that while variable, it is difficult material for knapping. The dominant chert at Al Wusta seems to be a particular poor quality form. The chert clasts are relatively small (always smaller than fist sized), have a very thick and undulating cortex, and frequently have inclusions. The material is also extremely hard, and has to be struck with considerable force to remove a flake. Despite these limitations, if knappers could access the inner part of the nodules, the chert is fine grained and appears to have reasonable properties of conchoidal fracture.

7.3 Core technology

The 74 residual Al Wusta cores are dominated by Levallois cores (54, 73%), with smaller numbers of non-Levallois multiple platform cores (12, 15.6%), double platform (bidirectional) cores (two, 2.7%), single platform cores (two, 2.7%), tested/minimally flaked cores (two, 2.7%), one radial core (i.e. a core flaked centripetally, but which is neither Levallois nor discoidal) (1.4%), and one core fragment (1.4%). It is clear, therefore, that reduction was dominated by use of the Levallois technique. Other methods appear more ad hoc in character. Multiple platform cores, for instance, tend to be small, having a mean weight of 19.3 g if one large outlier weighing 349.9 g is removed. This compares to the average weight for all cores of 34.9 g (with the same outlier also excluded). This suggests that multiple platform core reduction, the main non-Levallois reduction method, was employed on small clasts and/or on heavily reduced cores, which may have been Levallois cores earlier in reduction. There is no evidence of *façonnage* reduction, nor of blade production.

Of the 54 Levallois cores, 36 (66.7%) are preferential Levallois cores with centripetal preparation (e.g. Supplementary Figure 21a,b,d) and 13 (24.1%) are recurrent centripetal Levallois cores (e.g. Supplementary Figure 21C). The small number of pieces not in these categories are indeterminate Levallois cores where overshoot removals have removed the whole debitage surface (9.2%). From the geometry of the cores and the morphology of platform surface these are clearly centripetal Levallois cores, but it is not clear if their final phase of reduction was recurrent or preferential in character. Finally, one example each of recurrent unidirectional and recurrent bidirectional Levallois cores were identified. Striking platforms are faceted on 87% of the Levallois cores. In most cases preferential scars on Levallois cores demonstrate the product of parallel sided and sometimes oval Levallois flakes, with a small number producing pointed products.

Preferential and recurrent cores are similar in many features, for instance they have a similar number of scars (average of 12.9 and 10.9 respectively) and a similar percentage of cortical cover (23% and 27%). However, in terms of size recurrent cores are typically smaller (14.3 g, σ 9.0) than preferential cores (36.9 g, σ 42.7). These data indicate that, as a general pattern and perhaps within a situation of interchangeable reduction between Levallois methods, larger cores were reduced preferentially and a recurrent method was used late in reduction. The predominant focus on centripetal Levallois reduction demonstrates a level of homogeneity to core reduction at the site.

7.4 Debitage

Lithic debitage (n=284) at Al Wusta can be classified as complete flakes (185, 65.1%), broken flakes (44, 15.5%) and chips/chunks (55, 19.4%). Our description here focusses on complete flakes.

The average length (technological), width (mid-point), thickness (mid-point), and weight of complete flakes are 29.8 mm (σ 11.9), 24.5 mm (σ 9.2), 8.46 mm (σ 4.2), and 13.1 g (σ 16.3) respectively. To aid comparability, if we exclude flakes which are less than 20 mm in length, these values become 32.25 mm (σ 11.3), 25.7 mm (σ 9.4), 9.3 mm (σ 4.3), and 14.6 (σ 16.9).

Other basic features for complete flakes >20 mm in length include an average cortical cover of 16.5% of the debitage surface, an average of 2.8 dorsal scars (σ 1.7), striking platforms are prepared (faceted or dihedral) in 41.6% percent of cases and striking platforms have average external platform values of 74.9° (σ 6.2). Dorsal scar patterns can be classified as: 31.5% unidirectional, 2.4% unidirectional convergent, 1.6% perpendicular, 3.1% crossed, 11.8% bidirectional, 13.4% subcentripetal, and 36.2% centripetal.

Complete Al Wusta Levallois flakes (e.g. Supplementary Figure 21E,G-K) are on average 33 mm in length, 25.5 mm in width and 7.4 mm thick. They are generally broadly parallel sided in shape. They most commonly (64.5%) have centripetal scar patterns, with a further 12.9% having subcentripetal scar patterns. The remaining Levallois flakes are characterised by unidirectional scar patterns (16.1%) and unidirectional convergent (6.5%). 74.2% of the Al Wusta Levallois flakes have prepared platforms.

Summarising these data, Al Wusta flakes are therefore typically small, quite commonly cortical and with various scar patterns but particularly unidirectional and centripetal. Levallois flakes are also small, and indicate a tendency to produce Levallois flakes by centripetal preparation and hard hammer percussion from prepared platforms. These

characteristics are consistent with the evidence from the cores in indicating that lithic technology at Al Wusta is focussed on centripetal Levallois reduction.

7.5 Retouched flakes

A total of 21 retouched flakes were found at Al Wusta (15 complete). Three of these have only distal retouch (e.g. Supplementary Figure 21m), seven are retouched along one lateral (e.g. Supplementary Figure 21n), seven are retouched on both laterals (e.g. Supplementary Figure 21l), and two are retouched on both laterals and distally (two more are fragments). Where it can be determined, 87.5% of striking platforms are faceted or dihedral.

Of the complete retouched pieces, they are mostly parallel sided and 66.7% have either cortical surfaces or unidirectional scar patterns. Their average length, width, thickness and weight are 33.6 mm (σ 11.5), 24.34 mm (σ 12.4), 10.7 mm (σ 2.4), and 15.9 g (σ 2.4). These are similar values to the overall flake population.

In most cases retouch is 'regular', while 33% of retouched flakes have notches as well as regular retouch. In all but one case, the retouch is continuous rather than clustered, and virtually all of the retouch is semi-abrupt. Retouch is always found on either the dorsal surface or on both surfaces, and never exclusively on the ventral surface. Index of Invasiveness (I of I)¹⁰⁰, values range from 0.094 to 0.5, averaging at 0.25. Geometric Index of Unifacial retouch (GIUR)¹⁰¹ values range from 0.45 to 0.89 with an average of 0.69. The quite low values for I of I and high values for GIUR demonstrates that retouch was not horizontally extensive across the face of the flakes, but was vertically quite intensive.

The retouched component of Al Wusta therefore shows a focus on typical side retouched flakes (“scrapers”) with prepared platforms, as is commonly the case for Middle Palaeolithic assemblages.

7.6 Al Wusta lithic technology in comparative context

The lithic assemblage of Al Wusta demonstrates a consistent approach to lithic technology, focussed on centripetal Levallois technology. Here we will briefly consider this technology in relation to sites elsewhere in Arabia and surrounding regions.

The corpus of Arabian sites which can be related to later MIS 5 by chronometric dating techniques is small, but increasing. These include the ~75 ka assemblage at JQ-1 at Jubbah^{43,91}, the ~ 100-60 ka assemblage from JSM-1 at Jubbah^{91,95}, and the ~85 ka assemblage from Mundafan Al Buhayrah in southwestern Arabia⁹³. We emphasise the latter assemblage from Mundafan as the key example, with both good chronometric age estimates and a large and diagnostic lithic assemblage. Most Arabian Middle Palaeolithic assemblages remain undated, or have produced contradictory age estimates, which mean little can be said on their age with certainty. The available chronologically secure data indicate that mid to late MIS 5 sites demonstrate a focus on centripetal Levallois reduction, with both centripetally prepared preferential and recurrent centripetal Levallois methods employed. In all cases they lack bifacial *façonnage* technology and beaked (‘Nubian’) Levallois reduction. Levallois point production only occurs at marginal levels. Differences at these sites tend to correlate with raw material aspects. Both MDF-61 and JQ-1 appear to be located far (>10 km) from good raw material sources, and in both cases demonstrate highly reduced assemblages, both in terms of core reduction and high levels of retouch^{93,99}. In terms of size aspects, lithics at Al

Wusta are relatively small for MP/MSA assemblages. They are much smaller than many Arabian sites, but also not as small as sites with highly reduced assemblages elsewhere in southwest Asia, such as Tor Faraj in Jordan and Warwasi in Iran^{93,97,102}. We associate these small size aspects primarily with raw material factors at Al Wusta. The relatively high levels of retouch at Al Wusta likewise likely correlate with the limitations of raw material in the area, encouraging resharpener/edge rejuvenation of existing pieces.

In areas to the north and west of Arabia a number of assemblages date to mid to late MIS 5 and share similar technological characteristics with Al Wusta. In the Levant this is most clearly demonstrated by Qafzeh Cave, with its famous *Homo sapiens* fossils, dating to ~100-90 ka¹⁰¹. Moving into northeast Africa, sites such as 1017 and 34a have been attributed to later MIS 5 (~85-80 ka) on both stratigraphic grounds and with preliminary chronometric age estimates¹⁰⁴⁻¹⁰⁶. The technology of these sites again demonstrates a focus on centripetal Levallois reduction. The Bir Tarfawi and Bir Sahara palaeolakes feature broadly similar technology, and were repeatedly occupied during the wet phases of MIS 5¹⁰⁷. In the Horn of Africa key evidence comes from a series of dated assemblages in the Aduma area¹⁰⁸. These date to between ~100 and 80 thousand years ago, and as with sites such as Al Wusta demonstrate a focus on the centripetal reduction of small Levallois cores, and are also associated with *Homo sapiens* fossils. While these sites feature similar core reduction methods, in contrast to, for example, the unidirectional focus of MIS 4-3 Neanderthal assemblages in southwest Asia, further research is needed to understand similarities and differences in areas such as retouched tool technology.

While traditionally masked by variable analytical methodologies, differing regional nomenclatures and a lack of knowledge on areas such as Arabia, we interpret these findings

as indicating broad technological similarities across a large area in mid to late MIS 5¹¹⁰. However, East African sites appear to feature more of a focus on retouched point production (although this must be matched against these sites having large sample sizes compared to assemblages in places such as Arabia). In terms of core reduction methods, however, assemblages in East Africa, northeast Africa, the Levant and Arabia all appear to be similar in mid to late MIS 5. In line with available fossil evidence, we note that this pattern appears to be most ancient in East Africa^{108,109}. In contrast to East Africa, areas such as the Nile Valley, the Levant and Arabia demonstrate highly variable MP/MSA records. This pattern suggests that at least in terms of visible (archaeologically preserved) aspects of material culture, the earlier phases of dispersal into southwest Asia did not relate to radical technological innovation. Phases of dispersal correlate with windows of climatic amelioration in the Saharo-Arabian arid belt, while further research is needed to understand the nature of demographic, social and behavioural changes in sub-Saharan Africa which might have triggered dispersals. Beyond these broad observations of similarities in contemporaneous similarities, detailed comparative studies of chronometrically constrained lithic assemblages are needed to clarify patterns of similarities and differences in lithic assemblages.

Supplementary References

1. Martinez-Navarro, B., *et al.* Early Pleistocene “hominid remains” from southern Spain and the taxonomic assignment of the Cueva Victoria phalanx. *J. Hum. Evol.* **48**, 517-523 (2005).
2. Martinez-Navarro, B., *et al.* Reply to Gilbert et al. (2008) on the supposed human phalanx from Cueva Victoria (Cartagena, Spain). *J. Hum. Evol.* **54**, 157-161 (2008).
3. Ruff, C. B. Body size, body shape, and long bone strength in modern humans. *J. Hum. Evol.* **38**, 269-90 (2000).
4. Pearson O. M, Lieberman D. E. The aging of Wolff's "law": ontogeny and responses to mechanical loading in cortical bone. *Am. J. Phys. Anthropol.* **125**, 63-99 (2004)
5. Ruff, C., Holt, B., Trinkaus, E. Who's afraid of the big bad Wolff?: "Wolff's law" and bone functional adaptation. *Am. J. Phys. Anthropol.* **129**, 484-98 (2006).
6. Ruff, C. B., Trinkaus, E., Walker, A., Larsen, C. S. Postcranial robusticity in Homo. I: Temporal trends and mechanical interpretation. *Am. J. Phys. Anthropol.* **91**, 21-53 (1993).
7. Rasband, W. S. *ImageJ*. Bethesda, Maryland, USA: U.S. National Institutes of Health, <http://imagej.nih.gov/ij/> (1997-2016).
8. Doube, M. et al. BoneJ: Free and extensible bone image analysis in ImageJ. *Bone* **47**, 1076-9 (2010).
9. Ruff, C.B. in *Biological Anthropology of the Human Skeleton* (Katzenberg M.A., Saunders, S.R.) 183-206 (Wiley, 2007).
10. Ruff, C. Sexual dimorphism in human lower limb bone structure: relationship to subsistence strategy and sexual division of labor. *J. Hum. Evol.* **16**, 391-416 (1987).

11. Shaw, C. N, Stock, J. T. Intensity, repetitiveness, and directionality of habitual adolescent mobility patterns influence the tibial diaphysis morphology of athletes. *Am. J. Phys. Anth.* **140**, 149-59 (2009).
12. Grün, R., Eggins, S., Kinsley, L., Mosely, H., Sambridge, M. Laser ablation U-series analysis of fossil bones and teeth. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **416**, 150-167 (2014).
13. Benson, A., Kinsley, L., Defleur, A., Kokkonen, H., Mussi, M., Grün, R., 2013. Laser ablation depth profiling of U-series and Sr isotopes in human fossils. *J. Arch. Sci.* **40**, 2991-3000 (2013).
14. Grün, R. Methods of dose determination using ESR spectra of tooth enamel. *Radiat. Meas.* **32**, 767-772 (2000).
15. Duval, M., Grün, R., Falguères, C., Bahain, J.J., Dolo, J.M. ESR dating of Lower Pleistocene fossil teeth: Limits of the single saturating exponential (SSE) function for the equivalent dose determination. *Radiat. Meas.* **44**, 477-482 (2009).
16. Duval, M., Guilarte Moreno, V., Grün, R. ESR dosimetry of fossil enamel: some comments about measurement precision, long-term signal fading and dose–response curve fitting. *Rad. Prot. Dosim.* **157**, 463-476 (2013).
17. Duval, M., Grün, R. Are published ESR dose assessments on fossil tooth enamel reliable? *Quatern. Geochron.* **31**, 19-27 (2016).
18. Grün, R., Katzenberger-Apel, O. An alpha irradiator for ESR dating. *Ancient TL* **12**, 35-38 (1994).
19. Marsh, R. E. *Beta-gradient Isochrons Using Electron Paramagnetic Resonance: Towards a New Dating Method in Archaeology.* (MSc thesis, McMaster University, Hamilton, 1999).

20. Guérin, G., Mercier, N., Adamiec, G.. Dose rate conversion factors: update. *Ancient TL* **29**, 5-8 (2011).
21. Grün, R. The DATA program for the calculation of ESR age estimates on tooth enamel. *Quatern. Geochron.* **4**, 231-232 (2009).
22. Grün, R., Schwarcz, H.P., Chadam, J. ESR dating of tooth enamel: Coupled correction for U-uptake and U-series disequilibrium. *Int. J. Radiat. Appl. Instrum. Nucl. Tracks. Radiat. Meas.* **14**, 237-241 (1988).
23. Duval, M. in *Encyclopaedia of Scientific Dating* (eds Rink, W.J., Thompson, J.W.) 239-246 (Springer, 2015).
24. Ludwig, K. R. *User's manual for Isoplot 3.75*. Berkeley Geochronology Center, Special Publication **5**, 1-75 (2012).
25. Sambridge, M., Grün, R., Eggins, S. U-series dating of bone in an open system: The diffusion-adsorption-decay model. *Quatern. Geochron.* **9**, 42-53 (2012).
26. Bøtter-Jensen, L., Andersen, C. E., Duller, G. A. T., Murray, A. S. Developments in radiation, stimulation and observation facilities in luminescence measurements. *Radiat. Meas.* **37**, 535-541 (2003).
27. Duller, G. A. T., Bøtter-Jensen, L., Murray, A. S. Combining infrared- and green-laser stimulation sources in single-grain luminescence measurements of feldspar and quartz. *Radiat. Meas.* **37**, 543-550 (2003).
28. Duller, G. A. T., Bøtter-Jensen, L., Murray, A. S., Truscott, A. J. Single grain laser luminescence (SGLL) measurements using a novel automated reader. *Nucl. Instr. Meth. Phys. Res.* **155**, 506-514 (1999).
29. Armitage, S., Bailey, R. The measured dependence of laboratory beta dose rates on sample grain size. *Radiat. Meas.* **39**, 123-127 (2005).

30. Ballarini, M., Wintle, A. G., Wallinga, J. Spatial variation of dose rate from beta sources as measured using single grains. *Ancient TL* **24**, 1-7 (2006).
31. Armitage, S. J., Jasim, S. A., Marks, A. E., Parker, A. G., Usik, V. I., Uerpmann, H. - P. The Southern Route "Out of Africa": Evidence for an Early Expansion of Modern Humans into Arabia. *Science* **331**, 453-456 (2011).
32. Murray, A. S., Wintle, A. G. Luminescence dating of quartz using an improved single-aliquot regenerative-dose protocol. *Radiat. Meas.* **32**, 57-73 (2000).
33. Jacobs, Z. et al. Ages for the Middle Stone Age of southern Africa: Implications for human behavior and dispersal. *Science* **322**, 733-735 (2008).
34. Roberts, R. G., Galbraith, R. F., Olley, J. M., Yoshida, H., Laslett, G. M. Optical dating of single and multiple grains of quartz from Jinmium rock shelter, northern Australia: Part II, results and implications. *Archaeometry* **41**, 365-395 (1999).
35. Wallinga, J., Murray, A., Duller, G. A. T. Underestimation of equivalent dose in single-aliquot optical dating of feldspars caused by preheating. *Radiat. Meas.* **32**, 691-695 (2000).
36. Jacobs, Z., Duller, G.A.T., Wintle, A.G. Interpretation of single grain Distributions and calculation of De. *Radiat. Meas.* **41**, 264-277 (2006).
37. Thomsen, K.J., Murray, A.S., Bøtter-Jensen, L. Sources of variability in OSL dose measurements using single grains of quartz. *Radiat. Meas.* **39**, 47-61 (2005).
38. Duller, G. A. T. Assessing the error on equivalent dose estimates derived from single aliquot regenerative dose measurements. *Ancient TL* **25**, 15-24 (2007).
39. Duller, G. A. T., Bøtter-Jensen, L., Murray, A. S. Optical dating of single sand-sized grains of quartz: Sources of variability. *Radiat. Meas.* **32**, 453-457 (2000).
40. Jacobs, Z., Duller, G. A. T., Wintle, A. G. Optical dating of dune sand from Blombos Cave, South Africa: II - Single grain data. *J. Hum. Evol.* **44**, 613-625 (2003).

41. Thomsen, K. J., Murray, A. S., Bøtter-Jensen, L., Kinahan, J. Determination of burial dose in incompletely bleached fluvial samples using single grains of quartz. *Radiat. Meas.* **42**, 370-379 (2007).
42. Duller, G. A. T. Distinguishing quartz and feldspar in single grain luminescence measurements. *Radiat. Meas.* **37**, 161-165 (2003).
43. Petraglia, M. D. et al. Hominin Dispersal into the Nefud Desert and Middle Palaeolithic Settlement along the Jubbah Palaeolake, Northern Arabia. *PLoS ONE* **7**, e49840 (2012).
44. Galbraith, R. F., Roberts, R. G., Laslett, G. M., Yoshida, H., Olley, J. M. Optical dating of single and multiple grains of quartz from Jinmium rock shelter, northern Australia: Part I, experimental design and statistical models. *Archaeometry* **41**, 339-364 (1999).
45. Armitage, S. J., King, G. E. Optically stimulated luminescence dating of hearths from the Fazzan Basin, Libya: A tool for determining the timing and pattern of Holocene occupation of the Sahara. *Quat. Geochronol.* **15**, 88-97 (2013).
46. Tribolo, C., Mercier, N., Rasse, M., Soriano, S., Huysecom, E. Kobo 1 and L'Abri aux Vaches (Mali, West Africa): Two case studies for the optical dating of bioturbated sediments. *Quat. Geochronol.* **5**, 317-323 (2010).
47. Bøtter-Jensen, L., Mejdahl, V. Assessment of beta dose-rate using a GM multicounter system. *Int. J. Rad. Appl. Instrum. B.* **14**, 187-191 (1988).
48. De Corte, F. et al. Preparation and characterization of loess sediment for use as a reference material in the annual radiation dose determination for luminescence dating. *J. Radioanal. Nucl. Chem.* **272**, 311-319 (2007).
49. Hong, D. *Luminescence stimulated from quartz by green light: developments relevant to dating.* (PhD thesis, University of Edinburgh, 1988).

50. Mejdahl, V. Thermoluminescence dating: Beta-dose attenuation in quartz grains. *Archaeometry* **21**, 61-72 (1979).
51. Prescott, J. R., Hutton, J. T. Cosmic ray and gamma ray dosimetry for TL and ESR. *Int. J. Rad. Appl. Instrum.* **14**, 223-227 (1988).
52. Bennett K. D. Determination of the Number of Zones in a Biostratigraphical Sequence. *New Phytologist* **132**, 155-170 (1996).
53. Wagner B, *et al.* A 40,000-year record of environmental change from ancient Lake Ohrid (Albania and Macedonia). *J. Paleolimnology* **41**, 407-430 (2009).
54. Bennion, H., Simpson, G. The use of diatom records to establish reference conditions for UK lakes subject to eutrophication. *J. Paleolimnology* **45**, 1-20 (2011).
55. Rosenberg, T. M. *et al.* Middle and Late Pleistocene humid periods recorded in palaeolake deposits of the Nafud desert, Saudi Arabia. *Quatern. Sci. Rev.* **70**, 109-123 (2013).
56. Saros, J. E., Anderson, N. J. The ecology of the planktonic diatom *Cyclotella* and its implications for global environmental change studies. *Bio. Rev. Camb. Philos.* **90**, 522-541 (2015).
57. Wunsam, S., Schmidt, R., Klee, R. *Cyclotella*-taxa (Bacillariophyceae) in lakes of the Alpine region and their relationship to environmental variables. *Aquat. Sci.* **57**, 360-386 (1995).
58. Verschuren, D., Laird, K. R., Cumming, B. F. Rainfall and drought in equatorial east Africa during the past 1,100 years. *Nature* **403**, 410-414 (2000).
59. Wolin, J. A., Stone, J. R. in *The Diatoms Applications to the Environmental and Earth Sciences* (eds Stomer, F., Smol, J.P.) 174-185 (Cambridge University Press, Cambridge, 2010)

60. McGowan, S., Grauert, M., Anderson, N. J. A late Holocene record of landscape degradation from Heygsvatn, the Faroe Islands. *Palaeogeography Palaeoclimatology Palaeoecology* **264**(1-2) 11-24 (2008).
61. Phillips, G., Pietilainen, O. P., Carvalho, L., Solimini, A., Solheim, A. L., Cardoso, A. C Chlorophyll-nutrient relationships of different lake types using a large European dataset. *Aquatic Ecology* **42**(2) 213-226 (2008).
62. Alonso-Zarza, A. M. Palaeoenvironmental significance of palustrine carbonates and calcretes in the geological record. *Earth Sci. Rev.* **60**, 261-298 (2003).
63. Verrecchia, E. P. in *Geochemical Sediments and Landscapes* (eds Nash, D. J., McLaren, S. J.) 298-329 (Blackwell, Oxford, 2007).
64. Talbot, M. R. A review of the palaeohydrological interpretation of carbon and oxygen isotopic ratios in primary lacustrine carbonates. *Chem. Geol.* **80**, 261-279 (1990).
65. Li, H. C., Ku, T. L. $\delta^{13}\text{C}$ - $\delta^{18}\text{O}$ covariance as a paleohydrological indicator for closed-basin lakes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **133**, 69-80 (1997).
66. Leng, M.J. and Marshall, J.D. Palaeoclimate interpretation of stable isotope data from lake sediment archives. *Quatern. Sci. Rev.* **23**, 811-831 (2004).
67. Holmes, J.A., et al. Holocene palaeolimnology of Kajemamm Oasis, Northern Nigeria: an isotopic study of ostracodes, bulk carbonate and organic carbon. *J. Geol. Soc.*, **154**, 311-319 (1997).
68. Marshall, J.D., et al. A high resolution late glacial isotopic record from Hawes Water, Northwest England. Climatic oscillations: calibration and comparison of palaeotemperature proxies. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **185**, 25-40 (2002).
69. Rozanski, K., Araguas-Araguas, L., Gonfiantini, R. Isotopic Patterns in Modern Global Precipitation. In: Swart, P.K., Lohmann, K.C., McKenzie, J., Savin, S. (eds).

- Climate Change in Continental Isotopic Records*. Geophysical Monograph 78, American Geophysical Union, 1-36 (1993).
70. Parton, A., et al. Orbital-scale climate variability in Arabia as a potential motor for human dispersals. *Quatern. Int.* **382**, 82-97 (2015).
71. Fleitmann, D., Matter, A., Pint, J.J., Al-Shanti, M. The speleothem record of climate change in Saudi Arabia. Saudi Geological Survey report, SGS-OF-2004-8 (2004).
72. Eltringham, S. K. *The Hippos: Natural History and Conservation* (Academic Press, London, 1999)
73. Estres, R. D. *The Behaviour Guide to African Mammals*. (University of California Press, Berkeley, 1991).
74. Belmaker, M. In *Out of Africa I: The First Hominin Colonization of Eurasia* (eds. Fleagle, J. G., Shea, J. J., Grine, F. E., Baden, A. L., Leakey, R. E.) 183-205 (Springer, 2010).
75. Martínez-Navarro, B. in *Out of Africa I: The First Hominin Colonization of Eurasia* (eds. Fleagle, J. G., Shea, J. J., Grine, F. E., Baden, A. L. & Leakey, R. E) 207-224 (Springer, New York, 2010).
76. Martínez-Navarro, B., Rook, L., Segid, A., Yosief, D., Farretti, M. P., Shoshani, J., Techle, T. M., Libsekal, Y. The large fossil mammals from Buia (Eritrea): systematics, biochronology and paleoenvironments. *Riv. Ital. Paleontol. S.* **110**, 61-88 (2004).
77. Shreeve, D. C. A new record of Pleistocene hippopotamus from River Severn terrace deposits, Gloucester, UK-palaeoenvironmental setting and stratigraphical significance. *P. Geologist. Assoc.* **120**, 58-64 (2009).
78. van Kolfschoten, Th. The Eemian mammal fauna of central Europe. *Geol. Mijnbouw-N. J. G.* **79**, 269-281 (2000).

79. Tchernov, E. The Afro-Arabian Component in the Levantine Mammalian Fauna – A Short Biogeographical Review. *Isr. J. Zool.* **38**, 155-159 (1992).
80. Geraads, D. Plio-Pleistocene Mammalian Biostratigraphy of Atlantic Morocco. *Quaternaire.* **13**, 43-53 (2002).
81. Michel, P., Wengler, L. *Un Site Palaeontologique Avec Des Vestiges Archeologiques: La Carriere Doukkala II (Region De Temara – Maroc Atlantique).* **5**, 11-41 (1993).
82. Stimpson, C. et al. Middle Pleistocene vertebrate fossils from the Nefud Desert, Saudi Arabia: Implications for biogeography and palaeoecology. *Quatern. Sci. Rev.* **143**, 13-36 (2016).
83. McClure, H. A. *Late Quaternary Palaeoenvironments of the Rub' al Khali.* (PhD Thesis, Univeristy College, London, 1984).
84. O'Regan, H. J., Turner, A., Bishop, L. C., Elton, S., Lamb, A. L. Hominins without fellow travellers? First appearances and inferred dispersals of Afro-Eurasian large mammals in the Plio-Pleistocene. *Quatern. Sci. Rev.* **30**, 1343-1352 (2011).
85. Hadjouis, D. Présence du genera *Kobus* (Bovidae, Artiodactyla) dans le Pléistocène Supérieur D'Algérie. *L'Anthropologie.* **90**, 317-320 (1986).
86. Kowalski, K., Neer, W. V., Bockenski, Z., Mlynarsko, M., Rzabik-Kowalska, B., Szyndlar, Z., Gautier, A., Close, A. E., Wendorf, F. A Last Interglacial Fauna from the Eastern Sahara. *Quat. Res.* **32**, 335-341 (1989).
87. Blinkhorn, J., Achyuthan, H., Petraglia, M. D. Ostrich expansion into India during the Late Pleistocene: Implications for continental dispersal corridors. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **417**, 80-90 (2015).
88. Jain, S., Rai, N., Kumar, G., Pruthi, P. A., Thangaraj, K., Bajpai, S., Pruthi, V. Ancient DNA Reveals Late Pleistocene Existence of Ostriches in Indian Sub-Continent. *PLoS ONE.* **12**, e0164823 (2017).

89. Domínguez-Rodrigo, M., Egeland, C. P., Barba, R. in *Deconstructing Olduvai: A Taphonomic Study of the Bed I Sites* (eds. Domingues-Rodrigo, M., Barba, R. & Egeland, C. P.) 23-32 (Springer, New York, 2007)
90. Cruz-Urbe, K. Distinguishing hyena from hominid bone accumulations. *J. F. Archaeol.* **18**, 467–486 (1991).
91. Petraglia, M. D. et al. Middle Palaeolithic occupation on a Marine Isotope Stage 5 lakeshore in the Nefud Desert, Saudi Arabia. *Quatern. Sci. Rev.* **30**, 1555-1559 (2011).
92. Scerri, E. M. L. et al. Middle to Late Pleistocene human habitation in the western Nefud Desert. *Quatern. Int.* **382**, 200-214 (2015).
93. Groucutt, H. S. et al. Human occupation of the Arabian Empty Quarter during MIS 5: evidence from Mundafan al-Buhayrah. *Quatern. Sci. Rev.* **119**, 116-135 (2015).
94. Scerri, E. M. L., Drake, N. A., Jennings, R., Groucutt, H. S. Earliest evidence for the structure of *Homo sapiens* populations in Africa. *Quatern. Sci. Rev.* **101**, 207-2016 (2014).
95. Scerri, E. M. L., Groucutt, H. S., Jennings, R. P., Petraglia, M. D. Unexpected technological heterogeneity in northern Arabia indicates complex Late Pleistocene demography at the gateway to Asia. *J. Hum. Evol.* **75**, 125-142 (2014).
96. Scerri, E. M. L., Gravina, B., Blinkhorn, J., Delagnes, A. Can lithic attribute analyses identify discrete reduction trajectories? A quantitative study using refitted lithic sets. *J. Arch. Method Theory* **23**, 669-691.
97. Groucutt, H. S. et al. Late Pleistocene lakeshore settlement in Arabia: Middle Palaeolithic technology from Jebel Katefeh, Jubbah. *Quatern. Int.* **382**, 215-236.

98. Jennings, R. P., Parton, A., Clark-Balzan, L., White, T. S., Groucutt, H. S., Breeze, P., Parker, A. G., Drake, N. A., Petraglia, M. D. Human occupation of the northern Arabia interior during early Marine Isotope State 3. *J. Quat. Sci.* **31**, 953-966 (2017).
99. Groucutt, H. S. et al. Middle Palaeolithic raw material procurement and early stage reduction at Jubbah, Saudi Arabia. *Arch. Res. Asia.* **9**, 44-62 (2017).
100. Clarkson, C. An Index of Invasiveness for the measurement of unifacial and bifacial retouch: a theoretical, experimental, and archaeological verification. *J. Arch. Sci.* **29**, 65-75 (2002).
101. Kuhn, S. L., A geometric index of reduction for unifacial stone tools. *J. Arch. Sci.* **17**, 583-593 (1990).
102. Groucutt, H.S. Middle Palaeolithic point technology, with a focus on the site of Tor Faraj (Jordan, MIS 3). *Quatern. Int.* **350**, 205-226.
103. Hovers, E. *The Lithic Assemblages of Qafzeh Cave* (Oxford University Press, Oxford, 2009).
104. Marks, A. E. in *The Prehistory of Nubia, Vol. 1* (ed Wendorf, F.) 315-391 (SMU Press, Dallas, 1968).
105. Goder-Goldberger, M. The Khormusan: Evidence for an MSA East African industry in Nubia. *Quatern. Int.* **300**, 182-194 (2013).
106. Rose, J. I., Marks, A. E. "Out of Arabia" and the Middle-Upper Palaeolithic transition in the southern Levant. *Quartär* **61**, 49-85 (2014).
107. Wendorf, F., Schild, R., Close, A. E., and associates. *Egypt During the Last Interglacial: The Middle Paleolithic of Bir Tarfawi and Bir Sahara East.* (Plenum, New York, 1993).
108. Yellen, J. et al. The archaeology of Aduma Middle Stone Age sites in the Awash Valley, Ethiopia. *Paleoanthropology* **2005**, 25-100 (2005).

109. Shea, J. J. 2008. The Middle Stone Age archaeology of the Lower Omo Valley Kibish Formation: Excavations, lithic assemblages, and inferred patterns of early *Homo sapiens* behaviour. *J. Hum. Evol.* **55**, 448-485 (2008)
110. Groucutt, H. S. et al. Stone tool assemblages and models for the dispersal of *Homo sapiens* out of Africa. *Quatern. Int.* **382**, 8-30 (2015).
111. Horwitz, L.K., Smith, P., Faerman, M., Boaretto, E., Segal, I. The application of biometry LA-ICP-MS to provenance isolated bones: a study of hominin remains from Oumm-Qatafa Cave, Judean Desert. *Archaeol. Anthropol. Sci.* **3**, 245-62 (2011).
112. Walker, M. J., Ortega, J., López, M. V., Parmová, K. & Trinkaus, E. Neanderthal postcranial remains from the Sima de las Palomas del Cabezo Gordo, Murcia, Southeastern Spain. *Am. J. Phys. Anthropol.* **144**, 505-515 (2011).
113. Lorenzo, C, Arsuaga, J, Carretero, J.M. Hand and foot remains from the Gran Dolina Early Pleistocene site (Sierra de Atapuerca, Spain). *J. Hum. Evol.* **37**, 501-522 (1999).
114. Larson, S.G. et al. Descriptions of the upper limb skeleton of *Homo floresiensis*. *J. Hum. Evol.* **57**, 555-570 (2009).
115. Mersey, B., Jabbour, R.S., Brudvik, K., Defleur, A. Neandertal hand and foot remains from Moula-Guercy, Ardeche, France. *Am. J. Phys. Anthropol.* **152**, 516-529 (2013).
116. Semal, P., et al. New data on the late Neanderthals: direct dating of the Belgian Spy Fossils. *Am. J. Phys. Anthropol.* **138**, 421-428 (2009).
117. Trinkaus, E. *The Shanidar Neanderthals*. (Academic Press, New York, 1983).
118. Trinkaus E. *The Krapina Human Postcranial Remains: Morphology, Morphometrics and Paleopathology*. (University of Zagreb, Zagreb, 2016)

119. Sládek, V, Trinkaus, E, Hillson, S.W., Holliday, T.W. The people of the Pavlovian. Skeletal catalogue and osteometrics of the Gravettian fossil hominids from Dolni Vestonice and Pavlov. *Dolni Vestonice Studies, Svazek 5* (2000).
120. Trinkaus, E., Buzhilova, A.P., Mednikova, M.B., Dobrovolskaya, M.V. *The people of Sunghir: burials, bodies, and behavior in the earlier Upper Paleolithic*. (Oxford University Press, Oxford, 2014).
121. Shang, H., Trinkaus, E. *The early modern human from Tianyuan Cave, China* (Texas A & M University Press, 2010)
122. McCown, T.D., Keith, A. The Stone Age of Mount Carmel: The Fossil Human Remains from the Levallois-Mousterian, Vol. II. (Clarendon Press, Oxford, 1939).
123. Trinkaus, E., Svoboda, J.A., Wojtal, P., Nyvltova Fisakova, M., Wilczynski, J. Human remains from the Moravian Gravettian: morphology and taphonomy of additional elements from Dolni Vestonice II and Pavlov 1. *Int. J. Osteoarchaeol.* **20**, 645-669 (2010).
124. Trinkaus, E., Bailey, S.E., Zilhao, J. Upper Paleolithic human remains from the Gruta do Caldeirao, Tomar, Portugal. *Revista Portuguesa de Arquelologia* **4**, 5-17 (2001).

Supplementary Table 1. Calliper measurements of AW-1 phalanx following measurement scheme of Horwitz and colleagues¹¹¹, proximal radio-ulnar maximum breadth following¹¹².

Measurement	Measurement no. ¹¹¹	Value (mm)	Notes
Maximum length	1	32.25	
Inter-articular length	2	30.21	
Midshaft dorso-palmar breadth	3	6.25	Measured just distal to the pathology, which lies across midshaft (actual measurement location at <1 mm distal to true midshaft).
Midshaft radio-ulnar breadth	4	8.53	
Proximal joint surface dorso-palmar height	5	8.03	Proximal joint surface damaged, estimate.
Proximal joint surface radio-ulnar breadth	6	11.92	
Proximal radio-ulnar maximum breadth	n/a	14.98	
Distal joint surface dorso-palmar height	7	5.25	
Distal joint surface radio-ulnar breadth	8	8.72	Distal joint surface incomplete, estimate.
Midshaft circumference	9	14	

Supplementary Table 2. Comparative sample for linear metric analysis of Al Wusta-1 intermediate phalanx.

Taxon	Specimen	IP ray	total sample	Source of data
<i>A. afarensis</i>	AL 333x-18	IP2-4	n=5	measured directly from fossils by TLK
	AL 333-32	IP2-4		
	AL 333-46	IP2-4		
	AL 333-88	IP2-4		
	AL 333-149	IP2-4		
<i>A. africanus</i>	StW 331	IP2-4	n=1	measured directly from fossils by TLK
<i>A. sediba</i>	MH2	IP3, IP4	n=2	measured directly from fossils by TLK
Swartkrans (<i>A. robustus</i> /early <i>Homo</i>)	SKX 13476	IP2-4	n=6	measured directly from fossils by TLK
	SKX 5019	IP2-4		
	SKX 5021	IP2-4		
	SKX 9449	IP2-4		
	SKX 36712	IP2-4		
	SKX 35439	IP2-4		
<i>H. antecessor</i>	ATD6-28	IP3-4	n=2	published values from Lorenzo et al. ¹¹²
	ATD6-53	IP3-4		
<i>H. naledi</i>	Hand 1	IP2, IP3, IP4	n=5	measured directly from fossils by TLK
	UW 101-1646	IP3		
	UW 101-1647	IP4		
<i>H. floresiensis</i>	LB6/9	IP?	n=2	published values from Larson et al. ¹¹³
	LB1/48	IP?		
<i>H. neanderthalensis</i>	Amud 1	IP2, IP3?	min. n=12 individuals n=27 IPs	measured directly from fossils by TLK
	Kebara 2	IP2, IP3, IP4		
	Tabun 1	IP2, IP4?		
	Moula Guercy M-G1-154	IP2-4		
	Spy 430a	IP3		

	Spy 390a	IP3		
	Spy 484a	IP4		
	Shanidar 3	IP3, IP4		published values from Trinkaus ¹¹⁶
	Shanidar 4	IP2, IP3, IP4		
	Shanidar 5	IP3, IP4		
	Shanidar 6	IP3		
	Krapina 205.1	IP3-4		published values from Trinkaus ¹¹⁷
	Krapina 205.2	IP3-4		
	Krapina 205.3	IP3-4		
	Krapina 205.4	IP3-4		
	Krapina 205.5	IP3-4		
	Krapina 205.6	IP3-4		
	Krapina 205.7	IP3-4		
	Krapina 205.8	IP3-4		
	Krapina 205.10	IP2		
	Krapina 205.12	IP3-4		
	Krapina 205.13	IP3-4		
	Krapina 205.14	IP2		
	Krapina 205.15	IP2		
	Krapina 205.17	IP2		
	Krapina 205.18	IP2		
<hr/>				
early <i>H. sapiens</i>	Dolni Vestonice 3	IP2-3	n=17 individuals	published values from Sladek et al. ¹¹⁸
	Dolni Vestonice 13	IP2-3, IP4	n=33 IPs	
	Dolni Vestonice 14	IP3		
	Dolni Vestonice 15	IP2, IP3		
	Dolni Vestonice 16	IP2, IP3, IP4		
	Dolni Vestonice 34	IP2-3		
	Ohalo II H2	IP2, IP3, IP4		measured directly from fossils by TLK
	Qafzeh 9	IP2		measured directly from fossils by TLK
	Qafzeh 8	IP2, IP3, IP4		measured directly from fossils by TLK
	Barma Grande 2	IP3, IP4		measured directly from fossils by NBS

Arene Candide 2	IP2, IP3, IP4		measured directly from fossils by NBS
Sunghir 1	IP2, IP3, IP4, IP5		published values in Trinkaus et al. ¹¹⁹
Tianyuan 1	IP2?		published values in Shang and Trinkaus ¹²⁰
Skhul IV	IP3, IP4, IP5		published values in McCown and Keith ¹²¹
Pavlov 33	IP?		published values in Trinkaus et al. ¹²²
Caldeirao 9	IP2-4		published values in Trinkaus et al. ¹²³
Cueva Victoria CV-0*	IP2		published values in Martinez-Navarro et al. ¹
recent <i>H. sapiens</i>	IP2-4	n=22 individuals	measured directly from specimens by TLK
<i>Pan</i> (<i>P. troglodytes</i> & <i>P. paniscus</i>)	IP2-5	n=67 IPs	
		n=8 individuals	measured directly from specimens by TLK
		n=34 IPs	
<i>Gorilla</i> (<i>G. gorilla</i> & <i>G. beringei</i>)	IP2-5	n=8 individuals	measured directly from specimens by TLK
		n=34 IPs	
Cercopithecids	IP2-5	n=11 IPs	measured directly from specimens by TLK

*The taxonomic association of this specimen with *H. sapiens* is questionable^{1,2}

Supplementary Table 3. Results from Mann-Whitney U pairwise comparisons with Bonferroni correction ($\alpha = 0.003$) of all shape ratios (i.e. divided by the total length of the phalanx). All breadth measurements are radio-ulnar dimensions and all height measurements are dorsopalmar dimensions. All significant pairwise comparisons are in bold text.

			Cercopiths	Gorilla	Pan	Neandertal	early <i>H. sapiens</i>	<i>H. sapiens</i>
base breadth	above	Cercopiths	x	0.007	0.000	0.079	0.110	0.018
prox. articular breadth	below	Gorilla	0.001	x	0.000	0.000	0.133	0.869
		Pan	0.000	0.000	x	0.000	0.000	0.000
		Neandertal	0.196	0.000	0.000	x	0.000	0.000
		early <i>H. sapiens</i>	0.130	0.009	0.000	0.001	x	0.125
		<i>H. sapiens</i>	0.033	0.022	0.000	0.000	0.368	x
prox. shaft breadth	above	Cercopiths	x	0.001	0.000	0.000	0.000	0.000
midshaft breadth	below	Gorilla	0.526	x	0.000	0.180	0.000	0.481
		Pan	0.000	0.000	x	0.123	0.011	0.000
		Neandertal	0.886	0.418	0.000	x	0.011	0.404
		early <i>H. sapiens</i>	0.088	0.052	0.000	0.009	x	0.000
		<i>H. sapiens</i>	0.001	0.000	0.000	0.000	0.000	x
midshaft height	above	Cercopiths	x	0.000	0.000	0.866	0.014	0.000
distal shaft breadth	below	Gorilla	0.032	x	0.000	0.000	0.001	0.000
		Pan	0.000	0.000	x	0.000	0.000	0.000
		Neandertal	0.048	0.356	0.076	x	0.005	0.000
		early <i>H. sapiens</i>	0.000	0.000	0.002	0.003	x	0.359
		<i>H. sapiens</i>	0.005	0.625	0.000	0.481	0.000	x
distal shaft height	above	Cercopiths	x	0.000	0.000	0.004	0.000	0.020
prox. articular height	below	Gorilla	0.200	x	0.000	0.640	0.002	0.000
		Pan	0.000	0.000	x	0.087	0.010	0.000
		Neandertal	0.988	0.164	0.000	x	0.026	0.043
		early <i>H. sapiens</i>	0.318	0.885	0.000	0.120	x	0.000
		<i>H. sapiens</i>	0.217	0.674	0.000	0.083	0.876	x

Supplementary Table 4. Landmarks used in geometric morphometric analyses of AI Wusta-1 phalanx and the comparative samples.

Landmark	Description
1	Mid-point of distal articulation (distal view, dorsal up)
2	Furthest left point on distal head (dorsal view)
3	Furthest right point on distal head (dorsal view)
4	Furthest proximal point on midline of proximal articulation (proximal view, dorsal up)
5	Mid-point of ridge between two articulations on proximal face (proximal view, dorsal up)
6	Furthest left point on proximal base (dorsal view)
7	Furthest right point on proximal base (dorsal view)
8	Deepest point on left proximal articulation (proximal view, dorsal up)
9	Deepest point on right proximal articulation (proximal view, dorsal up)
10	Centre of trochlea
11	Place where triangular raised region merges with central ridge on proximal palmar surface (palmar view)
12	Most dorsal point on proximal base (dorsal view)
13	Most palmar point of left proximal articulation (proximal view, dorsal up)
14	Most palmar point of right proximal base (proximal view, dorsal up)
15	Furthest point left of the distal trochlea (palmar view)
16	Furthest left point of the proximal articular surface (proximal view, dorsal up)
17	Furthest right point of the proximal articular surface (proximal view, dorsal up)
18	Most palmar point at the mid-line of the proximal articular surface (proximal view, dorsal up)

Supplementary Table 5. Sample used in comparative primate geometric morphometric analyses.

Institutions: Duckworth: Duckworth Collection, University of Cambridge; NHM Vienna: Vienna Natural History Museum; Uni. of Florence: University of Florence; GAUG: Johann-Friedrich-Blumenbach-Institut für Zoologie und Anthropologie der Georg-August-Universität Göttingen; Uni. of Kent: University of Kent; Tel Aviv Uni.: Tel Aviv University; MPNBR: Museo Preistorico Nazionale dei Balzi Rossi, Italy; MAF: Museo Archeologico del Finale, Italy; NHM: Natural History Museum, London; MRAC: Musée Royal de l'Afrique Centrale, Tervuren; MPI-EVA: Max Planck Institute, Leipzig; Powell Cotton: Powell Cotton Collection.

Group	Sample number	Institution
Al Wusta	1	
<i>Colobus badius preussi</i> total	7	Powell Cotton
<i>Gorilla gorilla</i> total	20	Powell Cotton
<i>Mandrillus leucophaeus</i>	4	Powell Cotton
<i>Mandrillus sphinx</i>	2	Powell Cotton
<i>Mandrillus</i> total	6	
<i>Pan paniscus</i>	4	MRAC Powell Cotton, MPI- EVA
<i>Pan troglodytes</i>	4	
<i>Pan</i> total	8	
<i>Papio anubis neumanni</i> total	4	
Krapina	17	NESPOS
Regourdou	3	NESPOS
Kebara 2	3	Tel Aviv Uni.
Tabun C1	2	NHM
Neanderthal total	25	
Qafzeh	4	Tel Aviv Uni.
Ohalo	3	Tel Aviv Uni.
Barma Grande	2	MPNBR
Arene Candide	3	MAF
Early <i>H. sapiens</i> total	12	
Australian	3	Duckworth
Inuit	1	Duckworth
Kerma	3	Duckworth
Maiden Castle	3	Duckworth
Egyptian Nubian	19	NHM Vienna
Fuegian	3	Uni. of Florence
Siracusian	5	Uni. of Florence
German	13	GAUG
Canterbury	11	Uni. of Kent
Holocene <i>H. sapiens</i> total	61	

Supplementary Table 6. Procrustes distances from primate and hominin mean shapes to Al Wusta-1.

Mean Shape	Distance to AW-1
Holocene <i>H. sapiens</i>	0.080184
Early <i>H. sapiens</i>	0.084209
<i>Mandrillus</i>	0.093510
<i>Gorilla</i>	0.101452
<i>Papio</i>	0.106864
<i>H. neanderthalensis</i>	0.119294
<i>Pan</i>	0.119404
<i>Colobus</i>	0.150987

Supplementary Table 7. Details of the comparative sample of intermediate phalanges from Neanderthals and Pleistocene and Holocene *Homo sapiens* used for geometric morphometric analyses of known side and digit, and cross sectional geometric analyses.

Institutions: Duckworth: Duckworth Collection, University of Cambridge; NHM Vienna: Vienna Natural History Museum; Uni of Florence: University of Florence; GAUG: Johann-Friedrich-Blumenbach-Institut für Zoologie und Anthropologie der Georg-August-Universität Göttingen; Uni of Kent: University of Kent; Tel Aviv Uni: Tel Aviv University; MPNBR: Museo Preistorico Nazionale dei Balzi Rossi, Italy; MAF: Museo Archeologico del Finale, Italy; NHM: Natural History Museum, London.

Group	Specimen/Group	Institution	Lefts (n)			Rights (n)			Sample total
			2nd	3rd	4th	2nd	3rd	4th	
Holocene <i>H. sapiens</i>	Australian	Duckworth		1		1	1	2	5
Holocene <i>H. sapiens</i>	Kerma	Duckworth	1	1				1	3
Holocene <i>H. sapiens</i>	Inuit	Duckworth		1			1		2
Holocene <i>H. sapiens</i>	Maiden Castle	Duckworth		1		1		1	3
Holocene <i>H. sapiens</i>	Egyptian	NHM Vienna	2		2	5	6	4	19
	Nubian								
Holocene <i>H. sapiens</i>	Fuegian	Uni of Florence	1		1	1	1		4
Holocene <i>H. sapiens</i>	Siracusan	Uni of Florence	2	1	1	2	1	2	9
Holocene <i>H. sapiens</i>	German	GAUG	1	2	2	3	3	2	13
Holocene <i>H. sapiens</i>	Canterbury	Uni of Kent	1	1	2	2	3	2	11
Pleistocene <i>H. sapiens</i>	Ohalo 2	Tel Aviv Uni	1		1				2
Pleistocene <i>H. sapiens</i>	Qafzeh 8	Tel Aviv Uni	1	1	1	1	1	1	6
Pleistocene <i>H. sapiens</i>	Qafzeh 9	Tel Aviv Uni	1			1	1		3
Pleistocene <i>H. sapiens</i>	Barma Grande 2	MPNBR		1	1		1	1	4
Pleistocene <i>H. sapiens</i>	Arene Candide	MAF	1	1	1	1	1	1	6
	2								
Neanderthal	Kebara 2	Tel Aviv Uni	1	1	1	1		1	5
Neanderthal	Tabun C1	NHM	1		1				2
Total			14	12	14	19	20	18	91

Supplementary Table 8. Procrustes Distances between Al Wusta-1 and other hominin groups with sides pooled.

	Holocene <i>H. sapiens</i>	Pleistocene <i>H. sapiens</i>	Neanderthal
AW-1	0.078	0.084	0.099

Supplementary Table 9. Procrustes distances between Al Wusta-1 and its nearest neighbours with sides pooled. Specimen numbers refer to Figure 2.

Nearest neighbour	Side	Ray	Procrustes	
			Distance	
1) Maiden Castle EU.1.3.70_Sk20	Left	3 rd	0.084	
2) Egyptian NHMW K24_2	Right	3 rd	0.061	
3) Canterbury NGB_89_Sk15_1247	Left	3 rd	0.072	

Supplementary Table 10. Procrustes distances between Al Wusta-1 and comparative groups and ray numbers across groups, analysing left and right hands separately.

Phalanx side	Holocene <i>H.</i>	Pleistocene <i>H.</i>	
	<i>sapiens</i>	<i>sapiens</i>	Neanderthals
Right	0.080	0.090	0.119
Left	0.076	0.081	0.098
	2 nd Ray	3 rd Ray	4 th Ray
Right	0.095	0.076	0.083
Left	0.092	0.068	0.084

Supplementary Table 11. Procrustes distances between Al Wusta-1 and its nearest neighbours, analysing left and right hands separately.

Nearest neighbour	Side	Ray	Procrustes
			Distance
Egyptian NHMW K24 3 HP3MR	Right	3 rd	0.061
Canterbury NGB 89 Sk15 1247 HP3ML	Left	3 rd	0.072

Supplementary Table 12. U-series results on Al Wusta-1. All errors are 2- σ . Individual results do not contain errors of standard measurement (correlated errors), mean values incorporate errors of standard.

3675#1	U (ppm)	Th (ppb)	U/T h	²³⁰ Th/ ²³⁸ U U	²³⁰ Th/ ²³⁸ U error	²³⁴ U/ ²³⁸ U U	²³⁴ U/ ²³⁸ U error	Age (ka)	Age error (ka)
1	49.99	16	307	0.7293	0.0066	1.6887	0.0053	59.1	0.7
2	48.60	11	425	0.7310	0.0085	1.6817	0.0046	59.6	0.9
3	46.10	14	339	0.7537	0.0052	1.6798	0.0045	62.1	0.6
4	43.85	16	270	0.7724	0.0061	1.6899	0.0073	63.6	0.7
5	34.79	226	154	0.7744	0.0063	1.6801	0.0048	64.3	0.7
6	23.25	290	80	0.7950	0.0107	1.6829	0.0130	66.4	1.4
7	19.12	166	115	0.7851	0.0094	1.6861	0.0051	65.2	1.1
8	15.34	118	131	0.7791	0.0109	1.6895	0.0107	64.4	1.3
9	14.22	106	135	0.7522	0.0088	1.6807	0.0092	61.9	1.0
10	14.65	129	114	0.7574	0.0093	1.6882	0.0173	62.1	1.3
11	15.29	100	152	0.7622	0.0084	1.6794	0.0101	63.1	1.0
12	14.58	89	164	0.7559	0.0082	1.6895	0.0074	61.9	0.9
13	12.02	81	148	0.7474	0.0089	1.6828	0.0099	61.3	1.1
14	10.84	69	157	0.7516	0.0102	1.6798	0.0106	61.9	1.2
15	11.08	54	207	0.7598	0.0103	1.6808	0.0118	62.7	1.3
16	11.22	48	235	0.7585	0.0115	1.6772	0.0106	62.8	1.3
17	11.20	47	240	0.7558	0.0111	1.6871	0.0104	62.0	1.3
18	10.85	58	188	0.7606	0.0107	1.6933	0.0090	62.2	1.2
19	10.08	48	211	0.7558	0.0135	1.6884	0.0104	61.9	1.5
20	9.01	51	177	0.7572	0.0119	1.6952	0.0093	61.7	1.3
21	7.81	70	111	0.7489	0.0133	1.6825	0.0098	61.5	1.5
22	7.30	80	92	0.7311	0.0123	1.6858	0.0116	59.5	1.4
23	6.57	72	91	0.7275	0.0124	1.6769	0.0088	59.5	1.4
24	6.27	65	96	0.7251	0.0161	1.6833	0.0093	59.0	1.7
25	6.19	67	92	0.7361	0.0120	1.6666	0.0159	60.9	1.5
26	5.34	68	78	0.7338	0.0216	1.6691	0.0154	60.6	2.4
27	5.33	61	87	0.7405	0.0132	1.6833	0.0150	60.6	1.6
28	5.55	51	109	0.7351	0.0107	1.6971	0.0165	59.3	1.3
29	5.63	46	124	0.7527	0.0136	1.6880	0.0109	61.6	1.5
30	5.28	37	143	0.7333	0.0169	1.6821	0.0168	59.9	1.9
MEAN VALUES									
1-30				0.7547	0.0129	1.6843	0.0152	62.0	1.6

3675#2	U (ppm)	Th (ppb)	U/Th	²³⁰ Th/ ²³⁸ U U	²³⁰ Th/ ²³⁸ U error	²³⁴ U/ ²³⁸ U U	²³⁴ U/ ²³⁸ U error	Age (ka)	Age error (ka)
1	57.79	16	3687	0.5955	0.0200	1.6802	0.0107	46.3	1.9
2	57.29	9	6263	0.6742	0.0066	1.6675	0.0068	54.5	0.7
3	53.84	7	8207	0.7077	0.0114	1.6700	0.0061	57.8	1.2
4	47.64	5	9152	0.7705	0.0068	1.6852	0.0127	63.7	1.0
5	45.61	5	9414	0.7969	0.0088	1.6784	0.0077	66.9	1.1
6	42.56	4	1079	0.7910	0.0122	1.6714	0.0172	66.6	1.7
7	40.59	3	1267	0.7948	0.0084	1.6845	0.0094	66.3	1.1
8	38.07	2	1570	0.7810	0.0075	1.6844	0.0088	64.8	0.9
9	37.47	2	1760	0.7814	0.0125	1.6748	0.0196	65.4	1.7
10	37.71	2	1541	0.7671	0.0079	1.6749	0.0174	63.8	1.2
11	33.73	2	1364	0.7640	0.0067	1.6755	0.0094	63.5	0.9
12	30.04	3	1166	0.7767	0.0112	1.6806	0.0097	64.6	1.3
13	30.55	3	1109	0.7883	0.0082	1.6952	0.0125	65.1	1.1
14	29.63	3	1097	0.7787	0.0085	1.6844	0.0086	64.6	1.0
15	23.02	22	1060	0.7867	0.0099	1.6800	0.0076	65.7	1.2
16	17.81	33	540	0.8032	0.0098	1.6839	0.0093	67.3	1.2
17	14.64	22	660	0.8113	0.0091	1.6941	0.0118	67.6	1.2
18	12.94	30	425	0.7987	0.0076	1.6783	0.0112	67.1	1.0
19	10.54	39	272	0.7877	0.0106	1.6873	0.0097	65.4	1.3
20	8.24	48	171	0.7746	0.0125	1.6877	0.0069	64.0	1.4
21	6.20	61	102	0.7546	0.0143	1.6917	0.0102	61.6	1.6
22	5.00	31	160	0.7692	0.0162	1.6939	0.0166	63.1	1.9

23	4.40	17	256	0.7668	0.0163	1.6926	0.0146	62.9	1.9
24	3.90	12	319	0.7495	0.0165	1.6929	0.0151	61.0	1.9
25	3.57	11	329	0.7547	0.0183	1.6891	0.0145	61.8	2.1
26	2.97	10	296	0.7577	0.0177	1.6826	0.0146	62.4	2.0
27	2.63	9	282	0.7240	0.0209	1.6951	0.0197	58.3	2.3
28	2.34	8	299	0.6894	0.0234	1.6955	0.0151	54.8	2.4
29	2.16	7	300	0.6274	0.0259	1.6883	0.0174	49.0	2.6
30	1.85	6	331	0.6463	0.0358	1.6758	0.0322	51.3	3.7
MEAN VALUES									
5-26				0.7837	0.0135	1.6814	0.0154	65.3	1.7

3675#3	U	Th	U/Th	²³⁰ Th/ ²³⁸ U	²³⁰ Th/ ²³⁸ U	²³⁴ U/ ²³⁸ U	²³⁴ U/ ²³⁸ U	Age	Age error
	(ppm)	(ppb)		U	error	U	error	(ka)	(ka)
1	58.44	18	3325	0.6462	0.0124	1.6700	0.0073	51.6	1.3
2	56.93	8	7353	0.6569	0.0049	1.6696	0.0153	52.6	0.8
3	55.96	6	9673	0.6594	0.0066	1.6702	0.0096	52.9	0.8
4	47.14	5	9675	0.6976	0.0087	1.6696	0.0108	56.7	1.0
5	41.59	35	1201	0.7572	0.0132	1.6680	0.0139	63.1	1.6
6	37.40	11	3388	0.8065	0.0132	1.6908	0.0192	67.3	1.8
7	38.25	7	5596	0.7945	0.0094	1.6807	0.0074	66.5	1.1
8	38.32	5	7199	0.7931	0.0102	1.6773	0.0157	66.5	1.4
9	38.60	5	8198	0.8012	0.0106	1.6876	0.0156	66.9	1.4
10	37.61	2	1640	0.7850	0.0062	1.6900	0.0124	65.0	0.9
11	35.79	3	1383	0.7851	0.0086	1.6803	0.0087	65.5	1.0
12	35.59	2	1452	0.7800	0.0088	1.6783	0.0102	65.1	1.1
13	36.60	2	1522	0.7671	0.0057	1.6745	0.0088	63.8	0.8
14	32.65	2	1478	0.7566	0.0063	1.6792	0.0075	62.5	0.8
15	30.15	3	1155	0.7659	0.0103	1.6884	0.0125	63.0	1.3
16	27.97	3	1025	0.7703	0.0085	1.6835	0.0129	63.7	1.1
17	25.75	4	6415	0.7698	0.0120	1.6816	0.0171	63.8	1.6
18	24.69	3	8744	0.7781	0.0090	1.6691	0.0107	65.3	1.1
19	22.23	3	8589	0.7997	0.0127	1.6868	0.0087	66.8	1.5
20	19.33	3	6449	0.8015	0.0108	1.6820	0.0131	67.2	1.4
21	14.70	10	1482	0.7975	0.0083	1.6827	0.0059	66.7	1.0
22	8.99	19	471	0.7862	0.0162	1.6855	0.0115	65.3	1.9
23	5.23	11	497	0.7627	0.0172	1.6863	0.0158	62.8	2.0
24	5.05	6	827	0.7796	0.0105	1.7064	0.0110	63.5	1.2
25	5.18	8	667	0.7626	0.0118	1.6977	0.0141	62.2	1.4
26	5.43	8	653	0.7599	0.0196	1.6880	0.0132	62.4	2.2
27	5.14	5	1090	0.7318	0.0161	1.6956	0.0179	59.1	1.9
28	4.91	4	1297	0.7474	0.0144	1.6741	0.0193	61.7	1.8
29	4.70	3	1664	0.7497	0.0153	1.6738	0.0139	62.0	1.8
30	4.32	2	1859	0.7433	0.0144	1.6836	0.0160	60.8	1.7
MEAN VALUES									
5-30				0.7821	0.0135	1.6827	0.0154	65.0	1.7

3675#4	U	Th	U/Th	²³⁰ Th/ ²³⁸ U	²³⁰ Th/ ²³⁸ U	²³⁴ U/ ²³⁸ U	²³⁴ U/ ²³⁸ U	Age	Age error
	(ppm)	(ppb)		U	error	U	error	(ka)	(ka)
1	59.50	39	1524	0.6319	0.0102	1.6640	0.0054	50.4	1.0
2	61.70	14	4513	0.6370	0.0064	1.6770	0.0311	50.4	1.3
3	55.83	9	6523	0.6833	0.0087	1.6684	0.0049	55.3	0.9
4	46.15	6	8275	0.7477	0.0088	1.6755	0.0096	61.7	1.1
5	40.88	6	6811	0.8026	0.0100	1.6834	0.0081	67.3	1.2
6	37.30	5	6899	0.8437	0.0085	1.6773	0.0067	72.3	1.1
7	35.26	3	1049	0.8453	0.0054	1.6879	0.0048	71.8	0.7
8	36.35	3	1420	0.8305	0.0077	1.6833	0.0067	70.4	1.0
9	35.68	3	1253	0.8276	0.0079	1.6798	0.0080	70.3	1.0
10	33.76	2	1598	0.8366	0.0057	1.6846	0.0066	71.0	0.8
11	33.52	2	1678	0.8263	0.0069	1.6815	0.0104	70.0	1.0
12	32.93	2	1780	0.8117	0.0072	1.6814	0.0039	68.4	0.8
13	32.85	2	2134	0.8043	0.0070	1.6824	0.0086	67.5	0.9
14	31.36	2	1722	0.7961	0.0074	1.6893	0.0078	66.2	0.9

15	29.38	2	1603	0.7894	0.0074	1.6852	0.0093	65.7	1.0
16	28.26	1	1957	0.7801	0.0083	1.6719	0.0087	65.4	1.0
17	26.76	2	1245	0.7734	0.0080	1.6847	0.0066	64.0	0.9
18	24.71	2	1468	0.7655	0.0083	1.6735	0.0099	63.7	1.0
19	23.88	2	1269	0.7582	0.0097	1.6864	0.0156	62.3	1.3
20	22.21	7	3326	0.7556	0.0095	1.6836	0.0122	62.1	1.2
21	12.10	13	896	0.7489	0.0105	1.6782	0.0087	61.7	1.2
22	8.43	8	1063	0.7651	0.0101	1.6788	0.0119	63.4	1.2
23	7.16	5	1313	0.7567	0.0115	1.6846	0.0139	62.2	1.4
24	6.07	7	897	0.7675	0.0137	1.6901	0.0140	63.1	1.6
25	4.01	7	563	0.7555	0.0175	1.6847	0.0138	62.1	2.0
26	3.58	4	845	0.7533	0.0170	1.6764	0.0173	62.3	2.0
27	3.27	4	821	0.7348	0.0249	1.6763	0.0192	60.3	2.8
28	2.97	3	1072	0.7227	0.0224	1.6744	0.0174	59.1	2.5
29	2.59	2	1655	0.7398	0.0209	1.6773	0.0148	60.8	2.3
30	2.21	2	1242	0.7129	0.0255	1.6696	0.0139	58.3	2.7
MEAN VALUES									
5-30				0.8017	0.0137	1.6820	0.0152	67.2	1.7

3675#5	U	Th	U/Th	²³⁰ Th/ ²³⁸ U	²³⁰ Th/ ²³⁸ U	²³⁴ U/ ²³⁸ U	²³⁴ U/ ²³⁸ U	Age	Age error
	(ppm)	(ppb)		U	error	U	error	(ka)	(ka)
1	74.45	19	3962	0.5473	0.0042	1.6540	0.0049	42.7	0.4
2	68.19	8	8411	0.5574	0.0083	1.6487	0.0099	43.8	0.8
3	56.73	6	9114	0.6294	0.0134	1.6630	0.0063	50.2	1.3
4	47.30	6	7446	0.6985	0.0155	1.6621	0.0083	57.2	1.6
5	42.32	4	1050	0.7863	0.0120	1.6734	0.0149	66.0	1.6
6	38.37	3	1119	0.8324	0.0086	1.6852	0.0112	70.5	1.2
7	35.55	2	1465	0.8375	0.0101	1.6859	0.0067	71.0	1.2
8	33.52	2	1625	0.8548	0.0093	1.6904	0.0107	72.7	1.3
9	33.30	3	1090	0.8778	0.0072	1.6882	0.0107	75.5	1.1
10	32.28	2	1476	0.8746	0.0062	1.6817	0.0094	75.6	1.0
11	33.75	2	1990	0.8530	0.0068	1.6894	0.0165	72.6	1.3
12	32.58	2	2007	0.8570	0.0077	1.6872	0.0078	73.2	1.0
13	30.24	1	2053	0.8603	0.0106	1.6888	0.0099	73.5	1.4
14	27.27	2	1585	0.8684	0.0082	1.6860	0.0068	74.6	1.1
15	25.94	2	1567	0.8557	0.0082	1.6852	0.0112	73.2	1.2
16	24.93	1	2556	0.8451	0.0130	1.6847	0.0118	72.0	1.7
17	24.91	1	2502	0.8222	0.0100	1.6866	0.0063	69.3	1.2
18	23.05	1	3626	0.8358	0.0113	1.6888	0.0117	70.7	1.5
19	21.56	1	2173	0.8262	0.0080	1.6915	0.0091	69.4	1.0
20	20.31	1	1552	0.8322	0.0095	1.6917	0.0167	70.1	1.4
21	19.72	1	2411	0.8187	0.0146	1.7064	0.0278	67.7	2.2
22	19.69	1	2830	0.8080	0.0074	1.6861	0.0113	67.7	1.0
23	18.82	1	3367	0.8073	0.0079	1.6746	0.0132	68.3	1.2
24	18.26	1	1927	0.8028	0.0075	1.6792	0.0105	67.5	1.0
25	17.53	1	2329	0.8008	0.0102	1.6882	0.0064	66.8	1.2
26	16.43	1	1982	0.8079	0.0103	1.6917	0.0077	67.4	1.2
27	14.39	1	1501	0.8072	0.0094	1.6860	0.0132	67.6	1.3
28	10.81	1	8682	0.8119	0.0101	1.6874	0.0129	68.1	1.3
29	9.63	1	6644	0.8100	0.0107	1.6954	0.0113	67.4	1.3
30	9.14	1	8014	0.8081	0.0137	1.6825	0.0139	67.9	1.7
MEAN VALUES									
6-30				0.8392	0.0144	1.6874	0.0153	71.1	1.9

3675#6	U	Th	U/Th	²³⁰ Th/ ²³⁸ U	²³⁰ Th/ ²³⁸ U	²³⁴ U/ ²³⁸ U	²³⁴ U/ ²³⁸ U	Age	Age error
	(ppm)	(ppb)		U	error	U	error	(ka)	(ka)
1	76.83	11	7262	0.4753	0.0142	1.6530	0.0051	36.2	1.3
2	67.12	6	1086	0.5164	0.0068	1.6578	0.0123	39.7	0.7
3	63.03	7	9262	0.5360	0.0068	1.6572	0.0039	41.5	0.6
4	53.40	6	9030	0.6315	0.0155	1.6663	0.0084	50.3	1.6
5	47.14	4	1266	0.7257	0.0141	1.6735	0.0110	59.5	1.6
6	42.65	3	1293	0.7690	0.0090	1.6757	0.0050	64.0	1.0

7	40.02	3	1340	0.8297	0.0122	1.6803	0.0178	70.5	1.7
8	38.80	2	1670	0.8562	0.0078	1.6840	0.0100	73.3	1.1
9	36.41	2	2159	0.8708	0.0134	1.6868	0.0158	74.8	1.9
10	35.03	2	1790	0.8810	0.0087	1.6841	0.0106	76.2	1.2
11	34.68	2	1756	0.8966	0.0128	1.6925	0.0148	77.5	1.8
12	33.32	2	1818	0.9060	0.0089	1.6809	0.0091	79.4	1.3
13	30.97	2	1775	0.9079	0.0085	1.6863	0.0109	79.3	1.3
14	28.12	2	1582	0.8927	0.0089	1.6786	0.0083	78.0	1.2
15	25.78	2	1359	0.8953	0.0098	1.6778	0.0098	78.3	1.4
16	23.78	1	1965	0.8768	0.0109	1.6750	0.0130	76.3	1.6
17	21.87	1	2389	0.8852	0.0132	1.6860	0.0115	76.6	1.7
18	20.16	1	2144	0.8767	0.0125	1.6867	0.0105	75.5	1.6
19	19.88	1	2676	0.8574	0.0109	1.6881	0.0119	73.2	1.5
20	18.92	1	2179	0.8605	0.0097	1.6928	0.0105	73.2	1.3
21	17.80	1	1407	0.8454	0.0062	1.6846	0.0087	72.0	0.9
22	16.52	1	1433	0.8354	0.0126	1.6931	0.0158	70.4	1.7
23	16.42	1	2041	0.8389	0.0125	1.6902	0.0100	70.9	1.5
24	13.49	1	1810	0.8522	0.0131	1.6874	0.0123	72.6	1.7
25	12.28	1	1640	0.8742	0.0148	1.6841	0.0112	75.4	1.9
26	12.87	1	1602	0.8535	0.0161	1.7026	0.0164	71.9	2.1
27	11.94	1	8285	0.8409	0.0082	1.6995	0.0103	70.6	1.1
28	9.69	2	5389	0.8488	0.0108	1.6870	0.0103	72.2	1.4
29	8.40	2	4912	0.8500	0.0137	1.6845	0.0140	72.5	1.8
30	7.84	1	7437	0.8594	0.0136	1.6788	0.0122	74.0	1.8
MEAN VALUES									
5-30				0.8739	0.0150	1.6859	0.0154	75.2	2.0

3675#7	U	Th	U/Th	²³⁰ Th/ ²³⁸ U	²³⁰ Th/ ²³⁸ U	²³⁴ U/ ²³⁸ U	²³⁴ U/ ²³⁸ U	Age	Age error
	(ppm)	(ppb)		U	error	U	error	(ka)	(ka)
1	121.03	24	5024	0.3769	0.0033	1.6413	0.0025	28.0	0.3
2	115.97	15	7838	0.3748	0.0040	1.6423	0.0033	27.8	0.3
3	108.17	10	1073	0.4193	0.0081	1.6487	0.0030	31.4	0.7
4	86.57	9	9184	0.4735	0.0067	1.6484	0.0031	36.1	0.6
5	71.74	10	6855	0.5251	0.0146	1.6540	0.0038	40.6	1.3
6	57.44	13	4434	0.6336	0.0108	1.6660	0.0045	50.5	1.1
7	50.64	91	554	0.6722	0.0060	1.6690	0.0056	54.2	0.6
8	47.15	517	91	0.6413	0.0102	1.6627	0.0040	51.4	1.0
9	43.06	307	140	0.6922	0.0108	1.6750	0.0103	55.9	1.2
10	39.06	133	295	0.7215	0.0081	1.6787	0.0107	58.8	1.0
11	36.80	67	552	0.7133	0.0051	1.6666	0.0068	58.5	0.6
12	30.29	51	594	0.8452	0.0279	1.6882	0.0095	71.8	3.2
13	28.00	31	895	0.9252	0.0086	1.6917	0.0068	81.0	1.2
14	26.21	23	1162	0.9647	0.0104	1.7002	0.0094	85.3	1.5
15	23.79	17	1392	0.9813	0.0111	1.6944	0.0076	87.9	1.6
16	22.45	19	1199	0.9855	0.0112	1.6979	0.0067	88.1	1.5
17	21.92	22	982	0.9827	0.0079	1.6945	0.0066	88.0	1.2
18	21.01	17	1246	0.9752	0.0084	1.6878	0.0098	87.6	1.3
19	19.25	22	866	0.9940	0.0088	1.6967	0.0067	89.3	1.3
20	18.50	43	426	0.9641	0.0113	1.6841	0.0127	86.5	1.8
21	17.60	22	801	0.9799	0.0115	1.6883	0.0095	88.2	1.7
22	17.30	30	573	0.9743	0.0093	1.6958	0.0069	86.9	1.3
23	17.07	23	738	0.9737	0.0103	1.6907	0.0068	87.2	1.4
24	16.51	22	765	0.9703	0.0129	1.6869	0.0082	87.0	1.8
25	15.15	14	1054	0.9627	0.0098	1.6961	0.0058	85.4	1.3
26	13.61	8	1714	0.9810	0.0110	1.6945	0.0084	87.8	1.6
27	12.93	10	1267	0.9869	0.0123	1.6882	0.0095	89.1	1.8
28	11.39	19	614	0.9928	0.0109	1.6940	0.0073	89.4	1.5
29	7.08	76	93	0.9768	0.0133	1.6930	0.0112	87.4	1.9
30	5.92	53	112	0.9350	0.0144	1.6934	0.0148	82.1	2.1
MEAN VALUES									
15-30				0.9778	0.0168	1.6923	0.0153	87.6	2.5

Supplementary Table 13. U-series results on sample WU1601. All errors are 2- σ .

3672	U (ppm)	Th (ppb)	U/Th	²³⁰ Th/ ²³⁸ U	²³⁰ Th/ ²³⁸ U error	²³⁴ U/ ²³⁸ U	²³⁴ U/ ²³⁸ U error	Age (ka)	Age error (ka)
MEAN VALUES									
Enamel	0.83±0.28			0.8346	0.0533	1.5042	0.0329	83.5	8.1
Dentine	109±6			0.7295	0.0173	1.5742	0.0124	65.0	2.1

Supplementary Table 14. Combined US-ESR age results obtained for sample WU1601.

See Methods and Supplementary Information 2 for discussion.

Internal dose rate (μ Gy/a)	Beta dose rate (dentine) (μ Gy/a)	Beta dose rate sediment (μ Gy/a)	Gamma dose rate (μ Gy/a)	Cosmic dose rate (μ Gy/a)	Total dose rate (μ Gy/a)	Enamel thickness (μ m)	Removed surface layer (μ m) on each side	D _E (Gy)	p-parameter		US-ESR age (ka)
									enamel	dentine	
241±75	260±48	4±0	180±10	254±25	939±93	3244±100	Ext.: 176±50; Int.:181±50	97.1±1.9	-0.83	-0.53	103±10/-9

Supplementary Table 15. The number of single grains which were measured, rejected after application of the criteria outlined in Supplementary section 3.2, and accepted for inclusion in the calculation of D_b . Samples indicated (DR) represent dose recovery data.

Sample	PD15 (DR)	PD17 (DR)	PD15	PD17	PD40	PD41
<hr/> Total number of grains measured <hr/>						
	2500	2500	3600	3400	2800	3100
<hr/> Grains rejected for the following reasons <hr/>						
T_n signal	2358	2363	3373	3223	2606	2951
<hr/> <3*background <hr/>						
Poor recycling ratio	11	25	34	21	16	30
Depletion by IR	51	48	79	50	73	55
Recuperation	36	29	58	53	28	17
Oversaturation	3	1	0	5	12	7
$D_e < 2\sigma$ above 0 Gy	0	0	6	6	2	5
<hr/> Sum of rejected grains <hr/>						
	2459	2466	3550	3358	2737	3065
<hr/> Acceptable individual D_e values <hr/>						
	41	34	50	42	63	35

Supplementary Table 16. Sample depths, water content and dose rate for Al Wusta samples.

Sample	Depth (m)	Moisture (%)	Dose rate (Gy/ka)			Total dose rate, (Gy/ka)
			Beta	Gamma	Cosmic	
PD15	0.95±0.10	5±2.5	0.15±0.01	0.27±0.02	0.209±0.021	0.62±0.03
PD17	0.70±0.10	5±2.5	0.16±0.01	0.24±0.01	0.222±0.022	0.62±0.03
PD40	0.65±0.10	5±2.5	0.10±0.01	0.18±0.01	0.219±0.022	0.50±0.03
PD41	1.20±0.10	5±2.5	0.10±0.01	0.25±0.01	0.198±0.020	0.54±0.03

Supplementary Table 17. OSL Summary dating results and ages. Uncertainties in the age estimates are based on the propagation, in quadrature, of errors associated with individual errors for all measured quantities. In addition to uncertainties calculated from counting statistics, errors due to (1) beta source calibration (3 %); (2) single-grain instrument reproducibility (1.5 %); (3) dose rate conversion factors (3 %) and attenuation factors (3 %) have been included.

Sample	D_b calculation method (n)	D_b (Gy)	OD (%)	Total dose rate, D_r (Gy/ka)	Age (ka)
PD15	CAM (50)	57.4±3.0	22±4	0.62±0.03	92.0±6.3
PD17	CAM (42)	53.0±2.5	16±3	0.62±0.03	85.3±5.6
PD40	CAM (63)	49.6±2.5	26±3	0.50±0.03	98.6±7.0
PD41	CAM (35)	50.2±2.8	20±4	0.54±0.03	92.2±6.8

Supplementary Table 18. Quantification of bulk mineralogy from XRD data. Note that in all cases calcite makes up at least 85% of the crystalline mineral fraction. Gypsum and halite are below the detection limit of the instrumentation except in PD15-30. The absence of any saline tolerant diatom species at this level makes it unlikely that halite was precipitating in this system and it is more likely that this reflects aeolian dust blown onto the exposure. There is no evidence for Halite in any of the thin sections.



Sample ID	Calcite %	Quartz %	Gypsum %	Halite %
PD15_10	93.9	3.5	1.6	1
PD15_30	92.4	2	0.8	4.7
PD15_40	96.7	1.7	0.7	0.9
PD15_50	97.4	1.8	0.7	0.2
PD16_30	95.9	2.2	1.5	0.3
PD16_40	96.6	1.7	1.4	0.3
PD16_80	97.3	1.4	1.1	0.2
PD17_20	93.8	3.3	1.3	1.6
PD17_30	96.7	1.8	0.9	0.6
PD17_40	87.3	11	0.7	1

Supplementary Table 19. Taxonomy of Al Wusta vertebrate fossils.



Class	Order	Family	Taxon	Common name	
Reptilia	Squamata	Varanidae	<i>Varanus</i>	Monitor lizard	
Aves			Aves gen. et sp. indet. (small)	Small bird	
			Aves gen et sp. indet (medium)	Medium bird	
	Struthioniformes	Struthionidae	<i>Struthio</i> sp.	Ostrich	
Mammalia	Primates	Hominidae	<i>Homo sapiens</i>	Human	
	Rodentia		Myomorpha gen. et sp. indet.	Rodent	
	Artiodactyla	Hippopotamidae	<i>Hippopotamus</i> sp.	Hippopotamus	
			Bovidae	cf. <i>Pelorovis</i>	Pelorovis
				cf. <i>Kobus</i> sp.	Kobus
				Large bovid gen. et sp. indet.	Large bovid
				Medium bovid gen. et sp. indet.	Medium bovid
Small bovid gen. et sp. indet.	Small bovid				

Supplementary Table 20. Number of Identified Specimens for small, medium and large-sized fossil vertebrates from Al Wusta.

Element	Small	Medium	Large
Horn core	2	1	-
Crania	2	-	-
Mandible	2	4	-
Isolated teeth	5	10	2
Vertebrate	2	7	4
Ribs	7	12	4
Scapula	-	1	-
Pelvis	-	2	1
Humerus	2	3	2
Femur	-	-	-
Radius	-	2	-
Tibia	2	-	-
Carpals/Tarsals	2	4	1
Metapodials	1	-	2
Phalanges	1	2	3
Long bone shaft	11	46	7





| 263




- 1. AW-1 geometric morphometric analysis (see main text for methods and summary results)**


Supplementary Figure 1. Box-and-whisker plots of intermediate phalanx shape ratios of Al Wusta 1 and a sample of primates, including hominins. Al Wusta-1 is highlighted in red.




Supplementary Figure 2. Landmarks and wireframes used in geomorphometric analyses. A: Landmarks used to analyse phalanx shape using GMM. Landmarks numbered as in Supplementary Table 4 (AW-1 shown); B: Wireframes composed of straight lines connecting landmarks shown in A. Dorsal (left), lateral (middle) and proximal (right) views. Dotted lines connect landmarks not visible when bone is present, some lines omitted in proximal view for ease of visualisation.




Supplementary Figure 3. Scatterplot of the first two principal component (PC) scores of the geometric morphometric analysis of the Al Wusta-1 phalanx and a sample of primates, including hominins. Non-human primates in red; *Colobus*: triangles, *Pan*: stars, *Mandrillus*: squares, *Gorilla*: circles, *Papio*: diamonds. Hominins in blue; Neanderthals: diamonds, Holocene *H. sapiens*: squares, early *H. sapiens*: triangles. AW-1 in black. Data presented are the same as Figure 3, but filled convex hulls (for visualisation of data spread only) show hominins (blue) and non-human primates (red) generic groups of non-human primates and different groups of hominins. Circled shapes show means for groups (see Supplementary Table 6).




Supplementary Figure 4. Scatterplots of PC1 and PC2 scores from GMM analyses of left (top) and right (bottom) intermediate phalanges from a sample of Neanderthals and modern humans, and Al Wusta-1. Wireframes (see Supplementary Figure 2) show configurations at extremes of PC axes in dorsal (left), proximal (middle) and sagittal (right) views. Convex hulls added post-hoc for ease of visualisation.




Supplementary Figure 5. Scatterplots of cross-sectional geometric properties of Al Wusta-1 and comparative modern human and Neanderthal intermediate manual phalanges from rays 2-4. Plots against bone length: A = cortical area; B = total area; C = percent cortical area; D = J . Plot against I_{min} : E = I_{max} .




Supplementary Figure 6. AW-1 phalanx laser ablation sampling and data streams. A: Image of AW-1 (dating code 3675) with location of the laser ablation analysis holes; B: Data streams for ^{238}U , ^{234}U , ^{230}Th and ^{232}Th ; C: $^{230}\text{Th}/^{234}\text{U}$ and $^{234}\text{U}/^{238}\text{U}$ activity ratios and calculated closed system age estimates; D: Average age estimates for 30 bins of 33 cycles each; E: Plot of apparent age vs elemental U/Th.




Supplementary Figure 7. Uranium concentrations and age estimates. A: Plot of all average age estimates; B: U-concentrations (these are not corrected for diminishing U-yields from deeper domains in the hole). Note the different aspect compared to A; C: Age vs U-concentration for outside (bin 1); D: Age vs U-concentration for the age plateaux (bins 16-30).




Supplementary Figure 8. Image of sample WU1601 with location of the laser ablation analysis holes.




Supplementary Figure 9. ESR Dose response curve (DRC) obtained for WU1601. Final D_E values were calculated for each sample by pooling all the ESR intensities derived from the three repeated measurements in a single DRC. Fitting was performed with a SSE function and data weighting by $1/I^2$.




Supplementary Figure 10. OSL radial plots. Radial plots of the single-grain dose distributions for Al Wusta samples. For panels a and b the grey band is centred on the dose administered (49.8 Gy) in the dose recovery experiment. In panels c to f, the grey bar is centred on D_b determined with the Central Age Model. All points that lie within the grey band are consistent (at 2 standard errors) with either the administered dose (a,b) or D_b (c-f), and are shown as closed symbols. Open symbols denote equivalent doses which are greater than 2 standard errors from the administered dose (a,b) or D_b (c-f).




Supplementary Figure 11. Modelled ages for stratigraphic Units 1 (aeolian sand underlying the site) and 3 (waterlain sands and associated fossils overlying lacustrine marls). Two sequential phases were defined. Sample codes end with the age determination method in parentheses. The ages WU1601 (US-dentine), WU1601 (US-enamel) and AW-1 (US) are minimum age estimates, and the age model accounts for the fact that U-series ages are conventionally reported with 2 s uncertainties whereas OSL and ESR ages are reported with 1 s uncertainties. *A posteriori* densities are shown in darker shade while the likelihoods are shown in a lighter shade. Open circles underneath the *a posteriori* densities represent the mean age estimate, with 1 σ uncertainty bars, while the lower bar represents the 95.4 % range.





Supplementary Figure 12. Overview of the available age constraints for AW-1. In red, the modelled ages of the phase boundaries



Supplementary Figure 13. Stratigraphic sections from Al Wusta. A: stratigraphic logs of lacustrine marl deposits showing the main sedimentary units and stratigraphic variations in O and C isotopes and % CaCO₃. To avoid wasted space at the base of each stratigraphic section, OSL ages are not presented at their true depth below the sand-marl interface. B: Diagram summarising the difference in isotope values derived from aliquots of the same sample using different preparation techniques (dashed line = carbon isotope values, solid line = oxygen isotope values), although sieving of samples is often desirable the indurated nature of many samples makes such an approach impractical.




Supplementary Figure 14. Composite stratigraphic diagram showing the relationship between different sections across the site.




Supplementary Figure 15. XRD traces from PD15-50 (top) and PD17-40 (bottom).


Calcite peaks (orange) dominate with secondary quartz peaks (blue).




Supplementary Figure 16. Representative photo-micrographs of Al Wusta lake sediments. A) Finely laminated microsparite with silt-sized quartz grains (Allogenic). B) Massive microsparite with sand sized quartz grains (allogenic). C) freshwater sponge spicules in a microsparite/sparite massive matrix.




Supplementary Figure 17. Al Wusta carbonate isotopic data shown by sampling location (a) and as a single dataset (b). The overall r^2 value is 0.0521, while for each site the value is PD15 = 0.121, PD17 = 0.001, PD16 = 0.132, PD40 = 0.072. The three outlying samples are not shown but are included in the r^2 calculation.




Supplementary Figure 18. Summary diagram of Al Wusta Pit 1 PD 15 diatomite diatom assemblage. All taxa with relative abundances of $\geq 3\%$. The diatoms are ordered according to their weighted averaging distribution and divided up into assemblage zones derived from the optimal-sum-of squares partitioning using the program ZONE⁵². The statistically significant zones were deduced by comparison with the Broken-stick model using the program BSTICK version 1⁵². The planktonic: benthic ratio is shown with the habitat summary, Correspondence Analysis and Detrended Canonical Correspondence Analyses axis 1 sample scores (abbreviated to CA and DCCA respectively), log concentration, F-index which ranges from 0 (most dissolved) to 1 (most pristine).



Supplementary Figure 19. Summary diagram of Al Wusta Pit 2 PD 16 diatomite diatom assemblage. All taxa with relative abundances of $\geq 3\%$. The diatoms are ordered according to their weighted averaging distribution and divided up into assemblages zones derived from the optimal-sum-of squares partitioning using the program ZONE⁵². The statistically significant zones were deduced by comparison with the Broken-stick model using the program BSTICK version 1⁵². The planktonic: benthic ratio is shown with the habitat summary, Principal Component Analysis and Detrended Canonical Correspondence Analyses axis 1 sample scores (abbreviated to PCA and DCCA respectively), log concentration, F-index which ranges from 0 (most dissolved) to 1 (most pristine).



Supplementary Figure 20. Al Wusta vertebrate palaeontology. A: *cf. Kobus* sp. lower right M2-3; B: *Pelorovis* sp. mandibular fragment; C: *Hippopotamus* sp. ?upper canine fragment; D: *Hippopotamus* sp. medial upper incisor; E: *Varanus* sp. isolated tooth; F: Aves gen. et sp. indet. coracoid fragment; G: NISP for identified taxa. NISP is reported in parentheses; H: Evidence of carnivore gnawing on a bone fragment. Black scale bar = 10mm. White scale bar = 15mm.



Supplementary Figure 21. Selected Al Wusta lithic artefacts. A,B,D: Preferential Levallois cores with centripetal preparation, C: recurrent centripetal Levallois core, E,G-K: Levallois flakes, F: broken Levallois point, L: double side retouched flake, M: end retouched flake, N: side retouched flake.