

Molecular Brain

Naked mole-rat cortical neurons are resistant to acid-induced cell death

--Manuscript Draft--

Manuscript Number:	MBRJ-D-18-00046R2	
Full Title:	Naked mole-rat cortical neurons are resistant to acid-induced cell death	
Article Type:	Research	
Funding Information:	European Molecular Biology Organization (ALTF1565-2015)	Dr Zoé Husson
Abstract:	<p>Regulation of brain pH is a critical homeostatic process and changes in brain pH modulate various ion channels and receptors and thus neuronal excitability. Tissue acidosis, resulting from hypoxia or hypercapnia, can activate various proteins and ion channels, among which acid-sensing ion channels (ASICs) a family of primarily Na⁺ permeable ion channels, which alongside classical excitotoxicity causes neuronal death. Naked mole-rats (NMRs, <i>Heterocephalus glaber</i>) are long-lived, fossorial, eusocial rodents that display remarkable behavioral/cellular hypoxia and hypercapnia resistance. In the central nervous system, ASIC subunit expression is similar between mouse and NMR with the exception of much lower expression of ASIC4 throughout the NMR brain. However, ASIC function and neuronal sensitivity to sustained acidosis has not been examined in the NMR brain. Here, we show with whole-cell patch-clamp electrophysiology of cultured NMR and mouse cortical and hippocampal neurons that NMR neurons have smaller voltage-gated Na⁺ channel currents and more hyperpolarized resting membrane potentials. We further demonstrate that acid-mediated currents in NMR neurons are of smaller magnitude than in mouse, and that all currents in both species are fully blocked by the ASIC antagonist benzamil. We further demonstrate that NMR neurons show greater resistance to acid-induced cell death than mouse neurons. In summary, NMR neurons show significant cellular resistance to acidotoxicity compared to mouse neurons, contributing factors likely to be smaller ASIC-mediated currents and reduced NaV activity.</p>	
Corresponding Author:	Ewan St. John Smith, MPharmacol, PhD University of Cambridge Cambridge, Cambridgeshire UNITED KINGDOM	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	University of Cambridge	
Corresponding Author's Secondary Institution:		
First Author:	Zoé Husson, PhD	
First Author Secondary Information:		
Order of Authors:	Zoé Husson, PhD Ewan St. John Smith, MPharmacol, PhD	
Order of Authors Secondary Information:		
Response to Reviewers:	<p>Reviewer #1: Thanks for the response to reviewers.</p> <p>Minor comments:</p> <p>1, In the response to Reviewer 1 question 1, you mentioned that "we have however expanded the discuss on page 12...". Should be in page 13.</p> <p>That is simply a typo, nothing for us to address.</p> <p>2, question 6, I could not find reference 51 in the pubmed</p> <p>We can find the reference on PubMed, PMID 27964758 and so are not sure what the</p>	

	Reviewer means, although we appreciate that the formatting of online only journals for citations can be complicated and await advice from the copy editors.
Additional Information:	
Question	Response
Is this study a clinical trial?<hr><i>A clinical trial is defined by the World Health Organisation as 'any research study that prospectively assigns human participants or groups of humans to one or more health-related interventions to evaluate the effects on health outcomes'.</i>	No
Are you submitting to a Thematic Series?	No

[Click here to view linked References](#)1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 **Title:** Naked mole-rat cortical neurons are resistant to acid-induced cell
2 death

3 **Authors:** Zoé Husson¹, Ewan St John Smith¹

4 ¹ Department of Pharmacology, University of Cambridge, Tennis Court Road CB2 1PD, United
5 Kingdom

6 Corresponding author: Ewan St. John Smith, Department of Pharmacology, University of
7 Cambridge, Tennis Court Road, Cambridge, CB2 1PD, United Kingdom, Tel.: +44 1223 334048;
8 Fax: +44 1223 334100; E-mail: es336@cam.ac.uk

9
10 Co-author contact details: Zoé Husson, Department of Pharmacology, University of Cambridge,
11 Tennis Court Road, Cambridge, CB2 1PD, United Kingdom, Tel.: +44 1223 334006; Fax: +44
12 1223 334100; E-mail: zmah2@cam.ac.uk

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1

2 **Abstract**

3 Regulation of brain pH is a critical homeostatic process and changes in brain pH modulate various
4 ion channels and receptors and thus neuronal excitability. Tissue acidosis, resulting from hypoxia
5 or hypercapnia, can activate various proteins and ion channels, among which acid-sensing ion
6 channels (ASICs) a family of primarily Na⁺ permeable ion channels, which alongside classical
7 excitotoxicity causes neuronal death. Naked mole-rats (NMRs, *Heterocephalus glaber*) are long-
8 lived, fossorial, eusocial rodents that display remarkable behavioral/cellular hypoxia and
9 hypercapnia resistance. In the central nervous system, ASIC subunit expression is similar
10 between mouse and NMR with the exception of much lower expression of ASIC4 throughout the
11 NMR brain. However, ASIC function and neuronal sensitivity to sustained acidosis has not been
12 examined in the NMR brain. Here, we show with whole-cell patch-clamp electrophysiology of
13 cultured NMR and mouse cortical and hippocampal neurons that NMR neurons have smaller
14 voltage-gated Na⁺ channel currents and more hyperpolarized resting membrane potentials. We
15 further demonstrate that acid-mediated currents in NMR neurons are of smaller magnitude than
16 in mouse, and that all currents in both species are fully blocked by the ASIC antagonist benzamil.
17 We further demonstrate that NMR neurons show greater resistance to acid-induced cell death
18 than mouse neurons. In summary, NMR neurons show significant cellular resistance to
19 acidotoxicity compared to mouse neurons, contributing factors likely to be smaller ASIC-mediated
20 currents and reduced NaV activity.

21

22 **Key words**

23 ASIC, acid-induced currents, acidotoxicity, naked mole-rat, hippocampus

24

25

Background

Acid-sensing channels (ASICs) are ion channels of the ENaC/Deg superfamily and most subunits are activated by extracellular protons [1,2]. Six different ASIC subunits are encoded by 4 ASIC genes (ASIC1a, ASIC1b, ASIC2a, ASIC2b, ASIC3 and ASIC4), which assemble as homo- or heterotrimers [3]; neither ASIC2b nor ASIC4 form proton-sensitive homotrimers. ASICs are primarily permeable to Na⁺, although ASIC1a homomeric channels are also Ca²⁺ permeable [2]. In the central nervous system (CNS), neurons have been shown to primarily express ASIC1a homomers and heteromers of ASIC1a/2a and ASIC1a/2b [4–8], where they have been demonstrated to have key roles in synaptic plasticity [9–11] and fear conditioning [12–14], as well as being major players in neuronal death resulting from brain ischemia [5,15–17], and neurodegenerative diseases [18–20].

Regulation of brain pH is a highly complex and important process [21]. Brain tissue acidosis can result either from an increase in tissue partial pressure of carbon dioxide (PCO₂) during hypercapnia, or from the accumulation of the byproducts of anaerobic metabolism, such as lactate and protons, during hypoxia [22]. During periods of tissue acidosis, activation of ASICs by extracellular acidification is worsened by the release of allosteric modulators such as lactate [23], spermine [16] and arachidonic acid [24,25]. In addition to the activation of the Ca²⁺ permeable ASIC1a channel [15,16], a drop in pH also modulates the activity of numerous others ion channels, including voltage-gated ion channels [26–29] and glutamate receptors [30,31], therefore leading to disturbance in ion homeostasis, excitotoxicity and ultimately neuronal death [5,15,16].

Naked mole-rats (NMRs, *Heterocephalus glaber*) are subterranean rodents belonging to the Bathyergidae African mole-rat family found in East Africa [32]. Unusually for a mammal, NMRs are eusocial [33,34]. However, NMRs also display a range of remarkable physiological

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 peculiarities, which is beginning to make a significant impact on biomedical research [35]. The
2 unusual physiology of the NMR includes: extreme longevity with no increased risk of death with
3 ageing [36], an apparent absence of age-related neurodegenerative disorders [37,38], resistance
4 to cancer [39–41], insensitivity to certain noxious and irritant stimuli [42–45] and
5 hypoxia/hypercapnia resistance resulting from altered NMDA receptor function and an ability to
6 utilize fructose as an energy source [46–49]. It is striking that NMRs are resistant to many
7 pathological conditions known to involve ASICs. Recordings of ASIC-mediated currents in dorsal
8 root ganglion (DRG) sensory neurons demonstrated an increased frequency and magnitude of
9 ASIC responses in NMR neurons compared to mouse neurons [43], with APETx2, an inhibitor of
10 ASIC3-containing ASICs, demonstrating a key role for ASIC3, even though nmrASIC3 does not
11 appear to form functional homotrimers [50]. Previously we mapped out ASIC expression in
12 different NMR brain regions and observed similar expression between mouse and NMR, a key
13 exception being much lower ASIC4 levels throughout the NMR brain [51], however, no one has
14 yet studied the function of ASICs in NMR brain neurons.

15 In this study, we investigated acid-induced currents in mouse and NMR neurons using
16 whole-cell patch clamp recording of cultured neonatal hippocampal and cortical neurons. We find
17 that NMR neurons have ASIC-mediated currents of significantly smaller peak current amplitude
18 than those recorded from mouse neurons and that NMR neurons are resistant to acid-induced
19 cell death. Overall, these results suggest that the reduced acid-induced cell death in NMR
20 neurons may be neuroprotective.

22 **Methods**

23 *Animals*

24 All experiments were conducted in accordance with the United Kingdom Animal (Scientific
25 Procedures) Act 1986 Amendment Regulations 2012 under a Project License (70/7705) granted

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 to E. St. J. S. by the Home Office; the University of Cambridge Animal Welfare Ethical Review
2 Body also approved procedures. Breeding couples of 1 male and 2 female C57/bl6 mice were
3 conventionally housed with nesting material and a red plastic shelter; the holding room was
4 temperature-controlled (21 °C) and mice were on a normal 12-hour light/dark cycle with food and
5 water available *ad libitum*. Naked mole-rats were bred in house and maintained in a custom-made
6 caging system with conventional mouse/rat cages connected by different lengths of tunnel.
7 Bedding and nesting material were provided along with running wheels and chew blocks. The
8 room was warmed to 28 °C and humidified, with a heat cable to provide extra warmth running
9 under 2–3 cages, and red lighting (08:00–16:00) was used.

10 *Neuronal cultures*

11 P0-P2 mice and P0-P5 naked mole-rats were used to prepare cortical and hippocampal neuronal
12 cultures. Multiple pups (2-4) were used to prepare a single culture. Following decapitation, heads
13 were immediately placed in dishes containing ice-cold Hank's Balanced Salt Solution (HBSS)
14 solution (20 mM HEPES, 30 mM glucose in HBSS, Life Technologies). Brains were removed,
15 transferred to a new dish and the two hippocampi and cortices were isolated. Tissues were
16 subsequently incubated in an enzymatic digestion solution: 2 mg/mL papain (Worthington
17 Biochemical Corporation) in Hibernate-Ca²⁺ solution (Brain Bits), activated by 0.5 mM Glutamax
18 (Life Technologies) at 37 °C for 30 min in a 5 % CO₂ incubator. The digestion solution was then
19 replaced by HBSS solution supplemented with DNase I (250 Kunitz units/mL, Sigma Aldrich) and
20 tissues were slowly triturated (5-7 times) using a P1000 pipette. Neuronal suspensions were
21 filtered through a 100 µm nylon cell strainer (Corning) to remove non-dissociated pieces of tissues
22 before centrifugation for 5 mins at 1100 rpm at room temperature. Supernatants were discarded
23 and the pellets carefully resuspended in HBSS solution. After further centrifugation for 5 min 1100
24 rpm at room temperature, pellets were resuspended in MEM/HS solution: 10 % heat-inactivated
25 horse serum (Life Technologies), 2 mM glucose, 0.0025 % Glutamax, and 0.2 mg/mL primocin
26 (InVivogen). Hippocampal and cortical neurons were plated on 35 mm plastic dishes (Fisher

1
2
3
4 1 Scientific), previously coated with 100 mg/mL poly-L-lysine (Sigma-Aldrich), rinsed with water and
5
6 2 dried, at a density of 300 000 cells/mL (2 mL/dish). After a 4-hour incubation in a 37 °C / 5 % CO₂
7
8 3 incubator, the MEM/HS solution was removed and the dishes were flooded with Neurobasal/B27
9
10 4 solution (1X B27 Supplement, 0.0025 % Glutamax, and 0.2 mg/mL primocin). Naked mole-rat
11
12 5 neurons were kept at 33 °C in 5 % CO₂ incubator, whereas mouse neurons were kept at 37 °C in
13
14 6 a 5 % CO₂ incubator; this is due to NMRs being cold-blooded and NMR cells do not withstand 37
15
16 7 °C for long periods of time [52]. Half of the medium was exchanged for fresh medium every 2-3
17
18 8 days until the cultures were used for experiments.
19
20

21 9 *Electrophysiology*

22
23
24 10 Hippocampal and cortical neurons from mouse and NMR were used for whole-cell patch-clamp
25
26 11 recordings at 9-12 days *in vitro* (DIV9-12). Recordings were performed at room temperature using
27
28 12 the following solutions: extracellular (in mM) – 140 NaCl, 4 KCl, 2 CaCl₂, 1 MgCl₂, 4 glucose, 10
29
30 13 HEPES, adjusted to pH 7.4 with NaOH and 300–310 mOsm with sucrose; intracellular (in mM) –
31
32 14 110 KCl, 10 NaCl, 1 MgCl₂, 1 EGTA, 10 HEPES, 2 Na₂ATP, 0.5 Na₂GTP, adjusted to pH 7.3 with
33
34 15 KOH and to 310–315 mOsm with sucrose. Acidic extracellular solutions were made using MES
35
36 16 (pH 5.0). Patch pipettes were pulled (Model P-97, Flaming/Brown puller; Sutter Instruments,) from
37
38 17 borosilicate glass capillaries (Hilgenberg GmbH) and had a resistance of 6-10 MΩ. Data were
39
40 18 acquired using an EPC10 amplifier and Patchmaster software (HEKA). Whole-cell currents were
41
42 19 recorded at 20 kHz, pipette and membrane capacitance were compensated using Patchmaster
43
44 20 macros, and series resistance was compensated by >60 %. Cell capacitances and resting
45
46 21 membrane potentials were measured just after cell opening in whole-cell configuration. To study
47
48 22 macroscopic voltage-gated currents, a standard voltage-step protocol was used whereby cells
49
50 23 were held at -120 mV for 200 msec before stepping to the test potential (-80 mV - +65 mV in 5
51
52 24 mV increments) for 50 msec, returning to the holding potential (-60 mV) for 200 msec between
53
54 25 sweeps. In some experiments, tetrodotoxin (300 nM, Alomone Labs) was perfused for 30 seconds
55
56 26 before repeating the voltage-step protocol. To measure neuronal acid-sensitivity, cells were
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 exposed to the following protocol: 5 seconds of pH 7.4; 5-seconds of pH 5; and 5 seconds of pH
2 7.4. ASIC antagonists (100 μ M Benzamil, Sigma) were perfused during 30 seconds before
3 applying another 5 second pulse of pH 5. After 90 second wash time with pH 7.4 solution, a 5
4 second pulse of pH 5 was applied to check for reversal of any block observed. Current amplitude
5 was measured in Fitmaster (HEKA) by taking the maximum peak response and subtracting the
6 mean baseline amplitude in the preceding 50 msec (voltage-gated currents) or ~2.5 sec (ASIC
7 currents); current amplitude was normalized for cell size by dividing by cell capacitance. Using
8 Igor Pro, for each individual cell that underwent the voltage-step protocol, the following equation
9 was fitted to the normalized inward currents:

$$i(x) = \Gamma \cdot x \cdot \frac{1 - e^{-\frac{x-E_{rev}}{25mV}}}{1 - e^{-\frac{x}{25mV}}} \cdot \frac{1}{(1 - e^{-\frac{x-V_{Half}}{slope}})^3}$$

11 where E_{rev} is the reversal potential; V_{half} the half-activating potential; Γ a constant and x the
12 command potential. Similarly, a Boltzmann equation was fitted to the normalized outward
13 currents:

$$\frac{I}{I_{max}} = \frac{1}{1 + e^{\frac{(V_{Half}-V_m)}{slope}}}$$

15 where V_m is the membrane voltage and V_{Half} the voltage at half-maximal activation. To
16 determine the inactivation time of the ASIC-mediated currents, a single exponential was fitted.
17 Data are expressed as mean \pm standard error of the mean (SEM). Using Prism (GraphPad),
18 paired t-tests were used to compare the effects of ASIC antagonists on proton-gated currents
19 within both mouse and NMR neuron datasets; unpaired t-tests were used to compare parameters,
20 such as neuronal resting membrane potential and capacitance and ASIC-mediated current
21 amplitude, between mouse and NMR neuron datasets.

Acid-induced cell death assays

23 Mouse and NMR neurons were used to measure acid-induced cell death at DIV9-12. The pH 7.4
24 and pH 5 extracellular solutions used were the same as those described above for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 electrophysiology experiments. Neuronal cultures from both mouse and NMR were rinsed twice
2 with 37 °C solution (pH 7.4 or pH 5) and then incubated with pH 7.4 or pH 5 solution for 2 hours
3 at 37 °C. Cultures from both conditions were then rinsed with warm pH 7.4 solution and incubated
4 during 30 min with pH 7.4 solution containing 1.5 mM propidium iodide (PI, Sigma Aldrich) to stain
5 necrotic cells and Hoechst 33342 (dilution 1/2500, Sigma) to label all nuclei. Labelled cultures
6 were imaged using an epifluorescence microscope (Olympus) equipped with a 20X objective
7 (Olympus) and a QImaging camera. To determine the percentage of dead necrotic PI-positive
8 cells, we used the software Fiji to count the total number of cells per field of view by counting
9 nuclei on the Hoechst images, and subsequently counting the number of PI-positive nuclei on the
10 PI images. One to three dishes per condition were used, and three different images per dish were
11 taken. Data were collected from three different cultures and each culture was prepared from
12 multiple animals. A one-way ANOVA test (Prism, GraphPad) corrected for multiple comparisons
13 (Tukey test) was used to compare the percentage of cell death in each field of view at pH 7.4 and
14 pH 5 in both species. Data are expressed as mean ± standard error of the mean (SEM).

16 **Results**

17 *Basic electrophysiological properties and voltage-gated Na⁺ channel activity differ between NMR* 18 *and mouse neurons*

19 Electrophysiological recordings from NMR neurons have been performed in both DRG
20 sensory neurons [43,50] and CNS neurons [46,47,53]. However, neuronal activity from NMR CNS
21 has only been recorded in brain slices, in the form of field excitatory postsynaptic potentials, and
22 the basic electrophysiological properties of NMR neurons in hippocampal and cortical cultures
23 have not yet been described.

24 We first compared the capacitance and resting membrane potential of NMR and mouse
25 neurons from both cortical and hippocampal neuronal cultures (Fig. 1). The capacitance of NMR

1 neurons was significantly smaller than in mouse neurons in both cortical and hippocampal
2 cultures (cortex: 17.27 ± 1.02 pF versus 28.72 ± 2.32 pF for NMR (n = 30) and mouse (n = 24)
3 neurons, respectively; hippocampus: 17.41 ± 1.03 pF versus 38.13 ± 3.27 pF for NMR (n = 18)
4 and mouse (n = 26) neurons, respectively; unpaired two-sided t-tests, **** p < 0.0001, Fig. 1a).
5 Resting membrane potentials were measured as soon as the whole-cell configuration was
6 established and NMR neurons were significantly more hyperpolarized than mouse neurons, in
7 both cortical and hippocampal cultures (cortex: -57.03 ± 2.64 mV versus -44.05 ± 2.84 mV for
8 NMR (n = 30) and mouse (n = 21) neurons, respectively; hippocampus: -55.11 ± 4.79 mV versus
9 -43.85 ± 2.59 mV for NMR (n = 18) and mouse (n = 26) neurons, respectively; unpaired two-sided
10 t-tests; ** p < 0.01; * p < 0.05, Fig. 1b).

11 We then investigated macroscopic voltage-gated currents in NMR and mouse neurons,
12 using a voltage-step whereby cells were held at -120 mV for 200 msec before stepping to the
13 test potential (-80 mV to +65 mV in 5 mV increments) for 50 msec, and returning to the holding
14 potential (-60 mV) for 200 msec between sweeps (Fig. 1c). Both NMR and mouse neurons
15 showed inward and outward currents (Fig. 1c-e). In some experiments, the voltage-step protocol
16 was run twice, the second time after 300 nM tetrodotoxin (TTX) had been applied for 30 seconds
17 to investigate the contribution of TTX-sensitive voltage-gated Na⁺ channels (NaVs) to the
18 macroscopic voltage-gated inward currents recorded in NMR neurons (Fig. 1c, right panel). In
19 both cortical and hippocampal NMR neurons, the voltage-gated inward currents were fully blocked
20 by 300 nM TTX (n = 8 and n = 2 for cortical and hippocampal neurons, respectively). The fact that
21 no inward current remained after application of 300 nM TTX indicates that only NaVs were
22 activated with our voltage-step protocol and that there was no measurable contribution of voltage-
23 gated Ca²⁺ channels to the inward currents recorded. Moreover, these results indicate that cortical
24 and hippocampal NMR neurons only express TTX-sensitive NaVs.

25 Strikingly, NMR neurons had a significantly smaller inward peak current density in both
26 cortical and hippocampal cultures (cortex: 28.79 ± 3.83 pA/pF versus 202.40 ± 21.35 pA/pF for

1
2
3
4 1 NMR (n = 19) and mouse (n = 17) neurons, respectively; hippocampus: 39.00 ± 6.39 pA/pF versus
5
6 2 196.70 ± 28.28 pA/pF for NMR (n = 13) and mouse (n = 16) neurons, respectively; unpaired two-
7
8 3 sided t-tests, **** p < 0.0001, Fig. 1d.i-ii). Additionally, the voltage of half-activation (V_{half}) and
9
10 4 the peak inward current amplitude potential (peak V_m) were more depolarized in NMR neurons
11
12 5 compared to mouse neurons (cortex: V_{half} : -36.12 ± 2.80 mV versus -45.42 ± 1.99 mV and peak
13
14 6 V_m : -12.43 ± 3.02 mV versus -31.27 ± 2.61 mV for NMR (n = 19) and mouse (n = 17) neurons,
15
16 7 respectively; hippocampus: V_{half} : -37.10 ± 2.34 mV versus -45.71 ± 1.30 mV and peak V_m : -
17
18 8 17.62 ± 4.00 mV versus -30.68 ± 3.09 mV for NMR (n = 13) and mouse (n = 16) neurons,
19
20 9 respectively; unpaired two-sided t-tests, **** p < 0.0001, ** p < 0.01; * p < 0.05, Fig. 1d.iii-iv).
21
22 10 These results suggest that NMR neurons may be less excitable compared to mouse neurons,
23
24 11 with more hyperpolarized resting membrane potentials and smaller voltage-gated inward currents
25
26 12 that are activated at more depolarized potentials, i.e. a greater depolarizing stimulus is required
27
28 13 to activate NMR voltage-gated inward currents that produce much smaller currents.
29
30
31
32

33 14 By contrast, current-voltage curves for voltage-gated outward currents were similar
34
35 15 between NMR and mouse neurons from both cortical and hippocampal neurons (Fig. 1e.i) and
36
37 16 V_{halves} were not significantly different (cortex: 7.13 ± 1.91 mV versus 10.77 ± 2.18 mV for NMR (n
38
39 17 = 24) and mouse (n = 17) neurons, respectively; hippocampus: 5.15 ± 2.50 mV versus $11.59 \pm$
40
41 18 2.26 mV for NMR (n = 14) and mouse (n = 16) neurons, respectively; unpaired two-sided t-tests,
42
43 19 p = 0.221 and p = 0.094 for cortex and hippocampus respectively, Fig. 1e.ii).
44
45
46
47
48

49 21 *Acid-induced currents are mediated by ASICs in both NMR and mouse neurons*

50
51 22 The expression profile of ASIC subunits is similar in mouse and NMR brains, with the
52
53 23 exception of lower levels of ASIC4 throughout the NMR brain [51], results suggesting that
54
55 24 functional ASIC-mediated currents should be present in NMR as others have shown in mouse
56
57
58 25 [4,5,7,8,15].
59
60
61
62
63
64
65

1
2
3
4 1 A 5 second pulse of pH 5 was applied to NMR and mouse neurons from both hippocampal
5
6 2 and cortical neurons and rapidly activating and inactivating acid-induced responses were
7
8 3 recorded in every cell of both species (Fig. 2a). However, the peak current density of acid-
9
10 4 mediated responses recorded in NMR neurons was significantly smaller than in mouse neurons,
11
12 5 in both hippocampal and cortical neurons (cortex: 15.41 ± 1.82 pA/pF versus 85.66 ± 10.90 pA/pF
13
14 6 for NMR (n = 31) and mouse (n = 24) neurons, respectively; hippocampus: 20.54 ± 2.86 pA/pF
15
16 7 versus 100.90 ± 17.68 pA/pF for NMR (n = 22) and mouse (n = 26) neurons, respectively;
17
18 8 unpaired two-sided t-tests, **** p < 0.0001, *** p < 0.001, Fig. 2b). By contrast, the inactivation
19
20 9 time constant of the acid-mediated currents was similar between NMR and mouse neurons
21
22 10 (cortex: 0.35 ± 0.012 s versus 0.44 ± 0.056 s for NMR (n = 27) and mouse (n = 18) neurons,
23
24 11 respectively, unpaired two-sided t-test, p=0.0516; hippocampus: 0.47 ± 0.06 s versus 0.35 ± 0.04
25
26 12 s for NMR (n = 13) and mouse (n = 19) neurons, respectively; unpaired two-sided t-test, p =
27
28 13 0.0861, Fig. 2c).

29
30
31
32
33 14 The transient nature of the acid-mediated inward currents in both NMR and mouse
34
35 15 neurons is characteristic of ASIC-mediated currents [3] and to confirm the involvement of ASICs
36
37 16 we utilized the non-selective ASIC antagonist benzamil [54]. After a first application of pH 5 for 5
38
39 17 seconds, we applied 100 μ M benzamil for 30 seconds before a second pH 5 pulse, then followed
40
41 18 by a wash period of 90 seconds (Fig. 2d, 2g). In cortical and hippocampal mouse neurons, the
42
43 19 acid-induced currents were reversibly blocked by 100 μ M benzamil (cortex: pH 5: 133.60 ± 19.01
44
45 20 pA/pF; benzamil: 16.77 ± 5.28 pA/pF; wash: 133.80 ± 19.88 pA/pF (n = 9); hippocampus: pH 5:
46
47 21 151.00 ± 33.64 pA/pF; benzamil: 7.84 ± 2.67 pA/pF; wash: 89.26 ± 20.48 pA/pF (n = 10); one-
48
49 22 way paired ANOVA test, Tukey's multiple comparison test, *** p < 0.001, ** p < 0.01, Fig. 2e-f).
50
51 23 This result concurs with previous studies indicating that acid-evoked currents in mouse
52
53 24 hippocampal and cortical neurons are mediated by ASICs [4,8,15].

54
55
56
57
58 25 Similarly, in cortical and hippocampal NMR neurons, acid-induced currents were
59
60 26 reversibly inhibited by 100 μ M benzamil (cortex: pH 5: 9.66 ± 1.09 pA/pF; benzamil: 2.78 ± 0.49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 pA/pF; wash: 9.13 ± 0.89 pA/pF (n = 7); hippocampus: pH 5: 11.13 ± 1.88 pA/pF; benzamil: 3.50
2 ± 0.53 pA/pF; wash: 8.15 ± 1.30 pA/pF (n = 6); one-way paired ANOVA test, Tukey's multiple
3 comparison test, *** p < 0.001, ** p < 0.01, Fig. 2h-i). This is the first demonstration of functional
4 ASIC-mediated currents in CNS NMR neurons.

6 *NMR neurons are resistant to acid-induced cell death*

7 ASICs are involved in acid-induced cell death, so-called acidotoxicity, which can occur
8 during periods of ischemia [5,15,16]. Because NMR neurons exhibit significantly smaller ASIC
9 currents (Fig. 2), we hypothesized that this could be neuroprotective when neurons are in an
10 acidic environment. We exposed neuronal cultures from mouse and NMR cortices to a pH 7.4 or
11 a pH 5 solution for 2 hours at 37 °C. Nuclei were stained by Hoechst 33342 and necrotic (dead)
12 neurons were labeled using propidium iodide (PI) (Fig. 3). Percentages of dead neurons were
13 calculated by counting the number of necrotic PI-positive cells over the total number of Hoechst
14 33342-positive neurons in the field of view, and a one-way ANOVA test corrected for multiple
15 comparisons (see *Methods*) was used to compare neuronal death at pH 5 between both species
16 (Fig. 3c). At pH 7.4, percentages of dead cells were comparable between mouse and NMR cortical
17 cultures (mouse - pH 7.4: 20.31 ± 3.50 %, n = 18 fields of view; NMR – pH 7.4: 13.31 ± 1.53 %,
18 n = 27 fields of view; from 3 independent experiments), suggesting no differences in neuronal
19 death under basal conditions between NMR and mouse cultures. When incubated for 2 hours in
20 pH 5 solution, mouse cortical neurons exhibited a significantly increased percentage of cell death
21 compared to all other conditions (mouse – pH 5: 60.55 ± 4.87 %, n = 18 fields of view; NMR – pH
22 5: 22.29 ± 2.30 %, n = 28 fields of view, from 3 independent experiments, p < 0.0001, ****). By
23 contrast, when NMR cortical neurons were exposed to pH 5 for 2 hours, no difference in the level
24 of cell death was observed compared to incubation at pH 7.4. Similar results were obtained using
25 mouse and NMR hippocampal neurons (data not shown). This indicates that NMR neurons are

1
2
3
4 1 resistant to acid-induced cell death, possibly due to their reduced ASIC currents compared to
5
6 2 mouse neurons.
7
8
9 3

10 4 **Discussion**

11
12
13
14 5 In this study, we recorded for the first time from cultured NMR brain neurons and
15
16 6 described their basic electrophysiological properties (Fig. 1). We showed that NMR neuronal
17
18 7 capacitance was smaller than in mouse neurons and that the resting membrane potential of NMR
19
20 8 neurons is more hyperpolarized than in mouse neurons. One point to consider is that neurons
21
22 9 were recorded from blindly and so we can only broadly compare hippocampal and cortical
23
24 10 neurons between species and cannot comment on if these differences occur in all types of neuron
25
26 11 or specific neuronal subpopulations, such as interneurons. Although capacitance and resting
27
28 12 membrane potential values obtained from mouse neurons were similar to those reported by others
29
30 13 using cultured rodent neurons [55,56], for NMR neurons however, previous data from others
31
32 14 found their resting membrane potential to be more hyperpolarized (NMR cortical neurons: -57.03
33
34 15 ± 2.64 mV, NMR hippocampal neurons: -55.11 ± 4.79 mV, Fig. 1; NMR hippocampal pyramidal
35
36 16 neurons (4 months old): -70.3 ± 6.1 mV, NMR hippocampal dentate granule cells (4 months old):
37
38 17 -75.1 ± 3.6 mV, [53]). Explanations for differences observed in resting membrane potentials
39
40 18 between studies are the recording conditions (*in vitro* vs. *in vivo*), the developmental stage of the
41
42 19 animal from which neurons are obtained and the constituents of the intracellular and extracellular
43
44 20 solutions. When comparing our study with that of Penz and colleagues, who reported more
45
46 21 hyperpolarized resting membrane potentials, two factors that could contribute to the difference
47
48 22 observed is that we made recordings from cultured neurons, whereas they used brain slices, and
49
50 23 secondly, we isolated neurons from neonatal animals, whereas slice recordings were made from
51
52 24 animals aged at least 4-months.
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 1 A standard voltage-step protocol that has been already successfully used in NMR
5
6 2 sensory neurons to predominantly isolate NaV activity [43] was used to measure macroscopic
7
8 3 voltage-gated current activity in mouse and NMR neurons. NaV currents recorded from mouse
9
10 4 neurons were not different to what have been recorded in similar experimental conditions (for
11
12 5 example, mouse cortical neurons at DIV 10-12: V_{half} : -41.62 ± 1.46 mV in [57] versus this study
13
14 6 : -45.42 ± 1.99 mV). Similarly, inward currents recorded in NMR brain neurons were also very
15
16 7 similar to currents recorded in NMR sensory neurons (NMR cortex: V_{half} : -36.12 ± 2.80 mV; NMR
17
18 8 hippocampus: V_{half} : -37.10 ± 2.34 mV; NMR DRG neurons: V_{half} : from -34.2 ± 0.1 mV to -44.6
19
20 9 ± 0.5 mV [43]). However, NMR inward currents significantly differed from mouse currents: the
21
22 10 peak current density was significantly smaller and the V_{half} and peak V_m values were more
23
24 11 depolarized. These differences in NMR NaV activity likely result from differences in amino acid
25
26 12 sequence of NaV subunits and/or differential expression of accessory subunits and warrant
27
28 13 further investigation. We also found that addition of 300 nM TTX completely abolished all voltage-
29
30 14 gated inward currents demonstrating an absence of TTX resistant NaV subunits in the NMR brain,
31
32 15 this result aligns with the observation that the TTX-resistant NaV subunits, NaV1.5, NaV1.8 and
33
34 16 NaV1.9 are not expressed in the central nervous system [58]. It should also be noted that although
35
36 17 NMR neurons were cultured at 32 °C under hypoxic (3 % O₂) conditions, recordings under
37
38 18 normoxic, standard laboratory conditions, whereas NMRs live in a hypoxic and hypercapnic
39
40 19 environment [59,60], which may influence channel activity *in vivo*.

41
42
43
44
45
46 20 ASIC activity in brain neurons is now accepted as a key factor in numerous physiological
47
48 21 and pathological conditions [61]. However, nothing is known about acid-induced responses in
49
50 22 NMR brain neurons, which is of considerable interest considering the behavioral
51
52 23 hypoxia/hypercapnia resistance and lack of acid-induced nocifensive behavior display by NMR
53
54 24 that are likely adaptations to adapting to a safe, but relatively hypoxic and hypercapnic habitat
55
56 25 [35]. Recently, we described ASIC subunit expression in the NMR CNS [62], which is similar to
57
58 26 that in the mouse CNS, with the exception of lowered ASIC4 levels in the NMR brain. However,
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 evidence for functional ASIC activity is lacking and is of particular interest in light of our recent
2 finding that nmrASIC3 forms non-functional homomers [50]. Here, we find that NMR brain neurons
3 produce ASIC-mediated currents in response to acid stimulation, in both hippocampus and cortex,
4 as demonstrated by acid-induced responses being fully, reversibly blocked by 100 μ M benzamil
5 (Fig. 2). However, the peak current density of NMR ASIC-mediated responses was significantly
6 reduced compared to responses recorded in mouse neurons. The reasons for such reduction in
7 ASIC currents in NMR neurons is not known and additional research is needed to determine what
8 underpins this different, e.g. are regulators of ASIC plasma membrane trafficking different in
9 NMR? Is there a different developmental expression profile of ASICs between mouse and NMR?
10 With regard to the ASIC currents themselves, they have similar decay kinetics in both NMR and
11 mouse neurons (Fig. 2c), which suggests that a similar mixture of ASIC subunits are expressed,
12 as our previous mRNA based analysis suggested [62].

13 Incubation with a pH 5.0 solution showed that unlike mouse neurons, NMR neurons do
14 not undergo any significant acid-induced cell death (Fig. 3). This is the first demonstration of
15 resistance to acid-induced neuronal death in NMR neurons. In rodent neurons, some factors
16 shown to be protective against acid-induced neuronal injury, include lower temperature [63],
17 pharmacological blockade or genetic deletion of ASIC activity [15] or ASIC trafficking [64], i.e. it
18 is established that ASICs play a key role in acidotoxicity. Considering the similar prevalence of
19 ASIC currents, we suggest that the reduced ASIC-mediated current amplitude observed in NMR
20 neurons may be an additional neuroprotective mechanism in NMR brains, alongside the
21 previously described increased hypoxia-inducible transcription factor (HIF1- α) expression [65] or
22 more efficient *in vivo* CO₂ buffering [47]. One possible mechanism for the decreased amplitude
23 observed is reduced ASIC plasma membrane trafficking which is known to be modulated by an
24 extracellular acidic environment [64]. However, it is also possible that the reduced acid-induced
25 cell death observed is not ASIC-dependent because although ASIC activation appears to play a
26 major role in neuronal injury [61], several other molecular players are also modulated by a drop

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 in extracellular pH, such as NaVs [27,28] and glutamate receptors [31] and may contribute to the
2 lowered acid-induced cell death observed, for example, here we also show that NMR neurons
3 also have smaller NaV-mediated currents, which may also add a layer of neuroprotection.

5 **Conclusions**

6 In this work, we describe for the first time the basic electrophysiological properties of NMR
7 neurons in culture and showed that the resting membrane potential of NMR neurons is more
8 hyperpolarized, as well as the amplitude of NaVs being smaller than that of mouse neurons. We
9 then demonstrated that acid-induced currents are present in NMR neurons and are, as in mouse,
10 ASIC-mediated. The key result is that acid-induced cell death is virtually absent in NMR neurons,
11 with reduced ASIC and NaV amplitudes likely contributing to this observation, and thus this is a
12 further adaptation enabling NMR to live in a subterranean, hypercapnic/hypoxic environment.

14 **Abbreviations**

15 ASIC, acid-sensing ion channel; CNS, central nervous system; DRG, dorsal root ganglion, NaV,
16 voltage-gated Na⁺ channel; NMR, naked mole-rat; TTX, tetrodotoxin

18 **Declarations**

19 *Ethics*

20 All experiments were conducted in accordance with the United Kingdom Animal (Scientific
21 Procedures) Act 1986 Amendment Regulations 2012 under a Project License (70/7705) granted
22 to E. St. J. S. by the Home Office; the University of Cambridge Animal Welfare Ethical Review
23 Body also approved procedures.

24 *Consent for publication*

25 N/A

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 *Availability of data and material*

2 The datasets supporting the conclusions of this article are available in the University of Cambridge
3 Apollo repository: <https://doi.org/10.17863/CAM.18704>

4 *Competing interests*

5 The authors have no competing interests.

6 *Funding*

7 This work was supported by an Isaac Newton Trust Research Grant from the University of
8 Cambridge. ZH was funded by a EMBO Long-Term Fellowship (ALTF1565-2015).

9 *Authors' contributions*

10 ZH and ESS designed the study. ZH performed all the experiments and analysed the data. ZH
11 and ESS wrote the manuscript.

12 *Acknowledgements*

13 Thanks to all members of ESS lab for their useful discussions, and in particular to the assistant
14 staff in the Department of Pharmacology and Animal Facility for their technical help and animal
15 husbandry.

17
18 **References**

19 1. Gründer S, Pusch M. Biophysical properties of acid-sensing ion channels (ASICs).
20 Neuropharmacology. 2015;94:9–18.
21 2. Waldmann R, Champigny G, Bassilana F, Heurteaux C, Lazdunski M. A proton-gated cation
22 channel involved in acid-sensing. Nature. 1997. p. 173–7.
23 3. Hesselager M, Timmermann DB, Ahring PK. pH Dependency and Desensitization Kinetics of
24 Heterologously Expressed Combinations of Acid-sensing Ion Channel Subunits. J. Biol. Chem.
25 2004;279:11006–15.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 4. Baron A, Waldmann R, Lazdunski M. ASIC-like , proton-activated currents in rat hippocampal
2 neurons. *J. Physiol.* 2002;539:485–94.

3 5. Sherwood TW, Lee KG, Gormley MG, Askwith CC. Heteromeric Acid-Sensing Ion Channels
4 (ASICs) Composed of ASIC2b and ASIC1a Display Novel Channel Properties and Contribute to
5 Acidosis-Induced Neuronal Death. *J. Neurosci.* 2011;31:9723–34.

6 6. Wu L-J, Duan B, Mei Y-D, Gao J, Chen J-G, Zhuo M, et al. Characterization of Acid-sensing
7 Ion Channels in Dorsal Horn Neurons of Rat Spinal Cord. *J. Biol. Chem.* 2004;279:43716–24.

8 7. Gao J, Wu LJ, Xu L, Xu T Le. Properties of the proton-evoked currents and their modulation
9 by Ca²⁺ and Zn²⁺ in the acutely dissociated hippocampus CA1 neurons. *Brain Res.*
10 2004;1017:197–207.

11 8. Askwith CC, Wemmie J a., Price MP, Rokhlina T, Welsh MJ. Acid-sensing Ion Channel 2
12 (ASIC2) Modulates ASIC1 H⁺-activated Currents in Hippocampal Neurons. *J. Biol. Chem.*
13 2004;279:18296–305.

14 9. Wemmie J a., Chen J, Askwith CC, Hruska-Hageman AM, Price MP, Nolan BC, et al. The
15 acid-activated ion channel ASIC contributes to synaptic plasticity, learning, and memory.
16 *Neuron.* 2002;34:463–77.

17 10. Liu MG, Li HS, Li WG, Wu YJ, Deng SN, Huang C, et al. Acid-sensing ion channel 1a
18 contributes to hippocampal LTP inducibility through multiple mechanisms. *Sci. Rep.* 2016;6:1–
19 14.

20 11. Mango D, Braksator E, Battaglia G, Marcelli S, Mercuri NB, Feligioni M, et al. Acid-sensing
21 ion channel 1a is required for mGlu receptor dependent long-term depression in the
22 hippocampus. *Pharmacol. Res.* 2017;119:12–9.

23 12. Vralsted VC, Price MP, Du J, Schnizler M, Wunsch AM, Ziemann AE, et al. Expressing acid-
24 sensing ion channel 3 in the brain alters acid-evoked currents and impairs fear conditioning.
25 *Genes, Brain Behav.* 2011;10:444–50.

26 13. Taugher RJ, Lu Y, Fan R, Ghobbeh A, Kreple CJ, Faraci FM, et al. ASIC1A in neurons is

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 critical for fear-related behaviors. *Genes, Brain Behav.* 2017;16:745–55.

2 14. Wemmie JA, Coryell MW, Askwith CC, Lamani E, Leonard AS, Sigmund CD, et al.

3 Overexpression of acid-sensing ion channel 1a in transgenic mice increases acquired fear-

4 related behavior. *Proc. Natl. Acad. Sci.* 2004;101:3621–6.

5 15. Xiong Z-G, Zhu X-M, Chu X-P, Minami M, Hey J, Wei W-L, et al. Neuroprotection in

6 Ischemia: Blocking Calcium-Permeable Acid-Sensing Ion Channels. *Cell.* 2004;118:687–98.

7 16. Duan B, Wang Y-Z, Yang T, Chu X-P, Yu Y, Huang Y, et al. Extracellular Spermine

8 Exacerbates Ischemic Neuronal Injury through Sensitization of ASIC1a Channels to

9 Extracellular Acidosis. *J. Neurosci.* 2011;31:2101–12.

10 17. Gu L, Liu X, Yang Y, Luo D, Zheng X. ASICs aggravate acidosis-induced injuries during

11 ischemic reperfusion. *Neurosci. Lett.* 2010;479:63–8.

12 18. Friese MA, Craner MJ, Etzensperger R, Vergo S, Wemmie JA, Welsh MJ, et al. Acid-

13 sensing ion channel-1 contributes to axonal degeneration in autoimmune inflammation of the

14 central nervous system. 2007;13:1483–9.

15 19. Chu X-P, Xiong Z-G. Physiological and Pathological Functions of Acid-Sensing Ion

16 Channels in the Central Nervous System. *Curr. Drug Targets.* 2012;13:263–71.

17 20. Gonzales EB, Sumien N. Acidity and Acid-Sensing Ion Channels in the Normal and

18 Alzheimer’s Disease Brain. *J. Alzheimer’s Dis.* 2017;57:1137–44.

19 21. Chesler M. Regulation and Modulation of pH in the Brain. 2003;1183–221.

20 22. Rehncrona S. Brain acidosis. *Ann. Emerg. Med.* 1985;14:770–6.

21 23. Immke DC, McCleskey EW. Lactate enhances the acid-sensing Na⁺ channel on ischemia-

22 sensing neurons. *Nat. Neurosci.* 2001;4:869–70.

23 24. Allen NJ, Attwell D. Modulation of ASIC channels in rat cerebellar Purkinje neurons by

24 ischaemia-related signals. *J. Physiol.* 2002;543:521–9.

25 25. Smith ES, Cadiou H, McNaughton P a. Arachidonic acid potentiates acid-sensing ion

26 channels in rat sensory neurons by a direct action. *Neuroscience.* 2007;145:686–98.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 26. Tombaugh GC, Somjen GG. Effects of extracellular pH on voltage-gated Na⁺, K⁺ and Ca²⁺
2 currents in isolated rat CA1 neurons. *J. Physiol.* 1996;493:719–32.
3 27. Nakamura M, Jang I-S. Acid modulation of tetrodotoxin-resistant Na⁺ channels in rat
4 nociceptive neurons. *Neuropharmacology.* Elsevier Ltd; 2015;90:82–9.
5 28. Nakamura M, Kim DY, Jang IS. Acid modulation of tetrodotoxin-sensitive Na⁺ channels in
6 large-sized trigeminal ganglion neurons. *Brain Res.* 2016;1651:44–52.
7 29. Sepulveda F V., Pablo Cid L, Teulon J, Niemeyer MI. Molecular Aspects of Structure,
8 Gating, and Physiology of pH-Sensitive Background K₂P and Kir K⁺-Transport Channels.
9 *Physiol. Rev.* 2015;95:179–217.
10 30. Mcdonald JW, Bhattacharyya T, Sensi SL, Lobner D, Ying HS, Canzoniero LMT, et al.
11 Extracellular Acidity Potentiates AMPA Receptor-Mediated Cortical Neuronal Death.
12 *1998;18:6290–9.*
13 31. Traynelis SF, Cull-Candy SG. Proton inhibition of N-methyl-D-aspartate receptors in
14 cerebellar neurons. *Lett. to Nat.* 1990;345:347–50.
15 32. Bennett NC, Faulkes CG. African mole-rats: Ecology and Eusociality. Cambridge University
16 Press; 2000.
17 33. Brett RA. The ecology of naked mole-rat colonies: burrowing, food and limiting factors. In:
18 Sherman P, Jarvis J, Alexander R, editors. *Biol. Naked Mole-Rat.* Princeton University Press;
19 1991. p. 137–84.
20 34. Jarvis JUM. Reproduction of naked mole-rats. In: Sherman P, Jarvis J, Alexander M,
21 editors. *Biol. Naked Mole-Rat.* Princeton University Press; 1991. p. 384–425.
22 35. Schuhmacher L, Husson Z, Smith ES. The naked mole-rat as an animal model in biomedical
23 research: current perspectives. *Open Access Anim. Physiol.* 2015;7:137.
24 36. Ruby JG, Smith M, Buffenstein R. Naked Mole-Rat mortality rates defy gompertzian laws by
25 not increasing with age. *Elife.* 2018;7:1–18.
26 37. Edrey YH, Medina DX, Gaczynska M, Osmulski P a., Oddo S, Caccamo A, et al. Amyloid

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 beta and the longest-lived rodent: The naked mole-rat as a model for natural protection from
2 alzheimer's disease. *Neurobiol. Aging.* 2013;34:2352–60.

3 38. Orr ME, Garbarino VR, Salinas A, Buffenstein R. Sustained high levels of neuroprotective,
4 high molecular weight, phosphorylated tau in the longest-lived rodent. *Neurobiol. Aging. Elsevier*
5 *Inc;* 2015;36:1496–504.

6 39. Seluanov A, Hine C, Azpurua J, Feigenson M, Bozzella M, Mao Z, et al. Hypersensitivity to
7 contact inhibition provides a clue to cancer resistance of naked mole-rat. *Proc. Natl. Acad. Sci.*
8 *U. S. A.* 2009;106:19352–7.

9 40. Tian X, Azpurua J, Hine C, Vaidya A, Myakishev-Rempel M, Ablaeva J, et al. High-
10 molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. *Nature.*
11 2013;499:346–9.

12 41. Liang S, Mele J, Wu Y, Buffenstein R, Hornsby PJ. Resistance to experimental
13 tumorigenesis in cells of a long-lived mammal, the naked mole-rat (*Heterocephalus glaber*).
14 *Aging Cell.* 2010;9:626–35.

15 42. Park TJ, Lu Y, Jüttner R, Smith ESJ, Hu J, Brand A, et al. Selective inflammatory pain
16 insensitivity in the African naked mole-rat (*Heterocephalus glaber*). *PLoS Biol.* 2008;6:0156–70.

17 43. Smith ESJ, Omerbašić D, Lechner SG, Anirudhan G, Lapatsina L, Lewin GR. The molecular
18 basis of acid insensitivity in the African naked mole-rat. *Science.* 2011;334:1557–60.

19 44. Smith ESJ, Blass GRC, Lewin GR, Park TJ. Absence of histamine-induced itch in the
20 African naked mole-rat and “rescue” by Substance P. *Mol. Pain.* 2010;6:29.

21 45. Lavinka PC, Brand A, Landau VJ, Wirtshafter D, Park TJ. Extreme tolerance to ammonia
22 fumes in African naked mole-rats: Animals that naturally lack neuropeptides from trigeminal
23 chemosensory nerve fibers. *J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol.*
24 2009;195:419–27.

25 46. Larson J, Park TJ. Extreme hypoxia tolerance of naked mole-rat brain. *Neuroreport.*
26 2009;20:1634–7.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 47. Park TJ, Reznick J, Peterson BL, Blass G, Omerba D, Bennett NC, et al. Fructose-driven
2 glycolysis supports anoxia resistance in the naked mole-rat. 2017;311:307–11.

3 48. Peterson BL, Larson J, Buffenstein R, Park TJ, Fall CP. Blunted neuronal calcium response
4 to hypoxia in naked mole-rat hippocampus. PLoS One. 2012;7:1–8.

5 49. Peterson BL, Park TJ, Larson J. Adult naked mole-rat brain retains the NMDA receptor
6 subunit GluN2D associated with hypoxia tolerance in neonatal mammals. Neurosci. Lett.
7 2012;506:342–5.

8 50. Schuhmacher L-N, Callejo G, Srivats S, Smith ESJ. Naked mole-rat acid-sensing ion
9 channel 3 forms nonfunctional homomers, but functional heteromers. J. Biol. Chem.
10 2018;293:1756–66.

11 51. Schuhmacher L, Smith ESJ. Expression of acid-sensing ion channels and selection of
12 reference genes in mouse and naked mole rat. Mol. Brain. Molecular Brain; 2016;1–12.

13 52. Omerbašić D, Smith ESJ, Moroni M, Homfeld J, Eigenbrod O, Bennett NC, et al.
14 Hypofunctional TrkA Accounts for the Absence of Pain Sensitization in the African Naked Mole-
15 Rat. Cell Rep. 2016;17:748–58.

16 53. Penz OK, Fuzik J, Kurek AB, Romanov R, Larson J, Park TJ, et al. Protracted brain
17 development in a rodent model of extreme longevity. Sci. Rep. 2015;5:11592.

18 54. Baron A, Lingueglia E. Pharmacology of acid-sensing ion channels – Physiological and
19 therapeutical perspectives. Neuropharmacology. Elsevier Ltd; 2015;94:19–35.

20 55. Evans MS, Collings M a, Brewer GJ. Electrophysiology of embryonic, adult and aged rat
21 hippocampal neurons in serum-free culture. J. Neurosci. Methods. 1998;79:37–46.

22 56. Yang J, Thio LL, Clifford DB, Zorumski CF. Electrophysiological properties of identified
23 postnatal rat hippocampal pyramidal neurons in primary culture. Dev. Brain Res. 1993;71:19–
24 26.

25 57. Wang X, Zhang XG, Zhou TT, Li N, Jang CY, Xiao ZC, et al. Elevated neuronal excitability
26 due to modulation of the voltage-gated sodium channel Nav1.6 by Aβ1-42. Front. Neurosci.

1
2
3
4 1 2016;10:1–9.
5
6 2 58. Catterall WA. Voltage-gated sodium channels at 60 : structure , function and
7
8 3 pathophysiology. 2012;11:2577–89.
9
10 4 59. Shams I, Avivi A, Nevo E. Oxygen and carbon dioxide fluctuations in burrows of
11
12 5 subterranean blind mole rats indicate tolerance to hypoxic – hypercapnic stresses.
13
14 6 2005;142:376–82.
15
16 7 60. Mcnab BK. The Metabolism of Fossorial Rodents: A Study of Convergence. Ecol. Soc. Am.
17
18 8 1966;47:712–33.
19
20 9 61. Huang Y, Jiang N, Li J, Ji Y-H, Xiong Z-G, Zha X. Two aspects of ASIC function: Synaptic
21
22 10 plasticity and neuronal injury. Neuropharmacology. Elsevier Ltd; 2015;1–6.
23
24 11 62. Schuhmacher L, St E, Smith J. Expression of acid-sensing ion channels and selection of
25
26 12 reference genes in mouse and naked mole rat. Mol. Brain. Molecular Brain; 2016;1–12.
27
28 13 63. Ding D, Moskowitz SI, Li R, Lee SB, Esteban M, Tomaselli K, et al. Acidosis induces
29
30 14 necrosis and apoptosis of cultured hippocampal neurons. Exp. Neurol. 2000;162:1–12.
31
32 15 64. Zeng W-Z, Liu D-S, Buan B, Song X-L, Wang X, Wei D, et al. Molecular Mechanism of
33
34 16 Constitutive Endocytosis Of Acid-Sensing Ion Channel 1a and Its Protective Function in
35
36 17 Acidosis-Induced Neuronal Death. J Neurosci. 2013;33:7066–78.
37
38 18 65. Xiao B, Wang S, Yang G, Sun X, Zhao S, Lin L, et al. HIF-1 α contributes to hypoxia
39
40 19 adaptation of the naked mole rat. Oncotarget. 2017;8:109941–51.
41
42
43
44
45
46
47
48

49 **Figure Legends**

50
51
52 22 Fig. 1. Electrophysiological properties of NMR neurons differ from mouse neurons. NMR cortical
53
54 23 and hippocampal neurons have significantly smaller capacitance (**a**) and more hyperpolarized
55
56 24 resting membrane potential (**b**) than mouse neurons. **c**. Example of whole-cell patch-clamp
57
58 25 recordings made from a mouse (left, black) and a NMR (middle, blue) hippocampal neuron in
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 response to the voltage step protocol and the inhibition of response in NMR neurons by 300 nM
2 TTX (right, blue). Cortical and hippocampal NMR neurons have significantly smaller voltage-gated
3 inward currents (**a, b**), a more depolarized V_{half} (**c**) and a more depolarized peak activation
4 membrane potential V_m (**d**) compared to mouse neurons. **e**. Voltage-gated outward currents in
5 NMR cortical and hippocampal show no significant difference to those in mouse neurons. * $p <$
6 0.5 ; ** $p < 0.01$; **** $p < 0.0001$, unpaired t-tests within each structure. Numbers in brackets
7 indicate the number of recorded cells.

8
9 Fig. 2. ASICs mediate acid-induced currents in NMR and mouse CNS neurons. **a**. Both mouse
10 (black trace, left) and NMR (blue trace, right) neurons respond to a pH 5 solution with a transient
11 inward current. **b**. Acid-induced currents are of significantly smaller amplitude in NMR neurons
12 compared to mouse neurons. **c**. Inactivation time constants of the acid-induced responses are
13 similar between NMR and mouse neurons. **d**. Example trace of an acid-induced current elicited
14 by a pH 5 solution and the effect of 100 μ M benzamil in a mouse cortical neuron. **e-f**. Acid-induced
15 currents in both cortical and hippocampal mouse neurons were reversibly blocked by 100 μ M
16 benzamil ($n = 9$ and $n = 10$ cortical and hippocampal neurons, respectively). **g**. Example trace of
17 an acid-induced current evoked by a pH 5 solution in a cortical NMR neuron showing inhibition
18 by 100 μ M benzamil. **h-i**. In both cortical ($n = 7$) and hippocampal ($n = 6$) NMR neurons, acid-
19 induced currents were reversibly blocked by 100 μ M benzamil. ** $p < 0.01$; **** $p < 0.0001$, one-
20 way ANOVA paired tests, Tukey's multiple comparison tests. Numbers into brackets indicates the
21 number of recorded cells.

22
23 Fig. 3. NMR cortical neurons show resistance to acid-induced cell death. Mouse (**a**) and NMR (**b**)
24 cortical neurons were incubated for 2 hours with either a pH 7.4 or pH 5 solution and then stained
25 with Hoechst 33342 to label nuclei and PI to label necrotic dead cells. **c**. Percentages of PI-
26 positive neurons (i.e. dead) at pH 7.4 were similar between mouse and NMR neurons. The

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 percentage of mouse dead neurons was higher when cells were incubated in a pH 5 solution,
2 compared to pH 7.4 condition and compared to the percentage of cell death obtained with NMR
3 neurons at pH 5. NMR cortical neurons did not exhibit any significant difference with regard to cell
4 death when exposed to pH 5.0 compared to pH 7.4. **** p< 0.0001, one-way ANOVA test, Tukey's
5 multiple comparison test. Percentages of each field of view were used to compare acidotoxicity
6 between species. The number of fields of view analyzed for each condition are the following: n =
7 18 for mouse neurons (for both pH 7.4 and pH 5), n = 27 and 28 for NMR neurons at pH 7.4 and
8 at pH 5, respectively; obtained from 3 independent cultures for both species.





FIG. 1






FIG. 3