

The role of prenatal steroids at 34-36 weeks of gestation

Prof Gordon C.S. Smith, DSc

Department of Obstetrics and Gynaecology, University of Cambridge; NIHR Cambridge

Comprehensive Biomedical Research Centre, Box 223 The Rosie Hospital, Cambridge, UK.

Prof David H. Rowitch, ScD

Department of Paediatrics, Wellcome Trust-MRC Stem Cell Institute, University of Cambridge,

Cambridge, UK

Prof Ben W. Mol, PhD

Department of Obstetrics and Gynaecology, The Robinson Research Institute, School of Medicine,

University of Adelaide, 5000 SA Australia.

Correspondence to:

Prof Gordon C.S. Smith, Department of Obstetrics and Gynaecology, University of Cambridge; NIHR

Cambridge Comprehensive Biomedical Research Centre, Box 223 The Rosie Hospital, Cambridge, UK.

Tel: +44 (0)1223 336871, Fax: +44 (0)1223 215327, E-mail: gcss2@cam.ac.uk

Summary

A recent multi-centre, double blind, randomised, placebo controlled trial has demonstrated that administration of betamethasone to women with threatened preterm delivery at 34 to 36 weeks' of gestational age reduces the risk of neonatal respiratory morbidity. There is, however, no long term outcome data on the children and we believe that it is biologically plausible that this treatment may cause long term harm through effects on the infant's brain. Given this, we argue that steroids should not be used in the context of late preterm delivery until evidence of long term safety is available. This example illustrates some strengths and weaknesses of using "levels of evidence" to grade the empirical support for making clinical decisions.

One of the major advances in perinatal medicine in the last 30 years has been the administration of synthetic glucocorticoids to mothers who are likely to deliver at extreme preterm gestational ages, and this intervention clearly reduces perinatal mortality and severe morbidity.¹ A recent multi-centre, double blind, randomized controlled trial compared the effects of betamethasone versus placebo among women presenting between 34 and 36 weeks of gestational age with a high probability of delivery.² The primary outcome was need for respiratory support three days after delivery. The rate was 11.6% in steroid treated children and 14.4% with placebo, yielding a number needed to treat of 35 (95% CI 19 to 259). In addressing the question of whether this trial justifies immediate incorporation of glucocorticoids into the management of threatened delivery at 34 to 36 weeks, we need to consider both the science of glucocorticoids in pregnancy, and the science of clinical trials.

Physiologically, the fetus prepares for birth near term by an increased production of cortisol from the fetal adrenal. The glucocorticoids employed therapeutically to accelerate fetal lung maturation

are betamethasone and dexamethasone. The choice is purposeful as both are resistant to 11 β -hydroxysteroid dehydrogenase-2 (11 β -HSD-2),³ an enzyme which is highly expressed in the placenta and which limits transplacental passage of maternal cortisol, by converting it to inactive metabolites. Consequently, steroids which are metabolised by 11 β -HSD-2 (such as cortisol, hydrocortisone and prednisolone) cannot cross the placenta whereas 11 β -HSD-2-resistant steroids can. Importantly, the fetal brain has high levels of expression of 11 β -HSD-2, especially in areas which undergo rapid growth in the third trimester (e.g. the cerebellum).⁴ As a result, the brain is relatively protected from the effects of the physiological rise in endogenous cortisol in late gestation.⁵ It follows, however, that maternally administered betamethasone or dexamethasone near term will both cross the placenta and activate brain glucocorticoid receptors due to the resistance of these drugs to metabolism by 11 β -HSD-2. Hence, betamethasone and dexamethasone will cause un-physiological activation of glucocorticoid receptors in the fetal brain near term.

Should we be concerned? There are a number of observations that suggest we should. First, it has been shown 11 β -HSD-2 resistant glucocorticoids inhibited normal cerebellar growth when administered to neonatal mice⁶ and had profound and long lasting epigenetic effects on the brain of fetal guinea pigs following maternal administration in late gestation.⁷ Second, the one study that previously examined long term outcomes following betamethasone in late pregnancy (given prior to pre-labour caesarean section at 37 or 38 weeks¹ of gestational age) demonstrated that glucocorticoid-exposed children were twice as likely to be identified as being in the lowest achievement group at school compared with the control group (17.7% vs 8.5%, RR 2.1; 95% CI, 1.1-3.7).⁸ Interestingly, the same trial demonstrated no long term benefit of steroids. Part of the motivation for the original trial had been that prevention of short term respiratory morbidity around the time of birth may reduce the risk of respiratory disease in later childhood. However, the long term follow up study demonstrated no such benefits. Hence, the justification for glucocorticoids

prior to term pre-labour caesarean section is wholly based on reducing the risk of short term complications.

A number of studies have compared longer term follow up (in children¹ and adults⁹) in those exposed to antenatal glucocorticoids and placebo controls from the original RCTs of glucocorticoids in women presenting with threatened preterm birth. However, these studies generally assessed the effects of steroid exposure at earlier gestational ages. Overall, these results are reassuring and support the use of steroids in the management of preterm birth prior to 34 weeks.¹ However, the apparent long term safety of steroids following exposure at more extreme degrees of prematurity does not confirm their safety at later gestational ages. Potential direct harmful effects of steroids on the brain could be masked by the generally protective effect of improved neonatal cardiorespiratory function and reduced risks of specific acute perinatal complications, such as intra-ventricular haemorrhage. Moreover, activation of the glucocorticoid receptor by steroids is regulated by multiple aspects of the cellular and physiological context, termed "context-driven plasticity".¹⁰ It is possible – indeed likely – that the effect of a given concentration of a given steroid may differ at different gestational ages due to the multiple factors that change with advancing gestational age and the complex regulation of glucocorticoid receptor mediated responses.

In the context of extreme preterm birth (e.g. 24-28 weeks gestational age), the balance of risks and benefits favours the use of glucocorticoids, as they have clearly been shown to prevent death, and severe adverse outcome in the short and long term.¹ However, in our view, the case has not been established that the balance of risks and benefits favours their use after 33 weeks of gestational age. We know that use of steroids in women with threatened delivery at 34 to 36 weeks means that 35 children need to be treated to prevent one child requiring respiratory support three days after delivery. What we do not know is the long term effects of fetal exposure to these drugs at this

gestational age, i.e. whether the prevalence or severity of long term adverse effects – if there are any – outweigh the known benefit.

This example also illustrates a series of key points about randomized controlled trials in perinatal medicine: (i) there are multiple outcomes which may be affected by an intervention, (ii) the effect of an intervention might differ proportionally and directionally for different outcomes, (iii) the level of evidence around different effects may differ. In the case of late preterm glucocorticoids, their beneficial short term effect is supported by level 1 evidence. In contrast, the potential for adverse effects on the fetal brain is largely based on level 5 evidence, i.e. expert opinion and basic research. A relative lack of evidence on rare and long term outcomes is a general feature of trials, and reflects some practical realities: trials are expensive and the larger they are and the longer the follow up, the more expensive the trial. Consequently, trials are biased to be powered for outcomes which are common and occur in the short term. In contexts such as extreme preterm birth and cancer, the clinically most important outcomes, such as death, may be common and occur in the short term. However, in many other contexts, including late preterm delivery, common and short term complications are generally not severe, and the most important outcomes may be rare and/or long term, including harm done by the intervention. A danger of RCTs in obstetrics is that trials may not be funded or powered to detect long term adverse outcomes in children.

Ideally, the absence of level 1 evidence for all of the major outcomes of interest would not be a problem. The lack of information on long term outcomes, or underpowered analyses leading to wide confidence intervals, would be interpreted as indicating that there is no accurate information about whether these outcomes are affected by the intervention, and the absence of evidence would be a cause for concern. However, it has been suggested that the six most dangerous words in evidence-based medicine are: “there is no evidence to demonstrate”.¹¹ If the “absence of evidence” is conflated with “evidence of absence” we will be misled. The correct summary of using

glucocorticoids at 34 to 36 weeks is that there is evidence of short term benefit but an absence of evidence of long term safety. But there are those who might state that there is evidence of short term benefit and no evidence to demonstrate long term harm. The different wording yields a profound difference in inference.

Randomized trials have had a hugely positive effect on the practice of obstetrics and paediatrics. The strength of trials is that this study design has the highest internal validity, all other things being equal. But crucially, we cannot assume no effect for questions where there are limited or no data. Therefore, when we assess the evidence base we must consider all the issues. The problem is not trials. The problem is a simplistic approach to a complex question. Labelling evidence on the basis of study design alone, without reference to statistical power, particular attention to adequate length of follow up, harm, external validity, or biological plausibility is treating a complex question in a simplistic manner. We ignore complexity at our peril or, rather, at the peril of our patients.

Acknowledgements

We are very grateful to Professor Jim Thornton (Nottingham University, UK) for his helpful comments on an earlier draft of this manuscript.

Declaration of interests

GS receives/has received research support from GE, Roche and GSK. GS has been paid to attend advisory boards by GSK and Roche and to consult for GSK. GS has received support to attend a scientific meeting from Chiesi. GS is named inventor in a patent submitted by GSK for a novel application of an existing GSK compound for the prevention of preterm birth (PCT/EP2014/062602). GS is a member of a GSK Data Safety Monitoring Committee for a trial of RSV vaccination in pregnancy and infancy. BWM has been paid to attend advisory boards for ObsEva, Geneva and has presented at sponsored scientific meetings.

References

- (1) Roberts D, Dalziel S. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev* 2006; 3:CD004454.
- (2) Gyamfi-Bannerman C, Thom EA, Blackwell SC, Tita AT, Reddy UM, Saade GR et al. Antenatal Betamethasone for Women at Risk for Late Preterm Delivery. *N Engl J Med* 2016; 374:1311-1320.
- (3) Aiken CE, Fowden AL, Smith GC. Antenatal glucocorticoids prior to cesarean delivery at term. *JAMA Pediatr* 2014; 168(6):507-508.
- (4) Holmes MC, Sangra M, French KL, Whittle IR, Paterson J, Mullins JJ et al. 11beta-Hydroxysteroid dehydrogenase type 2 protects the neonatal cerebellum from deleterious effects of glucocorticoids. *Neuroscience* 2006; 137(3):865-873.
- (5) Wyrwoll CS, Holmes MC, Seckl JR. 11beta-hydroxysteroid dehydrogenases and the brain: from zero to hero, a decade of progress. *Front Neuroendocrinol* 2011; 32(3):265-286.
- (6) Heine VM, Rowitch DH. Hedgehog signaling has a protective effect in glucocorticoid-induced mouse neonatal brain injury through an 11betaHSD2-dependent mechanism. *J Clin Invest* 2009; 119(2):267-277.
- (7) Crudo A, Suderman M, Moisiadis VG, Petropoulos S, Kostaki A, Hallett M et al. Glucocorticoid programming of the fetal male hippocampal epigenome. *Endocrinology* 2013; 154(3):1168-1180.
- (8) Stutchfield PR, Whitaker R, Gliddon AE, Hobson L, Kotecha S, Doull IJ. Behavioural, educational and respiratory outcomes of antenatal betamethasone for term caesarean section (ASTECS trial). *Arch Dis Child Fetal Neonatal Ed* 2013; 98(3):F195-F200.
- (9) Dalziel SR, Lim VK, Lambert A, McCarthy D, Parag V, Rodgers A et al. Antenatal exposure to betamethasone: psychological functioning and health related quality of life 31 years after inclusion in randomised controlled trial. *BMJ* 2005; 331(7518):665.

- (10) Weikum ER, Knuesel MT, Ortlund EA, Yamamoto KR. Glucocorticoid receptor control of transcription: precision and plasticity via allostery. *Nat Rev Mol Cell Biol* 2017; 18(3):159-174.
- (11) Braithwaite RS. A piece of my mind. EBM's six dangerous words. *JAMA* 2013; 310(20):2149-2150.