

# SOME MINIMISATION ALGORITHMS IN ARITHMETIC INVARIANT THEORY

TOM FISHER AND LAZAR RADIČEVIĆ

**ABSTRACT.** We extend the work of Cremona, Fisher and Stoll on minimising genus one curves of degrees 2, 3, 4, 5, to some of the other representations associated to genus one curves, as studied by Bhargava and Ho. Specifically we describe algorithms for minimising bidegree (2, 2)-forms,  $3 \times 3 \times 3$  cubes and  $2 \times 2 \times 2$  hypercubes. We also prove a theorem relating the minimal discriminant to that of the Jacobian elliptic curve.

## 1. INTRODUCTION

Let  $F$  be a homogeneous polynomial in several variables with rational coefficients. Then making a linear change of variables and rescaling the polynomial by a rational number does not change the isomorphism class of the hypersurface defined by  $F$ . Thus a natural question is to find a change of variables and a rescaling of the polynomial so that its coefficients are small integers.

More generally we may consider the following situation. Let  $\mathcal{G}$  be a product of general linear groups, acting linearly on a  $\mathbb{Q}$ -vector space  $W$ . We fix a basis for  $W$ , and represent a vector  $w \in W$  by its vector of coordinates  $(w_1, \dots, w_N)$  relative to this basis. We refer to these co-ordinates as the *coefficients*. Then given  $w \in W$  we seek to find  $g \in \mathcal{G}(\mathbb{Q})$  such that  $g \cdot w$  has small integer coefficients.

An *invariant* is a polynomial  $I \in \mathbb{Z}[w_1, \dots, w_N]$  such that:

$$I(g \cdot w) = \chi(g)I(w)$$

for all  $g \in \mathcal{G}(\mathbb{C})$  and  $w \in W$ , where  $\chi$  is a rational character on  $\mathcal{G}$  (i.e. a product of determinants). In practice there will be an invariant  $\Delta$ , which we call the *discriminant*, and the elements  $w \in W$  of interest will be those with  $\Delta(w) \neq 0$ . We note that if  $w$  has integer coefficients then  $\Delta(w)$  is an integer. Our strategy is to first find  $g \in \mathcal{G}(\mathbb{Q})$  making this discriminant as small as possible (in absolute value). This is known as *minimisation*.

---

2010 *Mathematics Subject Classification.* 11G05, 11G07, 14H52.

*Key words and phrases.* elliptic curves, genus one curves, minimisation, coregular representations.

This is a local problem, in that for each prime  $p$  dividing  $\Delta(w)$  we seek to minimise the  $p$ -adic valuation  $v_p(\Delta(w))$ , without changing the valuations at the other primes. Once we've minimised the discriminant, the next step is to find a transformation in  $\mathcal{G}(\mathbb{Z})$ , making the coefficients as small as possible. This is known as *reduction*.

This strategy has been carried out in [3] and [4], for the models (i.e. collections of polynomials) defining genus one curves of degrees 2, 3, 4 and 5. In these cases the invariants give a Weierstrass equation for the Jacobian of the genus one curve. In this article, we extend these techniques to some of the other representation associated to genus one curves, as studied in [1]. Specifically we describe algorithms for minimising bidegree (2, 2)-forms,  $3 \times 3 \times 3$  cubes and  $2 \times 2 \times 2 \times 2$  hypercubes. In each of these cases the invariants define not only the Jacobian elliptic curve  $E$ , but also one or two marked points on  $E$ . One possible application of these algorithms is in computing the Cassels-Tate pairing (see [5]).

As explained below, each (2, 2)-form  $F$  determines a pair of binary quartics  $G_1, G_2$ , each  $3 \times 3 \times 3$  cube  $S$  determines a triple of ternary cubics  $F_1, F_2, F_3$ , and each  $2 \times 2 \times 2 \times 2$  hypercube  $H$  determines a quadruple of binary quartics  $G_1, \dots, G_4$ . Therefore a natural approach would be to minimise and reduce the corresponding binary quartics and ternary cubics, using the algorithms in [3], and then apply the transformations that arise in this way to  $F$ ,  $S$  or  $H$ . This strategy works for reduction (which we therefore do not study further in this article), but not for minimisation. For example if  $F \in \mathbb{Z}[x_1, x_2; y_1, y_2]$  is a (2, 2)-form with  $F \equiv x_2^2 y_2^2 \pmod{p^2}$  then the binary quartics  $G_1$  and  $G_2$  vanish mod  $p^2$ . The algorithm for minimising binary quartics says that we should divide each  $G_i$  by  $p^2$ . However this information on its own does not tell us how to minimise  $F$ .

Since minimisation is a local problem, we work in the following setting. Let  $K$  be a field with a discrete valuation  $v : K^\times \rightarrow \mathbb{Z}$ . We write  $\mathcal{O}_K$  for the valuation ring, and  $\pi$  for a uniformiser, i.e. an element  $\pi \in K$  with  $v(\pi) = 1$ . The residue field is  $k = \mathcal{O}_K/\pi\mathcal{O}_K$ . For example we could take  $K = \mathbb{Q}$  or  $\mathbb{Q}_p$ , and  $v = v_p$  the  $p$ -adic valuation. In these cases  $\mathcal{O}_K = \mathbb{Z}_{(p)}$  or  $\mathbb{Z}_p$ . We make no restrictions on the characteristics of  $K$  and  $k$ .

Since it serves as a prototype for our work, we briefly recall the algorithm for minimising binary quartics. See [3] for further details. A binary quartic is a homogeneous polynomial of degree 4 in two variables:

$$G(x_1, x_2) = ax_1^4 + bx_1^3x_2 + cx_1^2x_2^2 + dx_1x_2^3 + ex_2^4.$$

If  $R$  is any ring then there is an action of  $\mathcal{G}(R) = R^\times \times \mathrm{GL}_2(R)$  on the space of binary quartics over  $R$  via

$$(1) \quad \left[ \lambda, \begin{pmatrix} r & s \\ t & u \end{pmatrix} \right] : G(x_1, x_2) \mapsto \lambda^2 G(rx_1 + tx_2, sx_1 + ux_2).$$

We say that binary quartics are *R-equivalent* if they belong to the same orbit for this action. A polynomial  $I \in \mathbb{Z}[a, b, c, d, e]$  is an invariant of weight  $p$  if

$$I([\lambda, A] \cdot G) = (\lambda \det A)^p I(G)$$

for all  $[\lambda, A] \in \mathcal{G}(\mathbb{C})$ . The ring of invariants of a binary quartic is generated (in characteristics not 2 or 3) by

$$\begin{aligned} I &= 12ae - 3bd + c^2 \\ J &= 72ace - 27ad^2 - 27b^2e + 9bcd - 2c^3 \end{aligned}$$

of weights 4 and 6. The discriminant  $\Delta = (4I^3 - J^2)/27$  is an invariant of weight 12.

A binary quartic  $G$  is *integral* if it has coefficients in  $\mathcal{O}_K$ , and *non-singular* if  $\Delta(G) \neq 0$ . We write  $v(G)$  for the minimum of the valuations of the coefficients of  $G$ . Given a non-singular binary quartic, we seek to find a  $K$ -equivalent integral binary quartic  $G$  with  $v(\Delta(G))$  as small as possible.

We write  $\tilde{G}$  for the reduction of  $\pi^{-v(G)}G \bmod \pi$ . If an integral binary quartic  $G(x_1, x_2)$  is non-minimal, then it is  $\mathcal{O}_K$ -equivalent to a binary quartic with

$$G(x_1, \pi^s x_2) \equiv 0 \pmod{\pi^{2s+2}}$$

for some integer  $s \geq 0$ . The least such integer  $s$  is called the *slope*, and can only take values 0, 1 and 2. If  $v(G) \leq 1$  (i.e. the slope is positive) then  $\tilde{G}$  has a unique multiple root, and if we move this root to  $(1 : 0)$  then  $\pi^{-2}G(x_1, \pi x_2)$  is an integral binary quartic with the same invariants, but with smaller slope. After at most two iterations we reach a form of slope 0. We can then divide through by  $\pi^2$ , and repeat the process until a minimal binary quartic is obtained.

Our algorithms for minimising  $(2, 2)$ -forms,  $3 \times 3 \times 3$  cubes and  $2 \times 2 \times 2$  hypercubes are described in Sections 2, 3 and 4. We also give formulae for the Jacobian elliptic curve and the marked points that work in all characteristics. (In [1] the authors worked over a field of characteristic not 2 or 3, and the formulae were not always given explicitly.) In Section 5 we prove a theorem about the minimal discriminant, and describe how it is improved by our minimisation algorithms.

## 2. BIDEGREE $(2, 2)$ -FORMS

A  $(2, 2)$ -form is a polynomial in  $x_1, x_2, y_1, y_2$ , that is homogeneous of degree 2 in both sets of variables. We can view a  $(2, 2)$ -form  $F$  as a binary quadratic form in  $y_1, y_2$  whose coefficients are binary quadratic forms in  $x_1, x_2$ :

$$F(x_1, x_2; y_1, y_2) = F_1(x_1, x_2)y_1^2 + F_2(x_1, x_2)y_1y_2 + F_3(x_1, x_2)y_2^2.$$

The discriminant  $G_1 = F_2^2 - 4F_1F_3$  is then a binary quartic in  $x_1, x_2$ . Switching the two sets of variables we may likewise define a binary quartic  $G_2$  in  $y_1, y_2$ . It may be checked that  $G_1$  and  $G_2$  have the same invariants  $I$  and  $J$ . We define  $c_4(F) = I$  and  $c_6(F) = J/2$ . The discriminant is  $\Delta(F) = (c_4^3 - c_6^2)/1728$ .

A non-zero  $(2, 2)$ -form  $F$  over a field defines a curve in  $\mathbb{P}^1 \times \mathbb{P}^1$ . If  $\Delta(F) \neq 0$  then this curve  $\mathcal{C}_F$  is a smooth curve of genus one. It may be written as a double cover of  $\mathbb{P}^1$  (ramified over the roots of  $G_1$  or  $G_2$ ) by projecting to either factor.

Let  $R$  be a ring. There is an action of  $\mathcal{G}(R) = R^\times \times \mathrm{GL}_2(R) \times \mathrm{GL}_2(R)$  on the space of  $(2, 2)$ -forms over  $R$  given by

$$[\lambda, A, B] : F(x_1, x_2; y_1, y_2) \mapsto \lambda F((x_1, x_2)A; (y_1, y_2)B).$$

We say that  $(2, 2)$ -forms are *R-equivalent* if they belong to the same orbit for this action. If  $[\lambda, A, B] \cdot F = F'$  then the binary quartics  $G_1$  and  $G_2$  determined by  $F$ , and the binary quartics  $G'_1$  and  $G'_2$  determined by  $F'$ , are related by

$$(2) \quad \begin{aligned} G'_1 &= [\lambda \det B, A] \cdot G_1 \\ G'_2 &= [\lambda \det A, B] \cdot G_2 \end{aligned}$$

where the action on binary quartics is that defined in (1).

We may represent  $F$  by a  $3 \times 3$  matrix via:

$$(3) \quad F(x_1, x_2; y_1, y_2) = \begin{pmatrix} x_1^2 & x_1x_2 & x_2^2 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} y_1^2 \\ y_1y_2 \\ y_2^2 \end{pmatrix}.$$

A polynomial  $I \in \mathbb{Z}[a_{ij}]$  is an *invariant* of weight  $p$  if

$$I([\lambda, A, B] \cdot F) = (\lambda \det A \det B)^p I(F)$$

for all  $[\lambda, A, B] \in \mathcal{G}(\mathbb{C})$ . In particular the polynomials  $c_4, c_6$  and  $\Delta$  are invariants of weights 4, 6 and 12. Over a field of characteristic not 2 or 3, the invariants determine a pair  $(E, P)$  where  $E$  is an elliptic curve (the Jacobian of  $\mathcal{C}_F$ ) and  $P$  is a marked point on  $E$ . See [1, Section 6.1.2]. The next lemma gives formulae for

$$E : \quad y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$$

and  $P = (\xi, \eta)$  that work in all characteristics.

**Lemma 2.1.** *There exist  $\xi, \eta, a_1, a_2, a_3, a_4, a_6 \in \mathbb{Z}[a_{ij}]$  such that*

- (i) *We have  $c_4 = b_2^2 - 24b_4$  and  $c_6 = -b_2^3 + 36b_2b_4 - 216b_6$ , where  $b_2 = a_1^2 + 4a_2$ ,  $b_4 = a_1a_3 + 2a_4$  and  $b_6 = a_3^2 + 4a_6$ ,*

- (ii) The polynomials  $u = 12\xi + a_1^2 + 4a_2$  and  $v = 2\eta + a_1\xi + a_3$  are invariants of weights 2 and 3 satisfying  $(108v)^2 = (3u)^3 - 27c_4(3u) - 54c_6$ .
- (iii) We have  $\eta^2 + a_1\xi\eta + a_3\eta = \xi^3 + a_2\xi^2 + a_4\xi + a_6$ .

*Proof.* We put  $\xi = a_{11}a_{33} + a_{13}a_{31}$  and  $\eta = a_{11}a_{22}a_{33}$ .

(i) We put

$$\begin{aligned} a_1 &= -a_{22}, \\ a_2 &= -(a_{11}a_{33} + a_{12}a_{32} + a_{13}a_{31} + a_{21}a_{23}), \\ a_3 &= a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33}. \end{aligned}$$

Since we already defined  $c_4$  and  $c_6$ , we may solve for  $a_4$  and  $a_6$ . We find that these too are polynomials in the  $a_{ij}$  with integer coefficients.

(ii) The invariants  $u$  and  $v$  were denoted  $\delta_2$  and  $\delta_3$  in [1, Section 6.1.2]. In fact we have  $v = \det(a_{ij})$ .

(iii) This follows from (i) and (ii), exactly as in [6, Chapter III].  $\square$

Let  $(E, P)$  be a pair consisting of an elliptic curve  $E/K$  and a point  $0_E \neq P \in E(K)$ . On a minimal Weierstrass equation for  $E$ , the point  $P$  has co-ordinates  $(x_P, y_P)$ , where either  $x_P, y_P \in \mathcal{O}_K$  or  $v(x_P) = -2r$ ,  $v(y_P) = -3r$  for some integer  $r \geq 1$ . We define  $\kappa(P) = 0$  in the first case, and  $\kappa(P) = r$  in the second. We write  $\Delta_E$  for the minimal discriminant of  $E$ .

We say that a  $(2, 2)$ -form  $F$  is *integral* if it has coefficients in  $\mathcal{O}_K$ , and *non-singular* if  $\Delta(F) \neq 0$ .

**Lemma 2.2.** *Let  $F$  be a non-singular integral  $(2, 2)$ -form. Let  $(E, P)$  be the pair specified in Lemma 2.1. Then*

$$v(\Delta(F)) = v(\Delta_E) + 12\kappa(P) + 12\ell(F)$$

where  $\ell(F) \geq 0$  is an integer we call the level.

*Proof.* The formulae in Lemma 2.1 give an integral Weierstrass equation  $W$  for  $E$ , upon which  $P$  is a point with integral coordinates. The smallest possible discriminant of such an equation is  $v(\Delta_E) + 12\kappa(P)$ . Since the discriminant of  $F$  is equal to the discriminant of  $W$ , the result follows.  $\square$

In this section we give an algorithm for minimising  $(2, 2)$ -forms. That is, given a non-singular  $(2, 2)$ -form  $F$  over  $K$ , we explain how to find a  $K$ -equivalent integral  $(2, 2)$ -form with level (equivalently, valuation of the discriminant) as small as possible. In Section 5 we show that if  $\mathcal{C}_F(K) \neq \emptyset$  then the minimal level is zero.

By clearing denominators, we may start with an integral  $(2, 2)$ -form. If this form is  $K$ -equivalent to an integral form of smaller level, then our task

is to find such a form explicitly. Define  $v(F)$  to be the minimum of the valuations of the coefficients of  $F$ . If  $v(F) \geq 1$  then we can divide through by  $\pi$ , reducing the level of  $F$ . We may therefore assume that  $v(F) = 0$ .

Our algorithm for minimising  $(2, 2)$ -forms is described by the following theorem.

**Theorem 2.3.** *Let  $F$  be a non-minimal  $(2, 2)$ -form with  $v(F) = 0$ . Let  $f$  be the reduction of  $F \bmod \pi$ . Then we are in one of the following three situations.*

- (i) *The form  $f$  factors as a product of binary quadratic forms, both of which have a repeated root. By an  $\mathcal{O}_K$ -equivalence we may assume that  $f = x_2^2 y_2^2$ . Then at least one of the forms*

$$\begin{aligned} \pi^{-2}F(x_1, \pi x_2; y_1, y_2) \\ \pi^{-2}F(x_1, x_2; y_1, \pi y_2) \\ \pi^{-3}F(x_1, \pi x_2; y_1, \pi y_2) \end{aligned}$$

*is an integral  $(2, 2)$ -form of smaller level.*

- (ii) *The form  $f$  factors as a product of binary quadratic forms, exactly one of which has a repeated root. By an  $\mathcal{O}_K$ -equivalence, and switching the two sets of variables if necessary, we may assume that  $f = x_2^2 h(y_1, y_2)$ . Then  $\pi^{-1}F(x_1, \pi x_2; y_1, y_2)$  is an integral  $(2, 2)$ -form of the same level.*

- (iii) *The curve  $\mathcal{C}_f \subset \mathbb{P}^1 \times \mathbb{P}^1$  has a unique singular point. By an  $\mathcal{O}_K$ -equivalence, we may assume this is the point  $((1 : 0), (1 : 0))$ . Then  $\pi^{-2}F(x_1, \pi x_2; y_1, \pi y_2)$  is an integral  $(2, 2)$ -form of the same level.*

*Moreover the  $(2, 2)$ -form  $F$  computed in (ii) or (iii) either has  $v(F) \geq 1$  or has reduction mod  $\pi$  of the form specified in (i).*

**Remark 2.4.** Let  $F$  be an integral  $(2, 2)$ -form, with associated binary quartics  $G_1$  and  $G_2$ . It is clear by (2) that if either  $G_1$  or  $G_2$  is minimal then  $F$  is minimal. However the converse is not true. For example if  $F \equiv (x_1 y_1 + x_2 y_2)^2 \pmod{\pi^2}$ , then  $F$  is minimal by Theorem 2.3, yet we have  $G_1 \equiv G_2 \equiv 0 \pmod{\pi^2}$ .

Exactly as in the case of binary quartics, any non-minimal  $(2, 2)$ -form  $F$  is  $\mathcal{O}_K$ -equivalent to a form whose level can be reduced using diagonal transformations. Indeed, suppose that  $[\lambda, A_1, A_2] \in \mathcal{G}(K)$  is a transformation reducing the level. By clearing denominators, we may assume that the  $A_i$  have entries in  $\mathcal{O}_K$ , not all in  $\pi\mathcal{O}_K$ . Then writing these matrices in Smith normal form we have  $A_i = Q_i D_i P_i$  where  $P_i, Q_i \in \mathrm{GL}_2(\mathcal{O}_K)$  and

$$D_1 = \begin{pmatrix} 1 & 0 \\ 0 & \pi^a \end{pmatrix}, \quad D_2 = \begin{pmatrix} 1 & 0 \\ 0 & \pi^b \end{pmatrix},$$

for some integers  $a, b \geq 0$ . Replacing  $F$  by an  $\mathcal{O}_K$ -equivalent form, it follows that

$$\pi^{-a-b-1}F(x_1, \pi^a x_2; y_1, \pi^b y_2)$$

is an integral  $(2, 2)$ -form. We say that the pair  $(a, b)$  is *admissible* for  $F$ .

**Lemma 2.5.** *Let  $F$  be an integral  $(2, 2)$ -form. If some pair  $(a, b)$  is admissible for  $F$  then at least one of the following pairs is admissible:*

$$(0, 0), (1, 0), (0, 1), (1, 1), (2, 1), (1, 2).$$

*Proof.* The coefficients of  $F$ , arranged as in (3), have valuations satisfying

$$\begin{array}{lll} \geq a + b + 1 & \geq a + 1 & \geq a - b + 1 \\ \geq b + 1 & \geq 1 & \geq -b + 1 \\ \geq -a + b + 1 & \geq -a + 1 & \geq -a - b + 1. \end{array}$$

Conversely, if the valuations satisfy these inequalities then the pair  $(a, b)$  is admissible. If  $a = b = 0$  then we are done as  $(0, 0)$  is on the list. If  $a \geq 1, b = 0$  or  $a = 0, b \geq 1$ , then  $(1, 0)$  or  $(0, 1)$  is admissible. If  $a = b > 0$ , then  $(1, 1)$  is admissible. If  $a > b > 0$  or  $b > a > 0$ , then  $(2, 1)$  or  $(1, 2)$  is admissible.  $\square$

*Proof of Theorem 2.3.* For the proof we are free to replace the  $(2, 2)$ -form  $F$  by an  $\mathcal{O}_K$ -equivalent form. Indeed the transformations specified in the statement of the theorem induce well-defined maps on  $\mathcal{O}_K$ -equivalence classes, as may be verified using [3, Lemma 4.1]. We may therefore assume that one of the pairs  $(a, b)$  listed in Lemma 2.5 is admissible for  $F$ . Since  $v(F) = 0$  we cannot have  $a = b = 0$ . By switching the two sets of variables, we may assume that  $a \geq b$ . This leaves us with three cases. In considering each case, it is our running assumption that we are not in an earlier case.

**Case 1.** We assume  $(1, 0)$  is admissible for  $F$ . The coefficients of  $F$  have valuations satisfying

$$\begin{array}{lll} \geq 2 & \geq 2 & \geq 2 \\ \geq 1 & \geq 1 & \geq 1 \\ \geq 0 & \geq 0 & \geq 0 \end{array}$$

We have  $f = x_2^2 h(y_1, y_2)$  where  $h$  is a binary quadratic form. If  $h$  has a repeated root, then the first transformation in (i) decreases the level. Otherwise the transformation in (ii) gives a  $(2, 2)$ -form  $F$  with  $v(F) \geq 1$ .

**Case 2.** We assume  $(1, 1)$  is admissible for  $F$ . The coefficients of  $F$  have valuations satisfying

$$\begin{array}{ccc} \geq 3 & \geq 2 & \geq 1 \\ \geq 2 & \geq 1 & \geq 0 \\ \geq 1 & \geq 0 & \geq 0 \end{array}$$

We have

$$f = x_2 y_2 (\alpha x_1 y_2 + \beta x_2 y_1 + \gamma x_2 y_2)$$

for some  $\alpha, \beta, \gamma \in k$ . If  $\alpha = \beta = 0$  then the third transformation in (i) decreases the level. If exactly one of the coefficients  $\alpha$  and  $\beta$  is zero then the transformation in (ii) gives a  $(2, 2)$ -form whose reduction mod  $\pi$  is either zero, or of the form specified in (i). If  $\alpha$  and  $\beta$  are both non-zero then  $\mathcal{C}_f \subset \mathbb{P}^1 \times \mathbb{P}^1$  has a unique singular point at  $((1 : 0), (1 : 0))$ . The transformation in (iii) gives a  $(2, 2)$ -form  $F$  with  $v(F) \geq 1$ .

**Case 3.** We assume  $(2, 1)$  is admissible for  $F$ . The coefficients of  $F$  have valuations satisfying

$$\begin{array}{ccc} \geq 4 & \geq 3 & \geq 2 \\ \geq 2 & \geq 1 & = 0 \\ = 0 & \geq 0 & \geq 0 \end{array}$$

The two valuations indicated are zero, as we would otherwise be in Case 1 or Case 2. A calculation shows that  $\mathcal{C}_f \subset \mathbb{P}^1 \times \mathbb{P}^1$  has a unique singular point at  $((1 : 0), (1 : 0))$ . The transformation in (iii) gives a  $(2, 2)$ -form whose reduction mod  $\pi$  is of the form specified in (i).  $\square$

The following lemma will be needed in Section 4, in connection with our study of  $2 \times 2 \times 2 \times 2$  hypercubes.

**Lemma 2.6.** *Let  $F$  be a non-minimal  $(2, 2)$ -form, and let  $f = F \bmod \pi$ .*

(i) *If  $\mathcal{C}_f \subset \mathbb{P}^1 \times \mathbb{P}^1$  is singular at  $((1 : 0), (1 : 0))$ , then the coefficients of  $F$  have valuations satisfying*

$$\begin{array}{ccc} \geq 1 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & \geq 0 \\ \geq 0 & \geq 0 & \geq 0 \end{array} \quad \text{or} \quad \begin{array}{ccc} \geq 1 & \geq 1 & \geq 0 \\ \geq 1 & \geq 1 & \geq 0 \\ \geq 1 & \geq 0 & \geq 0 \end{array}$$

(ii) *If  $f = x_2^2 y_2^2$  then the coefficients of  $F$  have valuations satisfying*

$$\begin{array}{ccc} \geq 2 & \geq 2 & \geq 2 \\ \geq 1 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & = 0 \end{array} \quad \text{or} \quad \begin{array}{ccc} \geq 2 & \geq 1 & \geq 1 \\ \geq 2 & \geq 1 & \geq 1 \\ \geq 2 & \geq 1 & = 0 \end{array} \quad \text{or} \quad \begin{array}{ccc} \geq 3 & \geq 2 & \geq 1 \\ \geq 2 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & = 0 \end{array}$$

*Proof.* (i) The singular point forces  $a_{11} \equiv a_{12} \equiv a_{21} \equiv 0 \pmod{\pi}$ . The vanishing of the invariants  $u$  and  $v$  in Lemma 2.1 gives

$$8a_{13}a_{31} + a_{22}^2 \equiv a_{13}a_{22}a_{31} \equiv 0 \pmod{\pi}.$$



It follows that  $a_{22} \equiv 0 \pmod{\pi}$ . The same lemma shows that  $(\xi, \eta) = (a_{13}a_{31}, 0)$  is a singular point on the curve with Weierstrass equation  $y^2 \equiv x^2(x - a_{13}a_{31}) \pmod{\pi}$ . Therefore  $a_{13}a_{31} \equiv 0 \pmod{\pi}$ .

(ii) The proof of Theorem 2.3 shows that  $F$  is  $\mathcal{O}_K$ -equivalent to a  $(2, 2)$ -form  $F_1$  with

$$(4) \quad F_1(x_1, \pi^a x_2; y_1 \pi^b y_2) \equiv 0 \pmod{\pi^{a+b+1}}$$

for some  $(a, b) = (1, 0), (0, 1)$  or  $(1, 1)$ . Working mod  $\pi$  we have  $F_1 \equiv x_2^2 h(y_1, y_2), g(x_1, x_2) y_2^2$  or  $x_2 y_2 (\alpha x_1 y_2 + \beta x_2 y_1 + \gamma x_2 y_2)$ . In the last case it follows from our assumption  $F \equiv x_2^2 y_2^2 \pmod{\pi}$  that  $\alpha = \beta = 0$ . The equivalence relating  $F$  and  $F_1$  must now fix the points  $(x_1 : x_2) = (1 : 0) \pmod{\pi}$ ,  $(y_1 : y_2) = (1 : 0) \pmod{\pi}$ , or both. It follows that  $F$  also satisfies (4).  $\square$

### 3. $3 \times 3 \times 3$ RUBIK'S CUBES

We consider polynomials in  $x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3$  that are linear in each of the three sets of variables. Such a form may be represented as

$$\sum_{1 \leq i, j, k \leq 3} s_{ijk} x_i y_j z_k$$

where  $S = (s_{ijk})$  is a  $3 \times 3 \times 3$  cubical matrix. A *Rubik's cube*  $S$  may be partitioned into three  $3 \times 3$  matrices in three distinct ways:

- (i)  $M^1 = (s_{1jk})$  is the front face,  $N^1 = (s_{2jk})$  is the middle slice and  $P^1 = (s_{3jk})$  is the back face.
- (ii)  $M^2 = (s_{i1k})$  is the top face,  $N^2 = (s_{i2k})$  is the middle slice and  $P^2 = (s_{i3k})$  is the bottom face.
- (iii)  $M^3 = (s_{ij1})$  is the left face,  $N^3 = (s_{ij2})$  is the middle slice and  $P^3 = (s_{ij3})$  is the right face.

To each slicing  $(M^i, N^i, P^i)$ , we may associate a ternary cubic form

$$F_i(x, y, z) = \det(M^i x + N^i y + P^i z).$$

Following [3, Section 2] we scale the invariants  $c_4, c_6, \Delta$  of a ternary cubic so that  $c_4(xyz) = 1$ ,  $c_6(xyz) = -1$  and  $c_4^3 - c_6^2 = 1728\Delta$ . It may be checked that the  $F_i$  have the same invariants. We define  $c_4(S) = c_4(F_i)$ ,  $c_6(S) = c_6(F_i)$  and  $\Delta(S) = \Delta(F_i)$ .

If  $S$  is defined over a field and  $\Delta(S) \neq 0$  then each of the  $F_i$  defines a smooth curve of genus 1 in  $\mathbb{P}^2$ . These curves are isomorphic, although not in a canonical way. (See [1, Section 3.2] for further details.) We write  $\mathcal{C}_S$  to denote any one of them.

Let  $R$  be a ring. For each  $1 \leq i \leq 3$  there is an action of  $\mathrm{GL}_3(R)$  on the space of Rubik's cubes over  $R$  given by

$$A = (a_{ij}) : (M^i, N^i, P^i) \mapsto (a_{11}M^i + a_{12}N^i + a_{13}P^i, \\ a_{21}M^i + a_{22}N^i + a_{23}P^i, a_{31}M^i + a_{32}N^i + a_{33}P^i).$$

These actions commute, and so give an action of  $\mathcal{G}(R) = \mathrm{GL}_3(R)^3$ . We say that  $3 \times 3 \times 3$  cubes are *R-equivalent* if they belong to the same orbit for this action. If  $[A_1, A_2, A_3] \cdot S = S'$  then the associated ternary cubics are related by

$$(5) \quad F'_i(x, y, z) = \det(A_j A_k) F_i((x, y, z) A_i)$$

where  $\{i, j, k\} = \{1, 2, 3\}$ .

A polynomial  $I \in \mathbb{Z}[s_{ijk}]$  is an *invariant* of weight  $p$  if

$$I([A_1, A_2, A_3] \cdot S) = (\det A_1 \det A_2 \det A_3)^p I(S)$$

for all  $[A_1, A_2, A_3] \in \mathcal{G}(\mathbb{C})$ . In particular the polynomials  $c_4, c_6$  and  $\Delta$  are invariants of weights 4, 6 and 12. Over a field of characteristic not 2 or 3, the invariants determine a pair  $(E, P)$  where  $E$  is an elliptic curve (the Jacobian of  $\mathcal{C}_S$ ) and  $P$  is a marked point on  $E$ . See [1, Proposition 5.5]. The next lemma gives formulae for  $E$  and  $P$  that work in all characteristics.

**Lemma 3.1.** *There exist  $\xi, \eta, a_1, a_2, a_3, a_4, a_6 \in \mathbb{Z}[s_{ijk}]$  such that*

- (i) *We have  $c_4 = b_2^2 - 24b_4$  and  $c_6 = -b_2^3 + 36b_2b_4 - 216b_6$ , where  $b_2 = a_1^2 + 4a_2$ ,  $b_4 = a_1a_3 + 2a_4$  and  $b_6 = a_3^2 + 4a_6$ ,*
- (ii) *The polynomials  $u = 12\xi + a_1^2 + 4a_2$  and  $v = 2\eta + a_1\xi + a_3$  are invariants of weights 2 and 3 satisfying  $(108v)^2 = (3u)^3 - 27c_4(3u) - 54c_6$ .*
- (iii) *We have  $\eta^2 + a_1\xi\eta + a_3\eta = \xi^3 + a_2\xi^2 + a_4\xi + a_6$ .*

*Proof.* We define matrices  $A, B, C$  by the rule

$$(\mathrm{adj}(\lambda N^1 + \mu P^1))M^1 = \lambda^2 A + \lambda\mu B + \mu^2 C.$$

We put  $\xi = -\mathrm{tr}(AC)$  and  $\eta = -\mathrm{tr}(CBA)$ .

(i) We put

$$a_1 = \mathrm{tr}(B), \\ a_2 = \mathrm{tr}(AC) + \mathrm{tr}(A)\mathrm{tr}(C) - \mathrm{tr}(\mathrm{adj}(B)), \\ a_3 = \mathrm{tr}(ABC) + \mathrm{tr}(CBA) + \mathrm{tr}(AC)\mathrm{tr}(B).$$

Since we already defined  $c_4$  and  $c_6$ , we could now in principle solve for  $a_4$  and  $a_6$ . However it is simpler to argue as follows. Let  $a'_1, \dots, a'_6$  be the

$a$ -invariants (as defined in [3, Lemma 2.9]) of the ternary cubic  $F_1$ . We checked by computer algebra that there exist  $r, s, t \in \mathbb{Z}[s_{ijk}]$  satisfying

$$\begin{aligned} a'_1 &= a_1 + 2s, \\ a'_2 &= a_2 - sa_1 + 3r - s^2, \\ a'_3 &= a_3 + ra_1 + 2t. \end{aligned}$$

It follows by the transformation formulae for Weierstrass equations (see [6]) that  $a_4, a_6 \in \mathbb{Z}[s_{ijk}]$ . Note that our reason for working with  $a_1, \dots, a_6$ , in preference to  $a'_1, \dots, a'_6$ , is that this helped us find particularly simple expressions for  $\xi$  and  $\eta$ .

(ii) The invariants  $u$  and  $v$  were denoted  $4c_6$  and  $c_9$  in [1, Section 5.1.3]. In fact we have  $v = \text{tr}(ABC) - \text{tr}(CBA)$ .

(iii) This follows from (i) and (ii) exactly as in [6, Chapter III].  $\square$

A Rubik's cube  $S$  is *integral* if it has coefficients in  $\mathcal{O}_K$ , and *non-singular* if  $\Delta(S) \neq 0$ .

**Lemma 3.2.** *Let  $S$  be a non-singular integral Rubik's cube. Let  $(E, P)$  be the pair specified in Lemma 3.1. Then*

$$v(\Delta(S)) = v(\Delta_E) + 12\kappa(P) + 12\ell(S)$$

where  $\ell(S) \geq 0$  is an integer we call the level.

*Proof.* The proof is identical to that of Lemma 2.2.  $\square$

In this section we give an algorithm for minimising Rubik's cubes. In Section 5 we show that if  $\mathcal{C}_S(K) \neq \emptyset$  then the minimal level is zero.

We say that an integral cube  $S$  is *saturated* if for each  $i = 1, 2, 3$  the matrices  $M^i, N^i, P^i \in \text{Mat}_3(\mathcal{O}_K)$  are linearly independent mod  $\pi$ . If an integral cube is not saturated, then it is obvious how we may decrease the level.

Our algorithm for minimising  $3 \times 3 \times 3$  cubes is described by the following theorem.

**Theorem 3.3.** *Let  $S$  be a non-minimal saturated Rubik's cube. Let  $F_1, F_2, F_3$  be the associated ternary cubics, and  $f_1, f_2, f_3$  their reductions mod  $\pi$ . Then we are in one of the following two situations.*

- (i) *Two or more of the  $f_i$  are non-zero and have a repeated linear factor, say  $f_1$  and  $f_2$  are divisible by  $z^2$ . We apply a transformation*

$$\left[ \left( \begin{array}{ccc} 1 & & \\ & 1 & \\ & & \pi \end{array} \right), \left( \begin{array}{ccc} 1 & & \\ & 1 & \\ & & \pi \end{array} \right), A_3 \right]$$

where  $A_3 \in \text{GL}_3(K)$  is chosen such that the slices  $M^3, N^3, P^3 \in \text{Mat}_3(\mathcal{O}_K)$  of the transformed  $S$  are linearly independent mod  $\pi$ .

- (ii) Two or more of the  $f_i$  define a curve with a unique singular point, say  $f_1$  and  $f_2$  define curves with singular points at  $(1 : 0 : 0)$ . We apply a transformation

$$\left[ \begin{pmatrix} 1 & & \\ & \pi & \\ & & \pi \end{pmatrix}, \begin{pmatrix} 1 & & \\ & \pi & \\ & & \pi \end{pmatrix}, A_3 \right]$$

where  $A_3 \in \mathrm{GL}_3(K)$  is chosen such that the slices  $M^3, N^3, P^3 \in \mathrm{Mat}_3(\mathcal{O}_K)$  of the transformed  $S$  are linearly independent mod  $\pi$ .

The procedures in (i) and (ii) give an integral cube of the same or smaller level. Repeating these procedures either gives a non-saturated cube or decreases the level after at most three iterations.

**Remark 3.4.** Let  $S$  be an integral Rubik's cube, with associated ternary cubics  $F_1, F_2, F_3$ . It is clear by (5) that if any of the  $F_i$  are minimal then  $S$  is minimal. However the converse is not true. For example if  $S \equiv (\varepsilon_{ijk}) \pmod{\pi}$ , where  $\varepsilon_{ijk}$  is the Levi-Civita symbol (as appears in the definition of the cross product), then  $S$  is minimal by Theorem 3.3, yet we have  $F_1 \equiv F_2 \equiv F_3 \equiv 0 \pmod{\pi}$ .

Exactly as in the case of  $(2, 2)$ -forms, any non-minimal Rubik's cube  $S$  is  $\mathcal{O}_K$ -equivalent to a cube whose level can be reduced using diagonal transformations. Indeed, suppose that  $[\pi^{-s}A_1, A_2, A_3] \in \mathcal{G}(K)$  is a transformation reducing the level. By clearing denominators, we may assume that the  $A_i$  have entries in  $\mathcal{O}_K$ , not all in  $\pi\mathcal{O}_K$ . Then writing these matrices in Smith normal form we have  $A_i = Q_i D_i P_i$  where  $P_i, Q_i \in \mathrm{GL}_3(\mathcal{O}_K)$  and

$$D_i = \begin{pmatrix} \pi^{a_{1i}} & 0 & 0 \\ 0 & \pi^{a_{2i}} & 0 \\ 0 & 0 & \pi^{a_{3i}} \end{pmatrix}$$

with  $\min(a_{1i}, a_{2i}, a_{3i}) = 0$ . If this transformation reduces the level then  $\sum a_{ij} < 3s$ . In fact, by increasing one of the  $a_{ij}$ , we may assume that  $\sum a_{ij} = 3s - 1$ . We will from now on assume that  $a_{11} = a_{12} = a_{13} = 0$ . If the new cube has coefficients in  $\mathcal{O}_K$  then we say that the tuple  $(a_{21}, a_{31}; a_{22}, a_{32}; a_{23}, a_{33})$  is *admissible* for  $S$ .

**Lemma 3.5.** *Let  $S$  be a non-minimal Rubik's cube. Then after permuting the three slicings, and replacing  $S$  by an  $\mathcal{O}_K$ -equivalent cube, at least one of the following tuples is admissible.*

$$\begin{aligned} \tau_1 &= (1, 1; 0, 0; 0, 0), & \tau_2 &= (0, 1; 0, 1; 0, 0), & \tau_3 &= (1, 2; 0, 1; 0, 1), \\ \tau_4 &= (1, 1; 1, 1; 0, 1), & \tau_5 &= (1, 2; 1, 2; 1, 1), & \tau_6 &= (2, 3; 1, 2; 1, 2). \end{aligned}$$

*Proof.* We define the set of *weights*

$$\mathcal{W} = \left\{ (A, s) \in \text{Mat}_3(\mathbb{Z}) \times \mathbb{Z} \mid \begin{array}{l} a_{11} = a_{12} = a_{13} = 0, \\ a_{ij} \geq 0 \text{ for all } i, j, \\ \sum a_{ij} = 3s - 1 \end{array} \right\}.$$

If  $(A, s) \in \mathcal{W}$  then  $(a_{21}, a_{31}; a_{22}, a_{32}; a_{23}, a_{33})$  is admissible for  $S$  if and only if

$$v(s_{ijk}) \geq \max(s - a_{i1} - a_{j2} - a_{k3}, 0)$$

for all  $i, j, k \in \{1, 2, 3\}$ . We define a partial order on  $\mathcal{W}$  by  $(A, s) \leq (A', s')$  if

$$\max(s - a_{i1} - a_{j2} - a_{k3}, 0) \leq \max(s' - a'_{i1} - a'_{j2} - a'_{k3}, 0)$$

for all  $i, j, k \in \{1, 2, 3\}$ . A computer calculation, using Lemma 3.6 below, shows that  $(\mathcal{W}, \leq)$  has exactly 81 minimal elements. By an  $\mathcal{O}_K$ -equivalence we may assume that  $a_{2i} \leq a_{3i}$  for  $i = 1, 2, 3$ , and by permuting the three slicings of  $S$  we may assume that  $a_{31} \geq a_{32} \geq a_{33}$ . Only 8 of the 81 minimal elements satisfy these additional conditions. These are the 6 elements listed in the statement of the lemma, together with two more that are the same as  $\tau_4$  up to permuting the slicings.  $\square$

**Lemma 3.6.** *If  $(A, s) \in \mathcal{W}$  is minimal then  $s \leq 10$ .*

*Proof.* We suppose that  $(A, s)$  is minimal. Without loss of generality we have

$$(6) \quad a_{21} \leq a_{31}, \quad a_{22} \leq a_{32}, \quad a_{23} \leq a_{33} \quad \text{and} \quad a_{31} \geq a_{32} \geq a_{33}.$$

If  $a_{31} \leq 2$  then  $3s - 1 = \sum a_{ij} \leq 6a_{31} \leq 12$  and this gives the required bound on  $s$ . Otherwise we have  $a_{31} \geq 3$ . Let  $A'$  be the matrix obtained from  $A$  by replacing  $a_{31}$  by  $a_{31} - 3$ . Then  $(A', s - 1) \in \mathcal{W}$ , and by our minimality assumption  $(A', s - 1) \not\leq (A, s)$ . Therefore

$$\max(s - 1 - a'_{i1} - a'_{j2} - a'_{k3}, 0) > \max(s - a_{i1} - a_{j2} - a_{k3}, 0)$$

for some  $i, j, k \in \{1, 2, 3\}$ . Since we only changed the entry  $a_{31}$  we must have  $i = 3$  and  $s - 1 - (a_{31} - 3) > 0$ . Therefore

$$(7) \quad s + 1 \geq a_{31}.$$

The following inequalities are obtained in an entirely analogous way:

- (i) If  $a_{33} > 0$  then by considering  $(a_{21}, a_{31} - 1; a_{22}, a_{32} - 1; a_{23}, a_{33} - 1)$ , we have  $s \geq a_{32} + a_{33}$ .
- (ii) If  $a_{21}, a_{22}, a_{23} > 0$  then by considering  $(a_{21} - 1, a_{31} - 1, a_{22} - 1, a_{32} - 1, a_{23} - 1, a_{33} - 1)$ , we have  $s \geq a_{21} + a_{22} + a_{23}$ .
- (iii) If  $a_{22} > 0$  then by considering  $(a_{21}, a_{31} - 1; a_{22} - 1, a_{32} - 1; a_{23}, a_{33})$ , we have  $s \geq a_{31} + a_{22}$ .

- (iv) If  $a_{21}, a_{32} > 0$  then by considering  $(a_{21} - 1, a_{31} - 1; a_{22}, a_{32} - 1; a_{23}, a_{33})$ , we have  $s \geq a_{21} + a_{32}$ .
- (v) If  $a_{23} > 0$  then by considering  $(a_{21}, a_{31} - 1; a_{22}, a_{32}; a_{23} - 1, a_{33} - 1)$ , we have  $s \geq a_{31} + a_{23}$ .

We now claim that if  $a_{33} > 0$  then  $s \geq a_{21} + a_{22} + a_{23}$ . Indeed if  $a_{21}, a_{22}, a_{23} > 0$  then this is (ii). If  $a_{21} = 0$  then we instead use (i). If  $a_{21} > 0$  and  $a_{23} = 0$  then (noting that  $a_{32} \geq a_{33} > 0$ ) we instead use (iv). If  $a_{23} > 0$  and  $a_{22} = 0$  then we instead use (v).

To complete the proof of the lemma, we first suppose  $a_{33} > 0$ . Then the inequalities in (i) and (ii) hold without further hypothesis. We weaken the inequalities (iii), (iv) and (v) to

$$(8) \quad s + 1 \geq a_{31} + a_{22}$$

$$(9) \quad s + 1 \geq a_{21} + a_{32}$$

$$(10) \quad s + 1 \geq a_{31} + a_{23}$$

so that in cases where some of the  $a_{ij}$  are zero, these still hold by (6) and (7). Adding together all five inequalities gives

$$5s + 3 + a_{33} \geq 2 \sum a_{ij} = 2(3s - 1)$$

and hence  $a_{33} \geq s - 5$ . Using (i) again gives

$$s \geq a_{32} + a_{33} \geq 2a_{33} \geq 2(s - 5)$$

and hence  $s \leq 10$ , as required.

If  $a_{33} = 0$  then we still have (8) and (9) giving  $2(s + 1) \geq \sum a_{ij} = 3s - 1$ , and hence  $s \leq 3$ .  $\square$

*Proof of Theorem 3.3.* We represent  $S$  as a triple of matrices  $A, B, C$ , say.

$$\begin{array}{ccc|ccc|ccc} A_{11} & A_{12} & A_{13} & B_{11} & B_{12} & B_{13} & C_{11} & C_{12} & C_{13} \\ A_{21} & A_{22} & A_{23} & B_{21} & B_{22} & B_{23} & C_{21} & C_{22} & C_{23} \\ A_{31} & A_{32} & A_{33} & B_{31} & B_{32} & B_{33} & C_{31} & C_{32} & C_{33} \end{array}$$

The action of  $\mathcal{G}(K) = \mathrm{GL}_3(K)^3$  may be described as follows. The first factor replaces  $A, B, C$  by linear combinations of these matrices. The second factor acts by row operations (applied to  $A, B, C$  simultaneously), and the third factor acts by column operations.

We may assume that one of the tuples  $\tau_1, \dots, \tau_6$  in Lemma 3.5 is admissible for  $S$ . We therefore split into these 6 cases. In each case, it is our running assumption that we are not in an earlier case.

**Case 1.** We assume  $(1, 1; 0, 0; 0, 0)$  is admissible for  $S$ . Then the entries of  $A$  have valuation at least one, and so the cube  $S$  is not saturated.

**Case 2.** We assume  $(0, 1; 0, 1; 0, 0)$  is admissible for  $S$ . The entries of  $A, B$  and  $C$  have valuations satisfying

$$\begin{array}{ccccccc} \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 \\ \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 \\ \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \end{array}$$

Since  $S$  is saturated we may assume by column operations that  $v(C_{11}) = 0$ ,  $v(C_{12}) \geq 1$  and  $v(C_{13}) \geq 1$ . Subtracting a multiple of the first row from the second row gives  $v(C_{21}) \geq 1$ , and again by column operations  $v(C_{22}) = 0$  and  $v(C_{23}) \geq 1$ . Subtracting multiples of the first two rows from the third, the valuations now satisfy

$$\begin{array}{ccccccc} \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & = 0 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & = 0 & \geq 1 \\ \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 1 & \geq 1 & \geq 0 \end{array}$$

We compute  $f_1 = C_{11}C_{22}z^2(A_{33}x + B_{33}y + C_{33}z) \pmod{\pi}$ . Since  $S$  is saturated it follows that  $f_1$  is nonzero. The same argument shows that  $f_2$  has a repeated factor and is nonzero. On the other hand we have  $f_3 = 0$ . The procedure in (i) multiplies  $C$  and the third row by  $\pi$ , and then divides the cube by  $\pi$ . This transformation decreases the level.

**Case 3.** We assume  $(1, 2; 0, 1; 0, 1)$  is admissible for  $S$ . The entries of  $A, B$  and  $C$  have valuations satisfying

$$\begin{array}{ccccccc} \geq 2 & \geq 2 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \\ \geq 2 & \geq 2 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \\ \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \end{array}$$

Since  $S$  is saturated we have  $v(A_{33}) = 0$ . If  $B_{13} \equiv B_{23} \equiv 0 \pmod{\pi}$  then we are in Case 2, and likewise if  $B_{31} \equiv B_{32} \equiv 0 \pmod{\pi}$ . By operating on the first two rows and columns, and then subtracting a multiple of  $A$  from  $B$ , the valuations now satisfy

$$\begin{array}{ccccccc} \geq 2 & \geq 2 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 \\ \geq 2 & \geq 2 & \geq 1 & \geq 1 & \geq 1 & = 0 & \geq 0 & \geq 0 & \geq 0 \\ \geq 1 & \geq 1 & = 0 & \geq 1 & = 0 & \geq 1 & \geq 0 & \geq 0 & \geq 0 \end{array}$$

Working mod  $\pi$  we compute

$$\begin{aligned} f_1 &= -B_{23}B_{32}C_{11}y^2z + z^2(\dots) \\ f_2 &= -A_{33}B_{32}z^2(C_{11}x + C_{21}y + C_{31}z) \\ f_3 &= -A_{33}B_{23}z^2(C_{11}x + C_{12}y + C_{13}z) \end{aligned}$$

Since  $S$  is saturated, it is clear that  $f_2$  and  $f_3$  are nonzero.

We note that multiplying  $C$ , the last row and the last column by  $\pi$ , and then dividing the whole cube by  $\pi$ , gives an integral model of the same

level which is not saturated. These transformations are carried out by the procedure in (i), except possibly in the case where  $f_1$  has a repeated factor, and this factor is not  $z^2$ . In this remaining case  $v(C_{11}) = 0$ . We may assume by row and column operations that  $C_{12} \equiv C_{13} \equiv C_{21} \equiv C_{31} \equiv 0 \pmod{\pi}$ . Subtracting multiples of  $A$  and  $B$  from  $C$  gives  $C_{32} = C_{33} = 0 \pmod{\pi}$ . Now  $f_1 = C_{11}z(A_{33}C_{22}xz - B_{23}B_{32}y^2 - B_{32}C_{23}yz)$ , and so  $C_{22} \equiv C_{23} \equiv 0 \pmod{\pi}$ .

If the procedure in (i) picks  $f_1$  and  $f_2$  then we multiply  $B$  and the last row by  $\pi$ . Dividing the last two columns by  $\pi$  gives a model of the same level with valuations satisfying

$$\begin{array}{ccccccccccc} \geq 2 & \geq 1 & \geq 0 & \geq 2 & \geq 1 & \geq 1 & = 0 & \geq 0 & \geq 0 & & \\ \geq 2 & \geq 1 & \geq 0 & \geq 2 & \geq 1 & = 0 & \geq 1 & \geq 0 & \geq 0 & & \\ \geq 2 & \geq 1 & = 0 & \geq 3 & = 1 & \geq 2 & \geq 2 & \geq 1 & \geq 1 & & \end{array}$$

Since the first two columns of  $A$  and  $B$  are divisible by  $\pi$ , we are now in Case 2. The case where the procedure in (i) picks  $f_1$  and  $f_3$  works in the same way.

**Case 4.** We assume  $(1, 1; 1, 1; 0, 1)$  is admissible for  $S$ . The entries of  $A$ ,  $B$  and  $C$  have valuations satisfying

$$\begin{array}{ccccccccccc} \geq 2 & \geq 2 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 1 & \geq 1 & \geq 0 & & \\ \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & & \\ \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & & \end{array}$$

Working mod  $\pi$  we compute

$$f_1 = (B_{13}y + C_{13}z) \left| \begin{pmatrix} B_{21} & B_{22} \\ B_{31} & B_{32} \end{pmatrix} y + \begin{pmatrix} C_{21} & C_{22} \\ C_{31} & C_{32} \end{pmatrix} z \right|,$$

and

$$f_2 = (A_{23}y + A_{33}z) \left| \begin{pmatrix} B_{21} & B_{22} \\ C_{21} & C_{22} \end{pmatrix} y + \begin{pmatrix} B_{31} & B_{32} \\ C_{31} & C_{32} \end{pmatrix} z \right|.$$

Since  $S$  is saturated, the linear factors  $\ell_1 = B_{13}y + C_{13}z$  and  $\ell_2 = A_{23}y + A_{33}z$  cannot be identically zero. Let  $q_1$  and  $q_2$  be the quadratic factors. These are binary quadratic forms associated to the same  $2 \times 2 \times 2$  cube. In particular  $q_1$  and  $q_2$  have the same discriminant, say  $\delta$ . If this  $2 \times 2 \times 2$  cube is not saturated, it is easy to see we are in Case 1 or Case 2. Therefore  $f_1$  and  $f_2$  are nonzero.

Replacing  $B$  and  $C$  by suitable linear combinations, and likewise the last two rows, we may suppose that the linear factors  $\ell_1$  and  $\ell_2$  are multiples of  $z$ , i.e.

$$(11) \quad B_{13} \equiv A_{23} \equiv 0 \pmod{\pi}$$

Under this assumption  $f_3 = -A_{33}C_{13}z^2(B_{21}x + B_{22}y + B_{23}z)$ , and this is nonzero as we would otherwise be in Case 2.



If  $f_1$  and  $f_2$  don't have repeated factors, then each defines a curve with a unique singular point at  $(1 : 0 : 0)$ . The procedure in (ii) multiplies  $B$ ,  $C$  and the last two rows by  $\pi$ . The level is then reduced using column operations. The overall transformation applied in this case is exactly that suggested by the definition of admissibility, and the fact we are in Case 4.

Now suppose that at least one of the forms  $f_1$  and  $f_2$  has a repeated factor. Then the procedure in (i) is applied. We say we are in the *good situation* if the two of the  $f_i$  chosen are multiples of  $z^2$  and  $B_{21} \equiv B_{22} \equiv 0 \pmod{\pi}$ . Indeed in the good situation, the procedure in (i) reduces us to Case 1 or Case 2.

Suppose that  $f_1$  and  $f_3$  are chosen. Dropping the assumption (11) we may assume that  $f_1$  has repeated factor  $z^2$ . Then  $q_1$  has no  $y^2$  term and by row operations we reach the good situation. The case where  $f_2$  and  $f_3$  are chosen is similar. Finally we suppose that  $f_1$  and  $f_2$  are chosen. If  $q_1$  has a factor  $z$ , we may assume as above that  $B_{21} \equiv B_{22} \equiv 0 \pmod{\pi}$ . But then  $q_2$  has a factor  $z$ . So if  $\delta = 0$ , i.e.  $q_1$  and  $q_2$  each have a repeated factor, then we reach the good situation. Otherwise we make the assumption (11), and deduce that  $f_1$  and  $f_2$  are now multiples of  $z^2$ . The procedure in (i) multiplies  $C$  and the last row by  $\pi$ . The only coefficients not to vanish mod  $\pi$  are now those in the second row of  $B$ . It follows that after suitable column operations the level is preserved and we are reduced to Case 2 or Case 3.

**Case 5.** We assume  $(1, 2; 1, 2; 1, 1)$  is admissible for  $S$ . The entries of  $A, B$  and  $C$  have valuations satisfying

$$\begin{array}{ccccccccc} \geq 3 & \geq 2 & \geq 2 & \geq 2 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 \\ \geq 2 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \\ \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \end{array}$$

Since  $S$  is saturated, we may assume by column operations that  $v(A_{32}) \geq 1$  and  $v(A_{33}) = 0$ . Then  $v(B_{31}) = v(C_{12}) = v(C_{21}) = 0$ , otherwise we would be in Case 4. By row and column operations, and subtracting multiples of  $A$  from  $B$  and  $C$  we reduce to the case

$$\begin{array}{ccccccccc} \geq 3 & \geq 2 & \geq 2 & \geq 2 & \geq 1 & \geq 1 & \geq 1 & = 0 & \geq 1 \\ \geq 2 & \geq 1 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & = 0 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & = 0 & = 0 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 \end{array}$$

Working mod  $\pi$  we compute

$$\begin{aligned} f_1 &= C_{12}z(B_{31}B_{23}y^2 - A_{33}C_{21}xz) \\ f_2 &= -A_{33}z(B_{22}C_{21}y^2 - B_{31}C_{12}xz) \\ f_3 &= -A_{33}C_{12}yz(B_{22}y + B_{23}z) \end{aligned}$$

If  $B_{22} \not\equiv 0 \pmod{\pi}$  and  $B_{23} \not\equiv 0 \pmod{\pi}$  then  $f_1, f_2, f_3$  each define a curve with a unique singular point at  $(1 : 0 : 0)$ . If we multiply  $B, C$ , the last two rows and the last two columns by  $\pi$ , then the cube is divisible by  $\pi^2$ . From this we see that whichever two of the  $f_i$  are chosen by the procedure in (ii), the level is preserved and we are reduced to Case 2.

If  $B_{22} \not\equiv 0 \pmod{\pi}$  and  $B_{23} \equiv 0 \pmod{\pi}$  then  $f_1$  and  $f_3$  have repeated factors but  $f_2$  does not. The procedure in (i) multiplies  $C$  and the middle column by  $\pi$ . Then dividing the first two rows by  $\pi$  preserves the level and reduces us to Case 4 with  $\delta = 0$ . The observation that  $\delta = 0$  is needed to show that at most three iterations are required, as claimed in the statement of the theorem.

If  $B_{22} \equiv 0 \pmod{\pi}$  and  $B_{23} \not\equiv 0 \pmod{\pi}$  then we switch the first two slicings (i.e.  $A, B, C$  are replaced by the matrices formed from the first, second, third rows). Then switching the last two columns brings us to the situation considered in the previous paragraph.

Finally, if  $B_{22} \equiv B_{23} \equiv 0 \pmod{\pi}$  then we are already in Case 2.

**Case 6.** We assume  $(2, 3; 1, 2; 1, 2)$  is admissible for  $S$ . The entries of  $A, B$  and  $C$  have valuations satisfying

$$\begin{array}{ccccccc} \geq 4 & \geq 3 & \geq 2 & \geq 2 & \geq 1 & \geq 0 & \geq 1 & \geq 0 & \geq 0 \\ \geq 3 & \geq 2 & \geq 1 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \\ \geq 2 & \geq 1 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 & \geq 0 \end{array}$$

Since  $S$  is saturated, we have  $v(A_{33}) = 0$ . Then  $v(B_{22}) = 0$ , otherwise we would be in Case 3. We also have  $v(C_{12}) = v(C_{21}) = 0$ , otherwise we would be in Case 4, and  $v(B_{13}) = v(B_{31}) = 0$  otherwise we would be in Case 5. By row and column operations, and subtracting multiples of  $A$  from  $B$  and  $C$  we reduce to the case

$$\begin{array}{ccccccc} \geq 4 & \geq 3 & \geq 2 & \geq 2 & \geq 1 & = 0 & \geq 1 & = 0 & \geq 1 \\ \geq 3 & \geq 2 & \geq 1 & \geq 1 & = 0 & \geq 0 & = 0 & \geq 1 & \geq 1 \\ \geq 2 & \geq 1 & = 0 & = 0 & \geq 0 & \geq 1 & \geq 1 & \geq 1 & \geq 1 \end{array}$$

Working mod  $\pi$  we compute

$$\begin{aligned} f_1 &= -B_{31}B_{22}B_{13}y^3 - C_{12}C_{21}A_{33}xz^2 + (\dots)y^2z \\ f_2 &= A_{33}z(B_{31}C_{12}xz - C_{21}y(B_{22}y + B_{32}z)) \\ f_3 &= A_{33}z(B_{13}C_{21}xz - C_{12}y(B_{22}y + B_{23}z)) \end{aligned}$$

We see that  $f_1, f_2, f_3$  each define a curve with a unique singular point at  $(1 : 0 : 0)$ . If we multiply  $B, C$ , the last two rows and the last two columns by  $\pi$ , then the cube is divisible by  $\pi^2$ . From this we see that whichever two of the  $f_i$  are chosen by the procedure in (ii), the level is preserved and we are reduced to Case 3.  $\square$

4.  $2 \times 2 \times 2 \times 2$  HYPERCUBES

We consider polynomials in  $x_1, x_2, y_1, y_2, z_1, z_2, t_1, t_2$  that are linear in each of the four sets of variables. Such a polynomial may be represented as

$$(12) \quad \sum_{1 \leq i, j, k, l \leq 2} H_{ijkl} x_i y_j z_k t_l$$

where  $H = (H_{ijkl})$  is a  $2 \times 2 \times 2 \times 2$  hypercube. A hypercube  $H$  may be partitioned into two  $2 \times 2 \times 2$  cubes in four distinct ways:

- (i)  $A_1 = (H_{1jkl})$  and  $B_1 = (H_{2jkl})$
- (ii)  $A_2 = (H_{i1kl})$  and  $B_2 = (H_{i2kl})$
- (iii)  $A_3 = (H_{ij1l})$  and  $B_3 = (H_{ij2l})$
- (iv)  $A_4 = (H_{ijk1})$  and  $B_4 = (H_{ijk2})$

Let  $R$  be a ring. For each  $1 \leq i \leq 4$  there is an action of  $\mathrm{GL}_2(R)$  on the space of hypercubes over  $R$  via

$$\begin{pmatrix} r & s \\ t & u \end{pmatrix} : (A_i, B_i) \mapsto (rA_i + sB_i, tA_i + uB_i).$$

These actions commute, and so give an action of  $\mathrm{GL}_2(R)^4$ . We say that hypercubes are *R-equivalent* if they belong to the same orbit for this action.

For each  $1 \leq i < j \leq 4$  there is an associated  $(2, 2)$ -form  $F_{ij}$ . Indeed if we view (12) as a bilinear form in  $z_k$  and  $t_l$ , then the determinant of this form is a  $(2, 2)$ -form in  $x_i$  and  $y_j$ :

$$F_{12} = \left( \sum_{1 \leq i, j \leq 2} H_{ij11} x_i y_j \right) \left( \sum_{1 \leq i, j \leq 2} H_{ij22} x_i y_j \right) - \left( \sum_{1 \leq i, j \leq 2} H_{ij12} x_i y_j \right) \left( \sum_{1 \leq i, j \leq 2} H_{ij21} x_i y_j \right).$$

The other  $F_{ij}$  are defined similarly. If  $[M_1, M_2, M_3, M_4] \cdot H = H'$  then the  $(2, 2)$ -forms are related by

$$[\det(M_3) \det(M_4), M_1, M_2] \cdot F_{12} = F'_{12}.$$

As seen in Section 2, each  $(2, 2)$ -form determines a pair of binary quartics. It turns out that the binary quartics in  $x_1, x_2$  associated to  $F_{12}, F_{13}, F_{14}$  are all equal. Thus a hypercube  $H$  determines four binary quartics  $G_1, \dots, G_4$ , one in each of the four sets of variables. Each of these binary quartics has the same invariants  $I$  and  $J$ . Therefore the six  $(2, 2)$ -forms  $F_{ij}$  all have the same invariants  $c_4, c_6$  and  $\Delta$ . We define  $c_4(H) = c_4(F_{ij})$ ,  $c_6(H) = c_6(F_{ij})$  and  $\Delta(H) = \Delta(F_{ij})$ .

If  $H$  is defined over a field and  $\Delta(H) \neq 0$  then each of the  $F_{ij}$  defines a genus one curve in  $\mathbb{P}^1 \times \mathbb{P}^1$ . These curves are isomorphic, although not in

a canonical way. (See [1, Section 2.3] for further details.) We write  $\mathcal{C}_H$  to denote any one of them.

Let  $u$  and  $v$  be the invariants in Lemma 2.1. We find that  $u(F_{12}) = u(F_{34})$  and  $v(F_{12}) = v(F_{34})$ . Therefore  $F_{12}$  and  $F_{34}$  determine isomorphic pairs  $(E, P)$ . (A further calculation is needed to check this in characteristics 2 and 3, but we omit the details.) Repeating for the other  $F_{ij}$  gives a tuple  $(E, P_1, P_2, P_3)$  where  $E$  is an elliptic curve and  $0_E \neq P_1, P_2, P_3 \in E$  with  $P_1 + P_2 + P_3 = 0_E$ .

We say that a hypercube  $H$  is *integral* if it has coefficients in  $\mathcal{O}_K$ , and *non-singular* if  $\Delta(H) \neq 0$ .

**Lemma 4.1.** *Let  $H$  be a non-singular integral hypercube. Let  $(E, P_1, P_2, P_3)$  be the tuple determined by  $H$ . Then*

$$v(\Delta(H)) = v(\Delta_E) + 12 \max(\kappa(P_1), \kappa(P_2), \kappa(P_3)) + 12\ell(H)$$

where  $\ell(H) \geq 0$  is an integer we call the level.

*Proof.* This is immediate from Lemma 2.2. □

An integral hypercube is *saturated* if for all  $1 \leq i \leq 4$  the cubes  $A_i$  and  $B_i$  are linearly independent mod  $\pi$ . If an integral hypercube is not saturated, then it is obvious how we may decrease the level.

Our algorithm for minimising hypercubes is described by the following theorem.

**Theorem 4.2.** *Let  $H$  be a saturated hypercube with associated  $(2, 2)$ -forms  $F_{ij}$ . Suppose that all of the  $F_{ij}$  are non-minimal. Then after applying an  $\mathcal{O}_K$ -equivalence, and permuting the sets of variables, if necessary, we are in one of the following two situations:*

- (i) *The reduction of  $F_{12}$  mod  $\pi$  defines a curve in  $\mathbb{P}^1 \times \mathbb{P}^1$  with a unique singular point at  $((1 : 0), (1 : 0))$ , and the transformation*

$$(13) \quad \left[ \frac{1}{\pi} \begin{pmatrix} 1 & 0 \\ 0 & \pi \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & \pi \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right]$$

*gives an integral hypercube of the same level.*

- (ii) *We have  $F_{12} \equiv x_2^2 y_2^2 \pmod{\pi}$  and the transformation (13) gives a non-saturated hypercube of the same level.*

*Moreover, at most two iterations of the procedure in (i) are needed to give a non-saturated hypercube, or to reach the situation in (ii).*

It is clear that if any of the  $F_{ij}$  or  $G_i$  are minimal then  $H$  is minimal. We initially used the methods in Sections 2 and 3 to prove Theorem 4.2 under the apparently stronger hypothesis that  $H$  is non-minimal. The advantage of the theorem as stated here is that it has the following consequence.

**Corollary 4.3.** *Let  $H$  be a integral hypercube with associated  $(2, 2)$ -forms  $F_{ij}$ . Then  $H$  is minimal if and only if some  $F_{ij}$  is minimal.*

**Remark 4.4.** We may represent  $H = (H_{ijkl})$  as a  $4 \times 4$  matrix:

$$(14) \quad \left( \begin{array}{cc|cc} H_{1111} & H_{1211} & H_{1112} & H_{1212} \\ H_{2111} & H_{2211} & H_{2112} & H_{2212} \\ \hline H_{1121} & H_{1221} & H_{1122} & H_{1222} \\ H_{2121} & H_{2221} & H_{2122} & H_{2222} \end{array} \right).$$

If we write  $r_1, r_2, r_3, r_4$  for the rows, then the first copy of  $\mathrm{GL}_2$  acts by row operations simultaneously on  $\{r_1, r_2\}$  and  $\{r_3, r_4\}$ , the third copy of  $\mathrm{GL}_2$  acts by row operations on  $\{r_1, r_3\}$  and  $\{r_2, r_4\}$ , and the other two copies of  $\mathrm{GL}_2$  act by column operations.

**Remark 4.5.** Let  $H$  be an integral hypercube with associated binary quartics  $G_1, \dots, G_4$ . As noted above, if any of the  $G_i$  are minimal then  $H$  is minimal. However the converse is not true. For example if

$$H \equiv \left( \begin{array}{cc|cc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right) \pmod{\pi^2}$$

then  $H$  is minimal (since  $F_{12} \equiv (x_1y_1 + x_2y_2)^2 \pmod{\pi^2}$  and we saw in Remark 2.4 that this is minimal), yet we have  $G_1 \equiv \dots \equiv G_4 \equiv 0 \pmod{\pi^2}$ .

For the proof of Theorem 4.2 we need the following lemma.

**Lemma 4.6.** *Let  $H$  be an integral hypercube. Suppose that at least one of the associated  $(2, 2)$ -forms  $F_{ij}$  is non-minimal. Then by an  $\mathcal{O}_K$ -equivalence, and permuting the sets of variables, we may assume that  $H_{11kl} \equiv 0 \pmod{\pi}$  for all  $1 \leq k, l \leq 2$ .*

*Proof.* We suppose that  $F_{12}$  is non-minimal. If the reduction of  $F_{12} \pmod{\pi}$  is non-zero, then by Theorem 2.3 it defines a curve in  $\mathbb{P}^1 \times \mathbb{P}^1$  with singular locus a point, a line or a pair of lines. We may assume by an  $\mathcal{O}_K$ -equivalence that the curve is singular at  $((1 : 0), (1 : 0))$ . If  $H_{11kl} \not\equiv 0 \pmod{\pi}$  for some  $1 \leq k, l \leq 2$  then we may assume by an  $\mathcal{O}_K$ -equivalence that  $H_{1111} \not\equiv 0 \pmod{\pi}$ . A further  $\mathcal{O}_K$ -equivalence gives

$$H_{2111} \equiv H_{1211} \equiv H_{1121} \equiv H_{1112} \equiv 0 \pmod{\pi}.$$

Since the coefficients of  $x_1^2y_1^2$ ,  $x_1^2y_1y_2$  and  $x_1x_2y_1^2$  in  $F_{12}$  vanish mod  $\pi$ , we have

$$H_{1122} \equiv H_{1222} \equiv H_{2122} \equiv 0 \pmod{\pi}.$$

Lemma 2.6(i) now shows that either

$$H_{1221}H_{1212} \equiv 0 \pmod{\pi} \quad \text{or} \quad H_{2121}H_{2112} \equiv 0 \pmod{\pi}.$$

By switching the first two sets of variables and switching the last two sets of variables, as necessary, we may assume that  $H_{1212} \equiv 0 \pmod{\pi}$ . Now  $H_{1jk2} \equiv 0 \pmod{\pi}$  for all  $1 \leq j, k \leq 2$ , and this proves the lemma.  $\square$

*Proof of Theorem 4.2.* By Lemma 4.6 we may assume that  $H_{11kl} \equiv 0 \pmod{\pi}$  for all  $1 \leq k, l \leq 2$ . Applying Lemma 2.6(i) to  $F_{12}$ , and switching the first two sets of variables if necessary, we have

$$H_{1211}H_{1222} - H_{1212}H_{1221} \equiv 0 \pmod{\pi}.$$

By an  $\mathcal{O}_K$ -equivalence we may assume  $H_{1jkl} \equiv 0 \pmod{\pi}$  for all  $1 \leq j, k, l \leq 2$ , except  $(j, k, l) = (2, 2, 2)$ . Since  $H$  is saturated we have  $H_{1222} \not\equiv 0 \pmod{\pi}$ . Again by Lemma 2.6(i) we have  $H_{2111} \equiv 0 \pmod{\pi}$ .

We now split into cases, according as to whether

$$(15) \quad H_{2211} \equiv H_{2121} \equiv H_{2112} \equiv 0 \pmod{\pi}.$$

If this condition is not satisfied, then by permuting the last three sets of variables, we may suppose that  $H_{2211} \not\equiv 0 \pmod{\pi}$ . By an  $\mathcal{O}_K$ -equivalence we have

$$(16) \quad H \equiv \left( \begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ 0 & 1 & \beta & 0 \\ \hline 0 & 0 & 0 & 1 \\ \alpha & 0 & \gamma & 0 \end{array} \right) \pmod{\pi}$$

for some  $\alpha, \beta, \gamma \in k$ . We compute  $F_{12} \equiv x_1x_2y_2^2 + x_2^2(\alpha\beta y_1^2 + \gamma y_1y_2) \pmod{\pi}$ . The conclusions in (i) are satisfied unless  $\alpha\beta = \gamma = 0$ . In the remaining case we may assume, by switching the last two sets of variables if necessary, that  $\alpha = 0$ . Now switching the first and last sets of variables, and swapping over the third set of variables (i.e.  $z_1 \leftrightarrow z_2$ ), we may swap over  $\beta$  and  $\gamma$ . Therefore  $\beta = \gamma = 0$ , and this contradicts that  $H$  is saturated.

Now suppose the condition (15) is satisfied. Then by an  $\mathcal{O}_K$ -equivalence (and our assumption that  $H$  is saturated) we have

$$(17) \quad H \equiv \left( \begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 1 \\ 0 & -1 & -1 & 0 \end{array} \right) \pmod{\pi}.$$

We compute  $F_{12} \equiv x_2^2y_2^2 \pmod{\pi}$ . Let  $F_{12}$  have coefficients  $a_{ij}$  as labelled in (3). Lemma 2.6(ii) shows that either  $v(a_{12}) \geq 2$  or  $v(a_{21}) \geq 2$ . Therefore  $v(H_{1111}) \geq 2$ . Again by Lemma 2.6(ii) we have either  $v(a_{11}) \geq 3$ ,  $v(a_{13}) \geq 2$  or  $v(a_{31}) \geq 2$ . Therefore at least one of the coefficients  $H_{2111}$ ,  $H_{1211}$ ,  $H_{1121}$ ,  $H_{1112}$  has valuation at least two. By permuting the sets of variables we may suppose that  $v(H_{1112}) \geq 2$ . The conclusions in (ii) are now satisfied.

To prove the last part of the theorem, we need the following lemma.

**Lemma 4.7.** *Let  $H$  be a hypercube over a field  $k$  with associated  $(2, 2)$ -forms  $F_{ij}$ . We write*

$$\begin{aligned} F_{12} &= f_1(x_1, x_2)y_1^2 + f_2(x_1, x_2)y_1y_2 + f_3(x_1, x_2)y_2^2 \\ F_{13} &= g_1(x_1, x_2)z_1^2 + g_2(x_1, x_2)z_1z_2 + g_3(x_1, x_2)z_2^2 \end{aligned}$$

- (i) *We have  $g_2 = f_2 + 2h$  and  $g_1g_3 = f_1f_3 + f_2h + h^2$  for some  $h \in k[x_1, x_2]$ .*
- (ii) *If  $f_1 = f_2 = 0$  and  $g_1, g_2$  are multiples of  $x_2^2$ , then  $F_{13}$  is either zero or factors as a product of binary quadratic forms.*

*Proof.* (i) We have already remarked that  $f_2^2 - 4f_1f_3 = g_2^2 - 4g_1g_3$ . The result follows by considering the  $f_i$  and  $g_i$  as polynomials in  $\mathbb{Z}[H_{ijkl}][x_1, x_2]$ .

(ii) By (i) we have  $g_1 = \alpha x_2^2$ ,  $g_2 = 2\beta x_2^2$  and  $\alpha x_2^2 g_3(x_1, x_2) = \beta x_2^4$ . If  $\alpha = 0$  then  $g_1 = g_2 = 0$ , whereas if  $\alpha \neq 0$  then  $g_1, g_2, g_3$  are multiples of  $x_2^2$ .  $\square$

We say that a  $(2, 2)$ -form  $F$  is *slender* if  $F \bmod \pi$  is either zero, or factors as a product of binary quadratic forms. Theorem 2.3 shows that if  $F$  is non-minimal then either  $F \bmod \pi$  defines a curve with a unique singular point, or  $F$  is slender. These possibilities are mutually exclusive.

We now complete the proof of Theorem 4.2. Applying the transformation in (i) to  $H$  has the effect of applying the transformation in Theorem 2.3(iii) to  $F_{12}$ . The last sentence of Theorem 2.3 tells us that, after applying this transformation,  $F_{12} \bmod \pi$  is either zero, or factors as a product of binary quadratic forms both of which have a repeated root. In particular  $F_{12}$  is slender.

We claim that  $F_{13}$  is slender. If not then  $F_{13} \bmod \pi$  defines a curve with a unique singular point. By an  $\mathcal{O}_K$ -equivalence we may assume that this point is  $((1 : 0), (1 : 0))$ , and that  $F_{12} \equiv f_3(x_1, x_2)y_2^2 \pmod{\pi}$  for some binary quadratic form  $f_3$ . Lemmas 2.6(i) and 4.7(ii) now show that  $F_{13}$  is slender.

The same argument shows that all of the  $F_{ij}$  are slender, except possibly  $F_{34}$ . Since  $F_{34}$  was unchanged by the transformation (13), it follows that after at most two iterations, all of the  $F_{ij}$  are slender. In particular we cannot return to the situation in (i), and this completes the proof.  $\square$

## 5. MINIMISATION THEOREMS

The algorithms in [3] and [4] for minimising genus one curves of degree 2, 3, 4, 5 were complemented by a more theoretical result. This stated that if a genus one curve is soluble over  $K$  (or more generally over an unramified extension) then the discriminant of a minimal model is the same as that

for the Jacobian elliptic curve. In this section we prove the analogue of this result for  $(2, 2)$ -forms,  $3 \times 3 \times 3$  cubes and  $2 \times 2 \times 2 \times 2$  hypercubes.

In earlier papers, most notably [2, Lemmas 3,4,5], the minimisation algorithms and minimisation theorems were treated together. Following [3] we separate these out, and this leads to clean results that work the same in all residue characteristics. We phrase our result in terms of the level, as defined in Lemmas 2.2, 3.2 and 4.1.

**Theorem 5.1.** *Let  $\Phi$  be a non-singular  $(2, 2)$ -form,  $3 \times 3 \times 3$  cube, or  $2 \times 2 \times 2 \times 2$  hypercube defined over  $K$ . If  $\mathcal{C}_\Phi(K) \neq \emptyset$  then  $\Phi$  has minimal level 0.*

**Remark 5.2.** The algorithms in Sections 2, 3 and 4 show that the minimal level is unchanged by an unramified field extension. The hypothesis in Theorem 5.1 may therefore be weakened to solubility over an unramified field extension. We give examples below to show that this hypothesis cannot be removed entirely.

Let  $E/K$  be an elliptic curve and  $n \in \{2, 3\}$ . Let  $D$  and  $D'$  be  $K$ -rational divisors on  $E$  of degree  $n$ . The image of  $E$  in  $\mathbb{P}^{n-1} \times \mathbb{P}^{n-1}$  via  $|D| \times |D'|$  is defined by a  $(2, 2)$ -form in the case  $n = 2$ , and three bilinear forms in the case  $n = 3$ . The coefficients of the latter give a  $3 \times 3 \times 3$  cube. We note that the  $(2, 2)$ -form, respectively  $3 \times 3 \times 3$  cube, is uniquely determined up to  $K$ -equivalence by the triple  $(E, [D], [D'])$ , where  $[D]$  denotes the linear equivalence class of  $D$ . Moreover every  $(2, 2)$ -form, respectively  $3 \times 3 \times 3$  cube, defining a non-singular genus one curve with a  $K$ -rational point, arises in this way. Therefore the first two cases of Theorem 5.1 are immediate from the following theorem.

We write  $\text{sum} : \text{Div}_K(E) \rightarrow E(K)$  for the map that sends a formal sum of points to its sum using the group law on  $E$ . For a  $(2, 2)$ -form or  $3 \times 3 \times 3$  cube as constructed in the previous paragraph we have  $\text{sum}(D' - D) = P$  where  $P$  is the point described in Lemmas 2.1 and 3.1. See [1, Proposition 5.5 and Section 6.1].

**Theorem 5.3.** *Let  $E/K$  be an elliptic curve with integral Weierstrass equation*

$$(18) \quad y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x$$

*and let  $P = (0, 0) \in E(K)$ . Let  $D, D' \in \text{Div}_K(E)$  be divisors of degree  $n \in \{2, 3\}$  with  $\text{sum}(D' - D) = P$ . Then  $(E, [D], [D'])$  may be represented by an integral  $(2, 2)$ -form, or  $3 \times 3 \times 3$  cube, with the same discriminant as (18).*



We start by proving Theorem 5.3 in the case  $D \sim n \cdot 0_E$ . Since  $\text{sum}(D' - D) = P$  we have  $D' \sim (n-1) \cdot 0_E + P$ . We put

$$f = \frac{y + a_1x + a_3}{x} = \frac{x^2 + a_2x + a_4}{y}$$

and split into the cases  $n = 2$  and  $n = 3$ .

**Case  $n = 2$ .** The embedding  $E \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$  via  $|D| \times |D'|$  is given by

$$(x, y) \mapsto ((1 : x), (1 : f)).$$

The image is defined by the  $(2, 2)$ -form

$$F(x_1, x_2; y_1, y_2) = x_2^2 y_1^2 - x_1 x_2 y_2^2 + x_1 y_1 (a_1 x_2 y_2 + a_2 x_2 y_1 + a_3 x_1 y_2 + a_4 x_1 y_1),$$

with the same discriminant as (18).

**Case  $n = 3$ .** The embedding  $E \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$  via  $|D| \times |D'|$  is given by

$$(x, y) \mapsto ((1 : x : y), (1 : x : f)).$$

The image is defined by bilinear forms

$$\begin{aligned} B_1(x_1, x_2, x_3; y_1, y_2, y_3) &= x_2 y_1 - x_1 y_2, \\ B_2(x_1, x_2, x_3; y_1, y_2, y_3) &= x_3 y_1 + a_1 x_2 y_1 + a_3 x_1 y_1 - x_2 y_3, \\ B_3(x_1, x_2, x_3; y_1, y_2, y_3) &= x_2 y_2 + a_2 x_2 y_1 + a_4 x_1 y_1 - x_3 y_3. \end{aligned}$$

The coefficients of  $B_1, B_2, B_3$  give a  $3 \times 3 \times 3$  cube, and this has the same discriminant as (18).

**Lemma 5.4.** *Let  $S$  be a  $3 \times 3 \times 3$  cube corresponding to bilinear forms  $B_1, B_2, B_3$ , defining  $C \subset \mathbb{P}^2 \times \mathbb{P}^2$  a smooth curve of genus one, embedded via  $|D| \times |D'|$ .*

- (i) *If  $Q = ((0 : 0 : 1), (0 : 0 : 1)) \in C(K)$  then for  $i = 1, 2, 3$  we can write*

$$B_i = L_i(y_1, y_2)x_3 + M_i(x_1, x_2)y_3 + N_i(x_1, x_2; y_1, y_2).$$

- (ii) *The image of  $C$  in  $\mathbb{P}^1 \times \mathbb{P}^1$  via  $|D - Q| \times |D' - Q|$  is defined by the  $(2, 2)$ -form*

$$F(x_1, x_2; y_1, y_2) = \begin{vmatrix} L_1 & M_1 & N_1 \\ L_2 & M_2 & N_2 \\ L_3 & M_3 & N_3 \end{vmatrix}.$$

- (iii) *We have  $\Delta(F) = \Delta(S)$ .*

*Proof.* We map  $C \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$  via  $((x_1 : x_2), (y_1 : y_2))$ . The first two statements are clear. For (iii) we checked by a generic calculation that  $F$  and  $S$  have the same invariants  $c_4$  and  $c_6$ .  $\square$

**Lemma 5.5.** *Let  $F$  be a  $(2, 2)$ -form defining  $C \subset \mathbb{P}^1 \times \mathbb{P}^1$  a smooth curve of genus one, embedded via  $|D| \times |D'|$ .*

(i) *If  $Q = ((1 : 0), (1 : 0)) \in C(K)$  then we can write*

$$F(x_1, x_2; y_1, y_2) = \begin{pmatrix} x_1^2 & x_1x_2 & x_2^2 \end{pmatrix} \begin{pmatrix} 0 & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} y_1^2 \\ y_1y_2 \\ y_2^2 \end{pmatrix}.$$

(ii) *The image of  $C$  in  $\mathbb{P}^2 \times \mathbb{P}^2$  via  $|D + Q| \times |D' + Q|$  is defined by the  $3 \times 3 \times 3$  cube  $S$  with entries*

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & a_{12} & a_{13} \\ -1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & -1 \\ 0 & a_{21} & 0 \\ 0 & a_{31} & 0 \end{pmatrix}.$$

(iii) *We have  $\Delta(S) = \Delta(F)$ .*

*Proof.* We have  $D \sim Q + R$  and  $D' \sim Q + R'$  where  $R = ((1 : 0), (-a_{13} : a_{12}))$  and  $R' = ((-a_{31} : a_{21}), (1 : 0))$ . Choosing bases for the space of bilinear forms vanishing at  $R'$ , and the space of bilinear forms vanishing at  $R$ , we find that the map  $C \rightarrow \mathbb{P}^2 \times \mathbb{P}^2$  via  $|D + Q| \times |D' + Q|$  is given by

$$\begin{aligned} ((x_1 : x_2), (y_1 : y_2)) &\mapsto (((a_{21}x_1 + a_{31}x_2)y_1 : x_1y_2 : x_2y_2), \\ &\quad (x_1(a_{12}y_1 + a_{13}y_2) : x_2y_1 : x_2y_2)). \end{aligned}$$

The image is defined by

$$\begin{aligned} B_1 &= x_2y_1 + x_1y_2 + a_{22}x_2y_2 + a_{32}x_3y_2 + a_{23}x_2y_3 + a_{33}x_3y_3 \\ B_2 &= -x_3y_1 + a_{12}x_2y_2 + a_{13}x_2y_3, \\ B_3 &= -x_1y_3 + a_{21}x_2y_2 + a_{31}x_3y_2. \end{aligned}$$

The coefficients of these forms give the cube  $S$  in the statement of the lemma. Again we prove (iii) by a generic calculation.  $\square$

*Proof of Theorem 5.3.* We split into the cases  $n = 2$  and  $n = 3$ .

**Case  $n = 2$ .** We have  $D \sim 3.0_E - Q$  for some  $Q \in E(K)$ . By the special case of the theorem already established, there is an integral  $3 \times 3 \times 3$  cube representing  $(E, [D + Q], [D' + Q])$ , with the same discriminant as (18). We have  $E \subset \mathbb{P}^2 \times \mathbb{P}^2$ . Since  $\mathrm{SL}_3(\mathcal{O}_K)$  acts transitively on  $\mathbb{P}^2(K)$  we may assume that  $Q = ((0 : 0 : 1), (0 : 0 : 1))$ . Then Lemma 5.4 gives an integral  $(2, 2)$ -form representing  $(E, [D], [D'])$ , with the same discriminant as (18).

**Case  $n = 3$ .** We have  $D \sim 2.0_E + Q$  for some  $Q \in E(K)$ . By the special case of the theorem already established, there is an integral  $(2, 2)$ -form representing  $(E, [D - Q], [D' - Q])$ , with the same discriminant as (18). We have  $E \subset \mathbb{P}^1 \times \mathbb{P}^1$ . Since  $\text{SL}_2(\mathcal{O}_K)$  acts transitively on  $\mathbb{P}^1(K)$  we may assume that  $Q = ((1 : 0), (1 : 0))$ . Then Lemma 5.5 gives an integral  $3 \times 3 \times 3$  cube representing  $(E, [D], [D'])$ , with the same discriminant as (18).  $\square$

This completes the proof of Theorem 5.1 for  $(2, 2)$ -forms and  $3 \times 3 \times 3$  cubes. We now deduce the result for hypercubes from the result for  $(2, 2)$ -forms. Let  $H$  be a non-singular hypercube over  $K$ , with associated  $(2, 2)$ -forms  $F_{ij}$ . The genus one curve  $\mathcal{C}_H$  is that defined by any of the  $F_{ij}$ . So if  $\mathcal{C}_H(K) \neq \emptyset$  then the result for  $(2, 2)$ -forms shows that each  $F_{ij}$  has minimal level 0. By the definitions in Lemmas 2.2 and 4.1, we have  $\ell(H) = \min \ell(F_{ij})$ . It follows by Corollary 4.3 that  $H$  has minimal level 0.

**Remark 5.6.** We give some examples to show that the minimal level can be positive. We assume for convenience that  $\text{char}(k) \neq 2, 3$ . A binary quartic, or ternary cubic is called *critical* (see [3, Section 5]) if the valuations of its coefficients satisfy

$$= 1 \quad \geq 2 \quad \geq 2 \quad \geq 3 \quad = 3 \quad \text{or} \quad \begin{array}{ccc} & = 2 & \\ & \geq 2 & \geq 2 \\ \geq 1 & \geq 1 & \geq 2 \\ = 0 & \geq 1 & \geq 1 & = 1 \end{array}$$

We now define a *critical*  $(2, 2)$ -form,  $3 \times 3 \times 3$  cube or  $2 \times 2 \times 2 \times 2$  hypercube, to be one whose coefficients have valuations satisfying

$$\begin{array}{ccc} = 2 & \geq 2 & = 1 \\ \geq 2 & \geq 1 & \geq 1 \\ = 1 & \geq 1 & = 0 \end{array}$$

or

$$\begin{array}{ccccccc} \geq 2 & = 1 & \geq 1 & = 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & = 0 \\ = 1 & \geq 1 & \geq 1 & \geq 1 & \geq 1 & = 0 & \geq 1 & = 0 & \geq 0 \\ \geq 1 & \geq 1 & = 0 & \geq 1 & = 0 & \geq 0 & = 0 & \geq 0 & \geq 0 \end{array}$$

or

$$\begin{array}{ccc|ccc} \geq 2 & = 1 & & = 1 & \geq 1 & \\ = 1 & \geq 1 & & \geq 1 & = 0 & \\ \hline = 1 & \geq 1 & & \geq 1 & = 0 & \\ \geq 1 & = 0 & & = 0 & \geq 0 & \end{array}$$

Either by using our algorithms, or observing that the corresponding binary quartics and ternary cubics are critical, we see that any such model  $\Phi$  is minimal. However by applying the transformation

$$[\pi^{-2}, A_2, A_2], [\pi^{-4/3} A_3, A_3, A_3] \text{ or } [\pi^{-3/2} A_2, A_2, A_2, A_2],$$

where

$$A_2 = \begin{pmatrix} 1 & & \\ & \pi^{1/2} & \\ & & \end{pmatrix} \text{ and } A_3 = \begin{pmatrix} 1 & & \\ & \pi^{1/3} & \\ & & \pi^{2/3} \end{pmatrix},$$

we see that  $I(\Phi) \equiv 0 \pmod{\pi^p}$  for any invariant  $I$  of weight  $p$ . Therefore  $\Phi$  has positive level.

#### REFERENCES

- [1] M. BHARGAVA AND W. HO, *Coregular spaces and genus one curves*, Camb. J. Math. **4** (2016), no. 1, 1–119.
- [2] B.J. BIRCH AND H.P.F. SWINNERTON-DYER, *Notes on elliptic curves I*, J. reine angew. Math. **212** (1963) 7–25.
- [3] J.E. CREMONA, T.A. FISHER AND M. STOLL, *Minimisation and reduction of 2-, 3- and 4-coverings of elliptic curves*, Algebra & Number Theory **4** (2010), no. 6, 763–820.
- [4] T.A. FISHER, *Minimisation and reduction of 5-coverings of elliptic curves*, Algebra & Number Theory **7** (2013), no. 5, 1179–1205.
- [5] T.A. FISHER AND R.D. NEWTON, *Computing the Cassels-Tate pairing on the 3-Selmer group of an elliptic curve*, Int. J. Number Theory **10** (2014), no. 7, 1881–1907.
- [6] J.H. SILVERMAN, *The arithmetic of elliptic curves*, Graduate Texts in Mathematics **106**, Springer-Verlag, New York, 1986.

TOM FISHER, UNIVERSITY OF CAMBRIDGE, DPMMS, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, UK.

*E-mail address:* T.A.Fisher@dpmms.cam.ac.uk

*URL:* <https://www.dpmms.cam.ac.uk/~taf1000/>

LAZAR RADIČEVIĆ, UNIVERSITY OF CAMBRIDGE, DPMMS, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, UK.

*E-mail address:* lr397@cam.ac.uk