

RESEARCH ARTICLE

10.1029/2017JG004136

Key Points:

- Aquatic dynamics at Dove Lake are modulated by climate- and fire-driven terrestrial vegetation changes
- A period of high rainforest cover prior ca. 6 ka is linked to changing dystrophic conditions, lower light penetration depths, and anoxic conditions in the lake bottom waters
- Increasing sclerophyll cover after ca. 6 ka is associated with lower nutrient input, lower dystrophy, more oxic conditions, and higher light availability for aquatic organisms

Supporting Information:

- Figure S1

Correspondence to:

M. Mariani,
michela.mariani@unimelb.edu.au

Citation:

Mariani, M., Beck, K. K., Fletcher, M.-S., Gell, P., Saunders, K. M., Gadd, P., & Chisari, R. (2018). Biogeochemical responses to Holocene catchment-lake dynamics in the Tasmanian World Heritage Area, Australia. *Journal of Geophysical Research: Biogeosciences*, 123, 1610–1624. <https://doi.org/10.1029/2017JG004136>

Received 31 AUG 2017

Accepted 22 APR 2018

Accepted article online 30 APR 2018

Published online 19 MAY 2018

Biogeochemical Responses to Holocene Catchment-Lake Dynamics in the Tasmanian World Heritage Area, Australia

Michela Mariani¹ , Kristen K. Beck¹ , Michael-Shawn Fletcher¹ , Peter Gell², Krystyna M. Saunders³, Patricia Gadd³ , and Robert Chisari³

¹School of Geography, University of Melbourne, Parkville, Victoria, Australia, ²Faculty of Science and Technology, Federation University, Mt Helen, Victoria, Australia, ³Australian Nuclear Science and Technology Organization, Lucas Heights, New South Wales, Australia

Abstract Environmental changes such as climate, land use, and fire activity affect terrestrial and aquatic ecosystems at multiple scales of space and time. Due to the nature of the interactions between terrestrial and aquatic dynamics, an integrated study using multiple proxies is critical for a better understanding of climate- and fire-driven impacts on environmental change. Here we present a synthesis of biological and geochemical data (pollen, spores, diatoms, micro X-ray fluorescence scanning, CN content, and stable isotopes) from Dove Lake, Tasmania, allowing us to disentangle long-term terrestrial-aquatic dynamics through the last 12 kyear. We found that aquatic dynamics at Dove Lake are tightly linked to vegetation shifts dictated by regional hydroclimatic variability in western Tasmania. A major shift in the diatom composition was detected at ca. 6 ka, and it was likely mediated by changes in regional terrestrial vegetation, charcoal, and iron accumulation. High rainforest abundance prior ca. 6 ka is linked to increased terrestrially derived organic matter delivery into the lake, higher dystrophy, anoxic bottom conditions, and lower light penetration depths. The shift to a landscape with a higher proportion of sclerophyll species following the intensification of El Niño-Southern Oscillation since ca. 6 ka corresponds to a decline in terrestrial organic matter input into Dove Lake, lower dystrophy levels, higher oxygen availability, and higher light availability for algae and littoral macrophytes. This record provides new insights on terrestrial-aquatic dynamics that could contribute to the conservation management plans in the Tasmanian World Heritage Area and in temperate high-altitude dystrophic systems elsewhere.

1. Introduction

Terrestrial and aquatic ecosystem dynamics are influenced by climatic change (e.g., Ball et al., 2010; Fritz & Anderson, 2013; Massaferro et al., 2013), land-use/vegetation alteration (e.g., Cooper et al., 2015; Kissman et al., 2017), and fire regime shifts (e.g., Araneda et al., 2013; Bixby et al., 2015; Brown et al., 2014) at multiple scales of space and time. Given the importance and the interconnectedness of both climate and fire (e.g., Emelko et al., 2016; Fletcher et al., 2014; Power et al., 2016), and terrestrial and aquatic ecosystems (Beck, Fletcher, Kattel, et al., 2018; Kissman et al., 2017; Strock et al., 2017), the paucity of research on how climate and fire drive terrestrial and aquatic ecosystem dynamics constitutes a critical knowledge gap that potentially undermines effective management and conservation endeavors. This lack of cross-cutting research is particularly salient in Australia, where fires are key ecological agent that both has shaped the unique flora of the region (Bowman, 2000) and threatens relict fire-sensitive plant systems with extinction (Holz et al., 2014). Little is known about how fire-driven vegetation dynamics influence aquatic ecosystems in this region, exposing a concerning lack of understanding. This knowledge gap is important in the context of a predicted increase in fire activity over the upcoming centuries (Moritz et al., 2012), and it constitutes a significant impediment to effective management of aquatic systems in fire-prone environments. Here we integrate data sets of terrestrial and aquatic ecosystem dynamics spanning the past 12,000 years (12 kyear; kyear = duration in thousands of years) within the Tasmanian World Heritage Area, Australia, in an attempt to understand how these systems respond to inferred shifts in climate and fire activity through this time.

Fires can drive substantial changes in terrestrial vegetation and landscape processes, such as soil alteration and erosion, which can mobilize nutrients from soils, consequently increasing nutrient loading into streams, rivers, lakes, and reservoirs (Lane et al., 2008; Moody & Martin, 2001, 2009; Reneau et al., 2007; Sheridan et al.,

2007; Sherson et al., 2015; Wilkinson et al., 2009). Fire can also affect soil chemistry by altering organic matter content, cation exchange capacity, pH, electrical conductivity, and nutrient concentration (e.g., Bixby et al., 2015; Cerrato et al., 2016; Fletcher et al., 2014; González-Pérez et al., 2004; Knoepp et al., 2009; Kutiel & Inbar, 1993; Raison, 1979; Robinne et al., 2018). These fire-driven processes can influence the behavior of micronutrients, such as redox-sensitive iron (Fe) and manganese (Mn; Calvert & Pedersen, 2007; Davison, 1993; Engstrom & Wright, 1984; Mackereth, 1966; Och et al., 2012; Wersin et al., 1991), potentially leading to anoxia of bottom waters through reductive dissolution and (re)oxidation, which progressively leads to geochemical focusing. This process transfers and enriches these elements in deeper waters and can drive the development of anoxic bottom waters (Naehler et al., 2013; Schaller & Wehrli, 1996). Anoxia within aquatic systems is associated with biodiversity loss, nutrient availability, and food web alterations (Diaz, 2001; Hughes et al., 2015; North et al., 2014). While the influence of redox conditions over lake water oxygen dynamics is well understood (e.g., Balistrieri et al., 1992; Davison, 1993; Sigg, 2000), little is known about how long-term changes in terrestrial processes influence within lake dynamics, especially in temperate Australia.

Long-term disturbance from fire in fire-sensitive vegetation, such as rainforest, can result in substantial ecosystem transformation, such as altered species composition, soil formation, and nutrient loss that can have important implications for associated water bodies (Ball et al., 2010; Huvane & Whitehead, 1996; Korhola et al., 1996; Leys et al., 2016; Morris et al., 2015; Smith et al., 2011). Fires in temperate environments, such as western Tasmania, mostly occur in the period from September to March (from spring to early autumn) and are often followed by heavy rain events (Bridle et al., 2003; Pemberton, 1988), which can remove the soil layer into water bodies, altering water geochemistry and nutrient availability (Beck, Fletcher, Kattel, et al., 2018; Boerner, 1982) and precipitating an aquatic ecosystem response (Beck, Fletcher, Gadd, et al., 2018). Disturbance from repeated fire in the rainforest of Tasmania, for example, are associated with the destruction and complete erosion of highly organic soil profiles, localized plant species extinctions, and invasion by fire-promoting vegetation that can radically alter fire-vegetation-soil dynamics. Investigation on the effects of these changes in the dystrophic aquatic environments of Tasmania has revealed a tight coupling between fire-driven changes in the terrestrial environment and aquatic ecosystem dynamics, via the influence of organic soils over aquatic trophic status (Beck, Fletcher, Kattel, et al., 2018; Fletcher et al., 2014) and the influence of terrigenous input over lake water pH (Beck, Fletcher, Gadd, et al., 2018). While the recent work by Beck, Fletcher, Gadd, et al. (2018) reveals a tight coupling between terrestrial and aquatic ecosystem changes in response to fire, there is little empirical evidence for the influence of long-term changes in fire and vegetation over important within-lake processes, such as lake mixing, light availability changes, and redox conditions.

Here we present a new lake sediment geochemical (micro X-ray fluorescence [μ XRF] scanning, CN content, and stable isotopes) and biological (diatoms) proxy data set from a relatively large and deep dystrophic lake (Dove Lake) located in a cool temperate west of Tasmania, Australia. We compare our new proxy data set from Dove Lake to a recent pollen-based quantitative vegetation reconstruction from the same sediment sequence that reveals marked climate- and fire-driven changes in vegetation land-cover over the last 12 kyear (Mariani et al., 2017). These changes are characterized by shifts in the importance of high biomass rainforest vegetation and lower biomass sclerophyll-dominated vegetation. Peat soils predominate in this cool temperate and oligotrophic environment, with rainforest typically developing atop deep peat profiles, while sclerophyll vegetation typically produces shallow to skeletal peat profiles with a lower nutrient content (Beadle, 1966; Jackson, 2000). Like other temperate blanket peat environments (Bergström et al., 2001; Hansen, 1962; Steinberg et al., 2003), water bodies in western Tasmania receive substantial amounts of terrestrially derived organic matter and are consequently acidic, oligotrophic, and dystrophic (Buckney & Tyler, 1973; Tyler, 1992), with an elevated content of reactive Fe and Mn (Mackey et al., 1996). Given the high nutrient limitation in this region, the entire nutrient pool supporting terrestrial vegetation development only derives from the accumulation of organic debris. This implies that the nutrient input into water bodies is almost entirely derived from these acid peat soils, producing low-nutrient tea-colored waters (i.e., oligotrophic and dystrophic/humic lakes; Tyler, 1974, 1992). We hypothesize that (i) phases of maximum rainforest cover around Dove Lake will be associated with deeper catchment peat profiles that deliver elevated amounts of terrestrial organic matter, along with carbon, nitrogen, Fe, and Mn into the lake; (ii) the increase in sclerophyll vegetation cover will be




Figure 1. Introductory figure showing (a) site location (upper panel) and climatic diagram (lower panel). (b) 3D projection of Dove Lake watershed and bathymetric profile. Lower panel in (b) represents the depth profile from point A to point B (direction as indicated in the map). Red star in (b) indicates the location of the coring site. A detailed bathymetric map of Dove Lake is presented in supporting information Figure S1. (c) 3D projection of the Dove Lake watershed overlaid with the geological map. (d) 3D projection of the Dove Lake overlaid with the vegetation map (data from <https://www.thelist.tas.gov.au/app/content/home>).

associated with a shift toward shallower peat profiles and a concomitant decrease in nutrient supply into the lake; and (iii) changes in terrestrially derived organic matter will drive changes in within-lake redox conditions, light and oxygen availability, and diatom compositional changes.

1.1. Western Tasmania

Tasmania is a large cool temperate island located between 41–44°S (Figure 1) characterized by a temperate maritime climate with mild winters and cool summers (Gentili, 1972). A rain-shadow is produced by the prevailing westerly winds that rise over the northwest to southeast trending mountain range bisecting Tasmania. The regional geology of western Tasmania is dominated by highly resistant and inert quartz-dominated rock types that contribute little mineral matter to soils, resulting in the predominance of poor acidic soils (organosols/peats; di Folco, 2007; Tyler, 1992). Indeed, the ionic composition of Tasmanian lake water reflects a minimal geological contribution (Buckney & Tyler, 1973). The cool temperate climate, extreme oligotrophy, and predominance of organosols result in ubiquitously moderately to highly dystrophic and acidic waterbodies that has a thermal behavior from warm monomictic to polymictic, with some evidence of meromixis (or partial meromixis; Bowling & Tyler, 1984; Buckney & Tyler, 1973; King & Tyler, 1981; Tyler, 1974).

The modern vegetation landscape of western Tasmania is characterized by a mosaic of rainforest, moorland, and sclerophyll vegetation (Kirkpatrick & Dickinson, 1984) that has resulted from the long-term application of fire by people through the Late Pleistocene and Holocene (Fletcher & Thomas, 2010; Jackson, 1968; Mount, 1979; Thomas, 1995). The type of extant vegetation is intimately linked to peat development and nutrient availability for aquatic systems (Beck, Fletcher, Kattel, et al., 2018; Bowman & Jackson, 1981; Bridle et al., 2003). For instance, soils formed under sclerophyll vegetation in Tasmania are thinner and have significantly lower nutrient content compared to soils developed under rainforest vegetation (Bowman & Jackson, 1981; Jackson, 1968; Wood & Bowman, 2011), and fire-driven shifts from rainforest to more sclerophyll-dominant vegetation in Tasmania is associated with a shift to a more oligotrophic aquatic environment (Beck, Fletcher, Kattel, et al., 2018; Fletcher et al., 2014).

1.2. Study Site

Dove Lake (41°39′34.27″S, 145°57′35.14″E) is a large (90-ha) dystrophic subalpine lake (940 m a.s.l.) located within the Tasmanian World Heritage Area in Australia (Figure 1). Average total annual rainfall at the nearest meteorological station (Cradle Valley; 2.7 km from Dove Lake; 41°38′24.23″S, 145°56′24.28″E; 903 m a.s.l.) is approximately 2,700 mm/year (Figure 1a). The lake has a relatively steep bathymetric profile characterized by two deep basins (Figure 1b; supporting information Figure S1). Maximum depth was recorded approximately 60 m in the south basin, whereas a maximum depth of approximately 45 m was recorded in the northern basin (core location). The geology of the watershed is represented by two dominant types: quartzite and Quaternary glacial deposits (33.5% and 45.9%, respectively; Figure 1c). The modern catchment vegetation is dominated by wet sclerophyll forest on the eastern and western flanks, treeless fire-promoted moorland on the northern edge, and rainforest in the southwest corner (Figure 1d). Above approximately 1,000 m a.s.l., the landscape is dominated by highland and treeless vegetation, which represents 45.1% of the Dove Lake catchment. Important species in the modern landscape are *Eucalyptus coccifera*, *Lophozonia cunninghamii* (syn. *Nothofagus cunninghamii*), *Gymnoschenous sphaerocephalus*, *Athrotaxis selaginoides*, and a variety of ericaceous shrubs. Within the lake catchment and across the surrounding plateau, typical montane rainforest trees are also found, such as *Fuscospora gunnii* (syn. *Nothofagus gunnii*), *A. selaginoides*, and *Athrotaxis cupressoides* (Cupressaceae).

A quantitative vegetation reconstruction was previously carried out at this location (Mariani et al., 2017), which indicated the dominance of fire-promoted treeless moorland over the last 12 kyear. Forest cover was at approximately 40% between 10 and 4 ka (ka = specific point in time in thousands of years), peaking at approximately 50% of the surrounding landscape between 8 and 7 ka in response to a shift to a wetter hydroclimate and a reduction in regional fire activity (Fletcher & Moreno, 2011, 2012; Mariani & Fletcher, 2017; Wilkins et al., 2013; Xia et al., 2001). A shift to a drier and more variable hydroclimate after ca. 5.5 ka (Fletcher & Moreno, 2012; Mariani & Fletcher, 2017) drove an increase in regional biomass burning and a replacement of areas of rainforest by fire-promoting sclerophyll plant taxa, such as the fire-adapted *Eucalyptus* (Beck et al., 2017; Beck, Fletcher, Kattel, et al., 2018; Fletcher et al., 2014; Mariani et al., 2017; Stahle et al., 2016, 2017). Subsequently, during the late Holocene, a decrease in moisture availability and a synchronous phase of increased fire activity across western Tasmania (Fletcher et al., 2015; Mariani & Fletcher, 2017) resulted in a rainforest decline around Dove Lake (Mariani et al., 2017). These hydroclimatic changes are likely the long-term (millennial scale) effect of increasingly frequent El Niño events in the tropical Pacific (Donders et al., 2008; McGlone et al., 1992; Moy et al., 2002), which would have resulted in overall cooler than normal waters over the western tropical Pacific and high pressure systems driving negative moisture anomalies in eastern Australia. Today, El Niño events are associated with drier conditions and increased fire activity in southeast Australia today (Mariani et al., 2016; Nicholls & Lucas, 2007).

2. Methods

The 122-cm-long organic sediment core from Dove Lake was extracted in December 2015 using a 6.8-cm-diameter polycarbonate chamber attached to a Universal Gravity Corer (http://www.aquaticresearch.com/universal_core_head.htm). The core chronology was obtained using ^{210}Pb and ^{14}C dating methods (supporting information Tables S1a and S1b), and the age-depth model was previously published in Mariani et al. (2017; supporting information Figure S2).

2.1. Charcoal and Pollen Analysis

Macroscopic ($>125\ \mu\text{m}$) and microscopic ($<125\ \mu\text{m}$) charcoal content was analyzed to document the local fire history, and the results were previously presented in Mariani et al. (2017). Macroscopic charcoal content was measured using $1.25\ \text{cm}^3$ taken from 5-mm-thick (median time resolution = ca. 48 years) contiguous samples, and microscopic charcoal content was extracted from 0.5-cm^3 samples taken at 1-cm resolution. The macroscopic and microscopic charcoal counts were converted into accumulation rates (CHAR, particles cm^2/year), and for the remainder of this manuscript, they will be referred as macroscopic-CHAR and microscopic-CHAR, respectively.

On the same sequence, pollen analysis was performed at 1-cm resolution to investigate the vegetation history, and the results were previously shown in Mariani et al. (2017). The ratio between rainforest and

sclerophyll plant taxa (rainforest:sclerophyll) was calculated from the land-cover data published in Mariani et al. (2017) prior to the elaboration of statistics (see below). From the same data set, *Isoëtes*, a littoral macrophyte, spore abundances were considered as proxy for aquatic conditions, such as lake level and light penetration variation (Bogotá-A et al., 2016; Chappuis et al., 2015; Finkenbinder et al., 2014; Pesce & Moreno, 2014; Simi et al., 2017).

2.2. μ XRF Core Scanning

The μ XRF core scanning undertaken at the Australian Nuclear Science and Technology Organisation provided 0.5-mm-interval nondestructive elemental analyses of the Dove Lake core. Manganese:iron (Mn:Fe) and iron:titanium (Fe:Ti) ratios were calculated to document changes in redox conditions at the water-sediment interface (Corella et al., 2012; Cuven et al., 2010; Habertzettl et al., 2007).

2.3. Carbon and Nitrogen

Analysis for percent carbon (%C), percent nitrogen (%N), $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ for Dove Lake were performed at a resolution of 1 cm (median time resolution = ca. 96 years) at the Australian Nuclear Science and Technology Organisation facility for stable isotope analyses using standard procedures (Ohlsson & Wallmark, 1999). Briefly, the crushed and dried samples were weighed into tin capsules and introduced sequentially into an elemental analyzer (Thermo Fisher Flash 2000 HT EA) using an autosampler. No pretreatment was performed as no carbonates were detected in preliminary tests executed on the pollen samples from Mariani et al. (2017).

2.4. Diatoms

Diatom analysis was performed at an irregular resolution and volumes throughout the Dove Lake core due to sediment availability. Samples were processed using standard methods (Battarbee, 1986). Samples were placed in a hot bath, and a series of hydrogen peroxide additions were used to remove all organic matter. Sample dilutions were mounted with Naphrax[®], and at least 300 diatom valves were identified to genus level using an oil immersion differential interference contrast light microscope at 1,000X magnification. Diatoms were identified to genera rather than species level in order to observe the changes in diatom habitat to shifts in climate and terrestrial environmental dynamics, rather than a detailed ecological analysis of the lake.

2.5. Statistical Analyses

The μ XRF core scanning and diatom data from Dove Lake were summarized using principal curves (PCs) in R 3.3.1 (R Development Core Team, 2013). PCs (Hastie & Stuetzle, 1989) are a generalized form of the first principal component axis as a smooth, one-dimensional curve fitted through the data in multiple dimensions such that the curve best fits the data. Thus, distances of the samples to the PC are somewhat minimized (Simpson & Birks, 2012). PCs are considered to perform substantially better than principal component analysis and correspondence analysis in many cases, especially when data are dominated by a single dominant gradient (Simpson & Birks, 2012). Prior the PC analysis, μ XRF data were normalized by dividing all elemental counts by the total counts per second (Croudace & Rothwell, 2015) and binned to 50-year intervals to facilitate the statistical performance. All elements, but not ratios, were included in the PC analysis. Diatom data were square root transformed to correct for taxon skewness.

Generalized additive models (GAMs) were used to identify significant trends in the Dove Lake vegetation, charcoal influx, μ XRF core scanning, diatoms, and carbon and nitrogen data. The additive models do not involve a priori parameter setting, instead allowing the shape of the relationship to be determined from the data using penalized regression (Hastie & Tibshirani, 1990). All GAMs were fitted using the Mixed GAM Computation Vehicle (mgcv) package (Wood & Wood, 2007) for R 3.3.1 (R Development Core Team, 2013). To highlight periods of significant shifts in all time series analyzed with GAMs, the first derivative of the GAM splines were calculated (<https://github.com/gavinsimpson/tsgam>). This approach allows measurement of the slope at each point throughout a time series, thus enabling us to extract trends in the paleodata (Bunting et al., 2016). When slope exceeds confidence intervals, the relative time period is considered as "periods of significant change" of a certain time series. Significance level for the derivative calculations was set to 0.95.




Figure 2. Charcoal and geochemical results for the Dove Lake core. Black step-like curves are the results from microscopic and macroscopic charcoal analyses. Black scatter lines represent the results of carbon and nitrogen analyses; red curves represent selected micro X-ray fluorescence (μ XRF) elements and ratios (50-years binned). μ XRF elements counts were normalized by total counts per second (tcps). μ XRF principal curve includes all elements listed in supporting information Table S2.

3. Results

3.1. Charcoal and Pollen Analysis

The macroscopic-CHAR and microscopic-CHAR record reveal substantial changes in fire regimes throughout the record (Figure 2). In both records, relatively high accumulation rates are observed prior to ca. 6 ka with a peak occurring at 7.3 ka. A gradual decline is visible since ca. 6 ka and interrupted by an important peak at 3.7 ka, then the data reach minimum values between 2.5 and 0.5 ka. A total of 109 samples from Dove Lake were analyzed for pollen and spores content, and results were presented in Mariani et al. (2017). The ratio rainforest:sclerophyll vegetation calculated from the data of Mariani et al. (2017) shows high values between 10 and 6 ka, followed by a pronounced decline occurring between 6 and 4 ka. The littoral macrophyte *Isoetes* displays relatively low and stable abundances between 11.7 to 6.5 ka, then increases substantially (up to three times the initial values) during the period between 6.5 and 3 ka. A decline is evident since ca. 3 ka, but the percent abundance is maintained higher than the Early to mid-Holocene values.

3.2. μ XRF Core Scanning

A total of 4,578 data points was recorded on the Dove Lake core, and results are presented in Figure 2 (red curves). Selected elemental data obtained by this analysis revealed high counts of iron (Fe), the most abundant element in the spectrum. A gradual decline in Fe is detected between 11.7 and 5.2 ka with a sharper decline between ca. 7 and 5.2 ka, when Fe counts reach minimum values until >0.1 ka (Figure 2). Titanium (Ti) displays very low counts and the opposite trend of Fe, whereas manganese (Mn) counts are persistently low until ca. 3.5 ka, then gradually increase (Figure 2). The Mn:Fe ratio is an indicator for reducing conditions at the water-sediment interface (Corella et al., 2012; Cuven et al., 2010; Haberzettl et al., 2007) and shows fairly consistent high values prior to ca. 6 ka with a gentle decline evident between ca. 7.5 and 6 ka. The Fe:Ti ratio has been employed with the same scope as the Mn:Fe (Aufgebauer et al., 2012) as they show virtually identical trends, although specular (Figure 2).

3.3. Carbon and Nitrogen

A total of 92 samples were analyzed for %C, %N, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ (Figure 2). The percentages of C, N, and the C:N ratio show a slow gradual increase from ca. 11.7 to 8 ka and then become more stable until around 6 ka. At this point, a substantial, although gradual, decline occurs in these proxies, alongside a marked variation in the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values. The amount of C declines from 19% to 12%, whereas %N drops from 0.9% to 0.6%. Analogously, the ratio C:N follows a gradual decline from values clustered around 22 to around 18. The C:N




Figure 3. Results of the diatom analysis for selected taxa and diatom principal curve (PC). Red bars identify taxa with positive Pearson correlation values with the PC. Blue bars identify taxa with negative Pearson correlation values with the PC. Dashed lines represent the significant zonations (more detail in supporting information Figure S3 and Document S1). The asterisk (*) indicates that *Tabellaria* does not show a strong correlation with the PC ($r = 0.3$), but given its abundance in the record, its trend was considered important for the discussion.

has a mean value of 19.46 suggesting a high terrestrial component of this metric (Meyers & Teranes, 2002). $\delta^{13}\text{C}$ gradually increases between 6.2 to 3.6 ka from values around -26.7‰ to values around -26.3‰ . Given the extremely low variation in $\delta^{13}\text{C}$, this proxy was not an object of discussion. Meanwhile, $\delta^{15}\text{N}$ gradually increases from 6.1 to 3.5 ka with values ranging from 2‰ to 3.7‰ . Minerogenic flux was roughly estimated by mass difference using the summed C and N contents, then converted into accumulation rates (cm^2/year ; supporting information Figure S4).

3.4. Diatoms

A total of 37 diatom samples were counted, and selected genera are presented in Figure 3. The most abundant taxon in the spectrum is *Frustulia* spp., displaying higher abundances before ca. 9.8 ka and after ca. 5 ka (approximately 40%). Although less abundant in the record ($<5\%$), *Aulacoseira* spp. and *Gomphonema* spp. show important trends analogous to *Frustulia* spp. *Eunotia* spp., *Brachysira* spp., and *Actinella* spp. display relatively low abundances between ca. 11.7 and 9.8 ka and post ca. 5 ka, whereas higher percentages are recorded between ca. 9.8 and 5 ka. *Eunotia* spp. is the second most abundant taxon after *Frustulia* spp., reaching maximum values around 30% at ca. 6 ka. *Tabellaria* spp. shows relatively high percentages (8–10%) prior to ca. 5 ka, dropping to values around 5% afterward. An extended diatom stratigraphic plot is displayed in supporting information Figure S3, alongside the CONISS (stratigraphically constrained cluster analysis) zonation description (Document S1).

3.5. Statistical Analyses

Results from the PC analyses on the μXRF scanning and diatoms data are presented in Figures 2 and 3 (respectively). The μXRF data PC explains approximately 46% of the total variance in the data set. Iron (Fe)




Figure 4. Results from the application of generalized additive models and first derivative calculations on selected time series. Curve segments highlighted in red identify periods of significant increasing values, whereas blue highlights period of significant decrease in the time series. Significance levels for derivative calculations were set to 0.95.

shows the highest positive correlation with the PC ($r = 0.93$), whereas Ti displays a significant negative correlation with this curve ($r = -0.83$), alongside other detrital elements, such as potassium (K) and zirconium (Zr; supporting information Table S2). Fe:Ti and Mn:Fe show very high positive correlation values with the μ XRF data PC (0.85 and -0.49 , respectively).

The diatom PC explains approximately 31% of the variance of the assemblage data set. *Gomphonema* spp., *Aulacoseira* spp., and *Frustulia* spp. show the three highest negative correlations with the PC (-0.79 , -0.75 , and -0.65 , respectively). On the positive end of the correlation values spectrum, *Eunotia* spp., *Brachysira* spp., and *Actinella* spp. show the three highest values (0.77, 0.69, and 0.62 respectively). *Tabellaria* spp. display a significant positive correlation of 0.3 with the diatom PC. A table with the Pearson correlation values for all taxa is presented in supporting information Table S3.




Figure 5. Summary figure showing a schematic representation of the environmental changes recorded at Dove Lake based on the time series presented on the right: (a) macroscopic-CHAR; (b) carbon to nitrogen ratio (C:N); (c) rainforest to sclerophyll ratio (R:S); (d) diatoms principal curve; and (e) micro X-ray fluorescence (μ XRF) scanning data principal curve (PC). Drawings are not in scale.

The GAM splines produced for selected proxies from Dove Lake highlight the significant trends in the data by efficiently removing the noise embedded in the time series and, thus, allowing a better comparison of the paleorecords (Figure 4). Similar trends in the GAM splines are especially observed in the diatom PC, C:N, and rainforest:sclerophyll ratios. The analysis of the first derivatives allowed us to identify significant shifts in the time series and a better assessment of synchronicity between proxies (Figure 4). A significant increase in the rainforest:sclerophyll ratio between 11.7 and 9 ka was found at the same time as the significant increase in the diatom PC and the C:N ratio. A decrease in the rainforest:sclerophyll ratio, diatom PC, and C:N ratio between 6.5 and 3 ka is concurrent with an increase in the littoral macrophyte *Isoetes*. The μ XRF data PC follows a persistent declining trend from 11.7 to ca. 5 ka, reaching significant levels between 11.7–10 ka and 7–5 ka. Although there is a higher amount of noise in the data (i.e., peaks), macroscopic-CHAR appears to follow a trend analogous to the μ XRF data PC but reaches significance only between 6.5 and 1.7 ka.

4. Discussion

4.1. Vegetation Change and Macronutrient Dynamics at Dove Lake

Our data demonstrate that changes in vegetation and fire dynamics around Dove Lake in response to long-term changes in hydroclimate and their influence over fire activity (Mariani et al., 2017; Stahle et al., 2016, 2017) have played an important role in driving changes in within-lake dynamics at this site. The climate and fire-driven increase in sclerophyll-dominant vegetation cover (at the expense of rainforest) around Dove Lake at ca. 6 ka (Figure 5) is associated with a substantial decline in carbon and nitrogen content of lake sediment organic matter (Figure 4). This trend is mirrored elsewhere in Tasmania (Beck, Fletcher, Kattel, et al., 2018; Fletcher et al., 2014) and is consistent with a reduction in peat depth in response to a reduction of rainforest cover around Dove Lake (Figure 5c) and lower nutrient status of peat soils that develop under sclerophyll vegetation, relative to rainforest (Bowman & Jackson, 1981; Bridle et al., 2003).

A shift from high relative terrestrial nutrient input into the lake in response to the shift from high relative rainforest cover to high relative sclerophyll cover is supported by (1) the shift from high to low relative C:N ratios at ca. 6 ka (Meyers & Teranes, 2002) and (2) the concomitant shift from low to high relative $\delta^{15}\text{N}$ values

(Figure 2). Aquatic organisms preferentially uptake the lighter ^{14}N isotope, and N-enriched lake water results in the production of organic matter low in ^{15}N (so called Rayleigh Distillation; Talbot & Johannessen, 1992; Talbot, 2002). The high C:N values prior to ca. 6 ka indicate an increase in the delivery of terrestrial organic matter into Dove Lake from the well-developed peat soils within the catchment under high relative rainforest cover. This evidence is supported by an estimate of minerogenic flux (supporting information Figure S4), which shows relatively high fluxes prior to ca. 6 ka and a decline between 6 and 4 ka. The high relative input of terrestrial organic matter would have provided a constant input of ^{14}N and organic matter under an enriched ^{14}N regime, with the shift to increased sclerophyll vegetation and the resultant decrease in, and impoverishment of, terrestrial organic matter forcing the uptake of increased amounts of ^{15}N .

4.2. Changes in Acidity, Dystrophy, Redox Conditions, and Light Availability at Dove Lake

Our data reveal a covariance between millennial-scale trends in diatom community composition within Dove Lake (diatom PC; Figure 3) and trends in both the ratio of rainforest:sclerophyll vegetation and C:N throughout the last 12 kyear (Figures 4 and 5). This covariance indicates a tight coupling between millennial-scale terrestrial and aquatic ecosystem dynamics through this time. A positive correlation between C:N and dystrophy has been identified in cool temperate systems elsewhere (Hansen, 1962), and we interpret the coupling between C:N ratio and the diatom PC as an indication of high relative levels of dystrophy and acidity prior to ca. 6 ka when increased rainforest cover and associated peat profiles delivered high amounts of humic acids and leached terrestrially derived nutrients into the lake (Figure 4). Further, we contend that the high correlations found between the diatom PC with *Eunotia* spp., *Brachysira* spp., and *Actinella* spp. (positive) and with *Gomphonema* spp., *Aulacoseira* spp., and *Frustulia* spp. (negative; Figure 3) track dystrophy and acidity levels. While most diatom taxa in western Tasmania are adapted to high acidity and dystrophic/oligotrophic waters (Vyverman et al., 1996), the diatom compositional shifts represented by the diatom PC reflect shifts between more acidic/more dystrophic to less acidic/less dystrophic conditions (Figure 5). Indeed, *Eunotia* spp., *Brachysira* spp., and *Actinella* spp., which are important components of the diatom community prior to ca. 6 ka, are associated with generally high dystrophy and lower pH in temperate southeast Australia and Tasmania (Hodgson et al., 1996; Philibert et al., 2006; Sonneman et al., 1999; Vyverman et al., 1995, 1996), while taxa that increase in importance after ca. 6 ka, such as *Gomphonema*, *Aulacoseira*, and *Frustulia* species, are associated with slightly higher pH (Philibert et al., 2006; Vyverman et al., 1995, 1996).

Our inference of changes in dystrophy is consistent with changes in the diatom-type *Tabellaria* spp., which shifts gradually from high to low relative values at ca. 6 ka (Figure 3). *Tabellaria flocculosa* is the only member of the *Tabellaria* genus currently found in Dove Lake (<1% of modern composition; Vyverman et al., 1995). *T. flocculosa* displays a marked preference for low-light conditions during the photosynthetic process (Paul & Duthie, 1989), indicating that this species is sensitive to color changes in lake waters that influence light penetration, such as the results from changes in levels of dystrophy. Indeed, *T. flocculosa* and *Tabellaria quadrisepitata* are abundant in dystrophic environments (van Dam et al., 1981), indicating a competitive advantage in low light conditions by members of the *Tabellaria* genus. Thus, we interpret the relatively higher abundance of *Tabellaria* spp. prior to ca. 6 ka as reflecting increased inputs of terrestrially derived organic material into the lake (Figure 4). Moreover, our inferred changes in dystrophy resulting from hydroclimatic and fire-driven changes in terrestrial vegetation and organic soil development are consistent with trends in the littoral macrophyte, *Isoetes*, which shifts from low to high relative values at ca. 6 ka (Figure 4b). While comparatively little is known about the ecology of Tasmanian *Isoetes* species (Garrett & Kantivalis, 1992), evidence indicates an increase in mortality in response to a reduction in the penetration of light through the water column in the widespread European and North American species, *Isoetes lacustris* (Chappuis et al., 2015; Riera et al., 2017). Thus, in combination with the evidence for changes in dystrophy in Dove Lake, we interpret the low relative values of *Isoetes* prior to ca. 6 ka as reflecting a reduction in suitable habitat due to poor light penetration through the highly dystrophic lake waters, with an increase in this aquatic macrophyte after ca. 6 ka reflecting increased light penetration through less dystrophic waters (Figure 5).

We observe a high correlation between the μXRF data PC with Fe:Ti and Mn:Fe ratios that we interpret as indicating changes in redox conditions at the water-sediment interface. Ratios of Mn:Fe have been used repeatedly to reconstruct changing redox conditions in lakes (e.g., Dean & Doner, 2012; Koinig et al., 2003; Loizeau et al., 2001; Melles et al., 2012; Naeher et al., 2013; Wersin et al., 1991), with lower Mn:Fe ratios

associated with lower O₂ concentrations in the water column (e.g., Balistrieri et al., 1992; Davison, 1993; Loizeau et al., 2001; Mackereth, 1966; Neelson & Saffarini, 1994; O'Sullivan & Reynolds, 2008; Wersin et al., 1991). Thus, the substantially lower Mn:Fe ratios prior ca. 6.5 ka indicate lower oxygen abundance at the water-sediment interface (i.e., anoxic bottom waters) during the phase of high relative terrestrial organic matter input into the lake—more organic matter was available for remineralization that consumed more O₂ resulting in anoxia. The shift to lower terrestrial organic matter input after ca. 6 ka, thus, resulted in a reduction in O₂ consumption and the development of more oxic bottom waters (Figure 5).

4.3. Climate- or Biomass-Induced Charcoal Accumulation?

We observe high relative macroscopic charcoal influx into Dove Lake during the phase of high relative forest cover prior to ca. 6 ka, with an overall shift to lower charcoal influx between 6.2 and 1.7 ka in concert with a shift to increased sclerophyll vegetation and overall landscape openness (Figures 4 and 5; Mariani et al., 2017). While inconsistent with the fire ecology of rainforest (pyrophobic), and sclerophyll and open moorland (pyrophytic) vegetation in this region, our results are consistent with a primary control of biomass control over the production of macroscopic charcoal at Dove Lake. Fletcher et al. (2014) report similar reduction in sedimentary macroscopic charcoal in response to fire-driven shifts between pyrophobic and pyrophytic vegetation during the Holocene in Tasmania, concluding that the higher woody biomass and destructive nature of rainforest fires can produce high amounts of macroscopic charcoal relative to fire-promoting vegetation, where less woody material is usually burnt (Fletcher et al., 2014). Indeed, a reduction in macroscopic-CHAR was also observed in New Zealand following pyrophobic forest removal, where there was a decline in biomass availability following burning and deforestation after the Maori (McWethy et al., 2010). Further afield, our data showing high charcoal accumulation in line with high forest cover are analogous to studies from Kenya (Colombaroli et al., 2014), Spain (Gil-Romera et al., 2014), and Arizona (Brunelle et al., 2010), where biomass limitation was indicated as the most plausible explanation for the link between high forest cover and high charcoal influx. Nevertheless, the hypothesis of biomass-limited fire occurrence is unlikely in the generally high biomass environments of western Tasmania (McWethy et al., 2013); thus, it is more likely that a variation in the fuel type, from rainforest to sclerophyll vegetation, would have modulated charcoal influx in the Dove Lake record.

5. Conclusion

A tight coupling between the terrestrial-aquatic systems was found at Dove Lake, reflecting terrestrial organic material inputs into the lake from the vegetation-soil system over the last 12 kyear. The ratio between rainforest and sclerophyll vegetation cover shows synchronicity to shifts in carbon and nitrogen content and isotopic composition, alongside shifts in the diatoms assemblage. Phases of high rainforest cover with deep peat profiles are linked to high carbon, iron, and nitrogen delivery prior ca. 6 ka at Dove Lake. Contrastingly, an increase in sclerophyll abundance in the landscape after the intensification of ENSO since ca. 6 ka corresponds to a decline in nutrient supply to the lake. Moreover, we found that the changes in terrestrially derived organic matter inputs drove relatively synchronous changes in dystrophy and acidity levels, water redox conditions, light availability, and algal ecosystem dynamics.

Light was found to be an important driver of changes in algal composition and aquatic macrophyte abundance. Trends in *Tabellaria* spp. and *Isoëtes* spp. suggest a variation in light penetration depth resulting from dystrophy level changes after ca. 6 ka. Furthermore, the μ XRF core scanning results allowed us to reconstruct changes in redox conditions at the water-sediment interface: low Mn:Fe ratios prior ca. 6 ka probably indicate an oxygen depletion of the water column and anoxic lake bottom attributable to altered lake mixing and metabolism of nutrients. This record provides new insights on the importance of terrestrial inputs in driving light availability changes and aquatic ecosystem dynamics at Dove Lake in the World Heritage Area of Tasmania.

References

- Araneda, A., Jana, P., Ortega, C., Torrejón, F., Bertrand, S., Vargas, P., et al. (2013). Changes in sub-fossil chironomid assemblages in two northern Patagonian lake systems associated with the occurrence of historical fires. *Journal of Paleolimnology*, 50(1), 41–56. <https://doi.org/10.1007/s10933-013-9703-0>

Acknowledgments

This research was supported by Australian Research Council (ARC) grants DI110100019 and IN140100050. M. M. was also supported by an AINSE PGRA scholarship (12039) and the John and Allan Gilmour Science Award 2016 (Faculty of Science, University of Melbourne). Kristen K. Beck would like to thank the Albert Shimmins Fund for their financial support. We acknowledge that our work was conducted on Tasmanian Aboriginal lands and thank the Tasmanian Aboriginal community for their support. We thank Michael Comfort from the Department of Primary Industries, Parks, Water and Environment (DPIPWE) for granting the permit to core Dove Lake. We also thank Simon Connor, Valentina Vanghi, Anthony Romano, and Coralie Tate for assistance in the field. Data produced within the present work will be made available upon publication on NEOTOMA (<https://www.neotomadb.org/>).

- Aufgebauer, A., Panagiotopoulos, K., Wagner, B., Schaebitz, F., Viehberg, F. A., Vogel, H., et al. (2012). Climate and environmental change in the Balkans over the last 17 ka recorded in sediments from Lake Prespa (Albania/FYR of Macedonia/Greece). *Quaternary International*, *274*, 122–135. <https://doi.org/10.1016/j.quaint.2012.02.015>
- Balistrieri, L. S., Murray, J. W., & Paul, B. (1992). The cycling of iron and manganese in the water column of Lake Sammamish, Washington. *Limnology and Oceanography*, *37*(3), 510–528. <https://doi.org/10.4319/lo.1992.37.3.0510>
- Ball, B. A., Kominoski, J. S., Adams, H. E., Jones, S. E., Kane, E. S., Loecke, T. D., et al. (2010). Direct and terrestrial vegetation-mediated effects of environmental change on aquatic ecosystem processes. *Bioscience*, *60*(8), 590–601. <https://doi.org/10.1525/bio.2010.60.8.5>
- Battarbee, R. (1986). Diatom analysis. In B. E. Berglund (Ed.), *Handbook of Holocene palaeoecology and palaeohydrology* (pp. 527–570). London: John Wiley.
- Beadle, N. C. W. (1966). Soil phosphate and its role in molding segments of the Australian flora and vegetation, with special reference to xeromorphy and sclerophylly. *Ecology*, *47*(6), 992–1007. <https://doi.org/10.2307/1935647>
- Beck, K. K., Fletcher, M.-S., Gadd, P., Heijnis, H., & Jacobsen, G. (2017). An early onset of ENSO influence in the extra-tropics of the southwest Pacific inferred from a 14,600 year high resolution multi-proxy record from Paddy's Lake, northwest Tasmania. *Quaternary Science Reviews*, *157*, 164–175. <https://doi.org/10.1016/j.quascirev.2016.12.001>
- Beck, K. K., Fletcher, M. S., Gadd, P. S., Heijnis, H., Saunders, K. M., Simpson, G. L., & Zawadzki, A. (2018). Variance and rate-of-change as early warning signals for a critical transition in an aquatic ecosystem state: A test case from Tasmania, Australia. *Journal of Geophysical Research: Biogeosciences*, *123*, 495–508. <https://doi.org/10.1002/2017JG004135>
- Beck, K. K., Fletcher, M.-S., Kattel, G., Barry, L. A., Gadd, P. S., Heijnis, H., et al. (2018). The indirect response of an aquatic ecosystem to long-term climate-driven terrestrial vegetation in a subalpine temperate lake. *Journal of Biogeography*, *45*(3), 713–725. <https://doi.org/10.1111/jbi.13144>
- Bergström, S., Carlsson, B., Gardelin, M., Lindström, G., Pettersson, A., & Rummukainen, M. (2001). Climate change impacts on runoff in Sweden—Assessments by global climate models, dynamical downscaling and hydrological modelling. *Climate Research*, *16*(2), 101–112. <https://doi.org/10.3354/cr016101>
- Bixby, R. J., Cooper, S. D., Gresswell, R. E., Brown, L. E., Dahm, C. N., & Dwire, K. A. (2015). Fire effects on aquatic ecosystems: An assessment of the current state of the science. *Freshwater Science*, *34*(4), 1340–1350. <https://doi.org/10.1086/684073>
- Boerner, R. E. (1982). Fire and nutrient cycling in temperate ecosystems. *Bioscience*, *32*(3), 187–192.
- Bogotá-A, R., Hooghiemstra, H., & Berrio, J. (2016). North Andean environmental and climatic change at orbital to submillennial time-scales: Vegetation, water-levels and sedimentary regimes from Lake Fúquene between 284 and 130 ka. *Review of Palaeobotany and Palynology*, *226*, 91–107.
- Bowling, L., & Tyler, P. (1984). Endangered lakes of scientific and cultural value in the World Heritage Area of south-west Tasmania. *Biological Conservation*, *30*(3), 201–209. [https://doi.org/10.1016/0006-3207\(84\)90084-3](https://doi.org/10.1016/0006-3207(84)90084-3)
- Bowman, D. M. J. S. (2000). *Australian rainforests: Islands of green in a land of fire* (1st ed.). Cambridge: Cambridge University Press. <https://doi.org/10.1017/CBO9780511583490>
- Bowman, D. M. J. S., & Jackson, W. D. (1981). Vegetation succession in southwest Tasmania. *Search*, *12*(10), 358–362.
- Bridle, K., Cullen, P., & Russell, M. (2003). Peatland hydrology, fire management and Holocene fire regimes in southwest Tasmanian Blanket Bogs, Nature Conservation Branch, Department of Primary Industries, Water and Environment, Hobart, Tasmania.
- Brown, D. J., Nowlin, W. H., Ozel, E., Mali, I., Episcopo, D., Jones, M. C., & Forstner, M. R. (2014). Comparison of short term low, moderate, and high severity fire impacts to aquatic and terrestrial ecosystem components of a southern USA mixed pine/hardwood forest. *Forest Ecology and Management*, *312*, 179–192. <https://doi.org/10.1016/j.foreco.2013.10.006>
- Brunelle, A., Minkley, T., Blissett, S., Cobabe, S., & Guzman, B. (2010). A ~ 8000 year fire history from an Arizona/Sonora borderland cienega. *Journal of Arid Environments*, *74*(4), 475–481. <https://doi.org/10.1016/j.jaridenv.2009.10.006>
- Buckney, R. T., & Tyler, P. A. (1973). Chemistry of some sedgeland waters: Lake Pedder, south-west Tasmania. *Australian Journal of Marine and Freshwater Research*, *24*(3), 267–273. <https://doi.org/10.1071/MF9730267>
- Bunting, L., Leavitt, P., Simpson, G., Wissel, B., Laird, K., Cumming, B., et al. (2016). Increased variability and sudden ecosystem state change in Lake Winnipeg, Canada, caused by 20th century agriculture. *Limnology and Oceanography*, *61*(6), 2090–2107. <https://doi.org/10.1002/lno.10355>
- Calvert, S., & Pedersen, T. (2007). Chapter fourteen elemental proxies for palaeoclimatic and palaeoceanographic variability in marine sediments: Interpretation and application. *Developments in Marine Geology*, *1*, 567–644.
- Cerrato, J. M., Blake, J. M., Hirani, C., Clark, A. L., Ali, A.-M. S., Artyushkova, K., et al. (2016). Wildfires and water chemistry: Effect of metals associated with wood ash. *Environmental Science: Processes & Impacts*, *18*, 1078–1089. <https://doi.org/10.1039/c6em00123h>
- Chappuis, E., Lumbreras, A., Ballesteros, E., & Gacia, E. (2015). Deleterious interaction of light impairment and organic matter enrichment on *Isoetes lacustris* (Lycopodiophyta, Isoetales). *Hydrobiologia*, *760*(1), 145–158. <https://doi.org/10.1007/s10750-015-2321-2>
- Colombaroli, D., Ssemmanda, I., Gelorini, V., & Verschuren, D. (2014). Contrasting long-term records of biomass burning in wet and dry savannas of equatorial East Africa. *Global Change Biology*, *20*(9), 2903–2914. <https://doi.org/10.1111/gcb.12583>
- Cooper, S. D., Page, H. M., Wiseman, S. W., Klose, K., Bennett, D., Even, T., et al. (2015). Physicochemical and biological responses of streams to wildfire severity in riparian zones. *Freshwater Biology*, *60*(12), 2600–2619. <https://doi.org/10.1111/fwb.12523>
- Corella, J. P., Brauer, A., Mangili, C., Rull, V., Vegas-Vilarrúbia, T., Morellón, M., & Valero-Garcés, B. L. (2012). The 1.5-ka varved record of Lake Montcortés (southern Pyrenees, NE Spain). *Quaternary Research*, *78*(02), 323–332. <https://doi.org/10.1016/j.yqres.2012.06.002>
- Croudace, I. W., & Rothwell, R. G. (2015). *Micro-XRF studies of sediment cores: Applications of a non-destructive tool for the environmental sciences*. Netherlands: Springer. <https://doi.org/10.1007/978-94-017-9849-5>
- Cuven, S., Francus, P., & Lamoureux, S. F. (2010). Estimation of grain size variability with micro X-ray fluorescence in laminated lacustrine sediments, Cape Bounty, Canadian High Arctic. *Journal of Paleolimnology*, *44*(3), 803–817. <https://doi.org/10.1007/s10933-010-9453-1>
- Davison, W. (1993). Iron and manganese in lakes. *Earth-Science Reviews*, *34*(2), 119–163. [https://doi.org/10.1016/0012-8252\(93\)90029-7](https://doi.org/10.1016/0012-8252(93)90029-7)
- Dean, W. E., & Doner, L. A. (2012). A Holocene record of endogenic iron and manganese precipitation and vegetation history in a lake-fen complex in northwestern Minnesota. *Journal of Paleolimnology*, *47*(1), 29–42. <https://doi.org/10.1007/s10933-011-9544-7>
- Diaz, R. J. (2001). Overview of hypoxia around the world. *Journal of Environmental Quality*, *30*(2), 275–281. <https://doi.org/10.2134/jeq2001.302275x>
- di Folco, M. B. (2007). *Tasmanian organic soils*. Hobart, Australia: University of Tasmania.
- Donders, T. H., Wagner-Cremer, F., & Visscher, H. (2008). Integration of proxy data and model scenarios for the mid-Holocene onset of modern ENSO variability. *Quaternary Science Reviews*, *27*(5-6), 571–579. <https://doi.org/10.1016/j.quascirev.2007.11.010>

- Emelko, M. B., Stone, M., Silins, U., Allin, D., Collins, A. L., Williams, C. H., et al. (2016). Sediment-phosphorus dynamics can shift aquatic ecology and cause downstream legacy effects after wildfire in large river systems. *Global Change Biology*, 22(3), 1168–1184. <https://doi.org/10.1111/gcb.13073>
- Engstrom, D. R., & Wright, H. Jr. (1984). Chemical stratigraphy of lake sediments as a record of environmental change. In E. Y. Haworth & J. W. G. Lund (Eds.), *Lake sediments and environmental history: Studies in palaeolimnology and palaeoecology in honour of Winifred Tutin* (pp. 11–67). Minneapolis, MN: University of Minnesota Press.
- Finkenbinder, M. S., Abbott, M. B., Edwards, M. E., Langdon, C. T., Steinman, B. A., & Finney, B. P. (2014). A 31,000 year record of paleoenvironmental and lake-level change from Harding Lake, Alaska, USA. *Quaternary Science Reviews*, 87, 98–113. <https://doi.org/10.1016/j.quascirev.2014.01.005>
- Fletcher, M.-S., Benson, A., Heijnis, H., Gadd, P. S., Cwynar, L. C., & Rees, A. B. H. (2015). Changes in biomass burning mark the onset an ENSO-influenced climate regime at 42°S in southwest Tasmania, Australia. *Quaternary Science Reviews*, 122, 222–232. <https://doi.org/10.1016/j.quascirev.2015.05.002>
- Fletcher, M.-S., & Moreno, P. I. (2011). Zonally symmetric changes in the strength and position of the Southern Westerlies drove atmospheric CO₂ variations over the past 14 k.y. *Geology*, 39(5), 419–422. <https://doi.org/10.1130/G31807.1>
- Fletcher, M.-S., & Moreno, P. I. (2012). Have the Southern Westerlies changed in a zonally symmetric manner over the last 14,000 years? A hemisphere-wide take on a controversial problem. *Quaternary International*, 253, 32–46. <https://doi.org/10.1016/j.quaint.2011.04.042>
- Fletcher, M.-S., & Thomas, I. (2010). The origin and temporal development of an ancient cultural landscape. *Journal of Biogeography*, 37(11), 2183–2196. <https://doi.org/10.1111/j.1365-2699.2010.02363.x>
- Fletcher, M.-S., Wolfe, B. B., Whitlock, C., Pompeani, D., Heijnis, H., Haberle, S., et al. (2014). The legacy of mid Holocene fire on a Tasmanian montane landscape. *Journal of Biogeography*, 41(3), 476–488. <https://doi.org/10.1111/jbi.12229>
- Fritz, S. C., & Anderson, N. (2013). The relative influences of climate and catchment processes on Holocene lake development in glaciated regions. *Journal of Paleolimnology*, 49(3), 349–362. <https://doi.org/10.1007/s10933-013-9684-z>
- Garrett, M., & Kantvilas, G. (1992). Morphology, ecology and distribution of Isoetes L. in Tasmania. In *Papers and Proceedings of the Royal Society of Tasmania* (Vol. 126, pp. 115–122), Hobart.
- Gentilli, J. (1972). *Australian climate patterns*. Adelaide: The Griffin Press.
- Gil-Romera, G., González-Sampériz, P., Lasheras-Álvarez, L., Sevilla-Callejo, M., Moreno, A., Valero-Garcés, B., et al. (2014). Biomass-modulated fire dynamics during the Last Glacial–Interglacial Transition at the Central Pyrenees (Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 402, 113–124. <https://doi.org/10.1016/j.palaeo.2014.03.015>
- González-Pérez, J. A., González-Vila, F. J., Almendros, G., & Knicker, H. (2004). The effect of fire on soil organic matter—A review. *Environment International*, 30(6), 855–870. <https://doi.org/10.1016/j.envint.2004.02.003>
- Haberzettl, T., Corbella, H., Fey, M., Janssen, S., Lucke, A., Mayr, C., et al. (2007). Late glacial and Holocene wet-dry cycles in southern Patagonia: Chronology, sedimentology and geochemistry of a lacustrine record from Laguna Potrok Aike, Argentina. *The Holocene*, 17(3), 297–310. <https://doi.org/10.1177/0959683607076437>
- Hansen, K. (1962). The dystrophic lake type. *Hydrobiologia*, 19(2), 183–190. <https://doi.org/10.1007/BF00146361>
- Hastie, T., & Stuetzle, W. (1989). Principal curves. *Journal of the American Statistical Association*, 84(406), 502–516. <https://doi.org/10.1080/01621459.1989.10478797>
- Hastie, T., & Tibshirani, R. (1990). Generalized additive models. CRC Press. <https://doi.org/10.1002/0471667196.ess0297.pub2>
- Hodgson, D., Tyler, P., & Vyverman, W. (1996). The palaeolimnology of Lake Fidler, a meromictic lake in south-west Tasmania and the significance of recent human impact. *Journal of Paleolimnology*, 18(4), 313–333. <https://doi.org/10.1023/A:1007995614287>
- Holz, A., Wood, S. W., Veblen, T. T., & Bowman, D. M. (2014). Effects of high severity fire drove the population collapse of the subalpine Tasmanian endemic conifer *Athrotaxis cupressoides*. *Global Change Biology*. <https://doi.org/10.1111/gcb.12674>
- Hughes, B. B., Levey, M. D., Fountain, M. C., Carlisle, A. B., Chavez, F. P., & Gleason, M. G. (2015). Climate mediates hypoxic stress on fish diversity and nursery function at the land–sea interface. *Proceedings of the National Academy of Sciences of the United States of America*, 112(26), 8025–8030. <https://doi.org/10.1073/pnas.1505815112>
- Huvane, J. K., & Whitehead, D. R. (1996). The paleolimnology of North Pond: Watershed-lake interactions. *Journal of Paleolimnology*, 16(3). <https://doi.org/10.1007/BF00207576>
- Jackson, W. D. (1968). Fire, air, water and earth – An elemental ecology of Tasmania. *Proceedings of the Ecological Society of Australia*, 3, 9–16.
- Jackson, W. D. (2000). Nutrient stocks in Tasmanian vegetation and approximate losses due to fire. In *Papers and proceedings of the Royal Society of Tasmania* (Vol. 134, pp. 1–18), Hobart.
- King, R. D., & Tyler, P. A. (1981). Meromictic lakes of south-west Tasmania. *Marine and Freshwater Research*, 32(5), 741–756. <https://doi.org/10.1071/MF9810741>
- Kirkpatrick, J., & Dickinson, K. (1984). Vegetation of Tasmania 1:500 000, Forestry Commission Tasmania.
- Kissman, C. E., Williamson, C. E., Rose, K. C., & Saros, J. E. (2017). Nutrients associated with terrestrial dissolved organic matter drive changes in zooplankton: Phytoplankton biomass ratios in an alpine lake. *Freshwater Biology*, 62(1), 40–51. <https://doi.org/10.1111/fwb.12847>
- Knoepp, J. D., Elliott, K. J., Clinton, B. D., & Vose, J. M. (2009). Effects of prescribed fire in mixed oak forests of the southern Appalachians: Forest floor, soil, and soil solution nitrogen responses. *The Journal of the Torrey Botanical Society*, 136(3), 380–391. <https://doi.org/10.3159/08-RA-052.1>
- Koinig, K. A., Shoty, W., Lotter, A. F., Ohlendorf, C., & Sturm, M. (2003). 9000 years of geochemical evolution of lithogenic major and trace elements in the sediment of an alpine lake—The role of climate, vegetation, and land-use history. *Journal of Paleolimnology*, 30(3), 307–320. <https://doi.org/10.1023/A:1026080712312>
- Korhola, A., Virkanen, J., Tikkanen, M., & Blom, T. (1996). Fire-induced pH rise in a naturally acid hill-top lake, southern Finland: A palaeoecological survey. *Journal of Ecology*, 84(2), 257–265. <https://doi.org/10.2307/2261361>
- Kutiel, P., & Inbar, M. (1993). Fire impacts on soil nutrients and soil erosion in a Mediterranean pine forest plantation. *Catena*, 20(1–2), 129–139. [https://doi.org/10.1016/0341-8162\(93\)90033-L](https://doi.org/10.1016/0341-8162(93)90033-L)
- Lane, P. N., Sheridan, G. J., Noske, P. J., & Sherwin, C. B. (2008). Phosphorus and nitrogen exports from SE Australian forests following wildfire. *Journal of Hydrology*, 361(1–2), 186–198. <https://doi.org/10.1016/j.jhydrol.2008.07.041>
- Leys, B., Higuera, P. E., McLauchlan, K. K., & Dunnette, P. V. (2016). Wildfires and geochemical change in a subalpine forest over the past six millennia. *Environmental Research Letters*, 11(12), 125003. <https://doi.org/10.1088/1748-9326/11/12/125003>
- Loizeau, J.-L., Span, D., Coppee, V., & Dominik, J. (2001). Evolution of the trophic state of Lake Annecy (eastern France) since the last glaciation as indicated by iron, manganese and phosphorus speciation. *Journal of Paleolimnology*, 25(2), 205–214. <https://doi.org/10.1023/A:1008100432461>

- Mackereth, F. J. H. (1966). Some chemical observations on post-glacial lake sediments. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 250(765), 165–213. <https://doi.org/10.1098/rstb.1966.0001>
- Mackey, D. J., Butler, E. C., Carpenter, P. D., Higgins, H. W., O'Sullivan, J. E., & Plaschke, R. B. (1996). Trace elements and organic matter in a pristine environment: Bathurst Harbour, southwestern Tasmania. *Science of the Total Environment*, 191(1–2), 137–151. [https://doi.org/10.1016/0048-9697\(96\)05255-2](https://doi.org/10.1016/0048-9697(96)05255-2)
- Mariani, M., Connor, S. E., Fletcher, M. S., Theuerkauf, M., Kuneš, P., Jacobsen, G., et al. (2017). How old is the Tasmanian cultural landscape? A test of landscape openness using quantitative land-cover reconstructions. *Journal of Biogeography*, 44(10), 2410–2420. <https://doi.org/10.1111/jbi.13040>
- Mariani, M., & Fletcher, M.-S. (2017). Long-term climate dynamics in the extra-tropics of the South Pacific revealed from sedimentary charcoal analysis. *Quaternary Science Reviews*, 173, 181–192. <https://doi.org/10.1016/j.quascirev.2017.08.007>
- Mariani, M., Fletcher, M. S., Holz, A., & Nyman, P. (2016). ENSO controls interannual fire activity in southeast Australia. *Geophysical Research Letters*, 43, 10,891–10,900. <https://doi.org/10.1002/2016GL070572>
- Massaferro, J., Recasens, C., Larocque-Tobler, I., Zolitschka, B., & Maidana, N. (2013). Major lake level fluctuations and climate changes for the past 16,000 years as reflected by diatoms and chironomids preserved in the sediment of Laguna Potrok Aike, southern Patagonia. *Quaternary Science Reviews*, 71, 167–174. <https://doi.org/10.1016/j.quascirev.2012.07.026>
- McGlone, M. S., Kershaw, P., & Markgraf, V. (1992). El Niño/Southern Oscillation climatic variability in Australasian and South American paleoenvironmental records. In H. F. Diaz & V. Markgraf (Eds.), *El Niño: Historical and paleoclimatic aspects of the Southern Oscillation* (Chap. 21, pp. 435–462). Cambridge: Cambridge University Press.
- McWethy, D., Higuera, P., Whitlock, C., Veblen, T., Bowman, D., Cary, G., et al. (2013). A conceptual framework for predicting temperate ecosystem sensitivity to human impacts on fire regimes. *Global Ecology and Biogeography*, 22(8), 900–912. <https://doi.org/10.1111/geb.12038>
- McWethy, D., Whitlock, C., Wilmshurst, J. M., McGlone, M. S., Fromont, M., Li, X., et al. (2010). Rapid landscape transformation in South Island, New Zealand, following initial Polynesian settlement. *Proceedings of the National Academy of Sciences of the United States of America*, 107(50), 21,343–21,348. <https://doi.org/10.1073/pnas.1011801107>
- Melles, M., Brigham-Grette, J., Minyuk, P. S., Nowaczyk, N. R., Wennrich, V., DeConto, R. M., et al. (2012). 2.8 million years of Arctic climate change from Lake El'gygytgyn, NE Russia. *Science*, 337(6092), 315–320. <https://doi.org/10.1126/science.1222135>
- Meyers, P. A., & Teranes, J. L. (2002). Sediment organic matter. In *Tracking environmental change using lake sediments*. Volume 2: *Physical and Geochemical Methods* (pp. 239–269). Dordrecht, Netherlands: Springer. https://doi.org/10.1007/0-306-47670-3_9
- Moody, J. A., & Martin, D. A. (2001). Initial hydrologic and geomorphic response following a wildfire in the Colorado Front Range. *Earth Surface Processes and Landforms*, 26(10), 1049–1070. <https://doi.org/10.1002/esp.253>
- Moody, J. A., & Martin, D. A. (2009). Synthesis of sediment yields from wildland fire in different rainfall regimes in the western United States. *International Journal of Wildland Fire*, 18(1), 96–115. <https://doi.org/10.1071/WF07162>
- Moritz, M. A., Parisien, M.-A., Batllori, E., Krawchuk, M. A., Van Dorn, J., Ganz, D. J., & Hayhoe, K. (2012). Climate change and disruptions to global fire activity. *Ecosphere*, 3(6), art49.
- Morris, J. L., McLauchlan, K. K., & Higuera, P. E. (2015). Sensitivity and complacency of sedimentary biogeochemical records to climate-mediated forest disturbances. *Earth-Science Reviews*, 148, 121–133. <https://doi.org/10.1016/j.earscirev.2015.06.001>
- Mount, A. B. (1979). Natural regeneration processes in Tasmanian forests. *Search*, 10(5), 180–186.
- Moy, C. M., Seltzer, G. O., Rodbell, D. T., & Anderson, D. M. (2002). Variability of El Niño/Southern Oscillation activity at millennial timescales during the Holocene. *Nature*, 420(6912), 162–165. <https://doi.org/10.1038/nature01194>
- Naeher, S., Gilli, A., North, R. P., Hamann, Y., & Schubert, C. J. (2013). Tracing bottom water oxygenation with sedimentary Mn/Fe ratios in Lake Zurich, Switzerland. *Chemical Geology*, 352, 125–133. <https://doi.org/10.1016/j.chemgeo.2013.06.006>
- Nealson, K. H., & Saffarini, D. (1994). Iron and manganese in anaerobic respiration: Environmental significance, physiology, and regulation. *Annual Reviews in Microbiology*, 48(1), 311–343. <https://doi.org/10.1146/annurev.mi.48.100194.001523>
- Nicholls, N., & Lucas, C. (2007). Interannual variations of area burnt in Tasmanian bushfires: Relationships with climate and predictability. *International Journal of Wildland Fire*, 16(5), 540–546. <https://doi.org/10.1071/WF06125>
- North, R. P., North, R. L., Livingstone, D. M., Köster, O., & Kipfer, R. (2014). Long-term changes in hypoxia and soluble reactive phosphorus in the hypolimnion of a large temperate lake: Consequences of a climate regime shift. *Global Change Biology*, 20(3), 811–823. <https://doi.org/10.1111/gcb.12371>
- Och, L. M., Müller, B., Voegelin, A., Ulrich, A., Göttlicher, J., Steiniger, R., et al. (2012). New insights into the formation and burial of Fe/Mn accumulations in Lake Baikal sediments. *Chemical Geology*, 330, 244–259.
- Ohlsson, K. A., & Wallmark, P. H. (1999). Novel calibration with correction for drift and non-linear response for continuous flow isotope ratio mass spectrometry applied to the determination of $\delta^{15}\text{N}$, total nitrogen, $\delta^{13}\text{C}$ and total carbon in biological material. *Analyst*, 124(4), 571–577. <https://doi.org/10.1039/a900855a>
- O'Sullivan, P., & Reynolds, C. S. (2008). *The lakes handbook: Lake restoration and rehabilitation*. New York: John Wiley.
- Paul, B., & Duthie, H. (1989). Nutrient cycling in the epilithon of running waters. *Canadian Journal of Botany*, 67(8), 2302–2309. <https://doi.org/10.1139/b89-294>
- Pemberton, M. (1988). Soil erosion between Brichs Inlet and Elliot Bay, southwest Tasmania. *Papers and Proceedings of the Royal Society of Tasmania*, 122(2), 109–114.
- Pesce, O., & Moreno, P. (2014). Vegetation, fire and climate change in central-east Isla Grande de Chiloé (43 S) since the Last Glacial Maximum, northwestern Patagonia. *Quaternary Science Reviews*, 90, 143–157. <https://doi.org/10.1016/j.quascirev.2014.02.021>
- Philibert, A., Gell, P., Newall, P., Chessman, B., & Bate, N. (2006). Development of diatom-based tools for assessing stream water quality in south-eastern Australia: Assessment of environmental transfer functions. *Hydrobiologia*, 572(1), 103–114. <https://doi.org/10.1007/s10750-006-0371-1>
- Power, M., Whitney, B., Mayle, F., Neves, D., de Boer, E., & Maclean, K. (2016). Fire, climate and vegetation linkages in the Bolivian Chiquitano seasonally dry tropical forest. *Philosophical Transactions of the Royal Society B*, 371(1696), 20150165. <https://doi.org/10.1098/rstb.2015.0165>
- R Development Core Team (2013). *R: A language and environment for statistical computing*, edited. Vienna, Austria: R Foundation for Statistical Computing.
- Raison, R. J. (1979). Modification of the soil environment by vegetation fires, with particular reference to nitrogen transformations: A review. *Plant and Soil*, 51(1), 73–108. <https://doi.org/10.1007/BF02205929>
- Reneau, S. L., Katzman, D., Kuyumjian, G. A., Lavine, A., & Malmon, D. V. (2007). Sediment delivery after a wildfire. *Geology*, 35(2), 151–154. <https://doi.org/10.1130/G23288A1>

- Riera, J. L., Ballesteros, E., Pulido, C., Chappuis, E., & Gacia, E. (2017). Recovery of submersed vegetation in a high mountain oligotrophic soft-water lake over two decades after impoundment. *Hydrobiologia*, *794*(1), 139–151. <https://doi.org/10.1007/s10750-017-3087-5>
- Robinne, F.-N., Bladon, K. D., Miller, C., Parisien, M.-A., Mathieu, J., & Flannigan, M. D. (2018). A spatial evaluation of global wildfire-water risks to human and natural systems. *Science of the Total Environment*, *610*, 1193–1206.
- Schaller, T., & Wehrli, B. (1996). Geochemical-focusing of manganese in lake sediments—An indicator of deep-water oxygen conditions. *Aquatic Geochemistry*, *2*(4), 359–378.
- Sheridan, G. J., Lane, P. N., & Noske, P. J. (2007). Quantification of hillslope runoff and erosion processes before and after wildfire in a wet *Eucalyptus* forest. *Journal of Hydrology*, *343*(1-2), 12–28. <https://doi.org/10.1016/j.jhydrol.2007.06.005>
- Sherson, L. R., Van Horn, D. J., Gomez-Velez, J. D., Crossey, L. J., & Dahm, C. N. (2015). Nutrient dynamics in an alpine headwater stream: Use of continuous water quality sensors to examine responses to wildfire and precipitation events. *Hydrological Processes*, *29*(14), 3193–3207. <https://doi.org/10.1002/hyp.10426>
- Sigg, L. (2000). Redox potential measurements in natural waters: Significance, concepts and problems. In H. D. Schulz, et al. (Eds.), *Redox* (pp. 1–12). Berlin, Heidelberg: Springer.
- Simi, E., Moreno, P., Villa-Martínez, R., Vilanova, I., & de Pol-Holz, R. (2017). Climate change and resilience of deciduous *Nothofagus* forests in central–east Chilean Patagonia over the last 3200 years. *Journal of Quaternary Science*, *32*(6), 845–856. <https://doi.org/10.1002/jqs.2948>
- Simpson, G. L., & Birks, H. J. B. (2012). Statistical learning in palaeolimnology. In J. B. H. Birks, et al. (Eds.), *Tracking environmental change using lake sediments* (pp. 249–327). Netherlands: Springer.
- Smith, H. G., Sheridan, G. J., Lane, P. N. J., & Nyman, P. (2011). Wildfire effects on water quality in forest catchments: A review with implications for water supply. *Journal of Hydrology*, *396*(1-2), 170–192. <https://doi.org/10.1016/j.jhydrol.2010.10.043>
- Sonneman, J., Sincok, A., Fluin, J., Reid, M., Newall, P., Tibby, J., & Gell, P. (1999). *An illustrated guide to common diatom species from temperate Australia—Identification guide 33*. Albury: CRC for Freshwater Ecology.
- Stahle, L. N., Chin, H., Haberle, S., & Whitlock, C. (2017). Late-glacial and Holocene records of fire and vegetation from Cradle Mountain National Park, Tasmania, Australia. *Quaternary Science Reviews*, *177*, 57–77. <https://doi.org/10.1016/j.quascirev.2017.09.010>
- Stahle, L. N., Whitlock, C., & Haberle, S. G. (2016). A 17,000-year-long record of vegetation and fire from Cradle Mountain National Park, Tasmania. *Frontiers in Ecology and Evolution*, *4*, 82.
- Steinberg, C. E., Paul, A., Pflugmacher, S., Meinelt, T., Klöcking, R., & Wiegand, C. (2003). Pure humic substances have the potential to act as xenobiotic chemicals—A review. *Fresenius Environmental Bulletin*, *12*(5), 391–401.
- Strock, K. E., Theodore, N., Gawley, W. G., Ellsworth, A. C., & Saros, J. E. (2017). Increasing dissolved organic carbon concentrations in northern boreal lakes: Implications for lake water transparency and thermal structure. *Journal of Geophysical Research: Biogeosciences*, *122*, 1022–1035. <https://doi.org/10.1002/2017JG003767>
- Talbot, M. R. (2002). Nitrogen isotopes in palaeolimnology. *Tracking Environmental Change Using Lake Sediments*, 401–439. https://doi.org/10.1007/0-306-47670-3_15
- Talbot, M. R., & Johannessen, T. (1992). A high resolution palaeoclimatic record for the last 27,500 years in tropical West Africa from the carbon and nitrogen isotopic composition of lacustrine organic matter. *Earth and Planetary Science Letters*, *110*(1-4), 23–37. [https://doi.org/10.1016/0012-821X\(92\)90036-U](https://doi.org/10.1016/0012-821X(92)90036-U)
- Thomas, I. (1995). “Where have all the forests gone?” New pollen evidence from *Melaleuca Inlet* in southwestern Tasmania. Paper presented at IAG 1993 Conference Proceedings. Melbourne: Monash Publications in Geography.
- Tyler, P. (1974). III. Limnological studies. In *Biogeography and ecology in Tasmania* (Vol. 25, pp. 29–61). Dordrecht, Netherlands: Springer.
- Tyler, P. (1992). A lakeland from the dreamtime the second founders’ lecture. *British Phycological Journal*, *27*(4), 353–368. <https://doi.org/10.1080/00071619200650301>
- van Dam, H., Suurmond, G., & ter Braak, C. J. (1981). Impact of acidification on diatoms and chemistry of Dutch moorland pools. *Hydrobiologia*, *83*(3), 425–459.
- Vyverman, W., Vyverman, R., Hodgson, D., & Tyler, P. (1995). Diatoms from Tasmanian mountain lakes: A reference data-set for environmental reconstruction. The TASDIAT diatom training set: A systematic and autoecological study. *Bibliotheca Diatomol*, *33*, 1–198.
- Vyverman, W., Vyverman, R., Rajendran, V. S., & Tyler, P. (1996). Distribution of benthic diatom assemblages in Tasmanian highland lakes and their possible use as indicators of environmental changes. *Canadian Journal of Fisheries and Aquatic Sciences*, *53*(3), 493–508. <https://doi.org/10.1139/f95-220>
- Wersin, P., Höhener, P., Giovanoli, R., & Stumm, W. (1991). Early diagenetic influences on iron transformations in a freshwater lake sediment. *Chemical Geology*, *90*(3–4), 233–252. [https://doi.org/10.1016/0009-2541\(91\)90102-W](https://doi.org/10.1016/0009-2541(91)90102-W)
- Wilkins, D., Gouramanis, C., De Deckker, P., Fifield, L. K., & Olley, J. (2013). Holocene lake-level fluctuations in lakes Keilambete and Gnotuk, southwestern Victoria, Australia. *The Holocene*. <https://doi.org/10.1177/0959683612471983>
- Wilkinson, S., Wallbrink, P., Hancock, G., Blake, W., Shakesby, R., & Doerr, S. (2009). Fallout radionuclide tracers identify a switch in sediment sources and transport-limited sediment yield following wildfire in a eucalypt forest. *Geomorphology*, *110*(3-4), 140–151. <https://doi.org/10.1016/j.geomorph.2009.04.001>
- Wood, S. W., & Bowman, D. M. J. S. (2011). Alternative stable states and the role of fire–vegetation–soil feedbacks in the temperate wilderness of southwest Tasmania. *Landscape Ecology*, *27*, 13–28.
- Wood, S., & Wood, M. S. (2007). The mgcv package. Retrieved from www.r-project.org
- Xia, Y., Zhao, J., & Collerson, K. D. (2001). Early-mid Holocene climatic variations in Tasmania, Australia: Multi-proxy records in a stalagmite from Lynds Cave. *Earth and Planetary Science Letters*, *194*(1-2), 177–187. [https://doi.org/10.1016/S0012-821X\(01\)00541-6](https://doi.org/10.1016/S0012-821X(01)00541-6)