

Received: 5 June 2018 | Revised: 5 September 2018 | Accepted: 6 September 2018

DOI: 10.1111/jne.12644

ORIGINAL ARTICLE

WILEY 

Sex-dependent changes in neuroactive steroid concentrations in the rat brain following acute swim stress

Ying Sze^{1,2} | Andrew C. Gill^{2,3} | Paula J. Brunton^{1,2} 

¹Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh, UK

²The Roslin Institute, University of Edinburgh, Edinburgh, UK

³School of Chemistry, University of Lincoln, Lincoln, UK

Correspondence

Paula J. Brunton, Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh, UK.
Email: p.j.brunton@ed.ac.uk

Funding information

Biotechnology and Biological Sciences Research Council, Grant/Award Number: BB/J004332/1

Sex differences in hypothalamic-pituitary-adrenal (HPA) axis activity are well established in rodents. In addition to glucocorticoids, stress also stimulates the secretion of progesterone and deoxycorticosterone (DOC) from the adrenal gland. Neuroactive steroid metabolites of these precursors can modulate HPA axis function; however, it is not known whether levels of these steroids differ between male and females following stress. In the present study, we aimed to establish whether neuroactive steroid concentrations in the brain display sex- and/or region-specific differences under basal conditions and following exposure to acute stress. Brains were collected from male and female rats killed under nonstress conditions or following exposure to forced swimming. Liquid chromatography-mass spectrometry was used to quantify eight steroids: corticosterone, DOC, dihydrodeoxycorticosterone (DHDOC), pregnenolone, progesterone, dihydroprogesterone (DHP), allopregnanolone and testosterone in plasma, and in five brain regions (frontal cortex, hypothalamus, hippocampus, amygdala and brainstem). Corticosterone, DOC and progesterone concentrations were significantly greater in the plasma and brain of both sexes following stress; however, the responses in plasma were greater in females compared to males. This sex difference was also observed in the majority of brain regions for DOC and progesterone but not for corticosterone. Despite observing no stress-induced changes in circulating concentrations of pregnenolone, DHDOC or DHP, concentrations were significantly greater in the brain and this effect was more pronounced in females than males. Basal plasma and brain concentrations of allopregnanolone were significantly higher in females; moreover, stress had a greater impact on central allopregnanolone concentrations in females. Stress had no effect on circulating or brain concentrations of testosterone in males. These data indicate the existence of sex and regional differences in the generation of neuroactive steroids in the brain following acute stress, especially for the 5α -reduced steroids, and further suggest a sex-specific expression of steroidogenic enzymes in the brain. Thus, differential neurosteroidogenesis may contribute to sex differences in HPA axis responses to stress.

KEYWORDS

5α -reductase, glucocorticoids, hypothalamic-pituitary-adrenal (HPA) axis, neurosteroids, progestogens, sex differences

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Journal of Neuroendocrinology* published by John Wiley & Sons Ltd on behalf of British Society for Neuroendocrinology

1 | INTRODUCTION

Steroid hormones play a crucial role in regulating physiological responses to stress. They can exert their actions in target tissues by binding to classical steroid receptors, which then act as transcription factors in the nucleus to alter gene expression. Additionally, steroids can act in a nonclassical manner. Neuroactive steroids are active metabolites of classical steroid hormones that, independent of their source (ie, peripheral or central), exert rapid nongenomic effects on neuronal excitability by binding to membrane bound ion channel-linked receptors, thus influencing neurotransmission.¹ The brain itself can produce neuroactive steroids (referred to as “neurosteroids”), via local *de novo* synthesis from cholesterol or through the conversion of peripherally derived adrenal or gonadal steroids.^{1,2} In support of this, the brain possesses many steroidogenic enzymes, including 5 α -reductase and 3 α -hydroxysteroid dehydrogenase (3 α HSD) which are not only mainly expressed by glial cells,³⁻⁵ but also found in neurones.⁶ Furthermore, neurosteroids continue to be detected in the brain after combined adrenalectomy and gonadectomy,⁷ providing evidence for local neurosteroidogenesis. Nevertheless, the synthesis of neurosteroids is reliant upon the expression of the prerequisite steroidogenic enzymes, which exhibit significant regional differences in the brain.⁸

The major neuroendocrine response to stress is mediated via activation of the hypothalamic-pituitary-adrenal (HPA) axis, which results in a net increase in circulating glucocorticoids following activation of corticotrophin-releasing hormone (CRH) neurones in the medial parvocellular paraventricular nucleus (mpPVN) and adrenocorticotrophic hormone (ACTH) secretion from the anterior pituitary. Glucocorticoids, synthesised in the adrenal cortex, act at multiple targets in the body to facilitate the necessary metabolic

and behavioural adaptations required for stress coping and restoring physiological homeostasis. Moreover, glucocorticoids provide negative-feedback control of the HPA axis through actions on glucocorticoid and mineralocorticoid receptors in the brain and pituitary, crucial for terminating the stress response.

In addition to glucocorticoids, stress also results in a rapid increase in the secretion of other steroid hormones such as progesterone⁹ and 11-deoxycorticosterone (DOC)¹⁰ from the adrenal cortex. In the brain, these steroids can be further metabolised into neuroactive steroids, such as 5 α -dihydroprogesterone (DHP) and 5 α -dihydrodeoxycorticosterone (DHDOC) by 5 α -reductase, which in turn may be converted into allopregnanolone and tetrahydrodeoxycorticosterone (THDOC), respectively, by 3 α HSD (Figure 1), which are considered to fine-tune and aid cessation of the stress response. Indeed, levels of both allopregnanolone and THDOC are increased in the cerebral cortex and hypothalamus following exposure to acute stress¹¹ and both have been shown to negatively modulate stress-induced HPA axis activity *in vivo*.¹²⁻¹⁵ Moreover, allopregnanolone prevents the up-regulation of CRH mRNA in the mpPVN induced by adrenalectomy, attenuates stimulated CRH release from hypothalamic explants and inhibits the firing of mpPVN CRH neurones *in vitro*^{16,17}, while administration of the 5 α -reductase inhibitor, finasteride, leads to heightened HPA axis responses to stress in males and females.^{13,18} Allopregnanolone, DHDOC and THDOC are positive allosteric modulators of GABA_A receptors; by prolonging the opening time of chloride ion channels within GABA_A receptors, they enhance the inhibitory actions of GABA.¹⁹⁻²² Given that the PVN is substantially innervated by GABAergic neurones^{23,24} and the majority of mpPVN CRH neurones express GABA_A receptors,²⁵ this provides a means by which local neurosteroids may modulate inhibitory tone over the HPA axis.




FIGURE 1 Biosynthetic pathways involved in steroidogenesis. The major pathways and enzymes involved in the biosynthesis of corticosteroids, progestogens, androgens and oestrogens from cholesterol. 3 α -diol, 3 α -androstenediol; 3 α HSD, 3 α -hydroxysteroid dehydrogenase (*Akr1c*); 3 β -diol, 3 β -androstenediol; 3 β HSD, 3 β -hydroxysteroid dehydrogenase (*Hsd3b*); 5 α R, 5 α -reductase (*Srd5a*); 17 α HSD, 17 α -hydroxysteroid dehydrogenase (*Hsd17a*); 17 β HSD, 17 β -hydroxysteroid dehydrogenase (*Hsd17b*); DHDOC: 5 α -dihydrodeoxycorticosterone; DHEA, dehydroepiandrosterone; DHP, 5 α -dihydroprogesterone; DHT, 5 α -dihydrotestosterone; DOC, 11-deoxycorticosterone; p450scc, cholesterol side-chain cleavage enzyme (*Cyp11a1*); THDOC, tetrahydrodeoxycorticosterone

It is well established that there are sex differences in both basal and stress-induced activity of the HPA axis in rodents. Females generally display greater concentrations of circulating corticosterone under nonstress conditions and this is particularly evident at the onset of the dark phase.^{26,27} In addition, the HPA axis is more responsive to both physical and psychological stressors in females, than males, as reflected by greater secretion of ACTH and corticosterone and a larger increase in CRH gene transcription in the mpPVN.²⁷⁻³¹

Intriguingly, stress-induced activation of neurones in limbic brain regions, such as the prefrontal cortex and hippocampus, is greater in males than in females.³²⁻³⁴ Given that these regions project indirectly to the PVN (eg via a relay in the bed nucleus of the stria terminalis) and exert a net inhibitory effect on HPA axis activity,³⁵⁻³⁷ lower stress responses in males may reflect greater inhibitory signalling from limbic brain regions to the PVN.

The sex differences in HPA axis activity have been largely attributed to gonadal steroids, given that oestradiol typically has stimulatory actions, whereas androgens have inhibitory effects on HPA axis function. For example, ovariectomy decreases,³⁸⁻⁴⁰ whereas orchidectomy increases,^{27,41-47} the HPA axis response to stress and these effects of gonadectomy can be normalised with testosterone^{41,42,44,46} or oestradiol³⁸ replacement in males and females, respectively. However, the effect of testosterone on HPA axis activity is evidently mediated via the neuroactive metabolite of testosterone, 3 β -androstadiol (3 β -diol).⁴⁸ Likewise, studies indicate that progesterone may counteract some of the stimulatory effects of oestradiol on HPA axis activity,⁴⁹ although this is likely mediated via its 3 α -hydroxy A ring-reduced neuroactive metabolite, allopregnanolone.^{12,13} Despite these findings, it is not known whether there are sex differences in the stress-induced neuroactive steroid concentrations in the brain, which could potentially contribute to the differential HPA axis responses in males and females.

Although some studies have quantified the concentrations of a few steroids in the brain in response to foot-shock, forced swimming or CO₂ inhalation, previous work has focused on males and measurements have been largely limited to the cerebral cortex or hypothalamus.^{11,50-52} In studies where male and females have been directly compared, this has been under nonstress conditions.^{53,54} To date, no study has simultaneously measured stress-induced concentrations of multiple steroids in several different brain regions. Moreover, to the best of our knowledge, no study has compared changes in central steroid levels in response to stress in male and female rats in the same experiment. Traditional detection methods such as radioimmunoassays incur limitations: cross-reactivity issues may hinder accuracy, whereas low throughput and low sensitivity makes quantification of a panel of structurally related steroids in discrete brain regions difficult.^{55,56}

Accordingly, in the present study, we used a liquid chromatography-mass spectrometry (LC-MS) method to quantify a panel of steroids in the plasma, as well as in five separate brain regions known to be involved in regulating the activity of the HPA axis, from male and female rats under nonstress and acute stress conditions. We

aimed to establish: (i) which steroids (from our panel) are increased in the brain by acute stress; (ii) whether there are sex and/or regional differences in stress-induced central steroid concentrations; and (iii) whether circulating steroid concentrations correlate with those found in the brain following stress.

2 | MATERIALS AND METHODS

2.1 | Animals

Male and female Sprague-Dawley rats were bred in the rodent facility at the Roslin Institute. Rats were group housed by sex (four to six females, three or four males per cage) in open-top cages with ad libitum access to drinking water and a standard rodent diet (Harlan Teklad; Cambridgeshire, UK) and under a 12:12 hour light/dark cycle (lights on 8.00 AM) with controlled temperature (22 \pm 1 $^{\circ}$ C) and humidity (58 \pm 3%). Experiments were performed in rats aged 21 weeks and were approved by the local Animal Welfare and Ethical Review Body and performed in accordance with the UK Animals (Scientific Procedures) Act 1986 and the European Directive (2010/63/EU).

2.2 | Stress paradigm

All experiments and tissue collection were performed in male and female (randomly cycling) rats between 10.00 AM and 12.30 PM. Forced swimming was used because: (i) it is a robust activator of the HPA axis^{57,58}; (ii) sex differences in ACTH and corticosterone secretion have been demonstrated using this stressor⁵⁷; and (iii) it is a combined physical and psychological stressor, and thus activates stress-responsive nuclei in both the forebrain and hindbrain.⁵⁹ For the stressed groups (n = 7 per group per sex), rats were placed in a glass cylinder (diameter 25 cm; height 50 cm) filled with water (21-22 $^{\circ}$ C) to a depth of 30 cm and forced to swim for 2 minutes, after which they were gently dried in a towel and returned to their home cage. Thirty minutes after the onset of swimming stress, rats were rapidly transferred to a separate room and killed by conscious decapitation. This time-point has previously been shown to correspond with peak levels of immediate early gene induction⁵⁹ in the brain and elevated corticosterone secretion,⁶⁰ as well as with the peak in stress-induced allopregnanolone concentrations in the brain^{11,50} following swim stress. For the control group, rats (n = 7 per group per sex) remained undisturbed in their home cage prior to killing (as before). In each case, trunk blood was collected in tubes containing 0.5 mL of ice-cold 5% (w/v) EDTA and plasma separated by centrifugation (1500 g at 4 $^{\circ}$ C for 15 minutes). Brains were rapidly removed, and the regions of interest (frontal cortex, hypothalamus, amygdala, hippocampus, brainstem) were dissected (see Supporting information, Figure S1) and frozen on dry ice. Plasma was stored at -20 $^{\circ}$ C and brain samples were stored at -80 $^{\circ}$ C until further analysis.

2.3 | Steroid standards

All steroid standards were purchased from Steraloids Inc (Newport, RI, USA) and all solvents/chemicals used were high-performance

liquid chromatography (HPLC)-MS grade. Stock solutions of steroid standards (1 mg mL^{-1}) were prepared in methanol (Honeywell Riedel-de Haën, Seelze, Germany) and stored at -20°C . Steroids were combined and diluted in methanol into a working stock solution containing a concentration of $10 \text{ }\mu\text{g mL}^{-1}$ of each steroid (except for allopregnanolone and progesterone with a concentration of $25 \text{ }\mu\text{g mL}^{-1}$). On the day of sample processing, standards were further diluted 100-fold in a surrogate matrix 4% (w/v) bovine serum albumin (BSA) (VWR, Leicester, UK) in PBS, followed by a serial 2.5-fold dilution in 4% BSA to produce seven calibrants. The calibration standards used ranged from 41 to $10\,000 \text{ pg mL}^{-1}$ for corticosterone, DOC (4-pregnen-21-ol-3,20-dione), DHDOC (5 α -pregnan-21-ol-3,20-dione), testosterone, pregnenolone and DHP (5 α -pregnane-3,20-dione), and 102.4 to $25\,000 \text{ pg mL}^{-1}$ for progesterone and allopregnanolone (5 α -pregnan-3 α -ol-20-one). The deuterated internal standard progesterone-d9 was also dissolved in methanol and diluted to a final concentration of 25 ng mL^{-1} in 50% methanol.

2.4 | Sample processing

Samples from all four groups from any given region were processed in the same run, together with seven standard calibrants and a zero sample containing only 4% BSA. For standard calibrants and plasma, $100 \text{ }\mu\text{L}$ was used. Frozen brain regions were weighed prior to sample processing ($n = 7$ per group per sex, except for the frontal cortex, amygdala and hypothalamus where a sample from the female non-stressed group was lost).

2.4.1 | Tissue homogenisation

Brain samples were homogenised in $500 \text{ }\mu\text{L}$ of methanol containing 1% formic acid (Fisher Scientific, Loughborough, UK). For plasma and standards, $400 \text{ }\mu\text{L}$ of methanol containing 1% formic acid was added to $100 \text{ }\mu\text{L}$ of plasma/standards and vigorously vortexed. Next, $20 \text{ }\mu\text{L}$ of progesterone-d9 (25 ng mL^{-1}) was added and the homogenates were briefly sonicated. After incubation on dry ice for 30 minutes, the homogenates were centrifuged for 10 minutes ($13\,000 \text{ g}$ at 4°C) and the supernatant was decanted into a borosilicate tube and the pellet re-extracted again with $500 \text{ }\mu\text{L}$ of methanol containing 1% formic acid, and then sonicated and centrifuged as described above but without incubation. Supernatants were combined, then diluted with ultrapure water to a final concentration of 30% methanol (see Supporting information, Figure S2).

2.4.2 | Solid phase extraction

Steroids in both plasma and brain samples were extracted by solid phase extraction using DSC-Discovery C18 100 mg columns (#52602-U; Supelco, Belfont, PA, USA). Columns were activated with 1 mL of methanol and equilibrated with another 1 mL of 30% methanol. Diluted supernatants from homogenates (approximately 3 mL) were then loaded, followed by two 1-mL washes of 50% methanol. All steps were assisted by centrifugation at 50 g (average

flow rate of 0.5 mL min^{-1}). Steroids were eluted with 1 mL of 85% methanol by gravity flow. The collected eluate was dried in a vacuum concentrator (Savant SpeedVac; ThermoFisher Scientific, Waltham, MA, USA) overnight (see Supporting information, Figure S2). Dried samples were stored at -20°C until the day of analysis.

2.4.3 | Derivatisation

On the day of LC-MS analysis, $400 \text{ }\mu\text{L}$ of freshly prepared derivatisation agent (1 mg mL^{-1} of Girard's T reagent; #89397; Sigma-Aldrich, Gillingham, UK; dissolved in methanol containing 0.2% formic acid) was added to the dried samples. After incubation at 37°C for 30 minutes, the reaction was stopped by the addition of $50 \text{ }\mu\text{L}$ of 5% ammonium hydroxide (ACROS Organics, Morris Plains, NJ, USA) in methanol. Samples were dried in the SpeedVac then reconstituted in $50 \text{ }\mu\text{L}$ of 25 mmol L^{-1} phosphate buffer (pH 7.4) in 50% methanol to prevent acid hydrolysis of the derivatisation agent (see Supporting information, Figure S2). The sample was transferred to Chromacol vials (ThermoFisher Scientific) for analysis.

2.5 | LC-MS/MS analysis

Analysis of steroids was performed using an Ultimate 3000 Dionex HPLC system coupled to an AmaZon ETD ion trap mass spectrometer (Bruker Daltonics, Bremen, Germany). Separation of steroids on reverse phase HPLC was achieved on the ACE UltraCore $2.5 \text{ }\mu\text{m}$ Super C18 column (75 mm by 2.1 mm inner diameter; Advance Chromatography Technologies, Aberdeen, UK) maintained at 40°C . Gradient elution was performed (see Supporting information, Table S1) with the mobile phase consisting of 50 mmol L^{-1} ammonium formate (Thermo Fisher Scientific), pH 3 (solvent A), and methanol with 0.1% formic acid (solvent B). Steroids were analysed simultaneously using multiple reaction monitoring (see Supporting information, Table S2), with positive electrospray ionisation and collision-induced fragmentation. Injections for all samples, including calibrants, were carried out in duplicate. The method allowed us to simultaneously determine the concentrations of eight steroids: corticosterone, DOC, DHDOC, pregnenolone, progesterone, DHP, allopregnanolone and testosterone (Figure 2) with a total runtime of 19 minutes (see Supporting information, Table S1). Because Girard's T reagent was used as the derivatisation agent, only steroids with a carbonyl group could be examined; thus, oestradiol and the testosterone metabolites, 3 α - and 3 β -androstenediol, which do not have this functional group, were not detected using this method. THDOC and DHT did not pass the validation criteria and thus these data are not included.

Data were acquired using HYSTAR software (Bruker Daltonics, Bremen, Germany) and peak area under the curve was extracted and automatically integrated using QUANTANALYSIS, version 2.0 (Bruker Daltonics). The ratio of the peak area of the target analyte and the peak area of progesterone-d9 was used to construct the calibration curve. The seven-point calibration curves were linear for all analytes (see Supporting information, Table S3). Concentrations of samples were extrapolated and converted to ng mL^{-1} (for plasma) or

normalised to the wet weight of the tissues (ng g^{-1} ; for brain tissues). Performance characteristics (ie, recovery, sensitivity, accuracy and precision) were determined (see Supporting information, Table S4), and the concentrations of steroids detected in the plasma and brain were consistent with previous studies.^{61,62}

2.6 | Statistical analysis

Data are presented as group means \pm SEM. In each case, data were compared using a two-way ANOVA (SIGMAPLOT, version 11.0; Systat Software Inc., Chicago, IL, USA), with sex and stress as the two main factors followed by Student-Newman Keuls pairwise multiple comparison testing, except for the testosterone data, which were analysed using a two-tailed Student's *t*-test. The relationships between circulating and brain concentrations of steroids were determined using Pearson's correlation coefficients (SIGMAPLOT, version 11.0), and males and females were analysed separately. In each case, $P \leq 0.05$ was considered statistically significant.

3 | RESULTS

3.1 | Corticosterone

3.1.1 | Plasma

As expected, there was a main effect of stress ($F_{1,24} = 58.9, P < 0.001$) and sex ($F_{1,24} = 54.7, P < 0.001$) on corticosterone secretion and also a significant stress \times sex interaction ($F_{1,24} = 4.38, P = 0.047$). Plasma corticosterone concentrations were significantly greater in the stressed group compared to the control group in both sexes, with females displaying significantly greater plasma corticosterone

concentrations compared to males under both stress and nonstress conditions (Figure 3).

3.1.2 | Brain

There were no significant differences detected in basal corticosterone concentrations between males and females in any of the brain regions (Figure 3). There were significant main effects of stress in each of the five brain regions examined (frontal cortex, $F_{1,23} = 50.6, P < 0.001$; hypothalamus, $F_{1,23} = 26.7, P < 0.001$; amygdala, $F_{1,23} = 14.8, P < 0.001$; hippocampus, $F_{1,24} = 45.4, P < 0.001$; brainstem, $F_{1,24} = 21.3, P < 0.001$), with corticosterone concentrations being significantly greater in the stressed rats compared to the nonstressed rats in both sexes (Figure 3). There was a significant main effect of sex observed only in the brainstem ($F_{1,24} = 5.30, P = 0.03$), where corticosterone concentrations were 1.8-fold greater in stressed females compared to stressed males.

3.1.3 | Correlations between central and circulating corticosterone

There were significant positive correlations between corticosterone concentrations in the plasma and in all brain regions, except the amygdala in males and the hypothalamus in females (Table 1).

3.2 | Deoxycorticosterone

3.2.1 | Plasma

There was no significant difference in plasma DOC concentrations between males and females under nonstress conditions; however,




FIGURE 2 Representative chromatogram of the analytes. For some analytes, a double peak was observed as a result of the presence of both *syn* and *anti*-Girard's T derivatives, in which case the major product peak was used for quantification. The ratio between the major and minor peak was always consistent between runs. DHDOC, 5 α -dihydrodeoxycorticosterone; DHP, 5 α -dihydroprogesterone; DOC, 11-deoxycorticosterone




FIGURE 3 Effect of swim stress on corticosterone concentrations in the plasma and brain. Asterisks denote significant differences in the stressed group (S) compared to the nonstressed group (NS) within the same sex (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$); hashes denote significant sex differences within the same stress status (# $P < 0.05$; ### $P < 0.001$). Following stress, sex differences were detected for corticosterone concentrations in the plasma and brainstem. $n = 6-7$ rats per group. Note the difference in the scale of the y-axis for plasma

there was a significant effect of stress ($F_{1,24} = 67.6$, $P < 0.001$) and sex ($F_{1,24} = 26.0$, $P < 0.001$) and a significant stress \times sex interaction ($F_{1,24} = 20.7$, $P < 0.001$). Stress resulted in significantly greater circulating DOC concentrations in both sexes (Figure 4). However, under stressed conditions, females exhibited significantly greater plasma DOC concentrations compared to males (Figure 4).

3.2.2 | Brain

There were no differences in basal concentrations of DOC between males and females in any of the regions studied (Figure 4). There were significant main effects of stress in all five brain regions (frontal cortex: $F_{1,23} = 72.0$, $P < 0.001$; hypothalamus: $F_{1,23} = 37.1$, $P < 0.001$; amygdala: $F_{1,23} = 56.7$, $P < 0.001$; hippocampus: $F_{1,24} = 103.1$, $P < 0.001$; brainstem: $F_{1,24} = 105.1$, $P < 0.001$). Central DOC concentrations were significantly greater in the stressed groups compared to the basal groups for both sexes (Figure 4). There were additional main effects of sex observed for the hippocampus ($F_{1,24} = 5.34$, $P = 0.03$) and brainstem

($F_{1,24} = 14.4$, $P < 0.001$) and a stress \times sex interaction in the brainstem ($F_{1,24} = 12.9$, $P = 0.001$). Comparing the stressed groups, females had significantly greater DOC concentrations in the hippocampus and brainstem compared to males but not in the frontal cortex, hypothalamus or amygdala (Figure 4).

3.2.3 | Correlations between central and circulating DOC

There were significant positive correlations between DOC concentrations in the plasma and all five brain regions for both sexes (Table 1).

3.3 | Dihydrodeoxycorticosterone

3.3.1 | Plasma

There were no significant differences in plasma DHDOC concentrations in any of the groups (Figure 5).

TABLE 1 Correlation between central and circulating steroids

Steroid	Region	Males		Females	
		<i>r</i>	<i>P</i> value	<i>R</i>	<i>P</i> value
Corticosterone	Frontal cortex	0.89	<0.001***	0.88	<0.001***
	Hypothalamus	0.81	<0.001***	0.55	0.051*
	Amygdala	0.51	0.062	0.73	0.005**
	Hippocampus	0.82	<0.001***	0.62	0.017*
	Brainstem	0.62	0.019*	0.70	0.006**
DOC	Frontal cortex	0.80	0.001***	0.90	<0.001***
	Hypothalamus	0.76	0.002**	0.75	0.003**
	Amygdala	0.75	0.002**	0.85	<0.001***
	Hippocampus	0.87	<0.001***	0.88	<0.001***
	Brainstem	0.84	<0.001***	0.88	<0.001***
DHDOC	Frontal cortex	-0.07	0.826	0.23	0.459
	Hypothalamus	0.04	0.888	0.60	0.032*
	Amygdala	0.80	0.001***	0.92	<0.001***
	Hippocampus	0.19	0.525	0.68	0.008**
	Brainstem	-0.23	0.427	-0.16	0.591
Pregnenolone	Frontal cortex	-0.010	0.738	-0.02	0.942
	Hypothalamus	0.14	0.631	0.40	0.170
	Amygdala	0.53	0.051*	0.56	0.045*
	Hippocampus	0.06	0.828	0.48	0.079
	Brainstem	-0.25	0.388	-0.15	0.619
Progesterone	Frontal cortex	0.94	<0.001***	0.90	<0.001***
	Hypothalamus	0.98	<0.001***	0.82	0.001***
	Amygdala	0.99	<0.001***	0.83	<0.001***
	Hippocampus	0.94	<0.001***	0.89	<0.001***
	Brainstem	0.93	<0.001***	0.88	<0.001***
DHP	Frontal cortex	0.17	0.571	0.18	0.552
	Hypothalamus	-0.28	0.342	0.50	0.086
	Amygdala	0.83	<0.001***	0.94	<0.001***
	Hippocampus	0.01	0.747	0.66	0.011*
	Brainstem	-0.20	0.491	0.11	0.710
Allopregnanolone	Frontal cortex	-0.03	0.918	-0.02	0.952
	Hypothalamus	0.04	0.907	0.35	0.246
	Amygdala	0.59	0.026*	0.54	0.055
	Hippocampus	0.43	0.127	0.35	0.227
	Brainstem	0.41	0.143	-0.11	0.710
Testosterone	Frontal cortex	0.81	<0.001***		
	Hypothalamus	0.89	<0.001***		
	Amygdala	0.47	0.094		
	Hippocampus	0.49	0.075		
	Brainstem	0.80	0.001***		

Pearson's correlation coefficient (*r*) and probability (*P*) values between plasma and brain concentrations of steroids. Males and females were analysed separately and data from both stressed and nonstressed groups within the same sex were used for analysis (*n* = 13-14 per sex).

Asterisks denote significant correlations with plasma concentrations (**P* ≤ 0.05; ***P* ≤ 0.01; ****P* ≤ 0.001).




FIGURE 4 Effect of swim stress on DOC concentrations in the plasma and brain. Asterisks denote significant differences in the stressed group (S) compared to the nonstressed group (NS) within the same sex (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$); hashes denote significant sex differences with the same stress status (# $P < 0.05$; ## $P < 0.01$; ### $P < 0.001$). After stress exposure, sex differences were detected for DOC concentrations in the plasma, hippocampus and brainstem. $n = 6-7$ rats per group

3.3.2 | Brain

There were significant main effects of stress in all brain regions, except for the amygdala (frontal cortex, $F_{1,23} = 16.1$, $P < 0.001$; hypothalamus, $F_{1,23} = 8.46$, $P = 0.008$; hippocampus, $F_{1,24} = 7.67$, $P = 0.01$; brainstem, $F_{1,24} = 5.68$, $P = 0.025$) (Figure 5). There were significant main effects of sex on DHDOC concentrations in the frontal cortex ($F_{1,23} = 6.09$, $P = 0.02$) and brainstem ($F_{1,24} = 4.35$, $P = 0.048$). In the frontal cortex, DHDOC concentrations were greater in both the stressed males and females compared to their respective control groups, with stress-induced concentrations being significantly greater in females compared to males. However, in the hypothalamus, hippocampus and brainstem, stress resulted in greater DHDOC concentrations only in females.

3.3.3 | Correlations between central and circulating DHDOC

In males, only the DHDOC concentration in the amygdala was correlated with plasma levels, whereas, in females, there were positive correlations between DHDOC concentrations in the plasma and those in the amygdala, hippocampus and hypothalamus (Table 1).

3.4 | Pregnenolone

3.4.1 | Plasma

There were no significant differences in plasma pregnenolone concentrations between any of the groups (Figure 6).




FIGURE 5 Effect of swim stress on DHDOC concentrations in the plasma and brain. Asterisks denote significant differences in the stressed group (S) compared to the nonstressed group (NS) within the same sex (* $P < 0.05$; ** $P < 0.01$); hashes denote significant sex differences within the same stress status (# $P < 0.05$). Following stress, there was a significant sex difference in the frontal cortex and a tendency towards greater DHDOC concentrations in the brainstem of stressed females compared to stressed males ($P = 0.068$). $n = 6$ -7 rats per group. Note the difference in the scaling of the y-axis for the hypothalamus and brainstem

3.4.2 | Brain

There was a significant main effect of sex (frontal cortex, $F_{1,23} = 5.44$, $P = 0.029$; hypothalamus, $F_{1,23} = 7.75$, $P = 0.01$; amygdala, $F_{1,23} = 20.2$, $P < 0.001$; hippocampus $F_{1,24} = 27.3$, $P < 0.001$; brainstem, $F_{1,24} = 19.1$) and stress (frontal cortex, $F_{1,23} = 39.2$, $P < 0.001$; hypothalamus, $F_{1,23} = 10.4$, $P = 0.004$; amygdala, $F_{1,23} = 40.9$, $P < 0.001$; hippocampus; $F_{1,24} = 39.9$, $P < 0.001$; brainstem, $F_{1,24} = 26.7$) in all of the brain regions examined (Figure 6). There was also a significant sex \times stress interaction in the brainstem ($F_{1,24} = 4.89$), amygdala ($F_{1,23} = 12.8$, $P = 0.002$) and hippocampus ($F_{1,24} = 7.25$, $P = 0.01$). There was no difference in basal levels of pregnenolone in the brain between the sexes (Figure 6). Following stress, both males and females had greater pregnenolone concentrations than their respective control groups in all brain regions, except for the hypothalamus, where only females showed significantly greater pregnenolone concentrations compared to nonstressed females. For

each brain region examined, stress-induced pregnenolone concentrations were greater in females versus males (Figure 6).

3.4.3 | Correlations between central and circulating pregnenolone

For both males and females, the only significant correlation observed with plasma concentrations of pregnenolone was in the amygdala (Table 1). No other significant correlations were detected between circulating and brain pregnenolone concentrations.

3.5 | Progesterone

3.5.1 | Plasma

There were significant main effects of stress ($F_{1,24} = 49.1$, $P < 0.001$) and sex ($F_{1,24} = 49.7$, $P < 0.001$) and also a significant stress \times sex




FIGURE 6 Effect of swim stress on pregnenolone concentrations in the plasma and brain. Asterisks denote significant differences in the stressed group (S) compared to the nonstressed group (NS) within the same sex (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$); hashes denote significant sex differences with the same stress status (# $P < 0.05$; ## $P < 0.01$; ### $P < 0.001$). Concentrations of pregnenolone were greater in stressed females vs the stressed males in all of the brain regions examined. $n = 6-7$ rats per group. Note the difference in the scaling of the y-axis for the hypothalamus and brainstem

interaction ($F_{1,24} = 10.8$, $P = 0.003$) in plasma progesterone concentrations. Under basal conditions, circulating progesterone was significantly greater in females compared to males (Figure 7). In both sexes, stress resulted in increased progesterone secretion, with greater circulating concentrations in females than in males (Figure 7).

3.5.2 | Brain

There were significant main effects of stress and sex in all five brain regions examined (frontal cortex: stress $F_{1,23} = 62.6$, $P < 0.001$ and sex $F_{1,23} = 13.6$, $P = 0.001$; hypothalamus: stress $F_{1,23} = 48.3$, $P < 0.001$ and sex $F_{1,23} = 28.7$, $P < 0.001$; amygdala: stress $F_{1,23} = 50.7$, $P < 0.001$ and sex $F_{1,23} = 23.0$, $P < 0.001$; hippocampus: stress $F_{1,24} = 53.9$, $P < 0.001$ and sex $F_{1,24} = 38.9$, $P < 0.001$; brainstem: stress $F_{1,24} = 31.9$ and sex $F_{1,24} = 18.8$). There were also stress \times sex interactions observed for the hypothalamus ($F_{1,23} = 8.14$,

$P = 0.009$), amygdala ($F_{1,23} = 4.48$, $P = 0.045$) and hippocampus ($F_{1,24} = 3.94$, $P = 0.058$). There was a tendency for higher basal progesterone concentrations in females compared to males; however, this was only statistically significant in the hippocampus (Figure 7). Following stress, progesterone concentrations in all five brain regions were significantly higher than those measured under basal conditions in both sexes. Moreover, in each case, stress-induced progesterone concentrations were greater in females than in males (Figure 7).

3.5.3 | Correlations between central and circulating progesterone

There were significant positive correlations between circulating progesterone concentrations and those measured in all five brain regions for both males and females (Table 1).




FIGURE 7 Effect of swim stress on progesterone concentrations in the plasma and brain. Asterisks denote significant differences in the stressed group (S) compared to the nonstressed group (NS) within the same sex (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$); hashes denote significant sex differences within the same stress status (# $P < 0.05$; ## $P < 0.01$; ### $P < 0.001$). Following stress, sex differences in progesterone concentrations were detected in the plasma and all regions of the brain. $n = 6$ – 7 rats per group. Note the difference in the scaling of the y-axis for the brainstem

3.6 | Dihydroprogesterone

3.6.1 | Plasma

There was no significant difference in circulating DHP in any of the groups investigated (Figure 8).

3.6.2 | Brain

There was a significant main effect of stress on DHP concentration in the frontal cortex ($F_{1,23} = 13.1$, $P = 0.001$), hypothalamus ($F_{1,23} = 7.14$, $P = 0.01$), amygdala ($F_{1,23} = 5.73$, $P = 0.03$), hippocampus ($F_{1,24} = 9.77$, $P = 0.005$) and brainstem ($F_{1,24} = 22.0$, $P < 0.001$), as well as a main effect of sex for the hypothalamus ($F_{1,23} = 24.0$, $P < 0.001$) and brainstem ($F_{1,24} = 16.0$, $P < 0.001$). Under basal conditions, DHP concentrations in the hypothalamus were significantly greater in females than in males and there was a similar trend observed for the brainstem

($P = 0.06$) (Figure 8). Following stress, females had significantly higher DHP in each of the brain regions investigated compared to nonstress conditions, whereas, in males, stress-induced concentrations of DHP were greater than basal levels only in the frontal cortex and brainstem (Figure 8). There was a sex difference in stress-induced DHP concentrations, with stressed females exhibiting greater DHP than stressed males in the amygdala, hypothalamus and brainstem (Figure 8).

3.6.3 | Correlations between central and circulating DHP

There were significant positive correlations observed between DHP concentrations in the amygdala and circulation in both males and females (Table 1). In females, there was also a modest correlation between the DHP concentrations in the plasma and the hippocampus (Table 1). No other correlations were detected between central and circulating DHP levels.




FIGURE 8 Effect of swim stress on DHP concentrations in the plasma and brain. Asterisks denote significant differences in the stressed group (S) compared to the nonstressed group (NS) within the same sex (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$); hashes denote significant sex differences within the same stress status (# $P < 0.05$; ## $P < 0.01$). Under nonstress conditions, DHP concentrations were greater in the hypothalamus of females compared to males and a similar trend was detected in the brainstem ($P = 0.06$). Following stress, significant sex differences in DHP concentrations were detected in the hypothalamus, amygdala and brainstem. $n = 6$ -7 rats per group. Note the difference in the scaling of the y-axis for the brainstem

3.7 | Allopregnanolone

3.7.1 | Plasma

There was an effect of sex on circulating allopregnanolone concentrations ($F_{1,24} = 21.9$, $P < 0.001$). Basal allopregnanolone concentrations were significantly greater in females than in males (Figure 9). There was no main effect of stress on plasma allopregnanolone concentrations in either sex (Figure 9).

3.7.2 | Brain

There were significant main effects of sex on allopregnanolone concentrations for each of the brain regions examined (frontal cortex, $F_{1,23} = 14.9$, $P < 0.001$; hypothalamus, $F_{1,23} = 13.1$, $P = 0.001$;

amygdala, $F_{1,23} = 80.2$, $P < 0.001$; hippocampus, $F_{1,24} = 20.9$, $P < 0.001$; brainstem, $F_{1,24} = 40.2$, $P < 0.001$). A significant main effect of stress was observed for allopregnanolone concentrations in the frontal cortex ($F_{1,23} = 5.71$, $P = 0.03$), amygdala ($F_{1,23} = 27.6$, $P < 0.001$) and brainstem ($F_{1,24} = 9.08$, $P = 0.006$). A stress \times sex interaction was additionally observed in the amygdala ($F_{1,23} = 14.8$, $P < 0.001$). Basal allopregnanolone concentrations were significantly greater in females than in males in all of the brain regions, except the hypothalamus, where a tendency for higher levels was observed ($P = 0.07$) (Figure 9). Stress resulted in significantly greater allopregnanolone concentrations in the frontal cortex, amygdala and brainstem compared to nonstressed females (Figure 9). Similar trends were observed in the male brain, although this only reached statistical significance for the frontal cortex.




FIGURE 9 Effect of swim stress on allopregnanolone concentrations in the plasma and brain. Asterisks denote significant differences in the stressed group (S) compared to the nonstressed group (NS) within the same sex (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$); hashes denote significant sex differences with the same stress status (# $P < 0.05$; ## $P < 0.01$; ### $P < 0.001$). Under nonstress conditions, allopregnanolone concentrations were greater in females compared to males in the plasma and each of the brain regions examined, except the hypothalamus where there was a tendency for higher levels in females ($P = 0.07$). After stress exposure, significant sex differences in allopregnanolone concentrations were detected in the plasma and all regions of the brain. n/d, not detected. $n = 6-7$ rats/group. Note the difference in the scaling of the y-axis for the plasma, hypothalamus and brainstem

3.7.3 | Correlations between central and circulating allopregnanolone

In both sexes, only allopregnanolone concentrations in the amygdala were found to be significantly correlated with circulating allopregnanolone (Table 1).

3.8 | Testosterone

3.8.1 | Plasma

There was no significant effect of stress on plasma testosterone concentrations in males (1.05 ± 0.28 ng mL⁻¹ in nonstressed rats vs

0.77 ± 0.20 ng mL⁻¹ in stressed rats). Testosterone was below the lower limit of quantification (ie <41 ng mL⁻¹) in the plasma from females (see Supporting information, Table S4).

3.8.2 | Brain

Testosterone was below the lower limit of quantification in all of the brain regions in females. There was no significant effect of stress on central testosterone concentrations in male rats (frontal cortex, 2.20 ± 0.45 vs 2.20 ± 0.58 ng g⁻¹; hypothalamus, 2.79 ± 0.62 vs 3.15 ± 1.23 ng g⁻¹; amygdala, 0.88 ± 0.17 vs 1.03 ± 0.24 ng g⁻¹; hippocampus, 1.17 ± 0.21 vs 1.71 ± 0.61 ng g⁻¹; brainstem, 2.03 ± 0.19 vs 2.90 ± 0.35 ng g⁻¹ in the stressed and nonstressed groups, respectively).

3.8.3 | Correlations between central and circulating testosterone

Concentrations of testosterone in the frontal cortex, amygdala, hypothalamus and brainstem were positively correlated with circulating testosterone in male rats (Table 1).

4 | DISCUSSION

In the present study, we investigated the impact of an acute stressor on the circulating and central steroid profile in male and female rats. Several key findings emerged in the present study. We show for the first time there are clear sex differences in the brain's steroid response to acute swim stress in the rat, with females typically showing markedly greater concentrations in most cases. We also report that, although concentrations of corticosterone, progesterone and

DOC in the brain are positively correlated with those measured in the circulation, this is not the case for pregnenolone, DHP, DHDOC and allopregnanolone, supporting the concept of the de novo synthesis of these neurosteroids in the brain in response to stress.^{7,11} Lastly, although the effect of stress on the steroidal milieu of the brain was largely global, we did find evidence for regional differences (Figure 10), particularly for DHDOC and allopregnanolone, which may be of functional relevance.

A key feature of the endocrine response to stress is increased secretion of corticosterone from the adrenal cortex into the bloodstream. It is well established that females display enhanced HPA axis activity, characterised by a greater circulating concentrations of corticosterone under basal conditions and a larger increase in ACTH and corticosterone secretion following acute stress.⁶³⁻⁶⁵ Consistent with this, females in the present study exhibited greater basal and stress-induced plasma corticosterone concentrations. In the periphery, greater corticosterone secretion in females is associated with a




FIGURE 10 Summary of the key changes in steroid concentrations induced by stress. Changes in steroid concentrations in the plasma and brain 30 minutes after a forced swim stress. The source of steroids in the plasma is largely from peripheral steroidogenic organs, such as the adrenal glands and gonads, whereas steroids in the brain may be peripherally or centrally derived. Blue and red arrows represent stress-induced increases in steroid concentrations in males and females, respectively. Dashes indicate no significant changes following stress. Hashes denote the presence of a sex difference under stressed conditions. AP, allopregnanolone; CORT, corticosterone; DHDOC, dihydrodeoxycorticosterone; DHP, dihydroprogesterone; DOC, deoxycorticosterone; PREG, pregnenolone; P4, progesterone; T, testosterone

heightened sensitivity of the anterior pituitary to CRH and the adrenal gland to ACTH, as well as a greater capacity for glucocorticoid synthesis in the adrenal glands,⁶⁶ probably as a result of the actions of oestradiol.³⁹ In the brain, CRH mRNA expression in the mpPVN is increased following stress to a greater extent in females than in males,³¹ which may be further compounded by diminished GR-mediated negative-feedback control of the HPA axis in females.⁶⁷ Interestingly, despite sex differences in the plasma, corticosterone concentrations in the female brain were not different from those in males, except for in the brainstem. This is consistent with the finding of similar circadian corticosterone rhythms and concentrations in the brains of males and females, despite differences in the plasma.^{26,27,63,68} It is likely that greater concentrations of corticosteroid-binding globulin (which restricts the amount of free corticosterone in the blood that can enter the brain) in females^{64,69} contribute to this finding. It is not known whether sex differences exist in the expression of 11 β -hydroxysteroid dehydrogenase type 1 (which reactivates corticosterone from its inert metabolite, 11-dehydrocorticosterone) in the brain, which could play a role in modulating the local concentrations of corticosterone in the brain. However, sex and region-specific differences in 11 β -hydroxylase (Cyp11b1) expression have been reported in the brain,⁷⁰ suggesting that differences in local corticosterone synthesis could also potentially contribute.

Under nonstress conditions, circulating progesterone concentrations were greater in females than males, which is consistent with previous findings^{53,71}; however, there was no sex difference in plasma DOC concentrations. Following stress, circulating concentrations of DOC and progesterone were increased in both sexes, with significantly greater responses in females. Stress also resulted in greater levels of progesterone and DOC in both sexes in each of the brain regions investigated. Despite evidence of low levels of 21-hydroxylase gene expression and activity in the rodent brain,^{72,73} DOC is considered to be primarily synthesised from progesterone in the adrenal cortex and is secreted from the adrenal glands, together with progesterone, in response to stress.^{9,10} Consistent with this was our finding of a positive correlation between circulating DOC and progesterone concentrations and those in the brain, indicating that the likely source of DOC and progesterone in the brain is from the periphery, specifically adrenal derived. Although there was a robust sex difference in circulating DOC after stress, centrally, this was reflected only in the brainstem. This may be a result of sex differences in DOC metabolism in limbic brain regions in females or that circulating DOC more readily accesses the brainstem.

Importantly, DOC and progesterone mediate rapid nongenomic effects indirectly via their downstream neuroactive metabolites THDOC and allopregnanolone (Figure 1), which act as allosteric modulators of GABA_A receptor activity^{1,74} to enhance inhibitory GABA neurotransmission. THDOC and allopregnanolone are synthesised from DOC and progesterone via the intermediates DHDOC and DHP, respectively, and their conversion is catalysed by the same enzymes: 5 α -reductase and 3 α HSD (Figure 1). In the present study, stress resulted in higher concentrations of both DHDOC and DHP in discrete brain regions but not in the plasma, suggesting de novo

synthesis of these steroids in the brain. Moreover, differential regulation was observed depending on sex and brain region. For example, DHDOC concentrations were found to be increased only in the frontal cortex of males following stress, whereas, for females, DHDOC concentrations were increased in the frontal cortex, hypothalamus, hippocampus and brainstem. Similarly, although stress increased DHP concentrations in each of the five brain regions examined in females, in males, stress-induced elevations in DHP were only detected in the frontal cortex and brainstem. In addition to the presence of greater levels of precursors in the female brain, differential expression of the 5 α -reductase may contribute to the sex difference, given that expression has been shown to be higher in females, at least in the hippocampus.⁷⁵ Moreover, regional differences in expression and activity have been demonstrated for 5 α -reductase and 3 α -HSD in the adult brain,⁷⁶⁻⁷⁹ with sexual dimorphic expression being displayed after stress exposure.⁸⁰ Any direct effects of DHDOC and DHP are unclear, although DHP is known to have affinity for the progesterone receptor⁸¹ and DHDOC can potentiate GABA-activated chloride ion currents in neurones,²² suggesting that, similar to THDOC and allopregnanolone, DHDOC also acts as a positive allosteric modulator at GABA_A receptors.

Roles for allopregnanolone and THDOC in regulating HPA axis activity in response to stress via potentiating the inhibitory effects of GABA are well established.^{12-14,16,74,82,83} Unfortunately, we were unable to reliably quantify THDOC using our method; however, we were able to measure allopregnanolone. Under nonstress conditions, circulating and central allopregnanolone concentrations were greater in females than in males, which is consistent with previous findings.^{54,71} We did not observe a significant increase in circulating allopregnanolone concentrations 30 minutes after stress exposure in either sex, in contrast to previous studies in male rats.^{11,52} The reason for this is not clear; however, the different duration/intensity of the swim stress used (2 minutes in the present study versus 10 minutes) may contribute. Moreover, allopregnanolone concentrations are known to peak 30-60 minutes later in the plasma than in the brain.¹¹ Nevertheless, following stress, allopregnanolone concentrations were greater in the frontal cortex, amygdala and brainstem of females compared to controls, as well as in the frontal cortex of males, consistent with previous findings.^{11,52} Sex differences in central allopregnanolone concentrations may result from greater concentrations of the precursors, progesterone and DHP and/or sex differences in the activity of the converting enzymes in the brain. The strong positive correlation between central and peripheral progesterone concentrations, coupled with a lack of correlation between circulating and central allopregnanolone, and its precursor DHP, suggests that DHP and allopregnanolone are synthesised in the brain from progesterone following stress; however, it would be necessary to measure the activity of the converting enzymes in the brain to confirm this. Caruso et al⁵⁴ report a positive correlation between progesterone concentrations in the plasma and cerebral cortex in rats, which is consistent with our findings; however, in contrast to the present study, they also observed a direct correlation between pregnenolone, DHP and allopregnanolone levels in the

plasma and cerebral cortex. The reason for this difference is unclear; however, it is important to note that the correlations performed in the present study included data from both stressed and nonstressed rats within the same sex, whereas, in the previous study, correlations were analysed only under basal conditions with data from male and female rats combined. Thus, it is likely that the lack of correlation reported in the present study is a result of stress increasing central levels of pregnenolone, DHP and allopregnanolone, without affecting circulating levels.

It is not clear whether the sex differences in neurosteroids reported in the present study are of functional importance with respect to HPA axis regulation and further studies are needed to establish whether other stressors result in similar effects. Unlike the brainstem, the prefrontal cortex, amygdala and hippocampus do not innervate the PVN directly; rather, they modulate activity of the HPA axis via neural projections to GABAergic relay sites, such as the peri-PVN, the bed nucleus of the stria terminalis and the medial preoptic area, which exert considerable inhibitory tone over CRH neurones.^{24,36} Thus, it is conceivable that sex differences in brain concentrations of neurosteroids with GABA_A receptor agonist activity may lead to differential modulation of upstream inhibitory signalling pathways, resulting, for example, in a greater disinhibition of PVN CRH neurones in females, which could contribute to sex differences in HPA axis responses to stress.

As well as the increase in downstream metabolites of progesterone and DOC, the principal precursor for these steroid hormones, pregnenolone, was also increased throughout the brain following stress, and to a greater extent in the brains of females. This may indicate that, in females, either p450_{scc} is more highly expressed/more active or that pregnenolone is metabolised at a different rate in females; however, this requires further study. Consistent with previous studies, we found no stress-induced change in plasma pregnenolone concentrations.⁵²

One point to note is that the females used in the present study were randomly cycling, which may be expected to influence the steroid concentrations in the brain. However, there is very little variation in brain concentrations of pregnenolone, progesterone, DHP or allopregnanolone across the oestrous cycle when these are measured in the morning (as they were in the present study); indeed, circadian fluctuations in these steroids are far greater than those relating to any stage of the oestrous cycle.⁸⁴ Moreover, oestrous cycle stage does not affect basal corticosterone secretion⁸⁵ or the corticosterone response to swim stress⁸⁶ when sampling is performed in the morning.

The mechanisms involved in stress-induced increases in neuroactive steroid levels in the brain have yet to be established. One study has indicated that CRH increases concentrations of progesterone, DHP and allopregnanolone in the brain.⁸⁷ Steroidogenic enzymes such as 5 α -reductase are known to be regulated by androgens^{88,89} and, although adrenal-derived steroids have been shown to increase 3 α -HSD expression in the brain in both sexes, gonadal steroids regulate central 3 α -HSD expression in a sex-specific manner.⁹⁰ Given that testosterone was unaltered by stress, it is unlikely that it

contributes to the rapid changes in stress-induced neuroactive steroids seen in the present study; however, testosterone may underlie sex differences in steroidogenic enzyme expression in the brain, and therefore influence the brain's capacity to produce neurosteroids.

With respect to rapid regulation of neurosteroid production, GABA may play a role via an ultrashort feedback loop. Brain regions that express neurosteroidogenic enzymes are densely innervated by GABAergic neurones and express abundant GABA_A receptors.⁹¹⁻⁹³ Moreover, GABA or GABA_A receptor agonists inhibit the de novo synthesis of neurosteroids⁹⁴ and neurosteroidogenic neurones are evidently under tonic inhibition by GABA.⁹⁴ Given that some neuroactive steroids, such as allopregnanolone, are potent allosteric modulators of GABA_A receptor function,¹⁹⁻²¹ it is conceivable that neurosteroids may control their own production by regulating the activity of GABA_A receptors. Whether such a mechanism contributes to the sex differences in central neurosteroid concentrations requires further study; however, GABA receptor subunit expression does differ between males and females, under basal conditions and following stress.⁹⁵

In summary, the data obtained in the present study demonstrate robust sex differences in the steroid response to acute swim stress and indicate sex-specific expression of steroidogenic enzymes in the brain. Taken together, our data also support a role for the de novo synthesis of neurosteroids by the brain following stress, particularly those derived from 5 α -reductase activity (ie, DHDHC, DHP, allopregnanolone). The neurosteroid response to stress is likely to be adaptive and disparate neurosteroidogenesis may contribute to sex differences in HPA axis function; however, it is unclear whether greater neuroactive steroid levels in females confer an overall advantage or disadvantage. Stress-induced increases in neuroactive steroids are likely to have important neuroendocrine and/or behavioural effects to aid stress adaptation. For example, in addition to modulating HPA axis activity, allopregnanolone is known to have anxiolytic and analgesic actions.^{82,96} Sex differences in the brain's response to stress may account for differences in the propensity to develop stress-related disorders between males and females, further highlighting the importance of considering sex when developing therapies.

ACKNOWLEDGEMENTS

The authors thank Professor Ruth Andrew and Dr Shazia Khan (University of Edinburgh) for providing expert advice on derivitisation agents. Dr Yu-Ting Lai and Mrs Helen Cameron assisted with the animal experiments. This work was supported by Biotechnology and Biological Sciences Research Council (BBSRC) Institute Strategic funding (BB/J004332/1). YS was supported by a University of Edinburgh Principal's Career Development Scholarship.

CONFLICT OF INTERESTS

The authors declare that they have no conflicts of interest.

ORCID

Paula J. Brunton  <http://orcid.org/0000-0003-3827-6523>

REFERENCES

- Paul SM, Purdy RH. Neuroactive steroids. *FASEB J*. 1992;6:2311-2322.
- Baulieu EE. Neurosteroids: a new function in the brain. *Biol Cell*. 1991;71:3-10.
- Hu ZY, Bourreau E, Jung-Testas I, Robel P, Baulieu EE. Neurosteroids: oligodendrocyte mitochondria convert cholesterol to pregnenolone. *Proc Natl Acad Sci USA*. 1987;84:8215-8219.
- Jung-Testas I, Hu ZY, Baulieu EE, Robel P. Neurosteroids: biosynthesis of pregnenolone and progesterone in primary cultures of rat glial cells. *Endocrinology*. 1989;125:2083-2091.
- Krieger NR, Scott RG. Nonneuronal localization for steroid converting enzyme: 3 alpha-hydroxysteroid oxidoreductase in olfactory tubercle of rat brain. *J Neurochem*. 1989;52:1866-1870.
- Melcangi RC, Celotti F, Castano P, Martini L. Differential localization of the 5 alpha-reductase and the 3 alpha-hydroxysteroid dehydrogenase in neuronal and glial cultures. *Endocrinology*. 1993;132:1252-1259.
- Corpechot C, Young J, Calvel M, et al. Neurosteroids: 3alpha-Hydroxy-5alpha-pregnan-20-one and its precursors in the brain, plasma, and steroidogenic glands of male and female rats. *Endocrinology*. 1993;133:1003-1009.
- Mensah-Nyagan AG, Do-Rego JL, Beaujean D, Luu-The V, Pelletier G, Vaudry H. Neurosteroids: expression of steroidogenic enzymes and regulation of steroid biosynthesis in the central nervous system. *Pharmacol Rev*. 1999;51:63-81.
- Hueston CM, Deak T. On the time course, generality, and regulation of plasma progesterone release in male rats by stress exposure. *Endocrinology*. 2014;155:3527-3537.
- Reddy DS. Physiological role of adrenal deoxycorticosterone-derived neuroactive steroids in stress-sensitive conditions. *Neuroscience*. 2006;138:911-920.
- Purdy RH, Morrow AL, Moore P, Paul SM. Stress-induced elevations of gamma-aminobutyric acid type A receptor-active steroids in the rat brain. *Proc Natl Acad Sci USA*. 1991;88:4553-4557.
- Patchev VK, Hassan A, Holsboer F, Almeida O. The neurosteroid tetrahydroprogesterone attenuates the endocrine response to stress and exerts glucocorticoid-like effects on vasopressin gene transcription in the rat hypothalamus. *Neuropsychopharmacology*. 1996;15:533-540.
- Brunton PJ, McKay AJ, Ochedalski T, et al. Central opioid inhibition of neuroendocrine stress responses in pregnancy in the rat is induced by the neurosteroid allopregnanolone. *J Neurosci*. 2009;29:6449-6460.
- Brunton PJ, Donadio MV, Yao ST, et al. 5alpha-reduced neurosteroids sex-dependently reverse central prenatal programming of neuroendocrine stress responses in rats. *J Neurosci*. 2015;35:666-677.
- Owens MJ, Ritchie JC, Nemeroff CB. 5 alpha-pregnane-3 alpha, 21-diol-20-one (THDOC) attenuates mild stress-induced increases in plasma corticosterone via a non-glucocorticoid mechanism: comparison with alprazolam. *Brain Res*. 1992;573:353-355.
- Patchev VK, Shoaib M, Holsboer F, Almeida OF. The neurosteroid tetrahydroprogesterone counteracts corticotropin-releasing hormone-induced anxiety and alters the release and gene expression of corticotropin-releasing hormone in the rat hypothalamus. *Neuroscience*. 1994;62:265-271.
- Gunn BG, Cunningham L, Cooper MA, et al. Dysfunctional astrocytic and synaptic regulation of hypothalamic glutamatergic transmission in a mouse model of early-life adversity: relevance to neurosteroids and programming of the stress response. *J Neurosci*. 2013;33:19534-19554.
- Handa RJ, Kudwa AE, Donner NC, McGivern RF, Brown R. Central 5-alpha reduction of testosterone is required for testosterone's inhibition of the hypothalamo-pituitary-adrenal axis response to restraint stress in adult male rats. *Brain Res*. 2013;1529:74-82.
- Majewska MD, Harrison NL, Schwartz RD, Barker JL, Paul SM. Steroid hormone metabolites are barbiturate-like modulators of the GABA receptor. *Science*. 1986;232:1004-1007.
- Morrow AL, Suzdak PD, Paul SM. Steroid hormone metabolites potentiate GABA receptor-mediated chloride ion flux with nanomolar potency. *Eur J Pharmacol*. 1987;142:483-485.
- Lambert JJ, Cooper MA, Simmons RD, Weir CJ, Belelli D. Neurosteroids: endogenous allosteric modulators of GABA(A) receptors. *Psychoneuroendocrinology*. 2009;34(Suppl. 1):S48-S58.
- Reddy DS, Rogawski MA. Stress-induced deoxycorticosterone-derived neurosteroids modulate GABA(A) receptor function and seizure susceptibility. *J Neurosci*. 2002;22:3795-3805.
- Miklos IH, Kovacs KJ. GABAergic innervation of corticotropin-releasing hormone (CRH)-secreting parvocellular neurons and its plasticity as demonstrated by quantitative immunoelectron microscopy. *Neuroscience*. 2002;113:581-592.
- Cullinan WE, Ziegler DR, Herman JP. Functional role of local GABAergic influences on the HPA axis. *Brain Struct Funct*. 2008;213:63-72.
- Cullinan WE. GABA(A) receptor subunit expression within hypophysiotropic CRH neurons: a dual hybridization histochemical study. *J Comp Neurol*. 2000;419:344-351.
- Critchlow V, Liebelt RA, Bar-Sela M, Mountcastle W, Lipscomb HS. Sex difference in resting pituitary-adrenal function in the rat. *Am J Physiol*. 1963;205:807-815.
- Seale JV, Wood SA, Atkinson HC, et al. Gonadectomy reverses the sexually dimorphic patterns of circadian and stress-induced hypothalamic-pituitary-adrenal axis activity in male and female rats. *J Neuroendocrinol*. 2004;16:516-524.
- Frederic F, Oliver C, Wollman E, Delhaye-Bouchaud N, Mariani J. IL-1 and LPS induce a sexually dimorphic response of the hypothalamo-pituitary-adrenal axis in several mouse strains. *Eur Cytokine Netw*. 1993;4:321-329.
- Mitsushima D, Masuda J, Kimura F. Sex differences in the stress-induced release of acetylcholine in the hippocampus and corticosterone from the adrenal cortex in rats. *Neuroendocrinology*. 2003;78:234-240.
- Goel N, Bale TL. Organizational and activational effects of testosterone on masculinization of female physiological and behavioral stress responses. *Endocrinology*. 2008;149:6399-6405.
- Iwasaki-Sekino A, Mano-Otagiri A, Ohata H, Yamauchi N, Shibasaki T. Gender differences in corticotropin and corticosterone secretion and corticotropin-releasing factor mRNA expression in the paraventricular nucleus of the hypothalamus and the central nucleus of the amygdala in response to footshock stress or psychological stress in rats. *Psychoneuroendocrinology*. 2009;34:226-237.
- Figueiredo HF, Dolgas CM, Herman JP. Stress activation of cortex and hippocampus is modulated by sex and stage of estrus. *Endocrinology*. 2002;143:2534-2540.
- Bland ST, Schmid MJ, Der-Avakian A, Watkins LR, Spencer RL, Maier SF. Expression of c-fos and BDNF mRNA in subregions of the prefrontal cortex of male and female rats after acute uncontrollable stress. *Brain Res*. 2005;1051:90-99.
- Goel N, Bale TL. Sex differences in the serotonergic influence on the hypothalamic-pituitary-adrenal stress axis. *Endocrinology*. 2010;151:1784-1794.
- Herman JP, Figueiredo H, Mueller NK, et al. Central mechanisms of stress integration: hierarchical circuitry controlling

- hypothalamo-pituitary-adrenocortical responsiveness. *Front Neuroendocrinol.* 2003;24:151-180.
36. Ulrich-Lai YM, Herman JP. Neural regulation of endocrine and autonomic stress responses. *Nat Rev Neurosci.* 2009;10:397-409.
 37. Herman JP, McKlveen JM, Ghosal S, et al. Regulation of the hypothalamic-pituitary-adrenocortical stress response. *Comp Physiol.* 2016;6:603-621.
 38. Lesniewska B, Nowak M, Malendowicz LK. Sex differences in adrenocortical structure and function. XXVIII. ACTH and corticosterone in intact, gonadectomised and gonadal hormone replaced rats. *Horm Metab Res.* 1990;22:378-381.
 39. Figueiredo HF, Ulrich-Lai YM, Choi DC, Herman JP. Estrogen potentiates adrenocortical responses to stress in female rats. *Am J Physiol Endocrinol Metab.* 2007;292:E1173-E1182.
 40. Serova LI, Harris HA, Maharjan S, Sabban EL. Modulation of responses to stress by estradiol benzoate and selective estrogen receptor agonists. *J Endocrinol.* 2010;205:253-262.
 41. Handa RJ, Nunley KM, Lorens SA, Louie JP, McGivern RF, Bollnow MR. Androgen regulation of adrenocorticotropin and corticosterone secretion in the male rat following novelty and foot shock stressors. *Physiol Behav.* 1994;55:117-124.
 42. McCormick CM, Furey BF, Child M, Sawyer MJ, Donohue SM. Neonatal sex hormones have 'organizational' effects on the hypothalamic-pituitary-adrenal axis of male rats. *Dev Brain Res.* 1998;105:295-307.
 43. Seale JV, Wood SA, Atkinson HC, Harbuz MS, Lightman SL. Gonadal steroid replacement reverses gonadectomy-induced changes in the corticosterone pulse profile and stress-induced hypothalamic-pituitary-adrenal axis activity of male and female rats. *J Neuroendocrinol.* 2004;16:989-998.
 44. Spinedi E, Suescun MO, Hadid R, Daneva T, Gaillard RC. Effects of gonadectomy and sex hormone therapy on the endotoxin-stimulated hypothalamo-pituitary-adrenal axis: evidence for a neuroendocrine-immunological sexual dimorphism. *Endocrinology.* 1992;131:2430-2436.
 45. Viau V, Meaney MJ. The inhibitory effect of testosterone on hypothalamic-pituitary-adrenal responses to stress is mediated by the medial preoptic area. *J Neurosci.* 1996;16:1866-1876.
 46. Viau V, Chu A, Soriano L, Dallman MF. Independent and overlapping effects of corticosterone and testosterone on corticotropin-releasing hormone and arginine vasopressin mRNA expression in the paraventricular nucleus of the hypothalamus and stress-induced adrenocorticotropin hormone release. *J Neurosci.* 1999;19:6684-6693.
 47. Viau V, Lee P, Sampson J, Wu J. A testicular influence on restraint-induced activation of medial parvocellular neurons in the paraventricular nucleus in the male rat. *Endocrinology.* 2003;144:3067-3075.
 48. Handa RJ, Weiser MJ, Zuloaga DG. A role for the androgen metabolite, 5 α -androstane-3 β ,17 β -diol, in modulating oestrogen receptor beta-mediated regulation of hormonal stress reactivity. *J Neuroendocrinol.* 2009;21:351-358.
 49. Viau V, Meaney MJ. Variations in the hypothalamic-pituitary-adrenal response to stress during the estrous cycle in the rat. *Endocrinology.* 1991;129:2503-2511.
 50. Barbaccia ML, Roscetti G, Trabucchi M, et al. Time-dependent changes in rat brain neuroactive steroid concentrations and GABA_A receptor function after acute stress. *Neuroendocrinology.* 1996;63:166-172.
 51. Barbaccia ML, Roscetti G, Bolacchi F, et al. Stress-induced increase in brain neuroactive steroids: antagonism by abecarnil. *Pharmacol Biochem Behav.* 1996;54:205-210.
 52. Vallee M, Rivera JD, Koob GF, Purdy RH, Fitzgerald RL. Quantification of neurosteroids in rat plasma and brain following swim stress and allopregnanolone administration using negative chemical ionization gas chromatography/mass spectrometry. *Anal Biochem.* 2000;287:153-166.
 53. Caruso D, Pesaresi M, Maschi O, Giatti S, Garcia-Segura LM, Melcangi RC. Effect of short-and long-term gonadectomy on neuroactive steroid levels in the central and peripheral nervous system of male and female rats. *J Neuroendocrinol.* 2010;22:1137-1147.
 54. Caruso D, Pesaresi M, Abbiati F, et al. Comparison of plasma and cerebrospinal fluid levels of neuroactive steroids with their brain, spinal cord and peripheral nerve levels in male and female rats. *Psychoneuroendocrinology.* 2013;38:2278-2290.
 55. Taves MD, Ma C, Heimovics SA, Saldanha CJ, Soma KK. Measurement of steroid concentrations in brain tissue: methodological considerations. *Front Endocrinol.* 2011;2:39.
 56. Schumacher M, Guennoun R, Mattern C, et al. Analytical challenges for measuring steroid responses to stress, neurodegeneration and injury in the central nervous system. *Steroids.* 2015;103:42-57.
 57. Armario A, Gavaldà A, Martí J. Comparison of the behavioural and endocrine response to forced swimming stress in five inbred strains of rats. *Psychoneuroendocrinology.* 1995;20:879-890.
 58. Harbuz MS, Lightman SL. Responses of hypothalamic and pituitary mRNA to physical and psychological stress in the rat. *J Endocrinol.* 1989;122:705-711.
 59. Cullinan WE, Herman JP, Battaglia DF, Akil H, Watson SJ. Pattern and time course of immediate early gene expression in rat brain following acute stress. *Neuroscience.* 1995;64:477-505.
 60. Neumann ID, Johnstone HA, Hatzinger M, et al. Attenuated neuroendocrine responses to emotional and physical stressors in pregnant rats involve adenohipophysial changes. *J Physiol.* 1998;508:289-300.
 61. Sharp S, Mitchell SJ, Vallee M, et al. Isotope dilution-based targeted and nontargeted carbonyl neurosteroid/steroid profiling. *Anal Chem.* 2018;90:5247-5255.
 62. Fry JP, Li KY, Devall AJ, Cockcroft S, Honour JW, Lovick TA. Fluoxetine elevates allopregnanolone in female rat brain but inhibits a steroid microsomal dehydrogenase rather than activating an aldo-keto reductase. *Br J Pharmacol.* 2014;171:5870-5880.
 63. Handa RJ, Burgess LH, Kerr JE, O'Keefe JA. Gonadal steroid hormone receptors and sex differences in the hypothalamo-pituitary-adrenal axis. *Horm Behav.* 1994;28:464-476.
 64. Tinnikov AA. Responses of serum corticosterone and corticosteroid-binding globulin to acute and prolonged stress in the rat. *Endocrine.* 1999;11:145-150.
 65. Goel N, Workman JL, Lee TT, Innala L, Viau V. Sex differences in the HPA axis. *Comp Physiol.* 2014;4:1121-1155.
 66. Malendowicz LK. Sex differences in adrenocortical structure and function. III. The effects of postpubertal gonadectomy and gonadal hormone replacement on adrenal cholesterol sidechain cleavage activity and on steroids biosynthesis by rat adrenal homogenates. *Endokrinologie.* 1976;67:26-35.
 67. Burgess LH, Handa RJ. Chronic estrogen-induced alterations in adrenocorticotropin and corticosterone secretion, and glucocorticoid receptor-mediated functions in female rats. *Endocrinology.* 1992;131:1261-1269.
 68. Droste SK, de Groot L, Lightman SL, Reul JM, Linthorst AC. The ultradian and circadian rhythms of free corticosterone in the brain are not affected by gender: an in vivo microdialysis study in Wistar rats. *J Neuroendocrinol.* 2009;21:132-140.
 69. Gala RR, Westphal U. Corticosteroid-binding globulin in the rat: studies on the sex difference. *Endocrinology.* 1965;77:841-851.
 70. Mellon SH, Deschepper CF. Neurosteroid biosynthesis: genes for adrenal steroidogenic enzymes are expressed in the brain. *Brain Res.* 1993;629:283-292.
 71. Caruso D, D'Intino G, Giatti S, et al. Sex-dimorphic changes in neuroactive steroid levels after chronic experimental autoimmune encephalomyelitis. *J Neurochem.* 2010;114:921-932.

72. Lovelace M, Watson TG, Stephenson GL. Steroid 21-hydroxylase expression in cultured rat astrocytes. *Brain Res Bull.* 2003;61:609-615.
73. Kishimoto W, Hiroi T, Shiraishi M, et al. Cytochrome P450 2D catalyze steroid 21-hydroxylation in the brain. *Endocrinology.* 2004;145:699-705.
74. Morrow AL, Devaud LL, Purdy RH, Paul SM. Neuroactive steroid modulators of the stress response. *Ann NY Acad Sci.* 1995;771:257-272.
75. Rossetti MF, Varayoud J, Andreoli MF, Stoker C, Luque EH, Ramos JG. Sex- and age-associated differences in episodic-like memory and transcriptional regulation of hippocampal steroidogenic enzymes in rats. *Mol Cell Endocrinol* 2017;470:208-218.
76. Celotti F, Melcangi RC, Negri-Cesi P, Ballabio M, Martini L. Differential distribution of the 5-alpha-reductase in the central nervous system of the rat and the mouse: are the white matter structures of the brain target tissue for testosterone action? *J Steroid Biochem.* 1987;26:125-129.
77. Cheng KC, Lee J, Khanna M, Qin KN. Distribution and ontogeny of 3 alpha-hydroxysteroid dehydrogenase in the rat brain. *J Steroid Biochem Mol Biol.* 1994;50:85-89.
78. Li X, Bertics PJ, Karavolas HJ. Regional distribution of cytosolic and particulate 5alpha-dihydroprogesterone 3alpha-hydroxysteroid oxidoreductases in female rat brain. *J Steroid Biochem Mol Biol.* 1997;60:311-318.
79. Castelli MP, Casti A, Casu A, et al. Regional distribution of 5alpha-reductase type 2 in the adult rat brain: an immunohistochemical analysis. *Psychoneuroendocrinology.* 2013;38:281-293.
80. Sanchez P, Torres JM, Olmo A, O'Valle F, Ortega E. Effects of environmental stress on mRNA and protein expression levels of steroid 5alpha-reductase isozymes in adult rat brain. *Horm Behav.* 2009;56:348-353.
81. Wirth MM. Beyond the HPA Axis: progesterone-derived neuroactive steroids in human stress and emotion. *Front Endocrinol.* 2011;2:19.
82. Bitran D, Shiekh M, McLeod M. Anxiolytic effect of progesterone is mediated by the neurosteroid allopregnanolone at brain GABA(A) receptors. *J Neuroendocrinol.* 1995;7:171-177.
83. Gunn BG, Cunningham L, Mitchell SG, Swinny JD, Lambert JJ, Belelli D. GABAA receptor-acting neurosteroids: a role in the development and regulation of the stress response. *Front Neuroendocrinol.* 2015;36:28-48.
84. Corpechot C, Collins BE, Carey MP, Tsouros A, Robel P, Fry JP. Brain neurosteroids during the mouse oestrous cycle. *Brain Res.* 1997;766:276-280.
85. Guo AL, Petraglia F, Criscuolo M, et al. Acute stress- or lipopolysaccharide-induced corticosterone secretion in female rats is independent of the oestrous cycle. *Eur J Endocrinol.* 1994;131:535-539.
86. Ogle TF, Kitay JI. Ovarian and adrenal steroids during pregnancy and the oestrous cycle in the rat. *J Endocrinol.* 1977;74:89-98.
87. Torres JM, Ruiz E, Ortega E. Effects of CRH and ACTH administration on plasma and brain neurosteroid levels. *Neurochem Res.* 2001;26:555-558.
88. Sanchez P, Torres JM, del Moral RG, de Dios LJ, Ortega E. Steroid 5alpha-reductase in adult rat brain after neonatal testosterone administration. *IUBMB Life.* 2012;64:81-86.
89. Torres JM, Ortega E. Steroid 5alpha-reductase isozymes in the adult female rat brain: central role of dihydrotestosterone. *J Mol Endocrinol.* 2006;36:239-245.
90. Mitev YA, Darwish M, Wolf SS, Holsboer F, Almeida OF, Patchev VK. Gender differences in the regulation of 3 alpha-hydroxysteroid dehydrogenase in rat brain and sensitivity to neurosteroid-mediated stress protection. *Neuroscience.* 2003;120:541-549.
91. Tappaz ML, Wassef M, Oertel WH, Paut L, Pujol JF. Light- and electron-microscopic immunocytochemistry of glutamic acid decarboxylase (GAD) in the basal hypothalamus: morphological evidence for neuroendocrine gamma-aminobutyrate (GABA). *Neuroscience.* 1983;9:271-287.
92. Sakae M, Saito N, Taniguchi H, Baba S, Tanaka C. Immunohistochemical localization of gamma-aminobutyric acid in the rat pituitary gland and related hypothalamic regions. *Brain Res.* 1988;446:343-353.
93. Backberg M, Ultenius C, Fritschy JM, Meister B. Cellular localization of GABA receptor alpha subunit immunoreactivity in the rat hypothalamus: relationship with neurons containing orexigenic or anorexigenic peptides. *J Neuroendocrinol.* 2004;16:589-604.
94. Do-Rego JL, Mensah-Nyagan GA, Beaujean D, et al. gamma-Aminobutyric acid, acting through gamma-aminobutyric acid type A receptors, inhibits the biosynthesis of neurosteroids in the frog hypothalamus. *Proc Natl Acad Sci USA.* 2000;97:13925-13930.
95. Skilbeck KJ, Hinton T, Johnston GA. Sex-differences and stress: effects on regional high and low affinity [³H]GABA binding. *Neurochem Int.* 2008;52:1212-1219.
96. Kavaliers M, Wiebe JP. Analgesic effects of the progesterone metabolite, 3 alpha-hydroxy-5 alpha-pregnan-20-one, and possible modes of action in mice. *Brain Res.* 1987;415:393-398.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Sze Y, Gill AC, Brunton PJ. Sex-dependent changes in neuroactive steroid concentrations in the rat brain following acute swim stress. *J Neuroendocrinol.* 2018;30:e12644. <https://doi.org/10.1111/jne.12644>