

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE



Verhey, R; Gibson, L; Brakarsh, J; Chibanda, D; Seedat, S (2018)  
Prevalence and correlates of probable post-traumatic stress disorder  
and common mental disorders in a population with a high prevalence  
of HIV in Zimbabwe. *EUROPEAN JOURNAL OF PSYCHOTRAU-  
MATOLOGY*, 9 (1). ISSN 2000-8066 DOI: <https://doi.org/10.1080/20008198.2018.1536286>

Downloaded from: <http://researchonline.lshtm.ac.uk/4650473/>

DOI: [10.1080/20008198.2018.1536286](https://doi.org/10.1080/20008198.2018.1536286)

#### Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: Creative Commons Attribution Non-commercial  
<http://creativecommons.org/licenses/by-nc/3.0/>



# Prevalence and correlates of probable post-traumatic stress disorder and common mental disorders in a population with a high prevalence of HIV in Zimbabwe

R. Verhey<sup>a</sup>, L. Gibson<sup>b</sup>, J. Brakarsh<sup>c</sup>, D. Chibanda<sup>a</sup> and S. Seedat<sup>d</sup>

<sup>a</sup>Research Support Center, University of Zimbabwe, Harare, Zimbabwe; <sup>b</sup>Department of Epidemiology, London School of Hygiene and Tropical Medicine, London, UK; <sup>c</sup>Say and Play Therapy Centre, Harare, Zimbabwe; <sup>d</sup>Department of Psychiatry, Stellenbosch University, Stellenbosch, South Africa

## ABSTRACT

**Background:** We investigated the prevalence of and factors associated with post-traumatic stress disorder (PTSD) and common mental disorders (CMDs), which include depression and anxiety disorders, in a setting with a prevalence of high human immunodeficiency virus (HIV) within a primary care clinic, using the PTSD Checklist for DSM-5 and the 14-item Shona Symptom Questionnaire, both locally validated screening tools.

**Methods:** A cross-sectional survey was carried out with adult patients ( $n = 204$ ) from the largest primary care clinic facility in Harare, Zimbabwe, in June 2016.

**Results:** A total of 83 patients (40.7%) met the criteria for probable PTSD, of whom 57 (69.5%) had comorbid CMDs. Among people living with HIV, 42 (55.3%) had PTSD. Probable PTSD was associated with having experienced a negative life event in the past 6 months [adjusted odds ratio (OR) 3.73, 95% confidence interval (CI) 1.49–9.34] or screening positive for one or more CMD (adjusted OR 6.48, 95% CI 3.35–2.54).

**Conclusion:** People living with HIV showed a high prevalence of PTSD and CMD comorbidity. PTSD screening should be considered when the CMD screen is positive and there is a history of negative life events.

## Prevalencia y correlatos de probable trastorno de estrés postraumático y trastornos mentales comunes en una población con una alta prevalencia de VIH en Zimbabue

**Antecedentes:** Investigamos la prevalencia y los factores asociados con TEPT y trastornos mentales comunes (TMC) que incluyen depresión y trastornos de ansiedad en una zona de alta prevalencia del VIH, en un centro de atención primaria (APS), utilizando el cuestionario de trastorno por estrés postraumático (PCL-5) y el cuestionario de síntomas de Shona (SSQ-14), ambos instrumentos validados localmente.

**Métodos:** se realizó un estudio transversal con pacientes adultos ( $n = 204$ ) del centro de atención primaria de salud más grande de Harare en junio de 2016.

**Resultados:** Un total de 83 (40.7%) cumplieron los criterios para probable TEPT, de los cuales 57 (69.5%) presentaron TMC comórbido. Entre las personas que viven con el VIH (PLWH), 42 (55.3%) tenían TEPT. El TEPT probable, se asoció con haber experimentado un evento de vida negativo en los últimos seis meses (AOR 3,73; IC del 95%: 1,49 - 9,34) o con el screening positivo para TMC (AOR 6,48; IC del 95%: 3,35 - 2,54).

**Conclusión:** Las personas que viven con el VIH (PLWH) mostraron una alta prevalencia de TEPT y comorbilidad con TMC. Se debe considerar la detección de TEPT cuando el screening para TMC es positivo y hay un historial de eventos vitales negativos.

## 津巴布韦高艾滋感染率人群中潜在创伤后应激障碍和常见精神障碍的患病率和相关性

**背景:** 我们在高艾滋感染率初级保健诊所 (PHC) 中, 使用两个经过本土验证的筛查工  
**目标:** 创伤后应激障碍检查表 (PCL-5) 和 Shona 症状问卷 (SSQ-14), 调查了 PTSD 和常见精神障碍 (CMD; 包括抑郁和焦虑) 的患病率和相关因素。

**方法:** 在 2016 年 6 月, 对 Harare 最大 PHC 的成年患者 ( $n = 204$ ) 进行横断面调查。

**结果:** 共有 83 例 (40.7%) 符合可能的 PTSD 标准, 其中 57 例 (69.5%) 共病 CMD。在艾滋病毒感染者 (PLWH) 中, 42 人 (55.3%) 患有 PTSD。可能的 PTSD 与过去六个月经历过负性生活事件有关 (AOR 3.73, 95% CI 1.49-9.34), 或与 CMD 筛查阳性 (AOR 6.48, 95% CI 3.35-2.54) 有关。

**结论:** PLWH 群体中表现出 PTSD 和 CMD 合并症的高患病率。当 CMD 筛查阳性且存在负性生活事件的历史时, 应考虑对其进行 PTSD 筛查。

## ARTICLE HISTORY

Received 14 May 2018

Revised 2 August 2018

Accepted 23 September 2018

## KEYWORDS

Post-traumatic stress disorder; common mental disorders; HIV; low- and middle-income countries; Zimbabwe; primary health care

## PALABRAS CLAVE

Trastorno de estrés postraumático; Trastornos mentales comunes; VIH; Países de bajos y medios ingresos; Zimbabue; Atención primaria de salud.

## 关键词

创伤后应激障碍; 常见精神障碍; HIV; 低收入和中等收入国家; 津巴布韦; 初级卫生保健

## HIGHLIGHTS

- PTSD was found in almost 41% of primary health-care users in Zimbabwe's capital.
- Nearly 45% of participants were HIV positive.
- Participants with PTSD were four times more likely than those without PTSD to have experienced a negative life event in the past six months.
- Participants with PTSD were six times more likely than those without PTSD to have CMDs.

## 1. Introduction

Post-traumatic stress disorder (PTSD) is highly prevalent in populations with high human immunodeficiency virus (HIV) rates (Adewuya et al., 2009; Seedat, 2012). Common mental disorders (CMDs), which include depression and anxiety disorders, are equally prevalent in high HIV settings (Chibanda, Cowan, Gibson, Weiss, & Lund, 2016). These disorders are a leading cause of disability globally (Steel et al., 2014). It is estimated that less than one-third of individuals in the USA and Europe and less than one-fifth in low- and middle-income countries (LMICs) receive treatment for treatable psychiatric conditions (Group, 2007; Kessler et al., 2005; Wang et al., 2007).

Studies show high comorbidity between depression and PTSD in a variety of population samples, such as those from conflict or post-conflict settings (Miller & Rasmussen, 2010), refugee camps (Gerritsen et al., 2006; Momartin, Silove, Manicavasagar, & Steel, 2004) and combat situations, victims of interpersonal violence (Stein & Kennedy, 2001), or survivors of natural or human-made disasters (Fernando, Miller, & Berger, 2010; Kar & Bastia, 2006; Neria, Nandi, & Galea, 2008). Political and economic instability and disparity; poverty, with its consequences of food scarcity and lack of access to medical care and education; politically motivated violence; crime; and exposure to weather phenomena due to climate change, which impacts negatively on an agriculture-based country such as Zimbabwe, further contribute towards poor mental health outcomes (Lund et al., 2010).

Receiving an HIV diagnosis, stigma from one's community, which often includes acts of interpersonal violence, and being confronted with a scarcity of treatment options also contribute towards unfavourable mental health outcomes (Hosegood, Preston-Whyte, Busza, Moitse, & Timaeus, 2007; Rankin, Brennan, Schell, Laviwa, & Rankin, 2005) and reduced adherence to anti-retroviral treatment (ART) (Boarts, Sledjeski, Bogart, & Delahanty, 2006), consequently posing a threat to survival. Stressors, both acute and chronic, to which inhabitants of LMICs are exposed often occur on a daily basis.

The overlap of symptoms of PTSD and depression is clinically contextualized in the latest version (Fifth Edition) of the Diagnostic and Statistical Manual for Mental Disorders (DSM-5) (American Psychiatric Association [APA], 2013; (Friedman, Resick, Bryant, & Brewin, 2011). Criterion C of the DSM-5 consists of six items comprising a symptom cluster of negative alterations in mood and cognition that can be applied to both PTSD and depression. The diagnosis of PTSD depends on fulfilment of criterion A, a stressor which

varies in severity and subjective appraisal (Weathers & Keane, 2007). Several studies have found that continuous lower level stressors are cumulatively more predictive of psychological distress than single-event exposures, as documented in studies in conflict or post-conflict and non-conflict settings (Miller & Rasmussen, 2010; Al-Krenawi, Lev-Wiesel, & Mahmud, 2007).

Furthermore, exposure to armed conflict and organized violence is often experienced as distant and found to be less salient over time (Miller & Rasmussen, 2010) in comparison to daily stressors such as poverty, overcrowded housing, chronic illness, and failure to meet daily basic and medical needs (lack of access to clean water, education, medication and sufficient food, to name a few). These daily stressors are seen as being able to erode psychological resilience and effective coping mechanisms through their cumulative impact, thus impeding individuals' overall mental health (Kubiak, 2005).

There are gender differences in the prevalence of PTSD: although more men are exposed to traumatic events than women (Kessler et al., 1999), women are more likely to develop PTSD (Breslau & Anthony, 2007; Breslau, Peterson, Poisson, Schultz, & Lucia, 2004) after exposure to a traumatic event. Individuals with PTSD have a higher likelihood of suffering from chronic medical conditions, with the number of lifetime traumas mediating this correlation (Sledjeski, Speisman, & Dierker, 2008). Ill health, in turn, has a negative impact on resilience and coping with the daily stressors that are common in LMICs (Piazza, Charles, Sliwinski, Mogle, & Almeida, 2012; Thakkar & McCanne, 2000). Cumulative effects of multiple traumas are also found to be common and associated with worse psychiatric outcomes (Green et al., 2000; Sledjeski et al., 2008; Suliman et al., 2009).

Recent work from Zimbabwe revealed high levels of CMDs and PTSD in primary care users, necessitating adequate interventions to alleviate this burden. Psychosocial interventions such as the Friendship Bench (Chibanda et al., 2011, 2016) seem to reduce the perception of lack of control among affected populations (Miller & Rasmussen, 2010). The Zimbabwean Friendship Bench programme is a low-intensity psychological intervention that is successfully reducing the treatment gap for CMDs at the primary health-care (PHC) level. The task-shifted intervention is delivered by trained and supervised lay health workers (LHWs) (Chibanda et al., 2016). There is a dearth of data on cultural understandings of trauma as well as on the societal and psychological impacts of low- versus high-intensity stressors and their influence on potential treatment considerations

of subsequent PTSD, and therefore rigorous research is needed. The purpose of this study was to establish the prevalence and correlates of PTSD in a population sample with high HIV prevalence using PHC facilities in Zimbabwe.

## 2. Methods

We carried out a cross-sectional study using the PTSD Checklist for DSM-5 (PCL-5) (Blevins, Weathers, Davis, Witte, & Domino, 2015) as well as the 14-item Shona Symptom Questionnaire (SSQ-14) (Patel, Simunyu, Gwanzura, Lewis, & Mann, 1997), both validated in Zimbabwe (Chibanda et al., 2016; Verhey, Chibanda, Gibson, Brakarsh, & Seedat, 2018), at the largest clinic in the suburb of Mbare, near the central business district of Harare, Zimbabwe. This clinic has a catchment area of over 200,000 with an average attendance of 140 patients per day and a high HIV prevalence. This is a typical PHC clinic in Harare and is mostly staffed by nurses and LHWs.

### 2.1. Inclusion criteria

Adults aged 18 years and above attending the clinic were eligible for recruitment over a 2 week period in June 2016, provided they were able to give written consent and resided in the area. Pregnant women in their last trimester and women within the 3 month postnatal period were excluded, as were those who were unable to understand the purpose of the study.

### 2.2. PCL-5

The PCL-5 was validated in Zimbabwe to determine the adequate cut-off score for an adult population (Verhey et al., 2018). It was found to have good psychometric properties, with a cut-off score of  $\geq 33$ , and sensitivity and specificity of 74.5% [95% confidence interval (CI) 60.4–85.7 and 70.6% (95% CI 62.7–77.7)], respectively. Cronbach's alpha was 0.92, reflecting very good internal consistency.

The PCL-5 is a self-report measure that was developed at the National Center for PTSD (Weathers, Litz, Herman, Huska, & Keane, 1994). It is used in conjunction with the Life Events Checklist for DSM-5 (LEC-5) (Gray, Litz, Hsu, & Lombardo, 2004). The LEC-5 is a self-report questionnaire which assesses the prevalence of 16 potentially traumatic lifetime events as well as an added open category ('any other very stressful event or experience'), with five answer categories (Weathers et al., 2013). The original PCL was revised to match the adapted DSM-5 criteria for PTSD. It features an adapted answer scale ranging from 0 = 'not at all' to 4 = 'extremely', thus making the theoretical lowest score 0. It measures four symptom clusters: the original clusters, intrusion, avoidance and hyperarousal; and the

added cluster of negative alterations in cognition and mood, with three items (blame, negative emotions, and reckless or self-destructive behaviour). Item scores can be summed for an overall severity score as well as for individual symptom cluster sums. A PTSD diagnosis can be made provisionally considering items rated 2 = moderately or higher according to the DSM-5 diagnostic rule (at least one B, one C, two D and two E symptoms present).

### 2.3. SSQ-14

The SSQ (Patel et al., 1997) was developed and validated in Zimbabwe in 1997. We have revalidated it in a high HIV prevalence setting, where a cut-off score of 9, with sensitivity and specificity of 84% (95% CI 78–89%) and 73% (95% CI 63–81%), respectively, was found to be suitable for detecting CMDs (Chibanda et al., 2016). Most of the 14 items are common to those found in tools for depression worldwide, such as sleep disturbance and suicidal thoughts; others are local idioms of emotional distress, including 'thinking too much'. Participants are asked whether they have experienced a list of common mental health symptoms in the past week. Each of the items is scored dichotomously as yes (1) or no (0) (Patel et al., 1997).

### 2.4. Translation of tools

The PCL-5 and the LEC-5 were translated from English into the local language, Shona, by a bilingual clinical social worker and a bilingual psychiatrist (DC). The draft Shona version was reviewed by a team of five LHWs working for the Friendship Bench and five nurses working in the psychiatric ward of Harare Central Hospital, together with a psychologist. This phase focused on ensuring contextual equivalence to the original versions based on their understanding and use of local terms for trauma symptomatology.

An independent language expert back-translated the Shona version into English. The first author together with the social worker and the psychiatrist examined both original and back-translated versions, and any discrepancies were resolved by consensus. Translation and back-translation were carried out using a standard approach (Bontempo, 1993).

### 2.5. Ethical considerations

The study was approved by the Medical Research Council of Zimbabwe (MRCZ, reference MRCZ/A/1732) and by the Health Research Ethics Committee at Stellenbosch University (reference S14/05/102). Ethical considerations and confidentiality for all participants were respected in accordance with Medical Research Guidelines.

## 2.6. Data collection

During the study period, we randomly selected participants on a daily basis from a register of all adult patients waiting to be seen at the clinic. Participants were selected based on a computer-generated random number sequence. Fifteen selected participants were invited at a time to a quiet and private space where eligibility was determined. Informed written consent was sought from all those eligible.

Trained research assistants administered the SSQ-14 and the PCL-5 to participants in randomly assigned alternating questionnaire order, and also collected sociodemographic information such as age, gender, HIV status, and marital and employment status using Samsung Galaxy tablets. HIV status was established by self-report. We aimed to recruit between 180 and 200 participants over the 2 week period.

Interviews took 20–30 minutes and were conducted in a quiet space designated for the study team. Although the PCL-5, LEC-5 and SSQ-14 are self-report tools, we chose to have them administered by trained research assistants, as prior validation exercises using the same approach found that most of the clinic attendees were not familiar with tablet/computer use (Chibanda et al., 2010, 2016).

## 2.7. Statistical analysis

As in an earlier study (Chibanda et al., 2016), data were entered directly into the study desktop computer by a data-entry clerk using a predesigned data-entry program containing automated range checks, and data cleaning was carried out at the end of each day. Data were transferred to STATA version 13.0 for analysis. Analysis was based on outcome measures of the PCL-5. Following tests for effect modification of HIV status and factors associated with CMDs, results were presented stratified by PCL-5 outcome. Sociodemographic variables of the two stratified groups with respect to meeting PCL-5 criteria for PTSD and SSQ-14 criteria were initially compared to establish differences. Variables with  $p < 0.1$  on univariate logistic regression analyses were taken forward to be included in the multivariable regression. Those variables with a  $p$ -value of less than 0.05 were then included in the final regression model to estimate adjusted odds ratios (ORs) and 95% CIs.

## 3. Results

### 3.1. Sample description

A total of 204 adults aged 18 and above gave consent to participate in the study, of whom 174 (85.2%) were women. Most of the participants (42.2%) were aged

below 30 years, while those aged 30–39 years made up 30.4% of the participants, with the rest being 40 years and older. The mean age was 34 years. Most participants were married (69.1%). There were 91 (44.6%) HIV-positive participants. Of these, 72 (67.9%) were female. A total of 100 (49%) were HIV negative, while 13 (6.4%) did not know their HIV status.

### 3.2. Prevalence of PTSD by PCL-5

Table 1 shows the characteristics of study participants by PCL-5  $\geq 33$  versus  $< 33$ . In total, 83 participants (40.7%) scored  $\geq 33$  on the PCL-5. Of the 83 who scored  $\geq 33$  on the PCL-5, 75 (90.4%) reported a negative life event in the past 6 months (95% CI 81.7–95.2,  $p < 0.001$ ). A total of 57 (69.5%) of those meeting case-ness on the PCL-5 scored above 9 on the SSQ-14 (95% CI 58.5–78.7,  $p < 0.001$ ) (Table 1). The odds of experi-

**Table 1.** Characteristics of participants by PTSD Checklist for DSM-5 (PCL-5) score ( $n = 204$ ).

	PCL-5 < 33 ( $n = 121$ )		PCL-5 $\geq 33$ ( $n = 83$ )		$\chi^2$ $p$ -value
	$n$	%	$n$	%	
All participants					
<b>HIV status<sup>a</sup></b>					0.09
Negative	66	57.4	34	44.7	
Positive	49	42.6	42	55.3	
<b>Gender</b>					0.75
Male	17	14.1	13	15.7	
Female	104	86.0	70	84.3	
<b>Age group (years)</b>					0.12
< 30	58	47.9	28	33.7	
30–39	34	28.1	28	33.7	
$\geq 40$	29	24.0	27	32.5	
<b>Marital status</b>					0.25
Married	88	72.7	53	63.9	
Single	17	14.1	19	22.9	
Widowed	16	13.2	11	13.3	
<b>Education</b>					0.13
Less than 'O' level	44	36.4	39	47.0	
'O' level or more	77	63.6	44	53.0	
<b>Current employment status</b>					0.06
Unemployed	76	62.8	38	45.8	
Permanent full or part time	6	5.0	6	7.2	
Casual/self-employed	39	32.2	39	47.0	
<b>Main income source</b>					0.03
Own business/salary	56	46.7	53	65.4	
Partner/family	52	43.3	23	28.4	
No income	12	10.0	5	6.2	
<b>Suffer from chronic illness</b>					0.02
No	57	47.1	25	30.1	
Yes	64	52.9	58	69.9	
<b>Reason for clinic visit</b>					0.25
HIV-related	18	14.9	20	24.1	
Routine/family/antenatal	58	47.9	35	42.2	
Other	45	37.2	28	33.7	
<b>Negative life events in past 6 months</b>					< 0.001
No	45	37.2	8	9.6	
Yes	76	62.8	75	90.4	
<b>SSQ-14 <math>\geq 9</math></b>					< 0.001
No	95	78.5	25	30.5	
Yes	26	21.5	57	69.5	

HIV, human immunodeficiency virus; SSQ-14, 14-item Shona Symptom Questionnaire.

<sup>a</sup> $n=13$  with unknown HIV status.

**Table 2.** Factors associated with PTSD Checklist for DSM-5 (PCL-5) score  $\geq 33$  ( $n = 204$ ).

All participants	Total <i>n</i>	%	PCL-5 $\geq 33$					
			Multivariable analysis			Final model		
			OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>
<b>HIV status</b>					0.92			
Negative	100	34.0	1	–				
Positive	91	46.2	1.04	(0.45–2.44)				
<b>Gender</b>								
Male	30	43.3						
Female	174	40.2						
<b>Age group (years)</b>								
< 30	86	32.6						
30–39	62	45.2						
$\geq 40$	56	48.2						
<b>Marital status</b>								
Married	141	37.6						
Single	36	52.8						
Widowed	27	40.7						
<b>Education</b>								
Less than 'O' level	83	47.0						
'O' level or more	121	36.4						
<b>Current employment status</b>					0.97			
Unemployed	114	33.3	1					
Permanent full or part time	12	50.0	1.17	(0.24–5.68)				
Casual/self-employed	78	50.0	1.15	(0.38–3.46)				
<b>Main income source</b>					0.36			
Own business/salary	109	48.6	1					
Partner/family	75	30.7	0.45	(0.14–1.40)				
No income	17	29.4	0.44	(0.09–2.10)				
<b>Suffer from chronic illness</b>					0.83			
No	82	30.5	1					
Yes	122	47.5	1.10	(0.46–2.65)				
<b>Negative life events</b>					0.01			0.005
No	53	15.1	1			1		
Yes	151	49.7	4.17	(1.50–11.62)		3.73	(1.49–9.34)	
<b>SSQ-14 <math>\geq 9</math></b>					< 0.001			< 0.001
No	120	20.8	1			1		
Yes	83	68.7	6.57	(3.16–16.68)		6.48	(3.35–12.54)	

HIV, human immunodeficiency virus; SSQ-14, 14-item Shona Symptom Questionnaire; CI, confidence interval; OR, odds ratio.

encing a negative life event in the past 6 months among those who scored positively for probable PTSD was almost four times higher than in those who did not [odds ratio (OR) 3.73, 95% CI 1.49–9.34] (Table 2).

Participants who were found to suffer from probable PTSD were more likely to score  $\geq 9$  on the SSQ-14 than those who did not (OR 6.48, 95% CI 3.35–12.54) (Table 2).

### 3.3. LEC-5: self-experienced traumatic index events

All participants ( $n = 204$ ) responded to the 17 items describing potentially traumatic life events. The mean number of self-experienced lifetime traumatic events was 4.6 (*SD* 2.4). The most commonly experienced lifetime traumas are listed in Table 3. In total, 132 participants reported experienced physical assault (64.7%), followed by any other very stressful event/experience, indicated by 130 participants (63.7%), as well as sudden, unexpected death of someone close, named by 113 participants (55.4%), and life-threatening illness/injury, listed by 89 (43.6%). Participants could report as many items as they felt appropriate.

**Table 3.** Life Events Checklist for DSM-5 (LEC-5): self-experienced exposure ( $n = 204$ ).

Name of item	<i>n</i>	%
1. Natural disaster (for example, flood, hurricane, tornado, earthquake)	51	25.0
2. Fire or explosion	23	11.3
3. Transportation accident (for example, car accident, boat accident, train wreck, plane crash)	31	15.2
4. Serious accident at work, home, or during recreational activity	30	14.7
5. Exposure to toxic substance (for example, dangerous chemicals, radiation)	14	6.9
6. Physical assault (for example, being attacked, hit, slapped, kicked, beaten up)	132	64.7
7. Assault with a weapon (for example, being shot, stabbed, threatened with a knife, gun, bomb)	36	17.6
8. Sexual assault (rape, attempted rape, made to perform any type of sexual act through force or threat of harm)	20	9.8
9. Other unwanted or uncomfortable sexual experience	20	9.8
10. Combat or exposure to a war-zone (in the military or as a civilian)	12	5.9
11. Captivity (for example, being kidnapped, abducted, held hostage, prisoner of war)	39	19.1
12. Life-threatening illness or injury	89	43.6
13. Severe human suffering	86	42.2
14. Sudden, violent death (for example, homicide, suicide)	25	12.3
15. Sudden, unexpected death of someone close to you	113	55.4
16. Serious injury, harm, or death you caused to someone else	7	3.4
17. Any other very stressful event or experience	130	63.7

## 4. Discussion

This article highlights the substantial occurrence of PTSD in a high HIV prevalence population sample using a PHC facility in an urban setting in Zimbabwe.

A probable PTSD prevalence of 40.7% was reported in this sample of PHC users who live in a non-armed-conflict setting ( $n = 204$ ).

Of the self-experienced lifetime index events, the highest prevalence was for physical assault, often in the form of interpersonal violence. Other specific lifetime events mentioned included sudden and unexpected death of a family member, as well as illness and severe human suffering.

These findings are in line with earlier findings where interpersonal violence was reported by more than 40% of those presenting with CMDs and receiving the Friendship Bench intervention (Chibanda et al., 2016). Experiencing domestic violence has been found to be an important trauma type associated with PTSD and CMDs (Kaminer, Grimsrud, Myer, Stein, & Williams, 2008; Schumm, Briggs-Phillips, & Hobfoll, 2006). There is evidence showing high rates of PTSD among victims of domestic violence (Campbell et al., 2008; Dutton et al., 2006; Machtinger, Wilson, Haberer, & Weiss, 2012). Our results support findings that physical assault and interpersonal violence are common in a population with a high HIV prevalence (Davis, 2012; Kouyoumdjian et al., 2013).

Furthermore, sudden and unexpected death of a close person as well as illness which is often related to untimely death were listed as common traumatic events. In LMICs, the population is faced with reduced life expectancy and quality of life, as injuries and disease (chronic or acute) potentially cause long-term disability (Hofman, Primack, Keusch, & Hrynokow, 2005) or death that could be avoided with improved access to a well-functioning health system (Ozgediz, Jamison, Cherian, & McQueen, 2008; Samb et al., 2010).

This study shows a high rate of CMDs as measured by a locally validated tool, the SSQ-14 (Chibanda et al., 2016; Patel et al., 1997), among participants who met the criteria for probable PTSD as measured by the PCL-5. This comorbidity is not unusual as symptoms associated with CMDs, such as feelings of helplessness, hopelessness, poor functioning and sleep disturbances, can be found in PTSD (DeViva, Zayfert, Pigeon, & Mellman, 2005).

Our findings further reflect the results of earlier studies that show the impact of suffering from a chronic condition such as HIV/AIDS and the increased likelihood of the development of PTSD and comorbid CMDs (Martin & Kagee, 2011; Olley, Zeier, Seedat, & Stein, 2005). Despite more people living with HIV being started on ART and HIV/AIDS no longer being considered a death sentence, HIV-positive status is still associated with stigma (Lee, Kochman, & Sikkema, 2002; Logie & Gadalla, 2009), even in the absence of physical signs. Stigma continues to have a negative effect on adherence to medication (Katz et al., 2013).

The significance and meaning of having HIV as a chronic condition can be seen as important to the risk of developing PTSD in this context (Silove, 1999). An earlier study (Verhey et al., 2018) revealed traumatic experiences, referred to as 'njodzi' in the local language, often being related to HIV and associated with the experience of PTSD symptoms; however, these symptoms were seen within the context of CMDs by LHWs. LHWs in the aforementioned study (Verhey et al., *in press*) had not been trained in the identification of PTSD-related symptoms. Furthermore, 90.4% of all the participants who scored positive for PTSD on the PCL-5 reported having experienced a negative life event in past 6 months (Table 1), indicating that the occurrence of index traumatic events is common in their living environment.

As 'kufungisisa kwe njodzi' (the local term for PTSD or PTSD symptoms, literally translated as 'thinking too much due to having experienced a trauma') is pervasive but not recognized as a condition that can be treated, many people will not seek help, thereby increasing the likelihood of developing PTSD (Clark, 1997; Flannery, 2001). Our findings demonstrate an association of PTSD with a negative life event in the past 6 months, and symptoms of CMDs. These findings have also been reported in several independent research studies (Beck, Grant, Clapp, & Palyo, 2009; Momartin et al., 2004; Olley et al., 2005; Olley, Seedat, & Stein, 2006; Schumm et al., 2006).

To decrease the burden of mental illness, an evidence-based approach, such as the Friendship Bench programme, can easily be offered in all PHC facilities as it has been shown to offer an efficient and cost-effective as well as culturally acceptable way to bridge the existing treatment gap (Chibanda et al., 2016). However, there is a need to address the comorbidity that is often found (O'Donnell, Creamer, & Pattison, 2004), particularly at the PHC level (Campbell et al., 2007; Herman et al., 2009). Subthreshold presentations of PTSD are associated with similar levels of psychosocial impairment to those found in people diagnosed with PTSD (Cukor, Wyka, Jayasinghe, & Difede, 2010; Zlotnick, Franklin, & Zimmerman, 2002). To lower the impact of psychosocial impairment on PHC facility users, both PTSD and subthreshold PTSD have to be recognized.

Assessment of exposure to negative life events, whether acute or chronic, can become part of a comprehensive health-care package with the aim of addressing stress disorder symptoms timeously. The need to train LHWs in how to identify PTSD-specific symptoms, such as phobic avoidance, hypervigilance and recurring nightmares, adding to their mental health literacy, was highlighted recently (Chibanda, Verhey, Munetsi, et al., 2016b). Integrating a

common elements approach (Murray et al., 2014) in the management of conditions by LHWs is considered to be an acceptable way to bridge the treatment gap.

This study illustrates the high prevalence of PTSD-like symptoms among people with symptoms of CMDs. Further studies will need to examine how LHWs can be used resourcefully to address the burden of comorbidity in LMICs. There is growing evidence supporting the efficient and effective use of transdiagnostic approaches, for example, for anxiety disorders (McManus, Shafran, & Cooper, 2010). PTSD had been part of the Fourth Edition of the DSM (DSM-IV) (APA, 2000) anxiety disorders category until the release of DSM-5 (APA, 2013). With the Friendship Bench programme being a community-based intervention, we believe that an integrated, culturally acceptable task-shifting approach with an added focus on stress disorders can make a difference to beneficiaries. Furthermore, we propose rigorous research, focusing on its effectiveness in decreasing the burden of compromised mental health in a holistic manner.

This study has several limitations. It was carried out in a PHC clinic in one of Harare's townships. The majority of the sample was female, with the unequal gender distribution being a reflection of women displaying more active health-seeking behaviour in Zimbabwe. Therefore, our results may not be generalizable across gender. We did not stratify negative life events by specific type of trauma, and therefore it is not possible to make any firm conclusions as to which specific negative life events are more likely to lead to PTSD, or what mix of negative life events is important. There is a need for further work on this.

## 5. Conclusion

PTSD is highly prevalent in this setting and is associated with a wide range of negative life events and CMD symptoms. To sustainably decrease the burden of mental illness, stress-related disorder symptoms and their link to previous exposure to traumatic events have to be recognized by LHWs working for the Friendship Bench. There is a need to develop and strengthen community-based, cost-effective and efficient integrated interventions aimed at addressing this and other comorbidities. We furthermore propose that a differentiation between lower intensity stressors and their impact on mental health and those that hold potential traumatic salience as defined by DSM-5 (APA, 2013) is especially important in LMICs.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by Grand Challenges Canada (GCC) [Grant number: 0087-04].

## ORCID

R. Verhey  <http://orcid.org/0000-0002-5959-1891>

## References

- Adewuya, A. O., Afolabi, M. O., Ola, B. A., Ogundele, O. A., Ajibare, A. O., Oladipo, B. F., & Fakande, I. (2009). Post-traumatic stress disorder (PTSD) after stigma related events in HIV infected individuals in Nigeria. *Social Psychiatry and Psychiatric Epidemiology*, 44(9), 761–766.
- Al-Krenawi, A., Lev-Wiesel, R., & Mahmud, A. S. (2007). Psychological symptomatology among Palestinian adolescents living with political violence. *Child and Adolescent Mental Health*, 12(1), 27–31.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders*, 4th ed. Washington, DC:Author.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders*, 5th ed. Washington, DC:Author.
- Beck, J. G., Grant, D. M., Clapp, J. D., & Palyo, S. A. (2009). Understanding the interpersonal impact of trauma: Contributions of PTSD and depression. *Journal of Anxiety Disorders*, 23(4), 443–450.
- Blevins, C. A., Weathers, F. W., Davis, M. T., Witte, T. K., & Domino, J. L. (2015). The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): Development and initial psychometric evaluation. *Journal of Traumatic Stress*, 28(6), 489–498.
- Boarts, J. M., Sledjeski, E. M., Bogart, L. M., & Delahanty, D. L. (2006). The differential impact of PTSD and depression on HIV disease markers and adherence to HAART in people living with HIV. *AIDS and Behavior*, 10(3), 253–261.
- Bontempo, R. (1993). Translation fidelity of psychological scales an item response theory analysis of an individualism-collectivism scale. *Journal of Cross-Cultural Psychology*, 24(2), 149–166.
- Breslau, N., & Anthony, J. C. (2007). Gender differences in the sensitivity to posttraumatic stress disorder: An epidemiological study of urban young adults. *Journal of Abnormal Psychology*, 116(3), 607.
- Breslau, N., Peterson, E., Poisson, L., Schultz, L., & Lucia, V. (2004). Estimating post-traumatic stress disorder in the community: Lifetime perspective and the impact of typical traumatic events. *Psychological Medicine*, 34(5), 889–898.
- Campbell, D. G., Felker, B. L., Liu, C.-F., Yano, E. M., Kirchner, J. E., Chan, D., ... Chaney, E. F. (2007). Prevalence of depression-PTSD comorbidity: Implications for clinical practice guidelines and primary care-based interventions. *Journal of General Internal Medicine*, 22(6), 711–718.
- Campbell, J. C., Baty, M., Ghandour, R. M., Stockman, J. K., Francisco, L., & Wagman, J. (2008). The intersection of intimate partner violence against women and HIV/AIDS: A review. *International Journal of Injury Control and Safety Promotion*, 15(4), 221–231.



- Chibanda, D., Cowan, F., Gibson, L., Weiss, H. A., & Lund, C. (2016). Prevalence and correlates of probable common mental disorders in a population with high prevalence of HIV in Zimbabwe. *BMC Psychiatry*, *16*, 55.
- Chibanda, D., Mangezi, W., Tshimanga, M., Woelk, G., Rusakaniko, P., Stranix-Chibanda, L., ... Shetty, A. K. (2010). Validation of the Edinburgh Postnatal Depression Scale among women in a high HIV prevalence area in urban Zimbabwe. *Archives of Women's Mental Health*, *13*(3), 201–206.
- Chibanda, D., Mesu, P., Kajawu, L., Cowan, F., Araya, R., & Abas, M. A. (2011). Problem-solving therapy for depression and common mental disorders in Zimbabwe: Piloting a task-shifting primary mental health care intervention in a population with a high prevalence of people living with HIV. *BMC Public Health*, *11*, 828.
- Chibanda, D., Verhey, R., Gibson, L. J., Munetsi, E., Machando, D., Rusakaniko, S., ... Abas, M. (2016). Validation of screening tools for depression and anxiety disorders in a primary care population with high HIV prevalence in Zimbabwe. *Journal of Affective Disorders*, *198*, 50–55.
- Chibanda, D., Weiss, H. A., Verhey, R., Simms, V., Munjoma, R., Rusakaniko, S., ... Araya, R. (2016a). Effect of a primary care-based psychological intervention on symptoms of common mental disorders in Zimbabwe: A randomized clinical trial. *JAMA : the Journal of the American Medical Association*, *316*(24), 2618–2626.
- Chibanda, D., Verhey, R., Munetsi, E., Rusakaniko, S., Cowan, F., & Lund, C. (2016b). Scaling up interventions for depression in sub-Saharan Africa: lessons from Zimbabwe. *Global Mental Health*, *3*, doi: [10.1017/gmh.2016.8](https://doi.org/10.1017/gmh.2016.8). eCollection 2016.
- Clark, C. C. (1997). Posttraumatic stress disorder: How to support healing. *The American Journal of Nursing*, *97*(8), 27–32.
- Cukor, J., Wyka, K., Jayasinghe, N., & Difede, J. (2010). The nature and course of subthreshold PTSD. *Journal of Anxiety Disorders*, *24*(8), 918–923.
- Davis, K. B. (2012). Understanding and addressing stigma and attachment insecurity in HIV-positive women who experience intimate partner violence: A model of medical and psychosocial care. *Social Work in Health Care*, *51*(3), 213–231.
- DeViva, J. C., Zayfert, C., Pigeon, W. R., & Mellman, T. A. (2005). Treatment of residual insomnia after CBT for PTSD: Case studies. *Journal of Traumatic Stress*, *18*(2), 155–159.
- Dutton, M. A., Green, B. L., Kaltman, S. I., Roesch, D. M., Zeffiro, T. A., & Krause, E. D. (2006). Intimate partner violence, PTSD, and adverse health outcomes. *Journal of Interpersonal Violence*, *21*(7), 955–968.
- Fernando, G. A., Miller, K. E., & Berger, D. E. (2010). Growing pains: The impact of disaster-related and daily stressors on the psychological and psychosocial functioning of youth in Sri Lanka. *Child Development*, *81*(4), 1192–1210.
- Flannery, R. B., Jr. (2001). The employee victim of violence: Recognizing the impact of untreated psychological trauma. *American Journal of Alzheimer's Disease & Other Dementias*, *16*(4), 230–233.
- Friedman, M. J., Resick, P. A., Bryant, R. A., & Brewin, C. R. (2011). Considering PTSD for DSM-5. *Depression and Anxiety*, *28*(9), 750–769.
- Gerritsen, A. A., Bramsen, I., Devillé, W., Van Willigen, L. H., Hovens, J. E., & Van Der Ploeg, H. M. (2006). Physical and mental health of Afghan, Iranian and Somali asylum seekers and refugees living in the Netherlands. *Social Psychiatry and Psychiatric Epidemiology*, *41*(1), 18–26.
- Gray, M. J., Litz, B. T., Hsu, J. L., & Lombardo, T. W. (2004). Psychometric properties of the Life Events Checklist. *Assessment*, *11*(4), 330–341.
- Green, B. L., Goodman, L. A., Krupnick, J. L., Corcoran, C. B., Petty, R. M., Stockton, P., & Stern, N. M. (2000). Outcomes of single versus multiple trauma exposure in a screening sample. *Journal of Traumatic Stress*, *13*(2), 271–286.
- Group, L. G. M. H. (2007). Scale up services for mental disorders: A call for action. *The Lancet*, *370*(9594), 1241–1252.
- Herman, A. A., Stein, D. J., Seedat, S., Heeringa, S. G., Moomal, H., & Williams, D. R. (2009). The South African Stress and Health (SASH) study: 12-month and lifetime prevalence of common mental disorders. *SAMJ: South African Medical Journal*, *99*(5), 339–344.
- Hofman, K., Primack, A., Keusch, G., & Hrynokow, S. (2005). Addressing the growing burden of trauma and injury in low-and middle-income countries. *American Journal of Public Health*, *95*(1), 13–17.
- Hosegood, V., Preston-Whyte, E., Busza, J., Moitse, S., & Timaeus, I. M. (2007). Revealing the full extent of households' experiences of HIV and AIDS in rural South Africa. *Social Science & Medicine*, *65*(6), 1249–1259.
- Kaminer, D., Grimsrud, A., Myer, L., Stein, D. J., & Williams, D. R. (2008). Risk for post-traumatic stress disorder associated with different forms of interpersonal violence in South Africa. *Social Science & Medicine*, *67*(10), 1589–1595.
- Kar, N., & Bastia, B. K. (2006). Post-traumatic stress disorder, depression and generalised anxiety disorder in adolescents after a natural disaster: A study of comorbidity. *Clinical Practice and Epidemiology in Mental Health*, *2*(1), 17.
- Katz, I. T., Ryu, A. E., Onuegbu, A. G., Psaros, C., Weiser, S. D., Bangsberg, D. R., & Tsai, A. C. (2013). Impact of HIV-related stigma on treatment adherence: Systematic review and meta-synthesis. *Journal of the International AIDS Society*, *16*(3 Suppl 2), 18640.
- Kessler, R. C., Demler, O., Frank, R. G., Olfson, M., Pincus, H. A., Walters, E. E., ... Zaslavsky, A. M. (2005). Prevalence and treatment of mental disorders, 1990 to 2003. *New England Journal of Medicine*, *352*(24), 2515–2523.
- Kessler, R. C., Sonnega, A., Bromet, E., Hughes, M., Nelson, C. B., & Breslau, N. (1999). Epidemiological risk factors for trauma and PTSD. In R. Yehuda (Ed.), *Risk factors for posttraumatic stress disorder* (pp. 23–59). Arlington, VA: American Psychiatric Association.
- Kouyoumdjian, F. G., Calzavara, L. M., Bondy, S. J., O'campo, P., Serwadda, D., Nalugoda, F., ... Gray, R. (2013). Intimate partner violence is associated with incident HIV infection in women in Uganda. *AIDS*, *27*(8), 1331–1338.
- Kubiak, S. P. (2005). Trauma and cumulative adversity in women of a disadvantaged social location. *American Journal of Orthopsychiatry*, *75*(4), 451.
- Lee, R. S., Kochman, A., & Sikkema, K. J. (2002). Internalized stigma among people living with HIV-AIDS. *AIDS and Behavior*, *6*(4), 309–319.
- Logie, C., & Gadalla, T. (2009). Meta-analysis of health and demographic correlates of stigma towards people living with HIV. *AIDS Care*, *21*(6), 742–753.
- Lund, C., Breen, A., Flisher, A. J., Kakuma, R., Corrigall, J., Joska, J. A., ... Patel, V. (2010). Poverty and common mental

- disorders in low and middle income countries: A systematic review. *Social Science & Medicine*, 71(3), 517–528.
- Machtiger, E., Wilson, T., Haberer, J. E., & Weiss, D. (2012). Psychological trauma and PTSD in HIV-positive women: A meta-analysis. *AIDS and Behavior*, 16(8), 2091–2100.
- Martin, L., & Kagee, A. (2011). Lifetime and HIV-related PTSD among persons recently diagnosed with HIV. *AIDS and Behavior*, 15(1), 125–131.
- McManus, F., Shafran, R., & Cooper, Z. (2010). What does a transdiagnostic approach have to offer the treatment of anxiety disorders? *British Journal of Clinical Psychology*, 49(4), 491–505.
- Miller, K. E., & Rasmussen, A. (2010). War exposure, daily stressors, and mental health in conflict and post-conflict settings: Bridging the divide between trauma-focused and psychosocial frameworks. *Social Science & Medicine*, 70(1), 7–16.
- Momartin, S., Silove, D., Manicavasagar, V., & Steel, Z. (2004). Comorbidity of PTSD and depression: Associations with trauma exposure, symptom severity and functional impairment in Bosnian refugees resettled in Australia. *Journal of Affective Disorders*, 80(2), 231–238.
- Murray, L. K., Dorsey, S., Haroz, E., Lee, C., Alsiary, M. M., Haydary, A., ... Bolton, P. (2014). A common elements treatment approach for adult mental health problems in low- and middle-income countries. *Cognitive and Behavioral Practice*, 21(2), 111–123.
- Neria, Y., Nandi, A., & Galea, S. (2008). Post-traumatic stress disorder following disasters: A systematic review. *Psychological Medicine*, 38(04), 467–480.
- O'Donnell, M. L., Creamer, M., & Pattison, P. (2004). Posttraumatic stress disorder and depression following trauma: Understanding comorbidity. *American Journal of Psychiatry*, 161(8), 1390–1396.
- Olley, B., Zeier, M., Seedat, S., & Stein, D. (2005). Post-traumatic stress disorder among recently diagnosed patients with HIV/AIDS in South Africa. *AIDS Care*, 17(5), 550–557.
- Olley, B. O., Seedat, S., & Stein, D. J. (2006). Persistence of psychiatric disorders in a cohort of HIV/AIDS patients in South Africa: A 6-month follow-up study. *Journal of Psychosomatic Research*, 61(4), 479–484.
- Ozgediz, D., Jamison, D., Cherian, M., & McQueen, K. (2008). The burden of surgical conditions and access to surgical care in low-and middle-income countries. *Bulletin of the World Health Organization*, 86(8), 646–647.
- Patel, V., Simunyu, E., Gwanzura, F., Lewis, G., & Mann, A. (1997). The Shona Symptom Questionnaire: The development of an indigenous measure of common mental disorders in Harare. *Acta Psychiatrica Scandinavica*, 95(6), 469–475.
- Piazza, J. R., Charles, S. T., Sliwinski, M. J., Mogle, J., & Almeida, D. M. (2012). Affective reactivity to daily stressors and long-term risk of reporting a chronic physical health condition. *Annals of Behavioral Medicine*, 45(1), 110–120.
- Rankin, W. W., Brennan, S., Schell, E., Laviwa, J., & Rankin, S. H. (2005). The stigma of being HIV-positive in Africa. *PLoS Medicine*, 2(8), e247.
- Samb, B., Desai, N., Nishtar, S., Mendis, S., Bekedam, H., Wright, A., ... Patel, K. (2010). Prevention and management of chronic disease: A litmus test for health-systems strengthening in low-income and middle-income countries. *The Lancet*, 376(9754), 1785–1797.
- Schumm, J. A., Briggs-Phillips, M., & Hobfoll, S. E. (2006). Cumulative interpersonal traumas and social support as risk and resiliency factors in predicting PTSD and depression among inner-city women. *Journal of Traumatic Stress*, 19(6), 825–836.
- Seedat, S. (2012). Interventions to improve psychological functioning and health outcomes of HIV-infected individuals with a history of trauma or PTSD. *Current HIV/AIDS Reports*, 9(4), 344–350.
- Silove, D. (1999). The psychosocial effects of torture, mass human rights violations, and refugee trauma: Toward an integrated conceptual framework. *The Journal of Nervous and Mental Disease*, 187(4), 200–207.
- Sledjeski, E. M., Speisman, B., & Dierker, L. C. (2008). Does number of lifetime traumas explain the relationship between PTSD and chronic medical conditions? Answers from the National Comorbidity Survey-Replication (NCS-R). *Journal of Behavioral Medicine*, 31(4), 341–349.
- Steel, Z., Marnane, C., Iranpour, C., Chey, T., Jackson, J. W., Patel, V., & Silove, D. (2014). The global prevalence of common mental disorders: A systematic review and meta-analysis 1980–2013. *International Journal of Epidemiology*, 43(2), 476–493.
- Stein, M. B., & Kennedy, C. (2001). Major depressive and post-traumatic stress disorder comorbidity in female victims of intimate partner violence. *Journal of Affective Disorders*, 66(2), 133–138.
- Suliman, S., Mkabile, S. G., Fincham, D. S., Ahmed, R., Stein, D. J., & Seedat, S. (2009). Cumulative effect of multiple trauma on symptoms of posttraumatic stress disorder, anxiety, and depression in adolescents. *Comprehensive Psychiatry*, 50(2), 121–127.
- Thakkar, R. R., & McCanne, T. R. (2000). The effects of daily stressors on physical health in women with and without a childhood history of sexual abuse. *Child Abuse & Neglect*, 24(2), 209–221.
- Verhey, R., Chibanda, D., Gibson, L., Brakarsh, J., & Seedat, S. (in press). Validation of the posttraumatic stress disorder checklist - 5 (PCL-5) in a primary care population with high HIV prevalence in Zimbabwe. *BMC Psychiatry*, 18(1), 109.
- Wang, P. S., Aguilar-Gaxiola, S., Alonso, J., Angermeyer, M. C., Borges, G., Bromet, E. J., ... Gureje, O. (2007). Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. *The Lancet*, 370(9590), 841–850.
- Weathers, F. W., Blake, D. D., Schnurr, P. P., Kaloupek, D. G., Marx, B. P., & Keane, T. M. (2013). The Life Events Checklist for DSM-5 (LEC-5). Instrument available from the National Center for PTSD. 2013.
- Weathers, F., Litz, B., Herman, D., Huska, J., & Keane, T. (1994). *The PTSD Checklist-Civilian Version (PCL-C)*. Boston, MA: National Center for PTSD.
- Weathers, F. W., & Keane, T. M. (2007). The Criterion A problem revisited: Controversies and challenges in defining and measuring psychological trauma. *Journal of Traumatic Stress*, 20(2), 107–121.
- Zlotnick, C., Franklin, C. L., & Zimmerman, M. (2002). Does “subthreshold” posttraumatic stress disorder have any clinical relevance? *Comprehensive Psychiatry*, 43(6), 413–419.