



Miskolc Mathematical Notes
Vol. 8 (2007), No 1, pp. 23-30

HU e-ISSN 1787-2413
DOI: 10.18514/MMN.2007.161

Some new inverse-type Hilbert-Pachpatte inequalities

Zhao Changjian and Mihály Bencze



SOME NEW INVERSE-TYPE HILBERT-PACHPATTE INEQUALITIES

ZHAO CHANGJIAN AND MIHÁLY BENCZE

Received 2 June, 2006

Abstract. Inverses of some new inequalities similar to the Hilbert–Pachpatte inequality are established.

1991 *Mathematics Subject Classification:* 26D15

Keywords: Hilbert inequality, Hölder integral inequality, Jensen inequality

1. INTRODUCTION

In recent years several authors [2, 3, 5–10] have given considerable attention to the Hilbert inequalities and Hilbert type inequalities and their various generalizations. In particular, in 1988, B. G. Pachpatte [6] proved some new inequalities similar to Hilbert’s inequality [4, p. 226], The main purpose of this paper is to establish their inverses.

2. MAIN RESULTS

Our main results are given in the following theorems.

Theorem 1. *Let $0 < p \leq 1, 0 < q \leq 1$ and $\{a_m\}$ and $\{b_n\}$ be two nonnegative sequences of real numbers defined for $m = 1, 2, \dots, k$ and $n = 1, 2, \dots, r$, where k and r are the natural numbers and define $A_m = \sum_{s=1}^m a_s$ and $B_n = \sum_{t=1}^n b_t$. Then*

For the first author, the research is supported by Zhejiang Provincial Natural Science Foundation of China (Y605065), Foundation of the Education Department of Zhejiang Province of China (20050392), and the Academic Mainstay of Middle-age and Youth Foundation of Shandong Province of China (200203).

for $3n - 2m > 0$

$$\sum_{m=1}^k \sum_{n=1}^r \frac{A_m^p B_n^q}{3n-2m} \geq pqk^{-2}r^3 \left(\sum_{m=1}^k (k-m+1) (a_m A_m^p)^{-\frac{11}{3}} \right)^3 \times \left(\sum_{n=1}^r (r-n+1) (b_n B_n^{q-1})^{-\frac{1}{2}} \right)^{-2}. \quad (1)$$

Proof. By using the following inequality (see [4, p. 39])

$$rx^{r-1}(x-y) \leq x^r - y^r, \quad 0 < r \leq 1,$$

where x and y are positive real numbers, we obtain that

$$\sum_{m=0}^{k-1} pA_{m+1}^{p-1} (A_{m+1} - A_m) \leq \sum_{m=0}^{k-1} (A_{m+1}^p - A_m^p),$$

that is

$$A_k^p \geq p \sum_{m=1}^k a_m A_m^{p-1}. \quad (2)$$

From (2) and in view of Hölder's inequality [4, p. 24], we have

$$\begin{aligned} \frac{A_m^p B_n^q}{3n-2m} &\geq pq(3n-2m)^{-1} \left(\sum_{s=1}^m a_s A_s^{p-1} \right) \left(\sum_{t=1}^n b_t B_t^{q-1} \right) \\ &\geq pq(3n-2m)^{-1} n^3 m^{-2} \left(\sum_{s=1}^m (a_s A_s^{p-1})^{\frac{1}{3}} \right)^3 \\ &\quad \times \left(\sum_{t=1}^n (b_t B_t^{q-1})^{-\frac{1}{2}} \right)^{-2}. \end{aligned} \quad (3)$$

Using the following inequality [1, p. 15]

$$m^{\frac{1}{h}} n^{\frac{1}{l}} \geq \frac{m}{h} + \frac{n}{l}, \quad (4)$$

where $m > 0, n > 0, h^{-1} + l^{-1} = 1$, and $h < 1$, we easily get that

$$n^3 m^{-2} \geq 3n - 2m. \quad (5)$$

Taking the sum on both sides of (3) over n from 1 to r first and then the sum over m from 1 to k and in view of inequality (5) and using again Hölder's inequality, we

obtain

$$\begin{aligned}
 \sum_{m=1}^k \sum_{n=1}^r \frac{A_m^p B_n^q}{3n-2m} &\geq pq \left(\sum_{m=1}^k \left(\sum_{s=1}^m (a_s A_s^{p-1})^{\frac{1}{3}} \right)^3 \right) \\
 &\quad \times \left(\sum_{n=1}^r \left(\sum_{t=1}^n (b_t B_t^{q-1})^{-\frac{1}{2}} \right)^{-2} \right) \\
 &\geq pqk^{-2}r^3 \left(\sum_{m=1}^k \left(\sum_{s=1}^m (a_s A_s^{p-1})^{\frac{1}{3}} \right) \right)^3 \left(\sum_{n=1}^r \left(\sum_{t=1}^n (b_t B_t^{q-1})^{-\frac{1}{2}} \right) \right)^{-2} \\
 &= pqk^{-2}r^3 \left(\sum_{s=1}^k (a_s A_s^{p-1})^{\frac{1}{3}} \left(\sum_{m=s}^k 1 \right) \right)^3 \left(\sum_{t=1}^r (b_t B_t^{q-1})^{-\frac{1}{2}} \left(\sum_{n=t}^r 1 \right) \right)^{-2} \\
 &= pqk^{-2}r^3 \left(\sum_{s=1}^k (k-s+1) (a_s A_s^{p-1})^{\frac{1}{3}} \right)^3 \\
 &\quad \times \left(\sum_{t=1}^r (r-t+1) (b_t B_t^{q-1})^{-\frac{1}{2}} \right)^{-2} \\
 &= pqk^{-2}r^3 \left(\sum_{m=1}^k (k-m+1) (a_m A_m^{p-1})^{\frac{1}{3}} \right)^3 \\
 &\quad \times \left(\sum_{n=1}^r (r-n+1) (b_n B_n^{q-1})^{-\frac{1}{2}} \right)^{-2},
 \end{aligned}$$

which completes the proof. \square

Remark 1. Inequality (1) is just an inverse of the following Inequality A, which was proved by Pachpatte [6]:

Inequality A.

$$\begin{aligned}
 \sum_{m=1}^k \sum_{n=1}^r \frac{A_m^p B_n^q}{m+n} &\leq \frac{1}{2} pq (kr)^{\frac{1}{2}} \left(\sum_{m=1}^k (k-m+1) (a_m A_m^{p-1})^2 \right)^{\frac{1}{2}} \\
 &\quad \times \left(\sum_{n=1}^r (r-n+1) (b_n B_n^{q-1})^2 \right)^{\frac{1}{2}}.
 \end{aligned}$$

Theorem 2. Let $\{a_m\}, \{b_n\}, A_m, B_n$ be as defined in Theorem 1. Let $\{p_m\}$ and $\{q_n\}$ be two positive sequences for $m = 1, 2, \dots, k$ and $n = 1, 2, \dots, r$, where k and r are natural numbers and put $P_m = \sum_{s=1}^m p_s, Q_n = \sum_{t=1}^n q_t$. Let ϕ and ψ be two real-valued nonnegative, concave, and supermultiplicative functions* defined on \mathbb{R}_+ . Then for $3n - 2m > 0$

$$\sum_{m=1}^k \sum_{n=1}^r \frac{\phi(A_m)\psi(B_n)}{3n-2m} \geq M'(k, r) \left(\sum_{m=1}^k \left(p_m \phi \left(\frac{a_m}{p_m} \right) \right)^{\frac{1}{3}} (k-m+1) \right)^3 \times \left(\sum_{n=1}^r \left(q_n \psi \left(\frac{b_n}{q_n} \right) \right)^{-\frac{1}{2}} (r-n+1) \right)^{-2}, \quad (6)$$

where

$$M'(k, r) = \left(\sum_{m=1}^k \left(\frac{\phi(P_m)}{P_m} \right)^{-\frac{1}{2}} \right)^{-2} \left(\sum_{n=1}^r \left(\frac{\psi(Q_n)}{Q_n} \right)^{\frac{1}{3}} \right)^3.$$

Proof. From the hypotheses and by Jensen's inequality and Hölder's inequality, we obtain

$$\begin{aligned} \frac{\phi(A_m)\psi(B_n)}{3n-2m} &\geq (3n-2m)^{-1} \\ &\quad \times \phi(P_m)\psi(Q_n)\phi\left(\frac{\sum_{s=1}^m p_s a_s/p_s}{\sum_{s=1}^m p_s}\right)\psi\left(\frac{\sum_{t=1}^n q_t b_t/q_t}{\sum_{t=1}^n q_t}\right) \\ &\geq (3n-2m)^{-1} \frac{\phi(P_m)}{P_m} \frac{\psi(Q_n)}{Q_n} \sum_{s=1}^m p_s \phi\left(\frac{a_s}{p_s}\right) \sum_{t=1}^n q_t \psi\left(\frac{b_t}{q_t}\right) \\ &\geq (3n-2m)^{-1} n^3 m^{-2} \frac{\phi(P_m)}{P_m} \frac{\psi(Q_n)}{Q_n} \left(\sum_{s=1}^m \left(p_s \phi\left(\frac{a_s}{p_s}\right) \right)^{\frac{1}{3}} \right)^3 \\ &\quad \times \left(\sum_{t=1}^n \left(q_t \psi\left(\frac{b_t}{q_t}\right) \right)^{-\frac{1}{2}} \right)^{-2} \end{aligned}$$

* f is said to be a supermultiplicative function if $f(xy) \geq f(x)f(y)$ for $x, y \in \mathbb{R}_+ := [0, +\infty)$.

Taking the sum over n from 1 to r first and then the sum over m from 1 to k and using (5), in view of Hölder's inequality, we have

$$\begin{aligned}
 \sum_{m=1}^k \sum_{n=1}^r \frac{\phi(A_m) \psi(B_n)}{3n-2m} &\geq \sum_{m=1}^k \left(\frac{\phi(P_m)}{P_m} \left(\sum_{s=1}^m \left(p_s \phi \left(\frac{a_s}{p_s} \right) \right)^{\frac{1}{3}} \right)^3 \right) \\
 &\quad \times \sum_{n=1}^r \left(\frac{\psi(Q_n)}{Q_n} \left(\sum_{t=1}^n \left(q_t \psi \left(\frac{b_t}{q_t} \right) \right)^{-\frac{1}{2}} \right)^{-2} \right) \\
 &\geq \left(\sum_{m=1}^k \left(\frac{\phi(P_m)}{P_m} \right)^{-\frac{1}{2}} \right)^{-2} \left(\sum_{m=1}^k \sum_{s=1}^m p_s \phi \left(\frac{a_s}{p_s} \right)^{\frac{1}{3}} \right)^3 \\
 &\quad \times \left(\sum_{n=1}^r \left(\frac{\psi(Q_n)}{Q_n} \right)^{\frac{1}{3}} \right)^3 \left(\sum_{n=1}^r \sum_{t=1}^n q_t \psi \left(\frac{b_t}{q_t} \right)^{-\frac{1}{2}} \right)^{-2} \\
 &= M'(k, r) \left(\sum_{s=1}^k \left(p_s \phi \left(\frac{a_s}{p_s} \right) \right)^{\frac{1}{3}} \left(\sum_{m=s}^k 1 \right) \right)^3 \\
 &\quad \times \left(\sum_{t=1}^r \left(q_t \psi \left(\frac{b_t}{q_t} \right) \right)^{-\frac{1}{2}} \left(\sum_{n=t}^r 1 \right) \right)^{-2} \\
 &= M'(k, r) \left(\sum_{s=1}^k \left(p_s \phi \left(\frac{a_s}{p_s} \right) \right)^{\frac{1}{3}} (k-s+1) \right)^3 \\
 &\quad \times \left(\sum_{t=1}^r \left(q_t \psi \left(\frac{b_t}{q_t} \right) \right)^{-\frac{1}{2}} (r-t+1) \right)^{-2} \\
 &= M'(k, r) \left(\sum_{m=1}^k \left(p_m \phi \left(\frac{a_m}{p_m} \right) \right)^{\frac{1}{3}} (k-m+1) \right)^3 \\
 &\quad \times \left(\sum_{n=1}^r \left(q_n \psi \left(\frac{b_n}{q_n} \right) \right)^{-\frac{1}{2}} (r-n+1) \right)^{-2}.
 \end{aligned}$$

The proof is complete. \square

Remark 2. Inequality (6) is just an inverse of the following Inequality B, which was proved by Pachpatte [6]:

Inequality B.

$$\sum_{m=1}^k \sum_{n=1}^r \frac{\phi(A_m) \psi(B_n)}{m+n} \leq M(k, r) \left(\sum_{m=1}^k (k-m+1) \left(p_m \phi \left(\frac{a_m}{p_m} \right) \right)^2 \right)^{\frac{1}{2}} \\ \times \left(\sum_{n=1}^r (r-n+1) \left(q_n \psi \left(\frac{b_n}{q_n} \right) \right)^2 \right)^{\frac{1}{2}},$$

where

$$M(k, r) = \frac{1}{2} \left(\sum_{m=1}^k \left(\frac{\phi(P_m)}{P_m} \right)^2 \right)^{\frac{1}{2}} \left(\sum_{n=1}^r \left(\frac{\psi(Q_n)}{Q_n} \right)^2 \right)^{\frac{1}{2}}.$$

Similarly, the following two theorems can also be established.

Theorem 3. Let $\{a_m\}$ and $\{b_n\}$ be as in Theorem 1 and set $A_m = (1/m) \sum_{s=1}^m a_s$ and $B_n = (1/n) \sum_{t=1}^n b_t$, for $m = 1, 2, \dots, k$ and $n = 1, 2, \dots, r$, where k and r are natural numbers. Let ϕ and ψ be two real-valued, nonnegative, and concave functions defined on \mathbb{R}_+ . Then for $3n - 2m > 0$,

$$\sum_{m=1}^k \sum_{n=1}^r \frac{mn}{3n-2m} \phi(A_m) \psi(B_n) \geq k^{-2} r^3 \left(\sum_{m=1}^k (k-m+1) (\phi(a_m))^{\frac{1}{3}} \right)^3 \\ \times \left(\sum_{n=1}^r (r-n+1) (\psi(b_n))^{-\frac{1}{2}} \right)^{-2}. \quad (7)$$

Remark 3. Inequality (7) is just an inverse of the following Inequality C, which was proved by Pachpatte [6]:

Inequality C.

$$\sum_{m=1}^k \sum_{n=1}^r \frac{mn}{m+n} \phi(A_m) \psi(B_n) \leq \frac{1}{2} (kr)^{\frac{1}{2}} \left(\sum_{m=1}^k (k-m+1) (\phi(a_m))^2 \right)^{\frac{1}{2}} \\ \times \left(\sum_{n=1}^r (r-n+1) (\psi(b_n))^2 \right)^{\frac{1}{2}}.$$

Theorem 4. Let $\{a_m\}, \{b_n\}, \{p_m\}, \{q_n\}, P_m, Q_n$ be as in Theorem 2 and put $A_m = (1/p_m) \sum_{s=1}^m p_s a_s$, $B_n = (1/Q_n) \sum_{t=1}^n q_t b_t$, for $m = 1, 2, \dots, k$, $n = 1, 2, \dots, r$, where k and r are natural numbers. Let ϕ and ψ be as defined in Theorem 3. Then

for $3n - 2m > 0$,

$$\sum_{m=1}^k \sum_{n=1}^r \frac{P_m Q_n \phi(A_m) \psi(B_n)}{3n - 2m} \geq k^{-2} r^3 \left(\sum_{m=1}^k (k - m + 1) (p_m \phi(a_m))^{\frac{1}{3}} \right)^3 \times \left(\sum_{n=1}^r (r - n + 1) (q_n \psi(b_n))^{-\frac{1}{2}} \right)^{-2}. \quad (8)$$

Remark 4. Inequality (8) is just an inverse of the following Inequality D, which was proved by Pachpatte [6]:

Inequality D.

$$\sum_{m=1}^k \sum_{n=1}^r \frac{P_m Q_n \phi(A_m) \psi(B_n)}{m + n} \leq \frac{\sqrt{kr}}{2} \left(\sum_{m=1}^k (k - m + 1) (p_m \phi(a_m))^2 \right)^{\frac{1}{2}} \times \left(\sum_{n=1}^r (r - n + 1) (q_n \psi(b_n))^2 \right)^{\frac{1}{2}}.$$

The proofs of Theorems 3 and 4 can be completed by following the same steps as in the proof of Theorem 2 with suitable changes. Here, we omit the details.

REFERENCES

- [1] E. Beckenbach and R. Bellman, *An Introduction to Inequalities*, ser. New Mathematical Library. New York: Random House, 1961, vol. 3.
- [2] M. Gao, "On Hilbert's inequality and its applications," *J. Math. Anal. Appl.*, vol. 212, no. 1, pp. 316–323, 1997.
- [3] M. Gao and B. Yang, "On the extended Hilbert's inequality," *Proc. Amer. Math. Soc.*, vol. 126, no. 3, pp. 751–759, 1998.
- [4] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*. Cambridge: Cambridge University Press, 1952, 2d ed.
- [5] K. Jichang, "On new extensions of Hilbert's integral inequality," *J. Math. Anal. Appl.*, vol. 235, no. 2, pp. 608–614, 1999.
- [6] B. G. Pachpatte, "On some new inequalities similar to Hilbert's inequality," *J. Math. Anal. Appl.*, vol. 226, no. 1, pp. 166–179, 1998.
- [7] B. G. Pachpatte, "Inequalities similar to certain extensions of Hilbert's inequality," *J. Math. Anal. Appl.*, vol. 243, no. 2, pp. 217–227, 2000.
- [8] B. Yang, "On new generalizations of Hilbert's inequality," *J. Math. Anal. Appl.*, vol. 248, no. 1, pp. 29–40, 2000.
- [9] C.-j. Zhao, "Generalization on two new Hilbert type inequalities," *J. Math. (Wuhan)*, vol. 20, no. 4, pp. 413–416, 2000.
- [10] C.-J. Zhao and L. Debnath, "Some new inverse type Hilbert integral inequalities," *J. Math. Anal. Appl.*, vol. 262, no. 1, pp. 411–418, 2001.

*Authors' addresses***Zhao Changjian**

Department of Information and Mathematics Sciences, College of Science, China Jiliang University,
Hangzhou 310018, P. R. China

E-mail address: chjzhao@163.com

Mihály Bencze

6 Harmanului St., 505600 Sacele-Négyfalu, Jud. Brasov, Romania