



Miskolc Mathematical Notes
Vol. 15 (2014), No 2, pp. 615-623

HU e-ISSN 1787-2413
DOI: 10.18514/MMN.2014.1011

Best proximity point theorems for F -contractive non-self mappings

Mehdi Omidvari, S. M. Vaezpour, and Reza Saadati



BEST PROXIMITY POINT THEOREMS FOR F -CONTRACTIVE NON-SELF MAPPINGS

M. OMIDVARI, S. M. VAEZPOUR, AND R. SAADATI

Received 14 October, 2013

Abstract. In this article, we prove the existence of a best proximity point for F -contractive non-self mappings and state some results in the complete metric spaces. Also we define two kinds of F -proximal contraction and extend some best proximity theorems and improve the recent results.

2010 Mathematics Subject Classification: 46N40; 47H10; 54H25; 46T99

Keywords: metric space, best proximity point, fixed point, F -contractive non-self mapping, F -proximal contraction

1. INTRODUCTION

Let us assume that A, B be two nonempty subsets of a metric space (X, d) and $T : A \rightarrow B$. Clearly $T(A) \cap A \neq \emptyset$ is a necessary condition for the existence of a fixed point of T . Now if $T(A) \cap A = \emptyset$ then to find an element $x \in A$ such that $d(x, Tx) = d(A, B)$ which called best proximity point is the idea of best proximity point theorems. In the other words, we determine an approximate solution x such that the error of equation $d(x, Tx) = 0$ is minimum. Several authors such as Prolla [2], Reich [4], Sehgal and Singh [6, 7], Vertivel, Veermani and Bhattacharyya [8] and others [1] generalized and extended the best proximity point theorems in many directions. In 2011, Sadiq Basha [5] stated the best proximity points theorems for proximal contractions. On the other hand, Wardowski [9] introduced a new type of contraction which called F -contraction and proved a fixed point result in complete metric spaces. In this paper, by using Wardowski's contraction, we prove the existence of a best proximity point. Moreover we define F -proximal contractions of the first and second kind and establish the best proximity point theorems in spire of Wardowski's contraction.

2. PRELIMINARY

Let A, B be two nonempty subsets of a metric space X . The following notations will be used throughout this paper:

$$d(y, A) := \inf\{d(x, y) : x \in A\},$$

$$d(A, B) := \inf\{d(x, y) : x \in A \text{ and } y \in B\},$$

$$A_0 := \{x \in A : d(x, y) = d(A, B) \text{ for some } y \in B\},$$

$$B_0 := \{y \in B : d(x, y) = d(A, B) \text{ for some } x \in A\}.$$

We recall that $x \in A$ is a best proximity point of the mapping $T : A \rightarrow B$ if $d(x, Tx) = d(A, B)$. It can be observed that a best proximity reduces to a fixed point if the underlying mapping is a self-mapping.

Definition 1 ([3]). Let (A, B) be a pair of nonempty subsets of a metric space X with $A \neq \emptyset$. Then the pair (A, B) is said to have the P-property if and only if

$$\left. \begin{array}{l} d(x_1, y_1) = d(A, B) \\ d(x_2, y_2) = d(A, B) \end{array} \right\} \implies d(x_1, x_2) = d(y_1, y_2)$$

where $x_1, x_2 \in A_0$ and $y_1, y_2 \in B_0$.

It is clear that, for any nonempty subset A of X , the pair (A, A) has the P-property.

Definition 2 ([5]). A is said to be approximatively compact with respect to B if every sequence $\{x_n\}$ of A satisfying the condition that $d(y, x_n) \rightarrow d(y, A)$ for some y in B has a convergent subsequence.

It is easy to see that every set is approximatively compact with respect to itself.

Definition 3 ([5]). Given $T : A \rightarrow B$ and an isometry $g : A \rightarrow A$, the mapping T is said to preserve isometric distance with respect to g if

$$d(Tgx_1, Tgx_2) = d(Tx_1, Tx_2)$$

for all x_1 and x_2 in A .

Recently Wardowski [9] defined the following contraction which was called F -contraction.

Definition 4. Let $F : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a mapping satisfying:

(F1) F is strictly increasing, i.e. for all $a, b \in \mathbb{R}_+$ such that

$$\alpha < \beta \implies F(\alpha) < F(\beta);$$

(F2) For each sequence $\{\alpha_n\}_{n \in \mathbb{N}}$ of positive numbers

$$\lim_{n \rightarrow \infty} \alpha_n = 0 \text{ if and only if } \lim_{n \rightarrow \infty} F(\alpha_n) = -\infty;$$

(F3) There exists $k \in (0, 1)$ such that $\lim_{\alpha \rightarrow 0^+} \alpha^k F(\alpha) = 0$.

A mapping $T : X \rightarrow X$ is said to be an F -contraction if there exists $\tau > 0$ such that

$$\text{For all } x, y \in X, (d(Tx, Ty)) > 0 \implies \tau + F(d(Tx, Ty)) \leq F(d(x, y)). \quad (2.1)$$

Now, we consider the mapping F in the different types, and we obtain the variety of contractions.

Example 1 ([9]). Let $F : \mathbb{R}_+ \rightarrow \mathbb{R}$ be given by the formula $F(\alpha) = \ln \alpha$. It is clear that F satisfies the (F1)-(F3) ((F3) for any $k \in (0, 1)$) conditions of Definition 4. Each mapping $T : X \rightarrow X$ satisfying (2.1) is an F -contraction such that

$$d(Tx, Ty) \leq e^{-\tau} d(x, y), \quad \text{for all } x, y \in X, Tx \neq Ty.$$

It is clear that for $x, y \in X$ such that $Tx = Ty$ the inequality $d(Tx, Ty) \leq e^{-\tau} d(x, y)$ also holds, i.e. T is a Banach contraction.

3. MAIN RESULTS

Now, let us state our main result.

Theorem 1. *Let A and B be non-empty, closed subsets of a complete metric space X such that A_0 is nonempty. Let $T : A \rightarrow B$ be an F -contraction non-self mapping such that $T(A_0) \subseteq B_0$. Assume that the pair (A, B) has the P-property. Then there exists a unique x^* in A such that $d(x^*, Tx^*) = d(A, B)$.*

Proof. Choose $x_0 \in A_0$. Since $Tx_0 \in T(A_0) \subseteq B_0$, there exists $x_1 \in A_0$ such that $d(x_1, Tx_0) = d(A, B)$. Again, since $Tx_1 \in T(A_0) \subseteq B_0$, there exists $x_2 \in A_0$ such that $d(x_2, Tx_1) = d(A, B)$. Continuing this process, we can find a sequence $\{x_n\}$ in A_0 such that

$$d(x_{n+1}, Tx_n) = d(A, B), \quad \text{for all } n \in \mathbb{N}. \quad (3.1)$$

(A, B) satisfies the P-property, therefore from (3.1) we obtain

$$d(x_n, x_{n+1}) = d(Tx_{n-1}, Tx_n), \quad \text{for all } n \in \mathbb{N}. \quad (3.2)$$

We will prove that the sequence $\{x_n\}$ is convergent in A_0 . If there exists $n_0 \in \mathbb{N}$ such that $d(Tx_{n_0-1}, Tx_{n_0}) = 0$, then by (3.2) we have $d(x_{n_0}, x_{n_0+1}) = 0$ that implies $x_{n_0} = x_{n_0+1}$. Therefore

$$Tx_{n_0} = Tx_{n_0+1} \implies d(Tx_{n_0}, Tx_{n_0+1}) = 0 \quad (3.3)$$

From (3.2) and (3.3) we receive that

$$d(x_{n_0+2}, x_{n_0+1}) = d(Tx_{n_0+1}, Tx_{n_0}) = 0 \implies x_{n_0+2} = x_{n_0+1}$$

Therefore $x_n = x_{n_0}$, for all $n \geq n_0$ and $\{x_n\}$ is convergent in A_0 .

Now let $d(Tx_{n-1}, Tx_n) \neq 0$, for all $n \in \mathbb{N}$. T is a F -contraction and (3.2) holds, hence for any positive integer n we have

$$\tau + F(d(Tx_n, Tx_{n-1})) \leq F(d(x_n, x_{n-1}))$$

$$\begin{aligned} \implies F(d(x_{n+1}, x_n)) &\leq F(d(x_n, x_{n-1})) - \tau \\ &\vdots \\ &\leq F(d(x_1, x_0)) - n\tau. \end{aligned} \quad (3.4)$$

Put $\beta_n := d(x_{n+1}, x_n)$.

From (3.4), we obtain $\lim_{n \rightarrow \infty} F(\beta_n) = -\infty$ that together with (F2) gives

$$\lim_{n \rightarrow \infty} \beta_n = 0 \quad (3.5)$$

Also from (F3) we have

$$\exists k \in (0, 1) \text{ such that } \beta_n^k F(\beta_n) = 0 \quad (3.6)$$

By (3.3), the following holds for all $n \in \mathbb{N}$:

$$F(\beta_n) - F(\beta_0) \leq -n\tau.$$

Therefore

$$\beta_n^k F(\beta_n) - \beta_n^k F(\beta_0) \leq -n\beta_n^k \tau \leq 0.$$

Letting $k \rightarrow \infty$ in the above inequality and using (3.5), (3.6), we obtain

$$\lim_{n \rightarrow \infty} n\beta_n^k = 0.$$

Hence there exists $n_1 \in \mathbb{N}$ such that $n\beta_n^k \leq 1$ for all $n \geq n_1$. Therefore for any $n \geq n_1$,

$$\beta_n \leq \frac{1}{n^{\frac{1}{k}}}. \quad (3.7)$$

This means that series $\sum_{i=1}^{\infty} \beta_i$ is convergent.

Now let $m \geq n \geq n_1$. By the triangular inequality and (3.7), we have

$$d(x_m, x_n) \leq \beta_{m-1} + \beta_{m-2} + \dots + \beta_n \leq \sum_{i=n}^{\infty} \beta_i.$$

Therefore $\{x_n\}$ is a Cauchy sequence in A . Since (X, d) is complete and A is a closed subset of X , there exist $x^* \in A$ such that

$$\lim_{n \rightarrow \infty} x_n = x^*$$

Since T is continuous, we have $Tx_n \rightarrow Tx^*$. Hence continuity of the metric function d which implies that $d(x_{n+1}, Tx_n) \rightarrow d(x^*, Tx^*)$. From (3.1), $d(x^*, Tx^*) = d(A, B)$. So we show that x^* is a best proximity of T .

The uniqueness of the best proximity point follows from the condition that T is F -contraction. That is, suppose that

$$x_1, x_2 \in A \text{ such that } x_1 \neq x_2 \text{ and } d(x_1, Tx_1) = d(x_2, Tx_2) = d(A, B).$$

Then by the P-property of (A, B) , we have $d(x_1, x_2) = d(Tx_1, Tx_2)$. Also

$$x_1 \neq x_2 \implies d(x_1, x_2) \neq 0.$$

Therefore

$$F(d(x_1, x_2)) = F(d(Tx_1, Tx_2)) \leq F(d(x_1, x_2)) - \tau < F(d(x_1, x_2)),$$

which is a contraction. Hence the best proximity point is unique. \square

The following result is a special case of Theorem 1, obtained by setting $A = B$.

Corollary 1. *Let (X, d) be a complete metric space and A be a nonempty closed subset of X . Let $T : A \rightarrow A$ be a F -contractive self-map. Then T has a unique fixed point x^* in A .*

The next result is an immediate consequence of Theorem 1 by taking $F(\alpha) = \ln \alpha$.

Corollary 2 (Banach Contraction Principle). *Let (X, d) be a complete metric space and A be a nonempty closed subset of X . Let $T : A \rightarrow A$ be a contractive self-map. Then T has a unique fixed point x^* in A .*

Let F be the function as in Definition 4, we define the proximal contractions.

Definition 5. A mapping $T : A \rightarrow B$ is said to be a F -proximal contraction of the first kind if there exists a $\tau > 0$ such that

$$\left. \begin{array}{l} d(u_1, Tx_1) = d(A, B) \\ d(u_2, Tx_2) = d(A, B) \\ d(u_1, u_2), d(x_1, x_2) > 0 \end{array} \right\} \implies \tau + F(d(u_1, u_2)) \leq F(d(x_1, x_2))$$

where $u_1, u_2, x_1, x_2 \in A$.

Remark 1. If $T : A \rightarrow B$ is a F -proximal contraction of the first kind and (A, B) has the P-property then T is a F -contractive non-self mapping.

Definition 6. A mapping $T : A \rightarrow B$ is said to be a F -proximal contraction of the second kind if there exists a $\tau > 0$ such that

$$\left. \begin{array}{l} d(u_1, Tx_1) = d(A, B) \\ d(u_2, Tx_2) = d(A, B) \\ d(Tu_1, Tu_2), d(Tx_1, Tx_2) > 0 \end{array} \right\} \implies \tau + F(d(Tu_1, Tu_2)) \leq F(d(Tx_1, Tx_2))$$

where $u_1, u_2, x_1, x_2 \in A$.

The following theorem is a best proximity point theorem for non-self F -proximal contraction of the first kind.

Theorem 2. *Let A and B be non-empty, closed subsets of a complete metric space X such that A_0 is non-empty. Let $T : A \rightarrow B$ and $g : A \rightarrow A$ satisfy the following conditions:*

- (a) T is a continuous F -proximal contraction of the first kind.
- (b) g is an isometry.
- (c) $T(A_0) \subseteq B_0$.

(d) $A_0 \subseteq g(A_0)$.

Then, there exists a unique element $x \in A$ such that

$$d(gx, Tx) = d(A, B).$$

Proof. Choose $x_0 \in A_0$. Since $Tx_0 \in T(A_0) \subseteq B_0$ and $A_0 \subseteq g(A_0)$, there exists $x_1 \in A_0$ such that $d(gx_1, Tx_0) = d(A, B)$. If $x_0 = x_1$ then put $x_n := x_1$ for all $n \geq 2$. Also, since $Tx_1 \in T(A_0) \subseteq B_0$ and $A_0 \subseteq g(A_0)$, there exists $x_2 \in A_0$ such that $d(gx_2, Tx_1) = d(A, B)$. If $x_1 = x_2$ then put $x_n := x_2$ for all $n \geq 3$. Continuing this process, we can find a sequence $\{x_n\}$ in A_0 such that

$$d(gx_{n+1}, Tx_n) = d(A, B), \quad \text{for all } n \in \mathbb{N}. \quad (3.8)$$

We will prove the convergence of the sequence $\{x_n\}$ in A . If there exists $n_0 \in \mathbb{N}$ such that $d(gx_{n_0}, gx_{n_0+1}) = 0$ then it is clear that the sequence $\{x_n\}$ is convergent. Hence let $d(gx_n, gx_{n+1}) \neq 0$, for all $n \in \mathbb{N}$. T is a F -proximal contraction of the first kind and (3.8) holds, hence for any positive integer n we have

$$\begin{aligned} \tau + F(d(gx_n, gx_{n+1})) &\leq F(d(x_{n-1}, x_n)) \\ \implies F(d(x_n, x_{n+1})) &\leq F(d(x_{n-1}, x_n)) - \tau \\ &\vdots \\ &\leq F(d(x_0, x_1)) - n\tau. \end{aligned} \quad (3.9)$$

Similarly as the process in the proof of Theorem 1, $\{x_n\}$ is a Cauchy sequence in A . Since X is complete metric space and A is closed subset of X , there exists $x \in A$ such that $\lim_{n \rightarrow \infty} x_n = x$.

Since, T , g and d are continuous, therefore with letting $n \rightarrow \infty$ in (3.8), we obtain

$$d(gx, Tx) = d(A, B).$$

Now, x^* be in A such that

$$d(gx^*, Tx^*) = d(A, B).$$

We show that $x = x^*$. Suppose to the contrary, that $x \neq x^*$. Hence $d(x, x^*) \neq 0$. Since T is a F -proximal contraction of the first kind and g is an isometry,

$$F(d(x, x^*)) = F(d(gx, gx^*)) \leq F(d(x, x^*)) - \tau < F(d(x, x^*)),$$

which is a contraction. Therefore $x = x^*$ and this completes the proof of theorem. \square

The following result is a special case of Theorem 2, if g is the identity mapping.

Corollary 3. Let A and B be non-empty, closed subsets of a complete metric space X such that A is approximatively compact with respect to B . Further, suppose that A_0 is non-empty. Let $T : A \rightarrow B$ satisfies the following conditions:

- (a) T is a continuous F -proximal contraction of the first kind.
- (b) $T(A_0) \subseteq B_0$.

Then T has a unique best proximity point in A .

The following theorem is a best proximity point theorem for non-self F -proximal contraction of the second kind.

Theorem 3. Let A and B be non-empty, closed subsets of a complete metric space X such that A is approximatively compact with respect to B . Further, suppose that A_0 is non-empty. Let $T : A \rightarrow B$ and $g : A \rightarrow A$ satisfy the following conditions:

- (a) T is a continuous F -proximal contraction of the second kind.
- (b) g is an isometry.
- (c) $T(A_0) \subseteq B_0$.
- (d) $A_0 \subseteq g(A_0)$.
- (e) T preserves isometric distance with respect to g .

Then, there exists an element $x \in A$ such that

$$d(gx, Tx) = d(A, B).$$

Moreover, if x^* is another element of A such that $d(gx^*, Tx^*) = d(A, B)$ then $Tx = Tx^*$.

Proof. Choose $x_0 \in A_0$. Since $Tx_0 \in T(A_0) \subseteq B_0$ and $A_0 \subseteq g(A_0)$, there exists $x_1 \in A_0$ such that $d(gx_1, Tx_0) = d(A, B)$. If $Tx_0 = Tx_1$ then put $x_n := x_1$ for all $n \geq 2$. Otherwise again since $Tx_1 \in T(A_0) \subseteq B_0$ and $A_0 \subseteq g(A_0)$, there exists $x_2 \in A_0$ such that $d(gx_2, Tx_1) = d(A, B)$. If $Tx_1 = Tx_2$ then put $x_n := x_2$ for all $n \geq 3$. Continuing this process, we can find a sequence $\{x_n\}$ in A_0 such that

$$d(gx_{n+1}, Tx_n) = d(A, B), \quad \text{for all } n \in \mathbb{N}. \tag{3.10}$$

We will prove the convergence of the sequence $\{Tx_n\}$ in B . If there exists $n_0 \in \mathbb{N}$ such that $d(Tgx_{n_0}, Tgx_{n_0+1}) = 0$ then it is clear that the sequence $\{Tx_n\}$ is convergent. Hence let $d(Tgx_n, Tgx_{n+1}) \neq 0$, for all $n \in \mathbb{N}$. T is a F -proximal contraction of the second kind, T preserves isometric distance with respect to g and (3.10) holds, hence for any positive integer n we have

$$\begin{aligned} \tau + F(d(Tgx_n, Tgx_{n+1})) &\leq F(d(Tx_{n-1}, Tx_n)) \\ \implies F(d(Tx_n, Tx_{n+1})) &\leq F(d(Tx_{n-1}, Tx_n)) - \tau \\ &\vdots \\ &\leq F(d(Tx_0, Tx_1)) - n\tau. \end{aligned} \tag{3.11}$$

Similarly as the process in the proof of Theorem 1, we receive that $\{Tx_n\}$ is a Cauchy sequence in B . Since X is complete metric space and B is closed subset of X , there exists $y \in B$ such that $\lim_{n \rightarrow \infty} Tx_n = y$.

By the triangular inequality, we have,

$$\begin{aligned} d(y, A) &\leq d(y, gx_n) \leq d(y, Tx_{n-1}) + d(Tx_{n-1}, gx_n) \\ &= d(y, Tx_{n-1}) + d(A, B) \\ &\leq d(y, Tx_{n-1}) + d(y, A) \end{aligned}$$

Letting $k \rightarrow \infty$ in the above inequality, we obtain

$$\lim_{n \rightarrow \infty} d(y, gx_n) = d(y, A).$$

Since A is approximatively compact with respect to B , there exists a subsequence $\{gx_{n_k}\}$ of $\{gx_n\}$ such that converging to some $z \in A$. Therefore

$$d(z, y) = \lim_{k \rightarrow \infty} d(gx_{n_k}, Tx_{n_k-1}) = d(A, B).$$

This implies that $z \in A_0$. Since $A_0 \subseteq g(A_0)$, there exists $x \in A_0$ such that $z = gx$. As $\lim_{n \rightarrow \infty} g(x_{n_k}) = g(x)$ and g is an isometry, we have

$$\lim_{n \rightarrow \infty} x_{n_k} = x.$$

T is continuous and $\{Tx_n\}$ is convergent to y , Therefore

$$\lim_{n \rightarrow \infty} Tx_{n_k} = Tx = y.$$

Thus, it follows that

$$d(gx, Tx) = \lim_{n \rightarrow \infty} d(gx_{n_k}, Tx_{n_k}) = d(A, B).$$

Now let x^* be another element in A such that

$$d(gx^*, Tx^*) = d(A, B).$$

We will show that $Tx = Tx^*$. Suppose to the contrary, that $Tx \neq Tx^*$. Hence $d(Tx, Tx^*) \neq 0$. Since T preserves isometric distance with respect to g and T is a F -proximal contraction of the second kind,

$$F(d(Tx, Tx^*)) = F(d(Tgx, Tgx^*)) \leq F(d(Tx, Tx^*)) - \tau < F(d(Tx, Tx^*)),$$

which is a contraction. Therefore $Tx = Tx^*$. \square

The next result is an immediate consequence of the Theorem 3, if g is the identity mapping.

Corollary 4. *Let A and B be non-empty, closed subsets of a complete metric space X such that A is approximatively compact with respect to B . Further, suppose that A_0 is non-empty. Let $T : A \rightarrow B$ satisfies the following conditions:*

- (a) T is a continuous F -proximal contraction of the second kind.
- (b) $T(A_0) \subseteq B_0$.

Then, T has a best proximity point in A . Moreover, if x^ is another best proximity point of T then $Tx = Tx^*$.*

ACKNOWLEDGMENTS

The authors would like to thank referee for giving useful comments and suggestions for the improvement of this paper.

REFERENCES

- [1] M. R. Haddadi, "Best proximity point iteration for nonexpensive mapping in banach spaces," *J. Nonlinear Sci. Appl.*, vol. 7, no. 2, pp. 126–130, 2014.
- [2] J. B. Prolla, "Fixed-point theorems for set-valued mappings and existence of best approximants," *Numer. Funct. Anal. Optim.*, vol. 5, no. 4, pp. 449–455, 1982–1983.
- [3] V. S. Raj, "A best proximity point theorem for weakly contractive non-self-mappings," *Nonlinear Anal.*, vol. 74, no. 11, pp. 4804–4808, 2011.
- [4] S. Reich, "Approximate selections, best approximations, fixed points and invariant sets," *J. Math. Anal. Appl.*, vol. 62, no. 1, pp. 104–113, 1978.
- [5] S. Sadiq Basha, "Best proximity point theorems," *J. Approx. Theory*, vol. 163, no. 11, pp. 1772–1781, 2011.
- [6] V. M. Sehgal and S. P. Singh, "A generalization to multifunctions of fan's best approximation theorem," *Proc. Amer. Math. Soc.*, vol. 102, no. 3, pp. 534–537, 1988.
- [7] V. M. Sehgal and S. P. Singh, "A theorem on best approximations," *Numer. Funct. Anal. Optim.*, vol. 10, no. 1–2, pp. 181–184, 1989.
- [8] V. Vetrivel, P. Veeramani, and P. Bhattacharyya, "Some extensions of fan's best approximation theorem," *Numer. Funct. Anal. Optim.*, vol. 13, no. 3–4, pp. 397–402, 1992.
- [9] D. Wardowski, "Fixed points of a new type of contractive mappings in complete metric spaces," *Fixed Point Theory Appl.*, vol. 2012, no. 94, p. 6, 2012.

*Authors' addresses***M. Omidvari**

Department of Mathematics, Science and Research Branch, Islamic Azad University (IAU), Tehran, Iran

E-mail address: mehdi.omidvari@gmail.com

S. M. Vaezpour

Department of Mathematics and Computer Science, Amirkabir University of Technology, Hafez Ave., P. O. Box 15914, Tehran, Iran

E-mail address: vaez@aut.ac.ir

R. Saadati

Department of Mathematics, Iran University of Science and Technology, Tehran, Iran

E-mail address: rsaadati@iust.ac.ir