

TVERBERG PLUS MINUS

IMRE BÁRÁNY AND PABLO SOBERÓN

ABSTRACT. We prove a Tverberg type theorem: Given a set $A \subset \mathbb{R}^d$ in general position with $|A| = (r-1)(d+1) + 1$ and $k \in \{0, 1, \dots, r-1\}$, there is a partition of A into r sets A_1, \dots, A_r (where $|A_j| \leq d+1$ for each j) with the following property. There is a unique $z \in \bigcap_{j=1}^r \text{aff } A_j$ and it can be written as an affine combination of the element in A_j : $z = \sum_{x \in A_j} \alpha(x)x$ for every j and exactly k of the coefficients $\alpha(x)$ are negative. The case $k = 0$ is Tverberg's classical theorem.

1. INTRODUCTION AND MAIN RESULT

Assume $A = \{a_1, \dots, a_n\} \subset \mathbb{R}^d$ where $n = (r-1)(d+1) + 1$ and $r \geq 2$, $d \geq 1$ are integers. Suppose further that the coordinates of the a_i (altogether dn real numbers) are algebraically independent. A partition $\mathcal{A} = \{A_1, \dots, A_r\}$ of A is called *proper* if $1 \leq |A_j| \leq d+1$ for every $j \in [r]$. Here and in what follows $[r]$ stands for the set $\{1, \dots, r\}$. We will show later (Proposition 1.1) that in this case the intersection of the affine hull of the A_j s is a single point z , that is, $\{z\} = \bigcap_{j=1}^r \text{aff } A_j$. Equivalently, the following system of linear equations has a unique solution:

$$(1.1) \quad z = \sum_{x \in A_j} \alpha(x)x \quad \text{and} \quad 1 = \sum_{x \in A_j} \alpha(x) \quad \text{for all } j \in [r].$$

One form of Tverberg's classical theorem [Tve66] puts extra conditions on the coefficients $\alpha(x)$ (consult [Mat03] and the references therein for an introduction to the subject).

Theorem 1.1 (Tverberg's theorem). *Under the above conditions there is a proper partition of A into sets A_1, \dots, A_r such that $\alpha(x) \geq 0$ for all elements $x \in A$. In other words, $\{z\} = \bigcap_{j=1}^r \text{conv } A_j$.*

This means that the unique solution to (1.1) has $\alpha(x) > 0$ for all $x \in A$. Can we require here that exactly one (or two or more) of the $\alpha(x)$ are negative? A partial answer comes from the following theorem, which is the main result of this paper.

2000 *Mathematics Subject Classification.* Primary 52A37, secondary 52B05.

Key words and phrases. Tverberg's theorem, sign conditions.

Theorem 1.2. *Assume $k \in \{0, 1, \dots, r-1\}$. Under the conditions of Theorem 1.1 there is a (proper) partition of A into r parts so that in the unique solution to (1.1) $\alpha(x) < 0$ for exactly k elements $x \in A$.*

Of course the same holds for any set A of n points in \mathbb{R}^d , we only have to relax the condition $\alpha(x) < 0$ to $\alpha(x) \leq 0$ for k elements $x \in A$ and $\alpha(x) \geq 0$ for the rest. Actually, the general position condition (on A) is used in order to avoid cases when $\alpha(x) = 0$ for some $x \in A$.

It is not clear for what other values of k , $k \in [n]$, the theorem holds. Since the sum of the coefficients for each A_j is one, at least one is positive. This implies the upper bound $k \leq n - r$.

The case $d = 1$ is very simple. Then $n = 2r - 1$ and there is no r -partition with r or more negative coefficients, so the trivial bound $k \leq n - r = r - 1$ is tight. In the case $d = 2$, $r = 3$ and $n = 7$ Theorem 1.2 gives a suitable partition for $k = 0, 1, 2$. A careful case analysis shows that the statement holds for $k = 3$ as well, and an extensive computer aided search did not find any example where it fails to hold for $k = 4$.

The case of $r = 2$, that is, Radon (plus minus) partitions can be checked directly. Then $|A| = d + 2$ and the outcome is that for any $k \in \{0, 1, \dots, \lfloor \frac{d+2}{2} \rfloor\}$ there is a partition with exactly k negative $\alpha(x)$. Further, there are examples showing that this does not hold for $k > \lfloor \frac{d+2}{2} \rfloor$. In this case everything is governed by the unique affine dependence of the vectors in A , just as in the proof of Radon's theorem. We omit the details.

We will see in Corollary 3.1 in Section 3 that, for a strange reason, if both d and r are even, then Theorem 1.2 holds with $k = \frac{1}{2}[(r-1)(d+1) + 1]$ as well. This makes us wonder if Theorem 1.2 holds for all integers $k \leq \frac{1}{2}[(r-1)(d+1) + 1]$.

We are going to prove Theorem 1.2 in a stronger form: to some extent we can prescribe the subset of A where the coefficients in (1.1) are negative.

Theorem 1.3. *Under the conditions of Theorem 1.1 let $M \subset A$ be a set of size at most $r-1$ such that $\text{conv } M \cap \text{conv } (A \setminus M) = \emptyset$. Then there is a partition $\mathcal{A} = \{A_1, \dots, A_r\}$ of A such that in (1.1) $\alpha(x) < 0$ if and only if $x \in M$.*

We prove this theorem in Section 3 where we state a slightly stronger result whose proof is in Section 6. Examples showing the necessity of the condition on M are given in Section 2. In Section 4 we discuss coloured variations of Theorem 1.2. In Section 5 we prove the following fact.

Proposition 1.1. *Assume $A = \{a_1, \dots, a_n\} \subset \mathbb{R}^d$, the coordinates of the a_i are algebraically independent and $r \geq 2$, $d \geq 1$ are integers. If the partition $\mathcal{A} = \{A_1, \dots, A_r\}$ of A is proper and $n = (r-1)(d+1)+1$, then $\bigcap_{j=1}^r \text{aff } A_j$ is a single point. If $n \leq (r-1)(d+1)$, then $\bigcap_{j=1}^r \text{aff } A_j = \emptyset$.*

The last statement holds even if the partition is not proper. The first part must be known, see for instance [PS14] or [DV77] for similar statements. The second part is proved in [Tve66]. We give a simple proof in Section 5.

2. THE CONDITION ON M

The condition on M in Theorem 1.3 simply says that M and $A \setminus M$ can be separated by a hyperplane.

Example 1. We give an example showing the necessity of this condition. Let $V = \{v_1, \dots, v_{d+1}\}$ be the set of vertices of a regular simplex Δ , and let c be the centre of Δ and write F_h for its facet opposite to v_h . For every $h \in [d+1]$ let $U_h \subset v_h + \varepsilon B$ be an $(r-1)$ -element set. Here $\varepsilon > 0$ is small and B is the Euclidean unit ball in \mathbb{R}^d centred at the origin. Define $A = \{c\} \cup \bigcup_{h=1}^{d+1} U_h$. We assume A is in general position which can be clearly reached by choosing the sets U_h suitably. Set $M = \{c\}$ so the separation condition fails. We **claim** that there is no proper r -partition of A such that in (1.1) only $\alpha(c)$ is negative, $\alpha(x) > 0$ for all other $x \in A$.

Assume the contrary and let A_1, \dots, A_r be a proper partition with $z \in \bigcap_{j=1}^r \text{aff } A_j$ so that $\alpha(c) < 0$ and $\alpha(x) > 0$ for all other $x \in A$ in (1.1).

Given a convex compact set C in \mathbb{R}^d and a point $u \in \mathbb{R}^d \setminus C$ we let $S(u, C)$ denote the *shadow* of C from u which is the set of point $\{tu + (1-t)c : c \in C, t \leq 0\}$, see Figure 1.




FIGURE 1. Construction of $S(u, C)$. Notice that $C \subset S(u, C)$.

For every $h \in [d+1]$ there is a $j = j(h) \in [r]$ such that $A_{j(h)}$ and U_h are disjoint, simply because each U_h has $r-1$ elements and the number of sets A_j is r . It follows that $A_{j(h)} \subset F_h + \varepsilon B \subset S(c, F_h + \varepsilon B)$ if $c \notin A_{j(h)}$, and then $z \in \text{conv } A_{j(h)} \subset S(c, F_h + \varepsilon B)$. If $c \in A_{j(h)}$, then in the equation $z = \sum_{x \in A_{j(h)}} \alpha(x)x$ only the coefficient $\alpha(c)$ is negative, so $z \in S(c, F_h + \varepsilon B)$. Therefore

$$z \in \bigcap_{h=1}^{d+1} S(c, F_h + \varepsilon B).$$

However, $\bigcap_{h=1}^{d+1} S(c, F_h + \varepsilon B) = \emptyset$ as long as $\varepsilon < \frac{\text{diam}\Delta}{2d}$. See Figure 2 for an illustration. This follows from the fact that the shadows $S(c, F_h + \varepsilon B)$ for $h \in [d+1]$ are convex and their union covers the boundary of Δ . If they had a point in common, then their union would cover Δ . Since none of them contains c , this is impossible. This gives us the contradiction we were seeking.




FIGURE 2. The shadows of the sides of a triangle from its centre do not intersect.

The proof above does not use the fact that c is the only point near the centre of the simplex. If we consider $U_0 \subset c + \varepsilon B$ any set and declare $U_0 = M$, the same arguments as above show that there is no partition with the desired properties. Therefore, the condition $\text{conv } M \cap \text{conv } (A \setminus M) = \emptyset$ cannot be removed even if we allow A to have more than $(r-1)(d+1) + 1$ points.

Example 2. This example shows a construction where M satisfies the separation condition, $|M| = r$ and the conclusion of the theorem fails. We work with the same simplex Δ and $U_h \subset v_h + \varepsilon B$ is the same $(r-1)$ -element set as before for $h \in \{2, \dots, d+1\}$ but for $h=1$ it is an r -element set in $v_1 + \varepsilon B$. This time $A = \bigcup_{h=1}^{d+1} U_h$ and $M = U_1$. We assume of course that A is in general position. Now M is separated from $A \setminus M$ and has exactly r elements.

We **claim** that A has no partition into r parts with the required properties.

Proof. Assume the contrary and let A_1, \dots, A_r be a proper partition with $z \in \bigcap_{j=1}^r \text{aff } A_j$ such that in (1.1) $\alpha(x) < 0$ if $x \in M$ and $\alpha(x) > 0$ if $x \notin M$. For $h \in \{2, \dots, d+1\}$ let G_h be the convex hull of $V \setminus \{v_1, v_h\}$; this is a $(d-2)$ -face of Δ . Set $\beta = \sum_{x \in A_j \cap U_1} \alpha(x)$, and note that $\beta > 0$

if $A_j \cap U_1$ is nonempty. Define $u_j = \frac{1}{\beta} \sum_{x \in A_j \cap U_1} \alpha(x)x$ if $A_j \cap U_1$ is nonempty, and $u_j = v_1$ otherwise.

Note again that for each $h \in \{2, \dots, d+1\}$ there is a $j(h) \in [r]$ such that U_h and $A_{j(h)}$ are disjoint. Then $A_{j(h)} \setminus U_1 \subset G_h + \varepsilon B$. This and the sign condition in (1.1) imply that for every $h > 1$ with $A_{j(h)} \cap U_1 \neq \emptyset$,

$$z \in S(u_{j(h)}, G_h + \varepsilon B).$$

This holds even if $A_{j(h)} \cap U_1 = \emptyset$ since then $u_{j(h)} = v_1$ and $z \in \text{conv } A_{j(h)} \subset G_h + \varepsilon B \subset S(v_1, G_h + \varepsilon B)$. Thus

$$z \in \bigcap_{h=2}^{d+1} S(u_{j(h)}, G_h + \varepsilon B).$$

But again, the shadows on the right hand side have no point in common, as one can check easily. \square

3. PROOF OF THEOREM 1.3

We are going to use the colourful Carathéodory theorem [Bár82]. It says that given sets $S_1, \dots, S_{n+1} \subset \mathbb{R}^n$ with the condition that $0 \in \bigcap_{i=1}^{n+1} \text{conv } S_i$, there is a transversal, that is, a choice $s_i \in S_i$ for every $i \in [n+1]$, such that $0 \in \text{conv } \{s_1, \dots, s_{n+1}\}$.

Proof of Theorem 1.3. We use a modification of Sarkaria's argument [Sar92], in the form given by [BO97]. It starts with an artificial tool: let v_1, \dots, v_r be the vertices of a regular simplex in \mathbb{R}^{r-1} centred at the origin. The important property here is that, apart from scalar multiples, their unique linear dependence is $v_1 + \dots + v_r = 0$.

Assume $A = \{a_1, \dots, a_n\}$ where $n = (r-1)(d+1) + 1$. Recall that $M \subset A$, $|M| = k < r$, and M and $A \setminus M$ are separated by a hyperplane. Define

$$b_i = (a_i, 1) \text{ if } a_i \notin M \text{ and } b_i = (-a_i, -1) \text{ if } a_i \in M.$$

where $(a_i, 1) \in \mathbb{R}^{d+1}$ is vector a_i appended with an $(d+1)$ -th coordinate equal to one, and similarly for $(-a_i, -1)$. For $i \in [n]$ we set

$$S_i = \{v_1 \otimes b_i, v_2 \otimes b_i, \dots, v_r \otimes b_i\}.$$

Here $v_j \otimes b_i$ is the usual tensor product, which is the same as the matrix product of the $(r-1)$ -dimensional column vector v_j and the $(d+1)$ -dimensional row vector b_i^T : $v_j b_i^T$, where we consider our vectors as vertical matrices. So this product is an $(r-1) \times (d+1)$ matrix, or equivalently a vector in \mathbb{R}^{n-1} . Observe that $0 \in \text{conv } S_i$ for every i , so the colourful Carathéodory theorem applies and gives a transversal s_1, \dots, s_n whose convex hull contains the origin, that is, there are non-negative coefficients β_1, \dots, β_n whose sum is 1 such that $\sum_{i=1}^n \beta_i s_i = 0$. Here each s_i is of the form $v_j \otimes b_i$ for a unique $j = j(i) \in [r]$. We define

$A_j = \{a_i \in A : j(i) = j\}$ for all $j \in [r]$. The sets A_1, \dots, A_r form an r -partition of A . With the new notation

$$\begin{aligned} 0 &= \sum_{i=1}^n \beta_i s_i = \sum_{i=1}^n \beta_i v_{j(i)} \otimes b_i \\ &= \sum_{j=1}^r \sum_{a_i \in A_j} \beta_i v_j \otimes b_i = \sum_{j=1}^r v_j \otimes \left(\sum_{a_i \in A_j} \beta_i b_i \right). \end{aligned}$$

Define now $\alpha_i = -\beta_i$ if $a_i \in M$ and $\alpha_i = \beta_i$ otherwise. The last equation becomes

$$(3.1) \quad 0 = \sum_{j=1}^r v_j \otimes \left(\sum_{a_i \in A_j} \alpha_i(a_i, 1) \right).$$

There is a vector $u \in \mathbb{R}^{r-1}$, orthogonal to v_3, v_4, \dots, v_r with $\langle u, v_1 \rangle = 1$, where $\langle \cdot, \cdot \rangle$ denotes the dot product. The condition $v_1 + \dots + v_r = 0$ implies that $\langle u, v_2 \rangle = -1$. As (3.1) is a matrix equation, multiplying it from the left by the $(r-1)$ -dimensional row vector u^T gives $\sum_{a_i \in A_1} \alpha_i(a_i, 1) = \sum_{a_i \in A_2} \alpha_i(a_i, 1)$. By symmetry we have

$$(3.2) \quad z := \sum_{a_i \in A_1} \alpha_i(a_i, 1) = \sum_{a_i \in A_2} \alpha_i(a_i, 1) = \dots = \sum_{a_i \in A_r} \alpha_i(a_i, 1).$$

There are two cases to be considered.

Case 1: when $A_j = \emptyset$ for some $j \in [r]$. Then $z = 0$ and some A_h , say A_1 , is nonempty and not all coefficients α_i with $a_i \in A_1$ are zero. Thus $\sum_{a_i \in A_1} \alpha_i(a_i, 1) = 0$. Then $\alpha_i \leq 0$ for all $a_i \in A_1 \cap M$ and $\alpha_i \geq 0$ for all $a_i \in A_1 \setminus M$. Setting

$$\gamma := \sum_{a_i \in A_1 \cap M} \alpha_i = \sum_{a_i \in A_1 \setminus M} -\alpha_i,$$

it follows that $\gamma > 0$. Consequently $\text{conv}(A_1 \cap M)$ and $\text{conv}(A_1 \setminus M)$ have a point in common, namely

$$\frac{1}{\gamma} \sum_{a_i \in A_1 \cap M} \alpha_i a_i = \frac{1}{\gamma} \sum_{a_i \in A_1 \setminus M} (-\alpha_i) a_i,$$

contradicting the separation assumption.

Case 2: when A_j is nonempty for all $j \in [r]$. Reading the last coordinate of (3.2) gives that

$$\gamma := \sum_{a_i \in A_1} \alpha_i = \sum_{a_i \in A_2} \alpha_i = \dots = \sum_{a_i \in A_r} \alpha_i.$$

Since $|M| < r$, there is a $j \in [r]$ such that $\alpha_i > 0$ for all $a_i \in A_j$, implying that $\gamma > 0$. Then the point $\frac{1}{\gamma} z$ is in the affine hull of every A_j . The construction guarantees that $\alpha_i < 0$ if and only if $a_i \in M$. \square

Actually, this proof gives a stronger statement. In Case 2, the positivity of γ is guaranteed by the condition $|M| = k < r$. Not assuming $k < r$, γ can be negative or zero. When $\gamma < 0$ equation (3.2) implies again that $\frac{1}{\gamma}z$ is in the affine hull of every A_j , but this time $\alpha(x) > 0$ exactly when $x \notin M$. We will exclude the case $\gamma = 0$ using the general position condition. The proof of this is given in Section 6 because it uses the content of Section 5. So we have the following result.

Theorem 3.1. *Under the conditions of Theorem 1.1 let M be a subset A such that $\text{conv } M \cap \text{conv } (A \setminus M) = \emptyset$. Then there is a partition $\mathcal{A} = \{A_1, \dots, A_r\}$ of A such that in (1.1) either*

- $\alpha(x) < 0$ if and only if $x \in M$, or
- $\alpha(x) < 0$ if and only if $x \notin M$.

The second example in Section 2 shows that in some cases only the second alternative holds.

Corollary 3.1. *Assume r, d are both even and positive integers, $A \subset \mathbb{R}^d$ is in general position, $|A| = (r-1)(d+1) + 1$, and $k = \frac{1}{2}[(r-1)(d+1) + 1]$. Then A has a proper r -partition such that in (1.1) exactly k of the coefficients $\alpha(x)$ are negative.*

The **proof** is easy. Under the above conditions there is a subset M of A of size k that is separated from $A \setminus M$. According to Theorem 3.1, A has an r -partition such that in (1.1) either $\alpha(x) < 0$ if and only if $x \in M$, or $\alpha(x) < 0$ if and only if $x \in A \setminus M$. In both cases, exactly k coefficients are negative. \square

Remark. The same result can be proved using Tverberg's original method of *moving the points*. The main idea is to start with a set of points which have a partition with the required conditions. Then, as one moves one point continuously, if the partition stops working, one can show that points may be swapped in the partition in order to still satisfy the conclusion of the theorem. The proof given above is shorter and simpler.

4. COLOURFUL TVERBERG PLUS MINUS

Once a Tverberg type theorem with conditions on the signs of coefficients of the affine combinations has been established, it becomes natural to try to extend it to the coloured versions of Tverberg's theorem, as in [BL92].

Given disjoint sets F_1, \dots, F_n of r points each in \mathbb{R}^d , considered as colour classes, we say that A_1, \dots, A_r is a colourful partition of them if $|F_i \cap A_j| = 1$ for all $i \in [n], j \in [r]$. In such a case, we can denote the points by $x_{i,j} = F_i \cap A_j$. The coloured Tverberg theorem is concerned about the existence of colourful partitions for which the convex hulls

of the sets A_j intersect. In other words, we seek a colourful partition and a point $z \in \mathbb{R}^d$ for which there is a solution to the equations

$$(4.1) \quad z = \sum_{i=1}^n \alpha(x_{i,j}) x_{i,j} \text{ for all } j \in [r]$$

$$(4.2) \quad \text{subject to } 1 = \sum_{i=1}^n \alpha(x_{i,j}) \text{ for all } j \in [r], \text{ and}$$

$$(4.3) \quad \alpha(x_{i,j}) \geq 0 \text{ for all } i \in [n], j \in [r].$$

The question then becomes, given $M \subset [n]$, find a solution where we exchange (4.3) for

$$(4.4) \quad \alpha(x_{i,j}) \leq 0 \quad \text{for all } i \in M, j \in [r], \text{ and}$$

$$(4.5) \quad \alpha(x_{i,j}) \geq 0 \quad \text{for all } i \in [n] \setminus M, j \in [r].$$

In other words, we aim to prescribe negative coefficients, but we also require that the same restrictions hold across the colour classes. We obtain a partial result in this direction.

Theorem 4.1. *Let $n = (r - 1)d + 1$ and F_1, \dots, F_n be disjoint subsets of \mathbb{R}^d whose union is algebraically independent, each of cardinality r and $M \subset [n]$. Then, there is a colourful partition of F_1, \dots, F_n into r sets and solutions to equations (4.1) and (4.2) such that either*

- $\alpha(x_{i,j}) > 0$ for all $i \in M$ and $\alpha(x_{i,j}) < 0$ for all $i \in [n] \setminus M$, or
- $\alpha(x_{i,j}) < 0$ for all $i \in M$ and $\alpha(x_{i,j}) > 0$ for all $i \in [n] \setminus M$.

Moreover, the affine combinations use the same coefficients for the colour classes. In other words, for all $i \in [n]$ and $j, j' \in [r]$, $\alpha(x_{i,j}) = \alpha(x_{i,j'})$.

Proof. We use the main result of [Sob15]. It says that for $n = (r - 1)d + 1$ and the sets F_1, \dots, F_n , there are solutions to equations (4.1), (4.2) and (4.3) where $\alpha(x_{i,j}) = \alpha(x_{i,j'})$ for all $i \in [n], j \in [r], j' \in [r]$. Then, we apply this result to the sets

$$G_i = \begin{cases} F_i & \text{if } i \in M \\ -F_i & \text{otherwise.} \end{cases}$$

Let B_1, \dots, B_r be the colourful partition we obtain of G_1, \dots, G_n , with $y_{i,j} = G_i \cap B_j$ for all i, j . We denote by $\beta(y_{i,j})$ the coefficients we obtain satisfying equations (4.1), (4.2) and (4.3). We rename them as $\beta(y_{i,j}) = \beta_i$, since they do not depend on j . Let $x_{i,j} = \pm y_{i,j}$ and $\alpha_i = \pm \beta_i$ where the sign is positive (negative) if $i \in M$ ($i \notin M$), respectively. Let $\gamma = \sum_{i=1}^n \alpha_i$. Let us see what happens if $\gamma \neq 0$. By construction, if we consider A_1, \dots, A_r the partition induced by the points $x_{i,j}$ and $\alpha(x_{i,j}) = \alpha_i/\gamma$ for all i, j , they satisfy all the requirements for the conclusion of the theorem. The two cases in Theorem 4.1 correspond to the possibilities for the sign of γ .

We have to verify that the general condition assumption we have on F_1, \dots, F_n implies $\gamma \neq 0$. This part of the proof is technical, and it relies on the modification of Sarkaria's trick from [Sob15]. Given a set $F = \{z_1, \dots, z_r\} \subset \mathbb{R}^d$, a permutation $\sigma : [r] \rightarrow [r]$ and $v_1, \dots, v_r \in \mathbb{R}^{r-1}$ as in Section 3, we can define

$$F \otimes \sigma = \sum_{j=1}^r z_j \otimes v_{\sigma(j)} \in \mathbb{R}^{n-1}, \quad S(F) = \{F \otimes \sigma : \sigma \text{ is a permutation}\}$$

The existence of β_1, \dots, β_n follows from applying the colourful Carathéodory theorem to the sets $S(G_1), \dots, S(G_n)$ in \mathbb{R}^{n-1} . However, if the original set of points $\cup_{i=1}^n F_i$ is algebraically independent, then no transversal to $S(F_1), \dots, S(F_n)$ would have a non-trivial affine dependence in \mathbb{R}^{n-1} , so $\gamma \neq 0$, as required. \square

As mentioned, Theorem 4.1 has equal coefficients across the colour classes. Removing this condition leads to the following problem.

Open problem 4.1. *If we remove the equal coefficients condition, does Theorem 4.1 hold with $n = d + 1$?*

The answer is affirmative with $r = 2$. If $M = \emptyset$ this is the main conjecture from [BL92].

5. PROOF OF PROPOSITION 1.1

We write the equation (1.1) in matrix form $M\alpha = b$. The $(n + d) \times (n + d)$ matrix M is made up of blocks. The block corresponding to A_j is a $d \times |A_j|$ matrix N_j whose columns are the vectors in A_j . The row immediately below block N_j has a 1 in each column containing a vector from A_j and zeroes everywhere else. There are r further blocks, each one is $-I_d$, the negative $d \times d$ identity matrix. They are in the last d columns of M , with a row of zeroes between them. These submatrices are arranged in M as shown on Table 1. All other entries of M are zeroes. The i th column of M corresponds to the vector a_i . Note that $M = M(\mathcal{A})$ depends on A and on the partition $\mathcal{A} = \{A_1, \dots, A_r\}$ as well.

The variables are $\alpha = (\alpha_1, \dots, \alpha_n, z_1, \dots, z_d)^T \in \mathbb{R}^{n+d}$ and the right hand side vector is $b \in \mathbb{R}^{n+d}$ that has coordinate zero everywhere except in positions $d + 1, 2(d + 1), \dots, r(d + 1)$ where it has one. The original system (1.1) is the same as

$$(5.1) \quad M\alpha = b.$$

As we have seen, $\bigcap_{j=1}^r \text{aff } A_j$ is a single point if and only if the linear system (1.1), or what is the same, the equation (5.1) has a unique solution which happens if and only if $\det M \neq 0$. Here $\det M$ is a polynomial with integral coefficients in the coordinates of the a_i . If this polynomial is zero at some algebraically independent points a_1, \dots, a_n , then it is identically zero. So it suffices to show one example where it is

N_1				$-I_d$
1 1 ... 1				
	N_2			$-I_d$
	1 1 ... 1			
		\ddots		
			N_r	$-I_d$
			1 1 ... 1	

TABLE 1. The matrix M , the empty regions indicate zeros

non-zero or, what is the same, one example where $\bigcap_1^r \text{aff } A_j$ is a single point.

The example is simple. Suppose $|A_j| = d + 1 - m_j$ for all $j \in [r]$ and $m_1 \geq m_2 \geq \dots \geq m_r$. As \mathcal{A} is a proper partition, $0 \leq m_j \leq d$. Let H_j be the subspace of \mathbb{R}^d defined by equations $x_i = 0$ for $i = \sum_{h=1}^{j-1} m_h + 1, \dots, \sum_{h=1}^j m_h$. Since $n = (r - 1)(d + 1) + 1$, $\sum_1^r m_j = d$, implying that $\bigcap_1^r H_j$ is a single point, namely the origin. For each $p \in [r]$ choose $|A_j|$ affinely independent points in H_j . Their affine hull is exactly H_j , finishing the proof of the first part.

For the second part we can assume that A_j is nonempty for all j , and also that $|A_j| \leq d + 1$ as otherwise one can delete some elements of A_j while keeping its affine hull the same. We suppose further that $n = (r - 1)(d + 1)$ by adding extra (and algebraically independent) points to some suitable A_j s. Then $\bigcap_{j=1}^r \text{aff } A_j \neq \emptyset$ if and only if the corresponding linear system (5.1) has a solution. Now M is an $(n + 1) \times n$ matrix. Adding b to M as a last column we get a matrix that we denote by M^* . The system (5.1) has a solution if and only if $\text{rank } M = \text{rank } M^*$. The previous argument shows that $\text{rank } M = n - 1$ and so we have that, as a polynomial, $\det M^*$ is identically zero. Again it suffices to give a single example where $\bigcap \text{aff } A_j = \emptyset$. We use the same example as before except that this time $\sum_{j=1}^r m_j = d + 1$ so we can add the equation $\sum_{i=1}^d x_i = 1$ to the ones defining H_1 if $m_1 < d$ and then $\bigcap H_j = \emptyset$, indeed. If $m_1 = d$ then $H_1 = 0$ and $m_2 = 1$ and we define H_2 by the single equation $x_1 + x_2 = 1$, and again $\bigcap H_j = \emptyset$. The sets A_j are constructed the same way as above. \square

6. PROOF OF THEOREM 3.1

Proof. As we have seen we only have to show that $\gamma \neq 0$. Assume $\gamma = 0$. This happens if and only if the homogeneous version of equation (5.1), that is

$$(6.1) \quad A\alpha = 0$$

has a nontrivial solution, which happens again if and only if $\det M = 0$. This is impossible if the partition is proper (as we have seen in the previous section). Note that $z \neq 0$ and no A_j is the emptyset, this follows from Case 1 of the proof of Theorem 1.3. So assume the partition is not proper. Then A_j has more than $d+1$ elements for some j . Assume that $|A_1| > d+1$, say. This means that

$$\sum_{x \in A_1} \alpha(x)(x, 1) = (z, 0).$$

The vectors $(x, 1)$, $x \in A_1$ are affinely dependent, implying that there is a non-trivial affine dependence $\sum_{x \in A_1} \beta(x)(x, 1) = (0, 0)$. Then for all $t \in \mathbb{R}$

$$\sum_{x \in A_1} (\alpha(x) + t\beta(x))(x, 1) = (z, 0),$$

We choose here $t = t_0$ so that $\alpha(x) + t_0\beta(x) = 0$ for some $x = x_0 \in A_1$. Set $\alpha'(x) = \alpha(x) + t_0\beta(x)$ when $x \in A_1$ and $\alpha'(x) = \alpha(x)$ otherwise.

We change now the partition A_1, \dots, A_r to another one A'_1, \dots, A'_r as follows. Set $A'_1 = A_1 \setminus \{x_0\}$ and choose some A_j with $|A_j| \leq d$ and set $A'_j = A_j \cup \{x_0\}$. All other A_h remain the same. Let M' be the corresponding matrix.

We claim now that $\det M' = 0$. The linear system (6.1) has a non-trivial solution, namely $\alpha(x) = \alpha'(x)$ with $z \neq 0$ unchanged. So indeed, $\det M' = 0$.

Repeating this step finitely many times gives a proper partition such that (6.1) has a nontrivial solution. But the previous section shows that for a proper partition, (6.1) has no non-trivial solution. \square

Acknowledgements. This material is partly based upon work supported by the National Science Foundation under Grant No. DMS-1440140 while the first author was in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Fall 2017 semester. Support from Hungarian National Research Grants no K111827 and K116769 is acknowledged. We are also indebted to Attila Pór and Manfred Scheucher for useful discussions, and to an anonymous referee for careful reading and valuable comments.

REFERENCES

- [Bár82] I. Bárány, *A generalization of Carathéodory's theorem*, Discrete Math. **40** (1982), no. 2-3, 141–152.
- [BL92] I. Bárány and D. G. Larman, *A colored version of Tverberg's theorem*, J. London Math. Soc. (2) **45** (1992), no. 2, 314–320.
- [BO97] I. Bárány and S. Onn, *Colourful linear programming and its relatives*, Math. Oper. Res. **22** (1997), no. 3, 550–567.
- [DV77] J.-P. Doignon and G. Valette, *Radon partitions and a new notion of independence in affine and projective spaces*, Mathematika **24** (1977), no. 1, 86–96.
- [Mat03] J. Matoušek, *Using the Borsuk-Ulam theorem*, Universitext, Springer-Verlag, Berlin, 2003, Lectures on topological methods in combinatorics and geometry.
- [PS14] M. A. Perles and M. Sigron, *Strong general position*, arXiv:1409.2899, 2014.
- [Sar92] K. S. Sarkaria, *Tverberg's theorem via number fields*, Israel J. Math. **79** (1992), no. 2-3, 317–320.
- [Sob15] P. Soberón, *Equal coefficients and tolerance in coloured Tverberg partitions*, Combinatorica **35** (2015), no. 2, 235–252.
- [Tve66] H. Tverberg, *A generalization of Radon's theorem*, J. London Math. Soc. **41** (1966), 123–128.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES, H-1364 BUDAPEST PF. 127 HUNGARY

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON, WC1E 6BT, UK

E-mail address: `barany.imre@renyi.mta.hu`

MATHEMATICS DEPARTMENT, NORTHEASTERN UNIVERSITY, 360 HUNTINGTON AVE., BOSTON, MA 02115, USA

E-mail address: `p.soberonbravo@northeastern.edu`