

Title:

Effectiveness of dehydration on adaptation to heat

Authors:

ANDREW T. GARRETT^{1*}, NIELS G. GOOSENS¹, NANCY G. REHRER¹, MARK
J.PATTERSON² AND JAMES D.COTTER¹

* Present address: School of Life Science, University of Hertfordshire, College Lane,
Hatfield, U.K;

¹ School of Physical Education, University of Otago, Dunedin, New Zealand and

² Defence Science Technology Organisation (DSTO), Melbourne, Australia.

Corresponding author:

Dr Andrew Garrett
School of Life Science
Division of Sport, Health and Exercise
University of Hertfordshire
College Lane
Hatfield
Hertfordshire AL10 9AB
U.K.

Tel: 01707 281370

Fax: 01707 285046

Email: a.t.garrett@herts.ac.uk

Abstract:

Most advice for heat adaptation is to use long-term (>10 d) regimes, in which hydration status is maintained. We tested the hypothesis that short-term (5-day) heat acclimation would confer substantial improvements in physiological strain and exercise tolerance for exercise in the heat, and fluid regulatory strain provides a thermally-independent stimulus for such adaptations. Ten moderately-fit males were heat acclimated using controlled hyperthermia (rectal temperature 38.5°C) for 90 min on five consecutive days ($T_a = 40^\circ\text{C}$, 60% RH), on two occasions separated by a five-week washout, in a randomly assigned, cross-over design; one with euhydration (EUH) and one with dehydration (DEH) during acclimation bouts. One week before, then on the 2nd day after each acclimation regime, a heat stress test (HST) was completed, comprising cycling at 40% peak power output for 90 min ($T_a = 35^\circ\text{C}$, 60% RH), before incrementing to exhaustion. Plasma volume (PV) at rest was measured using CO rebreathing. Acclimation exercise-induced response of [aldo]_p became more pronounced across DEH ($\Delta 178 \text{ pg}\cdot\text{mL}^{-1}$; 95%CI: 33 to 324) but not EUH ($\Delta -47 \text{ pg}\cdot\text{mL}^{-1}$: -209 to 115) and this difference was significant ($P=0.02$). Compared to EUH, permissive DEH during acclimation bouts conferred larger acclimation-induced increases in resting PV (4.1%: -1.5 to 9.8%; $P=0.06$), \dot{Q}_F (4.2: 0.7 to 7.8 $\text{ml}\cdot\text{min}^{-1}\cdot 100 \text{ ml}^{-1}$; $P=0.009$), FVC (0.06: 0.02 to 0.10 $\text{ml}\cdot 100\text{ml Tissue}^{-1}\cdot\text{min}^{-1}\cdot\text{mmHg}^{-1}$; $P=0.006$) and decreased end-exercise f_c by 17% (19: -29 to 9 $\text{b}\cdot\text{min}^{-1}$; $P=0.05$). In conclusion, short-term (5-day) heat acclimation was effective with several adaptations more pronounced after fluid-regulatory strain from a dehydration acclimation regime.

Key words: hypohydration, fluid regulation, plasma volume

Introduction

The adaptive effects of medium to long-term heat acclimation (>8-12 d) have received much research attention (Nielsen et al. 1993; Regan et al. 1996; Nielsen et al. 1997; Cheung and McLellan 1998; Weller and Harrison 2001; Patterson et al. 2004).. However, many of the important adaptations to heat stress are cardiovascular, and occur relatively rapidly (<7 d). Therefore, the use of short-term heat acclimation regime (Weller and Harrison 2001; Patterson et al. 2004) has been adopted in this study. The use of permissive dehydration during acclimation contradicts the existing fluid replenishment guidelines for heat acclimation, which recommend the maintenance of good hydration status during exposure to heat stress conditions (Armstrong and Maresh 1991; Convertino et al. 1996; Cheung and McLellan 1998; Casa et al. 2000). However, the reality for many people undergoing acclimation bouts is that some level of dehydration is normal, if not frequently inevitable (Noakes et al. 1988; Greenleaf 1992).

It is well recognised that humans normally do not voluntarily replace all water lost during prolonged exercise in the heat (Szlyk et al. 1989; Greenleaf 1992; Armstrong et al. 1997), and in some individuals with high sweating rate, total body water loss may reach 8% of body mass (Armstrong et al. 1986). Heat-and-exercise induced hypohydration manifests itself as hyperosmotic hypovolaemia (Nadel et al. 1980), which impairs cutaneous blood flow and sweat rate, and raises heart rate, core temperature, glycogenolysis, perceived exertion and permeability of tight-membranes (Mack and Nadel 1996; Gonzalez-Alonso et al. 1998; Gonzalez-Alonso et al. 1999; Blatteis 2000; Gonzalez-Alonso and

Calbert 2003; Maughan 2003; Watson et al. 2005). The consequence is impaired exercise tolerance and performance in temperate conditions which has previously been debated (Sawka and Noakes 2007). Dehydration during prolonged exercise in the heat increases the response of the fluid- and stress-regulatory hormones aldosterone, AVP and cortisol, as well as thirst (Sawka et al. 1987), such that hydration is regulated at a lower level until after exercise and/or heat stress (Brandenberger et al. 1986; Brandenberger et al. 1989). The osmolality and volume effects of hypohydration incur fluid-regulatory responses that could partially mediate the hypervolaemia and improved fluid-regulatory efficiency that is observed with training and heat acclimation adaptations and which helps attenuate cardiovascular strain in exercise (Hopper et al. 1988). Further, there are two potential mechanisms explaining PV expansion during heat acclimation. First, greater renal and sweat related retention of electrolytes and fluid would expand the vascular compartment. Secondly, it has been reported that PV expansion after the movement of total plasma protein to the interstitial fluid in the vascular space. However, it is possible that both these mechanisms occur during heat acclimation and that interaction may contribute to PV expansion (Patterson et al. 2004). Therefore, we believe that in contrast to current recommendations of avoiding dehydration during exercise and heat acclimation, permissive dehydration independently facilitates acclimation by increased fluid-electrolyte retention, PV expansion and cardiovascular response to heat stress.

Methods

Experimental subjects and screening measures. Participants were 9 healthy, male, well-trained, volunteers (Mean \pm SD; Age 27 ± 7 y; Body mass 74.6 ± 4.4 kg; $\dot{V}O_{2\text{peak}}$ 60 ± 7 mL \cdot kg $^{-1}\cdot$ min $^{-1}$ and peak power output 340 ± 30 W). Acclimations were carried out in the winter-spring, to minimise seasonal acclimatisation effects. The study was conducted within the bounds of approval granted by University of Otago Human Ethics Committee.

Experimental design and overview

A randomly assigned, cross-over design was utilised (Figure 1). Nine participants undertook two, 5-d heat acclimation regimes, separated by a 5-week washout (Barnett and Maughan 1993; Pandolf 1998; Hayden et al. 2004); One acclimation with full rehydration (EUHydrated) and one with minimal fluid replenishment (DEHydrated) during each daily acclimation session. Heat acclimation consisted of 90-min exposure on 5 consecutive days (40°C, 60% RH), using controlled hyperthermia (rectal temperature (T_{re}) 38.5°C). Fluid regulatory measures of plasma aldosterone [aldo]_p, plasma AVP [AVP]_p and plasma total protein [TP]_p were recorded at rest and end exercise on day 1 and day 5 of acclimation. The nine participants completed an exercising heat stress test (HST) administered one week before and the 2nd day after EUH and DEH acclimation. On arrival at the laboratory before the HST participants' thermoregulatory, cardiovascular and fluid-regulatory status were measured during 60 min period of rest before cycling at 40% of pre-acclimation peak power output (PPO; $\sim 50\% \dot{V}O_{2\text{peak}}$) for 90 min in the heat (T_{db} , 35°C 60% RH, with a wind speed < 0.5 m \cdot s $^{-1}$). Participants then rested for 10 min before commencing a ramp protocol (2% PPO each 30 s) to volitional fatigue, or a $T_{re} \geq 39.5^\circ\text{C}$.

Before the HSTs participants consumed 250 ml of the 4% CHO fluid. At approximately 15 min intervals during HSTs, 150 ml of 4% CHO solution was consumed, for a total consumption of 900 ml in 90 min (Convertino et al. 1996; Casa et al. 2000). Blood volume (BV) was measured one week before the pre-acclimation HST and the 1st day after EUH and DEH acclimation using the carbon monoxide (CO) rebreathing technique (Burge and Skinner 1995). Plasma volume (PV) was derived by subtraction.

For logistical reasons regarding usage of the heat chamber facility and participant availability the HSTs were in the mornings, all beginning at the same time of the day (9.00 am). Participants were asked to refrain from strenuous exercise immediately before and 24 h prior to each HST, as it has been demonstrated that lower resting core temperature contributes to reduced physiological strain during acclimation (Kampmann et al. 2008). Acclimation bouts took place in the late afternoon/early evening. This consistency was important to control for changes in core temperature and sweating response that may occur if repeated heat exposures were to take place at different times of the day (Shido et al. 1999). Specifically, Shido and colleagues (1999) reported that the effect of acclimation on resting T_{re} was specific to the time of day of acclimation.

Procedures and Measurements

Acclimation: Heat acclimation sessions consisted of cycling (Monarch Ergomedic 824E, Sweden) for 90 min in hot and humid conditions (T_{db} 40°C 60% RH, with air velocity $<0.5 \text{ m}\cdot\text{s}^{-1}$). Since dehydration increases T_c under active heat stress, it was necessary to clamp T_c independently of hydration status to nullify its influence. Elevating T_{re} to the same level during exposure was intended to permit more work as acclimation

proceeded (Figure 2), and balance thermal strain between regimes. Modest hyperthermia (T_{re} of 38.5°C) was attained as rapidly as practicable, and was maintained by regular adjustment of workload to alter the physical stress imposed for the participant. Nominal fluid replacement (100 mL) was given immediately before the DEH acclimation bouts to limit perception of fluid deprivation. Fluid-regulatory measures and total protein were recorded on day 1 and day 5 of both (EUH and DEH) acclimation regimes. Blood volume was measured one week before the HST and on the 1st day after both (EUH and DEH) acclimations.




Figure 1: Experimental design for examining the adaptive effects of short-term heat acclimation with and without dehydration. Acclimation was staggered between participants for logistical reasons.

Fluid regulatory and stress hormones

A flexible 20-gauge catheter was placed in a suitable forearm vein before each HST and on days one and five of the EUH and DEH acclimation trials. Venous blood samples (15 mL) were taken without stasis, following a 1 mL discard. The catheter was flushed with saline after sampling. The samples were taken at rest, 30, 60 and 90 min in the HST, and at the same times on days one and five of the acclimation regimes, but with the omission of the 30 min sample (Morel and Doucet 1986; Creasy 2002). Plasma for aldosterone [aldo]_p, AVP and cortisol were analysed using the radioimmunoassay ¹²⁵I labeling technique. The intra-assay coefficient of variation for duplicate measures were as follows; [aldo]_p (9.9%), [AVP]_p (5.6%) and [cortisol]_p (12.1%). All samples for a given individual were analysed within the same assay.

Heat stress tests: The 90-min exercise bouts at 40% peak power output (T_{db} , 35°C 60% RH, with a wind speed $<0.5 \text{ m}\cdot\text{s}^{-1}$), were undertaken on an electro-magnetically braked cycle ergometer (Rodby Elektronik AB, Model RE 820/830, Sodertalje, Sweden). This protocol was adapted from that used by Patterson (1999). Body core temperature was measured using a rectal thermistor (Thermistor 400, Mallinckrodt Medical Inc., St Louis, USA) placed in the rectum 10 cm beyond the anal sphincter. Skin temperature was measured (Type EUS-U-V5-V2; Grant Instruments, Cambridge, England) at four, right-side sites: calf, upper thigh, chest and bicep. Mean skin temperature (\bar{T}_{sk}) and mean body temperature (\bar{T}_b) were calculated as: $\bar{T}_{sk} = 0.3 T_{\text{chest}} + 0.3 T_{\text{bicep}} + 0.2 T_{\text{thigh}} + 0.2 T_{\text{calf}}$ (Ramanathan 1964) and $\bar{T}_b = 0.9 T_{\text{re}} + 0.1 T_{\text{sk}}$ (Sawka et al. 1996). Temperatures were logged at 60-s intervals (1200 series, Squirrel Grant Instruments, Cambridge, England).

Sweat rate

Whole body sweat rate ($\text{ml}\cdot\text{h}^{-1}$) was calculated using measurements of pre- and post-exercise nude body mass using scales (accuracy 0.1 kg; resolution $\pm 20\text{g}$; Wedderburn Scales, Teraka Seiko, Tokyo, Japan). Continuous measurement of sweat rate from discrete skin sites was undertaken throughout HSTs using ventilated sweat capsules glued to the forehead and dorsal forearm (Brenzelmann et al. 1975; Graichen et al. 1982).

Forearm blood flow, blood pressure and vascular conductance

Forearm blood flow (\dot{Q}_F) was measured using venous occlusion plethysmography (Witney 1953; Joyner et al. 2001) during HSTs, with a custom built controller and commercial software for data capture and filtering (Matlab 4e and Chart 4.1 for Windows event Manager software by Power Lab, ADInstruments, USA). Blood pressure was measured by auscultation, as the average of three individual measurements taken following each \dot{Q}_F measurement series, using a sphygmomanometer. Forearm vascular conductance (FVC) was calculated as \dot{Q}_F divided by mean arterial pressure (MAP), where MAP was calculated as diastolic blood pressure plus one third of Pulse Pressure. Forearm blood flow (\dot{Q}_F), mean arterial pressure (MAP) and forearm vascular conductance (FVC) were measured at rest, 15, 45 and 75 min during the 90-min exercise stress test.

Blood measures

Haemoglobin (Hb) mass and blood volume were measured one week before the baseline HSTs and on the 1st day after each acclimation, using the carbon monoxide (CO) dilution/rebreathing technique (Burge and Skinner 1995; Ashenden et al. 1999). This

measurement was taken in the upright position with the participant seated for a 20 minute equilibrium period. This technique is based on the principle that CO has a strong affinity with Hb, yielding carboxyhaemoglobin (HbCO), which allows Hb mass to be determined from the extent of increase in [HbCO] due to rebreathing a known mass of CO. Venous blood samples were obtained after a priming dose and again after CO rebreathing for ten minutes to obtain [HbCO]. The dosage of CO for aerobically fit males is 20 mL for the priming dose and 1.5 mL CO/kg body mass for the main dose. Pilot testing (n=6) indicated that test-retest reliability for accuracy of repeated measurement had a coefficient of variation (CoV) of ~2.9% for blood volume. Blood, plasma and red cell volumes derived from Hb mass were measured before the baseline standardised heat stress tests and on the first day following each acclimation. Changes in plasma volume across HSTs and acclimations were calculated using a previously described method (Dill and Costill 1974).

Duplicate colorimetric analysis (Cobas Mira Plus, New Jersey, USA) was used to analyse albumin, total protein, Na^+ , K^+ and Cl^- . Osmolality was analysed using a Wescor Vapro Vapour Pressure Osmometer (5520), Wescor Inc., 459 South main Street, Logan, Utah 84321, USA.

Subjects' perceptions

Subjects rated their perceptions of exertion (16-point scale ranging 6-20: (Borg 1982), body temperature (1-13) and thermal comfort (1-5) (Gagge et al. 1967) at 20-30 min intervals during HSTs.

Data analysis

This research addressed the role of permissive dehydration in facilitating adaptations to the heat. Inferential analyses was performed on data collected at rest before (e.g., blood volumes) or within (e.g., heart rate) HSTs and at end-exercise in HSTs before and after each acclimation, as well as on the first and last days of each acclimation. Data was analysed using two-way analysis of variance (ANOVA) with repeated measures with nine participants. Factor one (time) had two levels for each variable, with factor two (acclimation type) having two levels. Paired t-test analysis has been used to isolate differences between means for significant two-way interactions. Estimates of population effects for acclimation status (pre vs post) and acclimation type (EUH vs DEH) are given and reported as means with 95% confidence intervals (95% CI) (Hopkins 2003). The relationship (r) between variables has been calculated using the Pearson Product Moment Correlation and expressed as r .

Results

All nine participants completed both acclimation regimes, the four heat stress tests (HSTs) and all procedures and measurements associated with them, except that one participant became exhausted in one heat stress test at 67 min, so his end-exercise data were taken as the 60-min measurement in all HSTs.

Acclimation

Thermal stress and strain

Thermal stress and strain from day 1 and day 5 of EUH and DEH acclimation regimes are shown in Table 1. Measures of dry bulb temperature (T_{db}) and relative humidity (RH) indicated that the thermal stress was similar within and between EUH and DEH acclimations, on day 1 and day 5. Similarly, the thermal strain was consistent between regimes illustrated by mean cardiac frequency (f_c) and rectal temperature (T_{re}). Time to T_{re} 38.5°C was longer on day 1 EUH than DEH but by day 5 the time to T_{re} 38.5°C was similar for EUH and DEH regimes. Therefore, less work was performed during the first acclimation day when comparing DEH and EUH in contrast with day 5, when work output performed in the 90 min was equivalent for DEH versus EUH (Figure 2).

Fluid regulatory hormones and total protein

Individual response data for blood measures are presented from DEH and EUH acclimation bouts for n=5, since a full cross-over of blood samples (i.e., rest and post exercise for day 1 and day 5, for both acclimations) was obtained from only five participants during the acclimation bouts (Table 2). The main effect for [aldo]_p did not change at rest but at 90-min exercise there was a significant main and interaction effect between EUH and DEH. However, [AVP]_p did not change across acclimation at rest or 90-min exercise. Similarly, [TP]_p did not change at rest or 90-min exercise but the interaction effect between EUH and DEH was significant.

Table 1 Thermal stress and strain.

	DAY 1		DAY 5		
	EUH	DEH	EUH	DEH	
T_{db} (°C)	39.5±0.1	39.6±0.1	39.6±0.1	39.5±0.1	
RH (%)	62 ±1.6	62 ±1.7	62 ±1.8	61 ±1.4	
Mean f_c (b·min ⁻¹)	123 ±13	122±16	123 ±14	121 ±10	
Mean T_{re} (°C)	38.2 ±0.1	38.3±0.1	38.2 ±0.1	38.2±0.2	
Time to T_{re} 38.5°C (min)	34 ±4.7	30.5±4.4	34±4.6	33±4.2	*†
Fluid consumed (mL)	1930±452	100 ±0	1861±333	100 ±0	†
Body mass change (%)	0.4 ±0.5	-1.7±0.5	0.1 ±0.7	-2.0±0.7	†

Dry Bulb Temperature (T_{db}), relative humidity (RH), rectal temperature (T_{re}), time to T_{re} 38.5°C, cardiac frequency (f_c), body mass change and fluid consumed on day one and five of acclimation undertaken with (EUH) or without (DEH) fluid rehydration. Data are mean ±SD for nine males for whom data were available across all sessions. Significant difference; day 1 versus day 5 * ($P<0.05$) and hydration regime † ($P<0.05$) analysed by two-way analysis of variance (ANOVA) with repeated measures.




Figure 2: Work output on the first day (1) to the last day (5) of euhydration and dehydration acclimation after 90 min heat exposure. Data are shown as mean \pm SE for nine males for whom data was available across all sessions. Significant difference * ($p < 0.05$) EUH cf. DEH acclimation analysed by two-way analysis of variance (ANOVA) with repeated measures. One-way analysis of variance (ANOVA) and Tukey's post-hoc test was used to isolate differences between days of EUH and DEH acclimation.

Blood, plasma and red cell volume

Plasma volume and thus also blood volume for nine males increased across acclimation ($P=0.02$ and $P=0.04$), irrespective of whether dehydration was permitted or not (PV: $P=0.52$; BV: $P=0.40$). Expansion of PV across the DEH acclimation was correlated ($r=0.65$) with the increase in $[\text{aldo}]_p$ measured across the final exercise and heat bout. Applying the Dill and Costill (1974) calculation to estimate the Δ PV across acclimations (i.e., assuming unchanged Hb mass), there was further tendency for PV to increase more in DEH ($8.3 \pm 3.2\%$) than in EUH ($4.2 \pm 3.4\%$) ($P=0.06$; difference of 4.1% : -1.5 to 9.8).

Table 2 Change in aldosterone, AVP and total protein response across day one and five EUHydration and DEHydration acclimation

	EUHydration	DEHydration	Main effect	Interaction
Aldosterone (pg.mL ⁻¹)				
Rest	-1.6: -30 to 33	-10: -51 to 71	<i>P</i> =0.65	<i>P</i> =0.74
90-min exercise	-47: -209 to 115	178: 33 to 324	<i>P</i>=0.05	<i>P</i>=0.02
AVP (p.mol.L ⁻¹)				
Rest	0.1: -3.5 to 5	0.9: -0.5 to 2	<i>P</i> =0.76	<i>P</i> =0.91
90-min exercise	-1.3: -7 to 4	3: -2 to 8	<i>P</i> =0.78	<i>P</i> =0.10
Total protein (mg.mL ⁻¹)				
Rest	-3: -13 to 8	-7: -20 to 6	<i>P</i> =0.14	<i>P</i> =0.23
90-min exercise	-5.9: -11.1 to 0.4	2.9: -4.2 to 9.9	<i>P</i> =0.46	<i>P</i>=0.03

Data are shown as mean and 95% confidence interval (95% CI) for five males for whom data was available across all sessions. Significant difference * (*P*<0.05) shown in **bold**.

Red cell volume (RCV), corrected for the volume of red cells removed during acclimation bouts did not have a significant main effect across acclimation (*P*=0.06) and the interaction effect between EUH and DEH was not significant (*P*=0.18). However, the mean change was 1.7 ±1.6% following EUH (Δ 0.6: -0.7 to 1.9 mL·kg⁻¹) but 4.1 ±0.9% after DEH (Δ 1.5: 0.7 to 2.3 mL·kg⁻¹). Therefore, this indicates that the difference in PV change between regimes may have been underestimated. Interestingly, 5 of 9 individuals RCV increased after EUH and all 9 of 9 participants following DEH acclimation.

Heat stress test

Cardiac frequency

End-exercise f_c (Figure 3) reduced across acclimation ($P=0.01$), and this was more pronounced for DEH (-19 $\text{b}\cdot\text{min}^{-1}$: -29 to -3) than EUH (-10 $\text{b}\cdot\text{min}^{-1}$: -17 to 2) ($P=0.05$). This reduction in f_c across DEH was moderately correlated with the extent of PV expansion ($r=0.72$).




Figure 3: Cardiac frequency during 90-min cycling at 40% peak power output in the heat (35°C , 60% RH) before (pre) and after (post) acclimation, undertaken with (EUH) or without (DEH) rehydration during daily heat sessions. Data are means \pm SE for nine males and expressed in $\text{b}\cdot\text{min}^{-1}$.

Body temperatures

The mean change for T_{re} at rest (Figure 4) was not lowered across EUH ($\Delta 0.15$: -0.40 to 0.10°C) or DEH ($\Delta 0.00$: -0.2 to 0.2°C). However, the main effect for T_{re} at 90-min exercise was significant ($P=0.004$) with no interaction effect between EUH and DEH

($P=0.52$). The mean change for T_{re} at 90-min exercise was reduced by EUH ($\Delta -0.2$: -0.6 to 0.05°C) and DEH ($\Delta -0.4$: -0.75 to -0.1°C).




Figure 4: Rectal temperature during 90-min cycling at 40% peak power output in the heat (35°C, 60% RH) before (pre) and after (post) acclimation, undertaken with (EUH) or without (DEH) rehydration during daily heat sessions. Data are means \pm SE for nine males and expressed in °C.

Mean resting \bar{T}_{sk} did not change across EUH ($\Delta 0.1$: -0.1 to 0.5°C) or DEH ($\Delta 0.2$: -0.1 to 0.5°C). However, the main effect in \bar{T}_{sk} at 90-min exercise was significant ($P=0.003$) but with no interaction effect between EUH and DEH ($P=0.15$). The mean change in \bar{T}_{sk} after 90-min exercise decreased across EUH ($\Delta -0.3$: -1.1 to 0.3°C) and DEH ($\Delta -0.4$: -0.7 to 0.1°C). Mean change in resting \bar{T}_b response was unaltered by EUH ($\Delta -0.2$: -0.4 to 0.1°C) or DEH ($\Delta 0.1$: -0.2 to 0.1°C). The main effect for \bar{T}_b at 90-min exercise was significant ($P=0.002$) but there was no interaction effect between EUH and

DEH ($p=0.17$). Therefore, at 90-min exercise mean change in \bar{T}_b reduced across EUH ($\Delta -0.2$; -0.4 to 0.1°C) and DEH ($\Delta -0.4$; -0.6 to 0.1°C).

Forearm blood flow, blood pressure and vascular conductance

The main effect for resting \dot{Q}_F (Figure 5) was significant across acclimation ($P=0.05$). There was a significant interaction effect for EUH and DEH ($P=0.009$). The mean change for resting \dot{Q}_F increased across DEH ($\Delta 4.2$: 0.7 to $7.8 \text{ ml}\cdot\text{min}^{-1}\cdot 100 \text{ ml}^{-1}$) in contrast to limited change for EUH ($\Delta -1.5$: -3.7 to $0.7 \text{ ml}\cdot\text{min}^{-1}\cdot 100 \text{ ml}^{-1}$). However, there was no main effect for exercise at 75 min \dot{Q}_F ($P=0.58$).




Figure 5: Forearm blood flow during 90-min cycling at 40% peak power output in the heat (35°C , 60% RH) before (pre) and after (post) acclimation, undertaken with (EUH) or without (DEH) rehydration during daily heat sessions. Data are means \pm SE for nine males and expressed in $\text{ml}\cdot\text{min}^{-1}\cdot 100 \text{ ml}^{-1}$.

Resting MAP did not change across acclimation ($P=0.56$) or at 75-min exercise ($P=0.14$). The main effect for resting FVC across acclimation was significant ($P=0.05$)

with an interaction effect between EUH and DEH ($P=0.006$). Therefore, the mean change in resting FVC increased across DEH (Δ 0.06: 0.02 to 0.10 ml \cdot 100ml Tissue $^{-1}\cdot$ min $^{-1}\cdot$ mmHg $^{-1}$), in contrast to no change across EUH (Δ -0.02: -0.05 to 0.01 ml \cdot 100ml Tissue $^{-1}\cdot$ min $^{-1}\cdot$ mmHg $^{-1}$). There was no main effect for FVC at 75-min exercise ($P=0.34$). In summary, the resting \dot{Q}_F and FVC response were elevated across DEH versus EUH but no such effect was seen at 75-min exercise \dot{Q}_F and FVC. Therefore, it appears that the resting response of \dot{Q}_F and FVC became more pronounced after the DEH versus EUH acclimation.

Sudomotor responses

Forehead and forearm sweat rate were measured at rest and 1-min intervals during the 90-min exercise stress test. Relative increases calculated from the mean data indicate that sweating on the forehead decreased after EUH (11%) and DEH (7%). However, the main effect was not significant for sweat rate at the forehead at 90 min ($P=0.72$) and there was no interaction between EUH and DEH ($P=0.822$). In contrast, sweat rate at the forearm was elevated after 90 min for EUH (19%) and DEH (28%). The main effect was significant ($P=0.01$) but there was no interaction effect between EUH and DEH [F ($P=0.77$)]. Mean changes in sweating at the forearm after 90-min exercise increased for EUH (Δ 0.3: -0.1 to 0.7 mg \cdot cm $^{-2}\cdot$ min $^{-1}$) and DEH (Δ 0.4: 0 to 0.8 mg \cdot cm $^{-2}\cdot$ min $^{-1}$).

Subjects' perceptions

Perceived body temperature did not change at rest across acclimation ($P=0.09$). However, the main effect at 90 min was significant ($P=0.02$) but there was no interaction effect between EUH and DEH ($P=0.18$). Thermal comfort improved at rest and had a significant main effect ($P=0.05$). There was no interaction effect between EUH and DEH

($P=0.43$). Similarly, after 90-min exercise thermal comfort improved across acclimation ($P=0.05$) but no interaction effect between EUH and DEH ($P=0.38$). Perceived exertion was reduced after 90-min exercise across acclimation and this was significant ($P=0.04$) but there was no interaction effect between EUH and DEH ($P=0.65$).

Fluid regulatory and stress hormones

Mean change at 90-min exercise $[\text{aldo}]_p$ substantially decreased ($P=0.001$) across EUH (Δ -144: -235 to -52 $\text{pg}\cdot\text{mL}^{-1}$) and DEH (Δ -55: -167 to -43 $\text{pg}\cdot\text{mL}^{-1}$) with no difference between regimes ($P=0.50$). Similarly, $[\text{AVP}]_p$ decreased across 90-min exercise for EUH (Δ -2.9: -6.1 to -0.2 $\text{p}\cdot\text{mol}\cdot\text{L}^{-1}$) and DEH (Δ -2.1: -4.1 to 0.1 $\text{p}\cdot\text{mol}\cdot\text{L}^{-1}$) and this was significant ($P=0.01$) but no differences between EUH and DEH ($P=0.31$). There was a decrease in $[\text{cortisol}]_p$ across 90-min exercise in EUH (Δ -3: -40 to 13 $\text{ug}\cdot\text{dL}^{-1}$) and DEH (Δ -2: -15 to 31 $\text{ug}\cdot\text{dL}^{-1}$) and this was significant ($P=0.02$) with no difference between regimes ($P=0.86$).

Blood measures

Mean change in $[\text{TP}]_p$ after 90-min exercise was reduced across EUH (Δ -1.8: -4.6 to 1.0 $\text{mg}\cdot\text{ml}^{-1}$) and DEH (Δ -1.2: -3.0 to -0.5 $\text{mg}\cdot\text{ml}^{-1}$) and this was significant ($P=0.02$) but with no difference between EUH and DEH ($P=0.99$). There was no change in plasma albumin $[\text{alb}]_p$ at rest ($P=0.43$) and there was no interaction effect between EUH and DEH ($P=0.10$). Similarly, after 90-min exercise there was no mean change ($P=0.36$) across EUH (Δ 0.3: -2.9 to 3.4 $\text{mg}\cdot\text{ml}^{-1}$), or DEH (Δ -1.5: -2.8 to -0.2 $\text{mg}\cdot\text{ml}^{-1}$) with no difference between regimes ($P=0.27$). There was no change in plasma sodium concentration $[\text{Na}^+]_p$ at rest ($P=0.38$) and no interaction effect between EUH and DEH ($P=0.11$). Similarly, after 90-min exercise there was no mean change ($P=0.16$) across

EUH (Δ -0.5: -1.5 to 1 mmol·L⁻¹) or DEH (Δ 1.0: 0.5 to 2 mmol·L⁻¹) with no interaction effect between EUH and DEH ($P=0.10$). Plasma osmolality at rest did not change ($P=0.90$) and there was no interaction between EUH and DEH ($P=0.28$). Similarly, after 90-min exercise there was no change ($P=0.85$) across EUH (Δ -4: -22 to 13 mmol·kg⁻¹) or DEH (Δ 4: -12 to 20 mmol·kg⁻¹) and there was no interaction effect between EUH and DEH ($P=0.66$).

Exercise performance capacity

Endurance exercise capacity was measured 10 min following the 90-min steady state exercise stress test, using an incremental protocol with 30-s step work increments to voluntary exhaustion. There was a significant main effect [F (1,14)=2.411, $p=0.001$] and a large effect size (eta squared=0.857) but no interaction effect between EUH and DEH [F (1,14)=2.411, $P=0.14$]. Exercise time significantly increased across acclimation by 14% in EUH (Δ 104: 57 to 150 s) and 19% for DEH (Δ 146: 101 to 191 s).

Discussion

The current project was designed to determine whether adaptations to short-term heat acclimation would be more pronounced when dehydration was permitted. This study demonstrated the effectiveness of short-term (90 min·d⁻¹ for 5-d) heat acclimation, using controlled hyperthermia (38.5°C rectal temperature), attenuating thermal strain and enhancing exercise capacity in the heat. Compared with the euhydration regime, permissive dehydration during acclimation bouts conferred larger acclimation-induced increases in resting plasma volume, \dot{Q}_F and decreased f_c at the end of the standardised exercise heat stress test.

We emphasise that the present findings were obtained using permissive dehydration to a modest level of hypohydration (~1.8%); a level at which additional fluid-regulatory stimuli are already induced (Table 2) (Brandenberger et al. 1986; Brandenberger et al. 1989; McConell et al. 1997) and thirst becomes stimulated to limit excessive dehydration (Engell et al. 1987; Sawka et al. 1987; Nadel et al. 1993). Whether greater dehydration and/or a longer acclimation regime would be more beneficial or detrimental is unknown and requires further investigation, but if exercise training is an important component of the acclimation then the training impulse would become reduced (especially if using the controlled hyperthermic regime).

Effectiveness of short-term acclimation to the heat

The adaptations from short-term (5-d) heat acclimation with euhydration, using the controlled hyperthermia technique, reduced exercising cardiovascular strain and enhanced exercise capacity. The cardiovascular stability was due to increased heat loss rather than lower heat content (~resting core temperature), at the time of day of testing HSTs and this concurs with previous work (Turk and Worsley 1974; Weller and Harrison 2001; Creasy 2002). It is suggested the increased capacity for work observed in the present study and the physiological adaptation to heat stress may have been enhanced by using the controlled hyperthermia technique during acclimation. This technique increases work stress progressively as the individual adapts to the heat stress conditions (Taylor 2000; Creasy 2002).

Fluid regulation and blood volume response to repeated heat stress

In the present study, there was no difference in environmental parameters between acclimation regimes and this was illustrated by similar T_{db} and RH measures (Table 1). Similarly, thermal strain was identical between regimes as evidenced by f_c and T_{re} . Therefore, participants experienced the same thermal load which is the basis of using the controlled hyperthermia technique for heat acclimation (Taylor 2000). Individual's hydration status was reflected on day five acclimation by total body water being maintained in EUH (~0.1% body mass) but after DEH the participants experienced a mild hypohydration of ~2% body mass (Table 1). In a review, on the fluid replacement of athletes Casa and colleagues (2000) reported that dehydration of 1-2% body weight increases physiological stress and decreases performance. However, in reality, it is often difficult and sometimes impossible to prevent at least some dehydration during repeated heat stress (Convertino et al. 1996) and this has been considered of little consequence by some authors (Noakes et al. 1988; Greenleaf 1991), on the basis that this readily occurs during repeated heat exposure. In summary, we postulate that the increased physiological strain of restricted fluid replenishment during DEH acclimation may have resulted in adaptation of the fluid-regulatory system.

Fluid regulatory hormones and electrolytes

The neuroendocrine system regulates the volume and composition of plasma volume and intracellular fluid of red blood cells, whose essential components for the retention of Na^+ and water are the renin-angiotensin-aldosterone system and AVP (Brandenberger et al. 1986). In the present study, after 90-min exercise $[aldo]_p$ increased across DEH bouts in contrast, to a limited response after EUH acclimation and this

difference was significant. However, there was no change in $[AVP]_p$ after rest and 90-min exercise across acclimation (Table 2). Therefore, it appears that the exercise-induced response of $[aldo]_p$ became more pronounced in the DEH compared with the EUH acclimation bouts.

The principal effects of aldosterone are the retention of Na^+ and therefore water from the urine output to maintain extracellular fluid volume and thus also blood volume. Increased Na^+ and water retention at the distal tubules (Morris 1981) are important mediators of the rapid PV expansion during the initial hours to days after exercise (Nagashima et al. 2001). However, in the present study, an exercise-induced response of increased $[Na^+]_p$ was not evident after acclimation for either the EUH or DEH regime, in the standardised exercising heat stress test. Therefore, this is in contrast with previous findings (Brandenberger et al. 1989; Francesconi et al. 1993; Allsopp et al. 1998) who reported a strong relationship between increased Na^+ with $[aldo]_p$ response.

Blood and PV responses showed similar findings with $[aldo]_p$ across DEH acclimation. There was a moderate relationship ($r=0.65$) between PV expansion and the increased $[aldo]_p$ after exercise in the DEH regime (Table 2). Blood volume expansion at rest increased after EUH (2.2%) and was elevated after the DEH regime (5.3%). This was consistent with the expansion of resting PV after EUH (5.3%) and DEH (9.0%). Patterson and colleagues (2004) used the Evans blue dye dilution technique for establishing resting PV with 12 participants. They reported a 9.8% PV expansion following 7-d heat acclimation, using the controlled hyperthermia technique. In the present study, the acclimation-induced resting PV was further supported by the ΔPV (Dill and Costill 1974), across HSTs and acclimations. Resting PV increased for EUH

($4.2 \pm 3.4\%$) but more so following DEH ($8.3 \pm 3.2\%$) and this was significantly different between regimes.

In summary, in the present study it is proposed that the adaptive response to fluid deficit during acclimation has resulted in greater cardiovascular stability. It is suggested that in comparison with EUH, the more pronounced biological action of $[\text{aldo}]_p$ observed during exercise in DEH had a major role in the more robust increase in PV response that was evident after that regime. This is further supported by the moderate relationship across DEH between PV expansion and $[\text{aldo}]_p$ response after exercise. Indicating that the increased $[\text{aldo}]_p$ response observed after DEH acclimation, may have contributed to the greater PV expansion in that regime. This further elucidates the notion that during exercise the fluid-regulatory hormone, $[\text{aldo}]_p$, maybe more systematically altered after a DEH acclimation regime and have a major role in acclimation-induced hypervolaemia. Therefore, it is surprising that the contribution of $[\text{aldo}]_p$ with PV expansion has received limited attention in the literature. Of the limited information available it has been demonstrated that renal and hormonal adaptations after intense exercise participate in the initial (first 24-48 h) process of PV expansion (Nagashima et al. 2001).

Blood, plasma and red cell volume

It is accepted that exercise-induced hypervolaemia mediated by PV expansion (Senay et al. 1976; Harrison 1985) has the beneficial effect of enhancing cardiovascular and thermoregulatory responses to exercise, resulting in greater cardiovascular stability (Fellman 1992). It has been established that there are two mechanisms of PV expansion to account for such a change; increased renal electrolyte and water retention (Wyndham 1973; Costill et al. 1975; Armstrong et al. 1987; Convertino et al. 1991) and elevated

plasma protein content (Harrison et al. 1981; Harrison 1985). Therefore, it has been suggested that both these mechanisms occur during heat acclimation and their interaction may contribute to PV expansion (Patterson et al. 2004). In the present study, resting $[TP]_p$ remained constant from the first to the last day of each of the acclimation regimes (Table 2). However, the blood volume (BV) expansion observed across EUH and DEH acclimation, indicated that resting $[TP]_p$ has increased and may well be a mechanism for the PV expansion that was evident. This is supported by previous research on the notion that PV expansion is mediated by elevated plasma protein content (Harrison et al. 1981; Mack and Nadel 1996). There was an increased exercise-induced response of $[TP]_p$ in the present study that became more pronounced within the DEH compared to the EUH acclimation bouts (Table 2). It is suggested that this differential response between regimes was indicative of the fluid loss from dehydration. In summary, our data suggests that elevated $[TP]_p$ rather than increased $[Na^+]_p$ has a major role in the mediation of the greater acclimation-induced PV expansion observed across acclimation.

In summary, the present data indicates that during exercise the fluid-regulatory hormone, $[aldo]_p$, may be more systematically altered following a DEH acclimation regime. Therefore, we suggest that the increased biological action of $[aldo]_p$ may be responsible for the greater increase in PV that was evident across the DEH regime. It is further suggested that the PV expansion observed after EUH and DEH acclimation may be mediated by elevated plasma protein content. The increased end-exercise responses of $[aldo]_p$ across DEH acclimation are consistent with the increased PV expansion and larger drops in f_c (Figure 3) in the standardised exercising heat stress after the DEH regime.

Adaptation to repeated heat stress

There was greater attenuation of end-exercise f_c after DEH (17%) compared with EUH (7%) acclimation (figure 3) and this difference was significant. The end-exercise T_{re} was lower for EUH (0.7%) and reduced after the DEH (0.8%) regime (Figure 4) with no difference between regimes. Similar findings have previously been reported using short-term heat acclimation regimes, with the controlled hyperthermia technique (Turk and Worsley 1974; Cotter et al. 1997; Patterson et al. 2004). Further, in the present study \bar{T}_{sk} and \bar{T}_b at end-exercise decreased across EUH and DEH acclimation with no difference between regimes. These physiological adaptations are characteristic features of heat acclimation (Armstrong and Maresh 1991) and reflected by greater cardiovascular stability (Wyndham et al. 1968; Strydom and Williams 1969; Senay et al. 1976; Sawka et al. 1985). In summary, these findings indicate that after acclimation per se, the cardiovascular system is not under as much physiological strain and more so after the DEH regime.

It is established that the control of V_s during exercise possibly reflects extramyocardial factors of increased blood volume (Hopper et al. 1988). It has previously been reported by Gonzalez-Alonso and colleagues (1999) that \dot{Q} reduces with dehydration and hyperthermia because of large decreases in V_s (19-27%), compared with parallel increases in f_c (5-10%). There is also a significant reduction in blood flow to the exercising muscle with dehydration during intense exercise in the heat, due to a lowering in perfusion pressure and systemic blood flow rather than increased vasoconstriction (Gonzalez-Alonso et al. 1998). More recently, it has been demonstrated

that in trained individuals, severe heat stress reduces $\dot{V}O_{2\max}$ by accelerating the decline in \dot{Q} and mean arterial pressure, leading to reduced exercising muscle blood flow, O_2 delivery and O_2 uptake (Gonzalez-Alonso and Calbert 2003). Therefore, we suggest that the greater PV expansion (Dill and Costill 1974) that was evident after the DEH acclimation (8.3%), across HSTs and acclimations. May have contributed to elevations in end-exercise V_s , although this measure was not taken in the present work. In a review by Sawka and colleagues (1996), V_s augmentation has been attributed to heat-acclimation-induced PV expansion (Hales et al. 1996).

Dehydration and hyperthermia are thought to reduce central blood volume during exercise in the upright position, affecting cardiopulmonary and arterial baroreceptors, which can inhibit forearm vascular conductance (FVC) to the skin and other tissues (Gonzalez-Alonso et al. 1999a). Gonzalez-Alonso and colleagues (1999) investigated whether the deleterious effects of dehydration and hyperthermia on cardiovascular function, during upright exercise, were attenuated by elevating central blood volume with supine exercise. It has been established that there is greater V_s , increased volume of venous return and central blood volume during supine exercise (Poliner et al. 1980). Gonzalez-Alonso and colleagues (1999) observed that the characteristic features of upright exercise during dehydration and hyperthermia of decreased mean arterial pressure, FVC and increased norepinephrine were absent during supine exercise, despite equal dehydration and hyperthermia. They concluded that the reduction in FVC and the increased plasma norepinephrine concentration, independent of hyperthermia, were associated with a reduction in central blood volume during dehydration. Therefore, the mechanisms mediating the reduction in central blood volume are not fully understood but

it has been hypothesised that decreased FVC may play a greater role than skin blood flow in reducing V_s and central blood volume during exercise in the heat (Gonzalez-Alonso et al. 1999a). In the present study, cutaneous blood flow [estimated from forearm blood flow (\dot{Q}_F)] (Figure 5) and FVC at rest were elevated across DEH but not EUH acclimation, with a significant differential response between regimes. This indicates elevated forearm perfusion across the DEH acclimation and this may be due to decreased arteriolar vascular resistance rather than increased perfusion pressure.

Conclusions

There were significant physiological adaptations and increased work capacity after short-term (5-day) heat acclimation. The nature of these adaptations were generally to increase heat loss rather than to lower resting body temperature, at least at the time of day at which testing took place. Several adaptations were generally more pronounced when acclimation bouts were undertaken with permissive dehydration versus a euhydration regime. This indicates that using permissive dehydration to a modest level of hypohydration (~1.8%) during acclimation, for enhanced adaptation requires further investigation.

Acknowledgements

Special thanks are given to the participants in this study and the technical expertise provided by Mrs R. McKay and Miss D. Wilson.

Grants

This work was supported by grants from the Australian Defence Science Technology Organisation and the School of Physical Education, University of Otago, New Zealand.

References

- Allsopp AJ, Sutherland R, Wood P, Wooton SA (1998) The effect of sodium balance on sweat sodium secretion and plasma aldosterone concentration. *Eur J Appl Physiol* 78:516-521
- Armstrong LE, Costill DL, Fink WJ (1987) Changes in body waters and electrolytes during heat acclimatization: effects of dietary sodium. *Aviat Space Environ Med* 58:143-148
- Armstrong LE, Hubbard RW, Jones BH, Daniels JT (1986) Preparing Alberto Salazar for the heat of the Olympic Marathon. *Phys Sports Med* 14:73-81
- Armstrong LE, Maresh CM (1991) The induction and decay of heat acclimatisation in trained athletes. *Sports Med* 12(5):302-312
- Armstrong LE, Maresh CM, Gabaree CV, Hoffman JR, Kavouras SA, Kenefik RW, Castellani JW, Ahlquist LE (1997) Thermal and circulatory responses during exercise: effects of hypohydration, dehydration, and water intake. *J Appl Physiol* 86:2028-2035
- Ashenden MJ, Gore CJ, Burge CM, Clough ML, Bourdon PC, Dobson GP, Hahn AG (1999) Skin-prick blood samples are reliable for estimating Hb mass with the CO-dilution technique. *Eur J Appl Physiol* 79:535-537
- Barnett A, Maughan R (1993) Response of unacclimatized males to repeated bouts of exercise in the heat. *Br J Sp Med* 27:39-44
- Blatteis CM (2000) Thermoregulation in complex situations: combined heat exposure, infectious fever and water deprivation. *Int J Biometeorol* 44:31-43

- Borg GAV (1982) Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 14:377-381
- Brandenberger G, Candas V, Follenius M, Kahn JM (1989) The influence of initial state of hydration on endocrine responses to exercise in the heat. *Eur J Appl Physiol* 58:674-679
- Brandenberger G, Candas V, Follenius M, Libert JP, Kahn JM (1986) Vascular fluid shifts and endocrine responses to exercise in the heat. *Eur J Appl Physiol* 55:123-129
- Bregelmann GL, McKeag M, Rowell LB (1975) Use of dew-point detection for quantitative measurement of sweating rate. *J Appl Physiol* 39:498-500
- Burge CM, Skinner SL (1995) Determination of hemoglobin mass and blood volume with CO: evaluation and application of a method. *J Appl Physiol* 79:623-631
- Casa DJ, Armstrong LE, Hillman SK, Montain SJ, Reiff RV, Rich BSE, Roberts WO, Stone JA (2000) National Athletic Trainers' Association: Position Statement: Fluid replacement for athletes. *J Athl Train* 35:121-224
- Cheung SS, McLellan TM (1998) Heat acclimation, aerobic fitness, and hydration effects on tolerance during uncompensable heat stress. *J Appl Physiol* 84:1731-1739
- Convertino VA, Armstrong LE, Coyle EF, Mack GW, Sawka MN, Senay LC, Sherman WM (1996) ACSM: Position stand on exercise and fluid replacement. *Med Sci Sports Exerc* 28:i-vii

- Convertino VA, Mack GW, Nadel ER (1991) Elevated central venous pressure: a consequence of exercise training-induced hypervolemia? *Am J Physiol*:R273-R277
- Costill DL, Cote R, Miller E, Miller T, Wynder S (1975) Fluid and electrolyte replacement during repeated days of exposure to heat. *Aviat Space Environ Med* 40:795-800
- Cotter JD, Patterson MJ, Taylor NAS (1997) Sweat distribution before and after repeated heat exposure. *Eur J Appl Physiol* 76:181-186
- Creasy R (2002) Post-exercise sauna bathing does not improve 2000-metre rowing performance. BPhEd, University of Otago, Dunedin, New Zealand
- Dill DB, Costill DL (1974) Calculation of percentage changes in volumes of blood, plasma and red cells in dehydration. *J Appl Physiol* 37(2):247-248
- Engell DB, Maller O, Sawka MN, Francesconi RN, Drolet L, Young AJ (1987) Thirst and fluid intake following graded hypohydration levels in humans. *Physiol Behav* 40:229-236
- Fellman N (1992) Hormonal and plasma volume alterations following endurance exercise. *Sports Med* 13:37-49
- Francesconi RN, Armstrong LE, Leva NM, Moore RJ, Szlyk PC, Matthew WT, Curtis WC, Hubbard RW, Askew EW (1993) Endocrinological Responses to Dietary Salt Restriction During Heat Acclimation. In: *Nutritional needs in hot environments*. National Academy Press, Washington, D.C., pp 259-275

- Gagge AP, Stolwijk JA, Hardy JD (1967) Comfort and thermal sensations and associated physiological responses at various ambient temperatures. *Environmental research* 1:1-20
- Gonzalez-Alonso J, Calbert JAL (2003) Reductions in systematic and skeletal muscle blood flow and oxygen delivery limit maximal aerobic capacity in humans. *Circ* 107:824-830
- Gonzalez-Alonso J, Calbert JAL, Nielsen B (1998) Muscle blood flow is reduced with dehydration during prolonged exercise in humans. *J Physiol* 513.3:895-905
- Gonzalez-Alonso J, Calbert JAL, Nielsen B (1999) Metabolic and thermodynamic responses to dehydration-induced reductions in muscle blood flow in exercising humans. *J Physiol* 520.2:577-589
- Gonzalez-Alonso J, Mora-Rodriguez R, Coyle EF (1999a) Supine exercise restores arterial blood pressure and skin blood flow despite dehydration and hyperthermia. *Am J Physiol*:H576-H583
- Graichen H, Rascati R, Gonzalez RR (1982) Automatic dew-point temperature sensor. *J Appl Physiol: Respirat Environ Exercise Physiol* 52:1658-1660
- Greenleaf JE (1991) The consequences of exercise on thirst and fluid intake. In: Ramsey DJ, Booth DA (eds) *Thirst: Physiological and Psychological aspects*. Springer-Verlag, London
- Greenleaf JE (1992) Problem: thirst, drinking behaviour, and involuntary dehydration. *Med Sci Sports Exerc* 24:645-646

- Hales JRS, Hubbard RW, Gaffin SL (1996) Limitation of heat tolerance. Handbook of Physiology: American Physiological Society. In: Fregly. MJ, Blatteis, C.M. (ed) Handbook of Physiology: American Physiological Society. Vol 1. Oxford University Press, pp 285-355
- Harrison MH (1985) Effects of thermal stress and exercise on blood volume in humans. *Physiol Rev* 65:149-209
- Harrison MH, Edwards RJ, Graveney MJ, Cochrane LA, Davies JA (1981) Blood volume and plasma protein responses to heat acclimatization in humans. *J Appl Physiol* 50:597-604
- Hayden G, Milne HC, Patterson MJ, Nimmo MA (2004) The reproducibility of closed-pouch sweat collection and thermoregulatory responses to exercise-heat stress. *Eur J Appl Physiol* 91:748-751
- Hopkins WG (2003) A spreadsheet for analysis of straightforward controlled trials. In: Hopkins WG (ed) *Sportsciorg*. Vol. 2005
- Hopper MK, Coggan AR, Coyle EF (1988) Exercise stroke volume in relation to plasma-volume expansion. *J Appl Physiol* 64:404-408
- Joyner MJ, Dietz NM, Shepard JT (2001) From Belfast to Mayo and beyond: the use and future of plethysmography to study blood flow in humans limbs. *J Appl Physiol* 91:2431-2441
- Kampmann B, Brode P, Schutte M, Griefahn B (2008) Lowering of resting core temperature during acclimation is influenced by exercise stimulus. *Eur J Appl Physiol* 104:321-327

- Mack GW, Nadel ER (1996) Body fluid balance during heat stress in humans. In: Fregly MJ, Blatteis, C.M. (ed) Handbook of Physiology. Vol 1. American Physiological Society, Bethesda, MD, pp 187-284
- Maughan RJ (2003) Impact of mild dehydration on wellness and on exercise performance. *European Journal of Clinical Nutrition* 57:S19-S23
- McConell GK, Burge CM, Skinner SL, Hargreaves M (1997) Influence of ingested fluid volume on physiological responses during prolonged exercise. *Acta Physiol Scand* 160:149-156
- Morel F, Doucet A (1986) Hormonal control of kidney functions at the cell level. *Physiol Rev* 66:377-468
- Morris DJ (1981) The metabolism and mechanism of action of aldosterone. *Endocr Rev* 2:234-247
- Nadel ER, Fortney SM, Wenger CB (1980) Effect of hydration state on circulatory and thermal regulations. *J Appl Physiol Respir Environ Exercise Physiol* 49:715-721
- Nadel ER, Hales J, Strange S, Christensen JW, Saltin B (1993) Thermoregulation, exercise and thirst: interrelationships in humans. In: Gisolfi CV, Lamb DR (eds) *Exercise, heat and thermoregulation*. Brown and Benchmark, Dubuque, I.A., pp 225-251
- Nagashima K, Jauchia W, Stavros A, Kavouras A, Mack GW (2001) Increased renal tubular sodium reabsorption during exercise-induced hypervolemia in humans. *J Appl Physiol* 91:1229-1236

- Nielsen B, Hales JRS, Strange S, Christensen JW, Saltin B (1993) Human circulatory and thermoregulatory adaptations with heat acclimation and exercise in a hot, dry environment. *J Physiol* 460:467-485
- Nielsen B, Strange S, Christensen NJ, Warberg J, Saltin B (1997) Acute and adaptive responses in humans to exercise in a warm, humid environment. *Pfluger's Archiv* 434:49-56
- Noakes TD, Adams BA, Myburgh KH, Greef C, Lotz T, Nathan M (1988) The danger of inadequate water intake during prolonged exercise. *Eur J Appl Physiol* 57:210-219
- Pandolf KB (1998) Time course of heat acclimation and its decay. *Int J Sport Med* 19:S157-S160
- Patterson MJ, Stocks JM, Taylor NAS (2004) Sustained and generalised extracellular fluid expansion following heat acclimation. *J Physiol* 559:327-334
- Poliner LR, Dehmer GJ, Lewis SE, Parksey RW, Blomqvist CG, Willerson JT (1980) Left ventricular performance in normal subjects: a comparison of the responses to exercise in the upright and supine positions. *Circulation* 62:528-534
- Ramanathan LN (1964) A new weighing system for mean skin temperature of the human body. *J Appl Physiol* 19:531-533
- Regan JM, Macfarlane DJ, Taylor NAS (1996) An evaluation of the role of skin temperature during heat adaptation. *Acta Physiol Scand* 158:365-375
- Sawka MN, Francesconi RP, Drolet L, Young AJ (1987) Thirst and fluid intake following graded hypohydration levels in humans. *Physiol Behav* 40:229-236

- Sawka MN, Noakes TD (2007) Does dehydration impair exercise performance? *Med Sci Sports Exerc* 39:1209-1217
- Sawka MN, Wenger CB, Pandolf KB (1996) Thermoregulatory responses to acute exercise-heat stress and heat acclimation. In: Fregly. MJ, Blatteis, C.M. (ed) *Handbook of Physiology: American Physiological Society. Vol 1. Oxford University Press*, pp 157-185
- Sawka MN, Young AJ, Caderette BS, Levine L, Pandolf KB (1985) Influence of heat stress and acclimation on maximal aerobic power. *Eur J Appl Physiol* 53:294-298
- Senay LC, Mitchell D, Wyndam CH (1976) Acclimatization in a hot, humid environment: body fluid adjustments. *J Appl Physiol* 40:786-796
- Shido O, Sugimoto N, Tanabe M, Sakurada S (1999) Core temperature and sweating onset in humans acclimated to heat at a fixed daily time. *Am J Physiol*:R1095-R1101
- Strydom NB, Williams CG (1969) Effect of physical conditioning on the state of heat acclimation on Banto labourers. *J Appl Physiol* 27:262-265
- Szlyk PC, Sils I, Francesconi RN, Hubbard RW, Armstrong LE (1989) Effects on water temperature and flavouring on voluntary dehydration in man. *Physiol Behav* 45:639-647
- Taylor N (2000) Principles and practices of heat adaptation. *J Human-Environ Sys* 4:11-22

Turk J, Worsley DE (1974) A technique for the rapid acclimatisation to heat for the army.

Army Personnel Research Establishment, Ministry of Defence, Farnborough,

U.K., pp 1-15

Watson G, Judelson DA, Armstrong LE, Yeargin SW, Casa DJ, Maresh CM (2005)

Influence of diuretic-induced dehydration on competitive sprint and power

performance. *Med Sci Sports Exerc* 37:1168-1174

Weller AS, Harrison MH (2001) Influence of heat acclimation on physiological strain

during exercise-heat stress in men wearing clothing of limited water vapour

permeability. *J Physiol* 531:51P

Witney R (1953) The measurement of volume changes in human limbs. *J Physiol*

121:1211-1227

Wyndham CH (1973) The physiology of exercise under heat stress. *Ann Rev Physiol*

35:193-220

Wyndham CH, Benade AJA, Williams CG, Strydom NB, Goldin A, Heynes AJA (1968)

Changes in central circulation and body fluid spaces during acclimatization to the

heat. *J Appl Physiol* 25:586-593