

Short Reports

Eur Neurol 2004;52:250–251
DOI: 10.1159/000082370

Stent Grafting Resolved Brachial Plexus Neuropathy due to Cervical Arteriovenous Fistula

Vera Carina Zingler^a, Michael Strupp^a, Thomas Brandt^a, Karin Herrmann^b, Thomas E. Mayer^c

Departments of ^aNeurology, ^bRadiology and ^cNeuroradiology, University of Munich, Munich, Germany

We report a rare case of an iatrogenic vertebrovenous fistula presenting with a brachial plexus neuropathy and a cervical bruit. This arteriovenous fistula (AVF) was successfully occluded by implanting a stent graft into the right vertebral artery. After the stenting, the cervical bruit ceased and the radiculopathy slowly improved. This case highlights the usefulness of such new devices as covered stent grafts, and the therapeutic options for effectively treating AVFs while maintaining patency of the vertebral artery.

Case Report

Spinal AVFs are characterized by the presenting symptoms of tinnitus, vertebrobasilar ischemia [1, 2], myelopathy or subarachnoid hemorrhage [3]. Only a few cases of cervical AVF associated with radiculopathy have been reported so far, because AVF rarely occurs at this site [4–7]. We describe a case of progressive brachial neuropathy that was caused by a vertebrovenous AVF. The patient was successfully treated by stent grafting of the vertebral artery.

A 63-year-old woman was admitted to our hospital with a 3-month history of progressive pain and fluctuating paresthesia of the right arm. The patient history revealed that right-sided breast carcinoma had been treated by surgery and local radiotherapy 4 years previously. Five months before onset of the brachial neuropathy, she had undergone a liver transplantation for fulminant hepatitis A and had received several central venous lines on the right side.

The neurological examination revealed hypesthesia and paresthesia of the right upper limb with predominant involvement of the dermatomes C₆–C₈. The muscle strength as well as the reflexes of both upper and lower limbs were normal. A loud, continuous bruit was detected during auscultation of the right lateral neck. Electromyography showed that the amplitude of the compound action potential of the right median nerve was reduced. Moreover, there were signs of chronic denervation of the right interosseus muscle. Motor and sensory electroneurography were normal. A recording of the somatosensory-evoked potentials after stimulation of the right median nerve revealed a loss of the peripheral responses (N13/14). Doppler ultrasound showed a pathological flow in the vertebral artery on the right side. MRI of the cervical spine demonstrated ectatic veins with flow




Fig. 1. T₂-weighted axial MRI at the level of C₆/C₇. Flow voids of dilated veins (arrows) around the nerve routes on the right side.

voids in the neuroforamina on the right (fig. 1). Conventional angiography disclosed a direct vertebral artery fistula draining into the paravertebral venous plexus at the level of the root C₇ (fig. 2a).

The AVF was occluded by implanting a Jomed stent graft (4 mm × 1.2 cm) into the right vertebral artery (by T.E.M.). The stent was dilated to 6 mm proximally with a Maverick balloon (fig. 2b). No neurological complications occurred during or after the endovascular treatment. The cervical bruit ceased, and the brachialgia and paresthesia of the right arm slowly improved after the stenting.

We had first thought that the right-sided brachial plexus syndrome was either caused by direct metastatic spread or was a delayed complication of the prior local radiation therapy [8, 9]. However, the cervical bruit pointed to an AVF that angiography confirmed. This case of AVF was most likely iatrogenic, having occurred after an unintentional puncture of the vertebral artery during insertion of a central venous line. Most cervical AVFs are caused by trauma, but they can also occur spontaneously or in association with other vascular dysplasias [5]. The brachial neuropathy of our patient was probably due to a direct compression of the cervical roots by dilated draining veins; this was documented by MRI and angiography (fig. 1, 2a).




Fig. 2. **a** Angiography of the right vertebral artery. Ectatic proximal vertebral artery, pseudoaneurysm and direct fistula (arrow) with early filling of the paravertebral venous plexus. **b** Remodeling of the vertebral artery and occlusion of the AVF by the polytetrafluoroethylene-covered stent graft (arrows).

Whereas in the past the most common treatment of AVF consisted of embolization with microcoils or surgery, nowadays successful stent repair has been reported in a few cases [10–12]. In our patient, a coronary polytetrafluoroethylene-covered stent graft repair was sufficient. It also successfully maintained patency of the vertebral artery (fig. 2b). This procedure appears superior to surgery or embolization, since patency of the vertebral artery is maintained.

References

- 1 Vinchon M, Laurian C, George B, et al: Vertebral arteriovenous fistula: A study of 49 cases and review of the literature. *Cardiovasc Surg* 1994;2:359–369.
- 2 Waitzman A, Anderson J, Willinsky R: Endovascular management of vertebral arteriovenous fistulas: The Toronto experience. *J Otolaryngol* 1996;25:322–328.
- 3 Willinsky R, Ter Brugge K, Lasjaunias P, et al: The variable presentations of craniocervical and cervical dural arteriovenous malformations. *Surg Neurol* 1990;34:118–123.
- 4 Kohno M, Takahashi H, Ide K, et al: A cervical dural arteriovenous fistula in a patient presenting with radiculopathy. *J Neurosurg* 1996;84:119–123.
- 5 Taylor CG, Husami Y, Colquhoun IR, Byrne JV: Direct cervical vertebral arteriovenous fistula with radiculopathy and MRI changes resolving after successful endovascular embolisation: A report of two cases. *Neuroradiology* 2001;43:1118–1122.
- 6 Cahan LD, Higashida RT, Halbach VV, et al: Variants of radiculomeningeal vascular malformations of the spine. *J Neurosurg* 1987;66:333–337.
- 7 Beaujeux R, Reizine D, Casasco A, et al: Endovascular treatment of vertebral arteriovenous fistula. *Radiology* 1992;183:361–367.
- 8 Rubin DI, Schomberg PJ, Shepherd RF, Panneton JM: Arteritis and brachial plexus neuropathy as delayed complications of radiation therapy. *Mayo Clin Proc* 2001;76:849–852.
- 9 Fardin P, Lelli S, Negrin P, Maluta S: Radiation-induced brachial plexopathy: Clinical and electromyographical (EMG) considerations in 13 cases. *Electromyogr Clin Neurophysiol* 1990;30:277–282.
- 10 Priestley R, Bray P, Bray A, Hunter J: Iatrogenic vertebral arteriovenous fistula treated with a hemobahn stent-graft. *J Endovasc Ther* 2003;10:657–663.
- 11 Surber R, Werner GS, Cohnert TU, Wahlers T, Figulla HR: Recurrent vertebral arteriovenous fistula after surgical repair: Treatment with a self-expanding stent-graft. *J Endovasc Ther* 2003;10:49–53.
- 12 Gonzalez A, Mayol A, Gil-Peralta A, Gonzalez-Marcos JR: Endovascular stent-graft treatment of an iatrogenic vertebral arteriovenous fistula. *Neuroradiology* 2001;43:784–786.

Vera Zingler, MD
 Department of Neurology, University of Munich, Klinikum Grosshadern
 Marchioninistrasse 15, DE–81377 Munich (Germany)
 Tel. +49 89 7095 2585, Fax +49 89 7095 5584
 E-Mail vzingler@nro.med.uni-muenchen.de