

t a

d

o n

Submitted by Dmitry Konstantinovich Polyushkin

to the University of Exeter

as a thesis for the degree of

Doctor of Philosophy in Physics June 2013

This thesis is available for library use on the understanding that it is copyright material and that no quotation from the thesis may be published without proper acknowledgement.

I certify that all material in this thesis which is not my own work has been identified and that no material has previously been submitted and approved for the award of a degree by this or any other university. I lovingly dedicate this thesis to all my family, especially to my wonderful wife Yulia, who fills my life with joy and happiness; to my father Konstantin, mother Irina and sister Daria who have always provided me with their unconditional love and support through the years; to my grandparents Gennady and Nina and my aunt Ekaterina who with my parents have brought me up in love and care; and in memory of my dear grandparents Ivan and Nadezhda.

Abstract

In this thesis the interaction of ultrashort laser pulses with metal nanostructures is investigated via two different phenomena: coherent acoustic oscillations of nanoparticles and generation of THz pulses on metal surfaces.

Both of these effects rely on the collective oscillations of free conduction electrons in metal surfaces, plasmons. The field of plasmonics gained a great interest in the last twenty years due to the unique properties of these surface modes. It is the effects of the resonant response of plasmonic structures to incident electromagnetic wave, in particular, in visible and infrared bands and the concentration of the electromagnetic field in small subwavelength regions with significant enhancement of the incident field that make plasmonics so attractive for various applications, such as biochemical sensing, enhanced fluorescence, surface-enhanced Raman scattering, and second harmonic generation, amongst others.

Investigation of the coherent particle vibrations is performed using the pumpprobe technique which allows measurement of the transient transmission signals. The expansion and subsequent contraction of the nanoparticle following the ultrashort laser pulse excitation lead to a shift of the plasmon band which can be traced by transient spectroscopy. We have investigated the effect of the particle thickness on the frequency of the fundamental vibrational mode. In addition, we measured the vibrational particle response during the particle shape deformation, both symmetrical and asymmetrical.

Exploration of the THz generation phenomena on plasmonic structures was performed using THz time-domain spectroscopy, the method which allows tracing of the generated THz field in the time-domain. We were able for the first time to measure the THz pulses generated from arrays of metal nanoparticles. Our observations verify the role of the particle plasmon mode in the generation of THz pulses. In addition, by exploring the dependence of the THz emission on the femtosecond pulse intensity we showed a high nonlinearity in the THz generation mechanism. The experimental results were assessed in the context of a recently proposed model where the THz radiation is generated via the acceleration of the ejected electrons by ponderomotive forces.

3

To reveal another proposed mechanism of the THz generation from plasmonic structures, namely optical rectification, we investigated the THz generation and electron emission from the arrays of nanoparticles and nanoholes. Our results suggest that both mechanisms may contribute to generation of THz pulses from the same sample under different illumination conditions.

In addition to periodic arrays of nanoparticles and nanoholes, THz generation from random metal-dielectric films was investigated. The microstructuring of such films allowed selective THz frequency generation which was explained by a model of dipole THz emitters. In addition, the effects of low temperature and pressure on the THz generation efficiency were investigated.

Acknowledgements

I would like to thank a number of people who have given me help and support over the past four years during my PhD at the University of Exeter.

First, I would like to express my deepest gratitude to my supervisor Prof. Bill Barnes for his persistent support in many aspects of my research. Bill offers a unique blend of the personal and scientific qualities which could rarely be met. His patient encouragement, sage and helpful advices were so indispensable to me during my PhD. Unfortunately, there are no words to properly appreciate all that he has done for me for these years. Many thanks for everything!

I am also very grateful to Prof. Euan Hendry. Although not a formal supervisor of mine, he has spent so much time helping me in my work! Every time I contacted him he inspired me with his enthusiasm, encouraging further work and providing me with bright ideas and comprehensive support both experimentally and theoretically. Many ideas I have implemented belong to him.

Before acknowledging other people I would like to mention the contribution of Prof. Alex Savchenko. To him and to Bill I owe the opportunity to study in Exeter. It was Alex who initially invited me to do a PhD research in his group and after a problem with a scholarship offered my candidacy to Bill who kindly accepted it. Alex's passing was very tragic for all of us. Having occasionally a chance to contact him as my mentor, I have seen an image of a true scientist fascinated with his work and very attentive to colleges and especially to his students. I would also like to thank my second mentor Dr. Saverio Russo for his friendly student-mentor meetings and for our fruitful collaboration.

I am indebted as well to Dr. Chris Burrows for his help in the first two years of my PhD. He was the person who intensively supported me during the most difficult time of my study, the first year. I am very grateful to him for his introduction to the most of the equipment I have used during my work in Exeter. Many thanks to him also for our conversations and for his desire to improve my English. Another person with whom I was constantly contacted during my work is Dr. Edmund Stone. His help with THz setup and also contribution to the project, both experimental and through the computer modelling, are invaluable. I would also like to thank Dr. Pete Hale for his introduction to the laser lab facilities. I am grateful to Dr. Ehsan Ahmad for his great deal of help with NSL technique and the plasma etching system and to Dr. David Horsell for his support in AFM measurements.

I would like to thank the whole Electromagnetic Materials group for their input during the group meetings. Especially, I am grateful to Prof. Alastair Hibbins who was organising these meetings and Prof. Roy Sambles, the bright person and talented scientist, for his insightful suggestions. My gratitude also goes to those who shared an office with me during my PhD, Dr. Chris Burrows, Dr. Tomasz Trzeciak, Dr. Melita Taylor, Dr. Celia Butler and Rostislav Mikhaylovskiy. Thank you Slava for supporting me when I was doing physics problems demonstrating.

It is my pleasure as well to thank all the collaborators who contributed to my research, in particular Dr. Ivan Khrapach, James Milton and Dr. Salvatore Santandrea with whom I was working on the graphene projects; Dr. Péter Dombi, István Márton and Péter Rácz from Budapest who provided me with electron emission data; Emiliano Martins and Calvyn Howells from the University of St Andrews for their warm welcome and for accomodating me during my visit there; and Dr. Nicolas Vogel for his crescent samples.

In addition, I would like to thank the clean room and workshop technicians, especially Dave Anderson and Nick Cole, and also Chris Forrest, our system administrator, for his computer related issues support.

I would also like to acknowledge many people who contributed both personally and professionally during my stay in Exeter, especially Dr. Ivan Khrapach, Dr. Tim Khodkov, Dr. Alexander Markevich, and Dr. Volodymyr Kruglyak. Many thanks to Dr. Ivan Khrapach for keeping the clean room in properly clean condition.

Finally, and extremely importantly, I would like to thank those whose contribution to the thesis is not so visible but whose love and support were invaluable, my beloved wife, Mum, Dad, sister and everyone in my family. In addition, I very much appreciate the support of my friends, parishioners and clergy from Exeter Orthodox Church, especially Dr. Ivan Khrapach, Protodeacon Peter Scorer, Fr. John Marks and Nadya Keenan. Thanks to all of them!

6

Contents

Abstrac	t	. 3		
Acknowledgements				
List of Figures				
List of T	ables	20		
Abbrevi	ations	21		
Chapter	¹ Background physics	23		
1.1	Maxwell's equations	23		
1.2	The Drude model	25		
1.3	Volume and surface plasmons	26		
1.4	Excitation of SPP	29		
1.5	Localized surface plasmons: mathematical techniques	32		
1.6	Particles small compared with light wavelength	35		
Chapter	2 Methods	40		
2.1	Fabrication techniques	40		
2.1	.1 Electron-beam lithography	40		
2.1	.2 Nanosphere lithography	43		
2.2	Sample characterization	45		
2.2	.1 Scanning electron microscopy	45		
2.2	.2 Atomic force microscopy	46		
2.2	.3 Optical characterization	47		
2.3	Experimental techniques: time-resolved measurements	52		
2.3	.1 Pump-probe spectroscopy	53		
2.3	.2 THz time-domain spectroscopy	56		
Chapter 3 Acoustic oscillations59				
3.1	Two-temperature model	59		

3.2	Acc	oustic oscillations	51
3.3	Exp	perimental results6	\$4
3	.3.1	Intensity dependence6	5
3	.3.2	Thickness dependence	6
3	.3.3	Particle shape	0
3	.3.4	Asymmetrical NSL particles	'2
3	.3.5	Surrounding medum7	'4
3.4	Sur	nmary and conclusions7	'7
Chapt	ter 4 T	Hz generation from plasmonic structures7	'9
4.1	TH	z generation from metal surfaces7	'9
4.2	Ser	nicontinuous metal films8	10
4.3	TH	z generation from semicontinuous metal films8	3
4	.3.1	Film thickness dependence	34
4	.3.2	SERS measurements on silver SMFs8	39
4.4	TH	z generation from arrays of nanoparticles9)2
4	.4.1	LSPR resonance position, experimental results)2
4	.4.2	Particle thickness, computer modelling9)4
4.5	TH	z intensity dependence9)7
4.6	TH	z generation model9	9
4.7	Ang	gle dependence)3
4.8	Sur	nmary and conclusions10)4
Chapt	ter 5 T	Hz generation and electron emission from plasmonic structures 10)6
5.1	Intr	oduction10)6
5.2	Exp	perimental results)9
5	.2.1	Electron emission 11	1
5	.2.2	THz generation	3
5	.2.3	Computer modelling11	7

Contents

5.3	Discussion of the results	
5.4	Summary and conclusions 120	
Chapte	r 6 Temperature and pressure effects in THz generation from plasmonic	
structur	es	
6.1	Temperature effects 122	
6.1	.1 Temperature dependence of metal losses	
6.1	.2 THz generation efficiency as a function of temperature	
6.1	.3 Temperature dependence of the order of the THz generation	
pro	ocess	
6.1	.4 Pressure effects	
6.2	Summary and conclusions 133	
Chapte	r 7 Selective THz frequency generation from microstructured SMF 135	
7.1	Introduction 135	
7.2	Dipole model for THz emission pattern 137	
7.3	Angle dependence of THz generation	
7.4	Unstructured silver film142	
7.5	Structured silver film	
7.6	Summary and conclusions 145	
Conclu	sions and outlook146	
Publications and presentations150		
Referer	nces	