


Contents lists available at ScienceDirect

Physics Letters A

www.elsevier.com/locate/plaShort-time quantum propagator and Bohmian trajectories Maurice de Gosson ^{a,*}, Basil Hiley ^b^a Universität Wien, Fakultät für Mathematik, NuHAG, Wien 1090, Austria^b University of London, Birkbeck College, Theoretical Physics Unit, London WC1E 7HX, United Kingdom

ARTICLE INFO

Article history:

Received 13 May 2013

Received in revised form 16 August 2013

Accepted 19 August 2013

Available online 10 September 2013

Communicated by P.R. Holland

Keywords:

Generating function

Short-time action

Quantum potential

Bohmian trajectories

ABSTRACT

We begin by giving correct expressions for the short-time action following the work Makri–Miller. We use these estimates to derive an accurate expression modulo Δt^2 for the quantum propagator and we show that the quantum potential is negligible modulo Δt^2 for a point source, thus justifying an unfortunately largely ignored observation of Holland made twenty years ago. We finally prove that this implies that the quantum motion is classical for very short times.

© 2013 The Authors. Published by Elsevier B.V. All rights reserved.

1. Introduction

In exploring the WKB limit of quantum theory, Bohm [2] was the first to notice that although one starts with all the ambiguities about the nature of a quantum system, the first order approximation fits the ordinary classical ontology. By that we mean that the real part of the Schrödinger equation under polar decomposition of the wave function becomes the classical Hamilton–Jacobi equation in the limit where terms involving \hbar are neglected. In contrast to this approach, in this Letter we show that the classical trajectories arise from a short-time quantum propagator when terms of $O(\Delta t^2)$ can be neglected. This fact was actually already observed by Holland some twenty years ago: In page 269 of his book [6] infinitesimal time intervals are considered whose sequence constructs a finite path. It is shown that along each segment the motion is classical (negligible quantum potential), and that it follows that the quantum path may be decomposed into a sequence of segments along each of which the classical action is a minimum. The novel contribution of the present Letter is an improved proof of Holland's result using an improved version of the propagator due to Makri and Miller [9,10]. (See also de Gosson [3] for a further discussion.)

* This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

* Corresponding author. Tel.: +43 0 6767877995.

E-mail addresses: maurice.degosson@gmail.com, maurice.de.gosson@univie.ac.at (M. de Gosson).

Now it is well known that explicit approximate expressions for the short-time action already play an essential role in various aspects of quantum mechanics (for instance the Feynman path integral, or semi-classical mechanics), and so does the associated Van Vleck determinant. Unfortunately, as already observed by Makri and Miller [9,10], these expressions, while giving the correct results for long time behavior, are not accurate enough to allow us to explore the short-time propagator rigorously. It is actually worse, the literature seems to be dominated by formulas which Makri and Miller show are wrong even to the first order of approximation!

These results have enabled us to provide precise estimates for the short-time Bohmian quantum trajectories for an initially sharply located particle. We will see that these trajectories are classical to the second order in time, due to the vanishing of the quantum potential for small time intervals.

In this Letter we sidestep the philosophical and ontological debate around the “reality” of Bohm's trajectories and rather focus on the mathematical issues.

2. Bohmian trajectories

Consider a time-dependent Hamiltonian function

$$H(x, p, t) = \sum_{j=1}^n \frac{p_j^2}{2m_j} + U(x, t) \quad (1)$$

and the corresponding quantum operator

$$\hat{H}(x, -i\hbar\nabla_x, t) = \sum_{j=1}^n \frac{-\hbar^2}{2m_j} \frac{\partial^2}{\partial x_j^2} + U(x, t). \quad (2)$$

The associated Schrödinger equation is

$$i\hbar \frac{\partial \Psi}{\partial t} = \widehat{H}(x, -i\hbar \nabla_x, t) \Psi, \quad \Psi(x, 0) = \Psi_0(x). \quad (3)$$

Let us write Ψ in polar form $Re^{iS/\hbar}$; here $R = R(x, t) \geq 0$ and $S = S(x, t)$ are real functions. On inserting $Re^{iS/\hbar}$ into Schrödinger's equation and separating real and imaginary parts, one sees that the functions R and S satisfy, at the points (x, t) where $R(x, t) > 0$, the coupled system of non-linear partial differential equations

$$\frac{\partial S}{\partial t} + \sum_{j=1}^n \frac{1}{2m_j} \left(\frac{\partial S}{\partial x_j} \right)^2 + U(x, t) - \sum_{j=1}^n \frac{\hbar^2}{2m_j R} \frac{\partial^2 R}{\partial x_j^2} = 0, \quad (4)$$

$$\frac{\partial R^2}{\partial t} + \sum_{j=1}^n \frac{1}{m_j} \frac{\partial}{\partial x_j} \left(R^2 \frac{\partial S}{\partial x_j} \right) = 0. \quad (5)$$

The crucial step now consists in recognizing the first equation as a Hamilton–Jacobi equation, and the second as a continuity equation. In fact, introducing the quantum potential

$$Q^\Psi = - \sum_{j=1}^n \frac{\hbar^2}{2m_j R} \frac{\partial^2 R}{\partial x_j^2} \quad (6)$$

(Bohm and Hiley [2]) and the velocity field

$$v^\Psi(x, t) = \left(\frac{1}{m_1} \frac{\partial S}{\partial x_1}, \dots, \frac{1}{m_n} \frac{\partial S}{\partial x_n} \right) \quad (7)$$

Eqs. (4) and (5) become

$$\frac{\partial S}{\partial t} + H(x, \nabla_x S, t) + Q^\Psi(x, t) = 0, \quad (8)$$

$$\frac{\partial \rho}{\partial t} + \operatorname{div}(\rho v^\Psi) = 0, \quad \rho = R^2. \quad (9)$$

The main postulate of the Bohmian theory of motion is that particles follow quantum trajectories, and that these trajectories are the solutions of the differential equations

$$\dot{x}_j^\Psi = \frac{\hbar}{m_j} \operatorname{Im} \frac{1}{\Psi} \frac{\partial \Psi}{\partial x_j}. \quad (10)$$

The phase space interpretation is that the Bohmian trajectories are determined by the equations

$$\dot{x}_j^\Psi = \frac{1}{m_j} p_j^\Psi, \quad \dot{p}_j^\Psi = - \frac{\partial U}{\partial x_j}(x^\Psi, t) - \frac{\partial Q^\Psi}{\partial x_j}(x^\Psi, t). \quad (11)$$

It is straightforward to check that these are just Hamilton's equations for the Hamiltonian function

$$H^\Psi(x, p, t) = \sum_{j=1}^n \frac{p_j^2}{2m_j} + U(x, t) + Q^\Psi(x, t) \quad (12)$$

which can be viewed as a perturbation of the original Hamiltonian H by the quantum potential Q^Ψ (see Holland [7,8] for a detailed study of quantum trajectories in the context of Hamiltonian mechanics).

The Bohmian equations of motion are *a priori* only defined when $R \neq 0$ (that is, outside the nodes of the wave function); this will be the case in our constructions since for sufficiently small times this condition will be satisfied by continuity if we assume that it is case at the initial time.

An important feature of the quantum trajectories defined above is that they cannot cross; thus there will be no conjugate points like those that complicate the usual Hamiltonian dynamics.

3. The short-time propagator

The solution Ψ of Schrödinger's equation (3) can be written

$$\Psi(x, t) = \int K(x, x_0; t) \Psi_0(x_0) dx_0 \quad (13)$$

where the kernel K is the quantum propagator:

$$K(x, x_0; t) = \langle x | \exp(-i\widehat{H}t/\hbar) | x_0 \rangle. \quad (14)$$

Schrödinger's equation (3) is then equivalent to

$$i\hbar \frac{\partial K}{\partial t} = \widehat{H}(x, -i\hbar \nabla_x, t) K, \quad K(x, x_0; 0) = \delta(x - x_0) \quad (15)$$

where δ is the Dirac distribution. Physically this equation describes an isotropic source of point-like particle emanating from the point x_0 at initial time $t_0 = 0$. We want to find an asymptotic formula for K for short time intervals Δt . Referring to the usual literature (see, e.g., Schulman [11]), such approximations are given by expressions of the type

$$K(x, x_0; \Delta t) = \left(\frac{1}{2\pi i\hbar} \right)^{n/2} \sqrt{\rho(x, x_0; \Delta t)} \exp\left(\frac{i}{\hbar} S(x, x_0; \Delta t) \right)$$

where $S(x, x_0; \Delta t)$ is the action along the classical trajectory from x_0 to x in time Δt and

$$\rho(x, x_0; \Delta t) = \det \left(- \frac{\partial^2 S(x, x_0; \Delta t)}{\partial x_j \partial x_k} \right)_{1 \leq j, k \leq n}$$

is the corresponding Van Vleck determinant. We will need precise short-time behavior of the action. In this regard Makri and Miller [9,10] have shown that the asymptotic expression for the generating function is given by

$$S(x, x_0; \Delta t) = \sum_{j=1}^n \frac{m_j}{2\Delta t} (x_j - x_0)^2 - \widetilde{U}(x, x_0) \Delta t + O(\Delta t^2) \quad (16)$$

where $\widetilde{U}(x, x_0, 0)$ is the average value of the potential over the straight line joining x_0 at time t_0 to x at time t with constant velocity:

$$\widetilde{U}(x, x_0) = \int_0^1 U(\lambda x + (1 - \lambda)x_0, 0) d\lambda. \quad (17)$$

For instance when

$$H(x, p) = \frac{1}{2m} (p^2 + m^2 \omega^2 x^2)$$

is the one-dimensional harmonic oscillator formula (16) yields the correct expansion

$$S(x, x_0; t) = \frac{m}{2\Delta t} (x - x_0)^2 - \frac{m\omega^2}{6} (x^2 + xx_0 + x_0^2) \Delta t + O(\Delta t^2), \quad (18)$$

the latter can of course be deduced directly from the exact value

$$S(x, x_0; t, t_0) = \frac{m\omega}{2 \sin \omega \Delta t} ((x^2 + x_0^2) \cos \omega \Delta t - 2xx_0) \quad (19)$$

by expanding $\sin \omega \Delta t$ and $\cos \omega \Delta t$ for $\Delta t \rightarrow 0$.

Introducing the following notation,

$$\widetilde{S}(x, x_0; \Delta t) = \sum_{j=1}^n m_j \frac{(x_j - x_0)^2}{2\Delta t} - \widetilde{U}(x, x_0) \Delta t, \quad (20)$$

leads us to the Makri and Miller approximation (formula (17c) in [9]) for the short-time propagator:

$$K(x, x_0; \Delta t) = \left(\frac{1}{2\pi i\hbar}\right)^{n/2} \sqrt{\rho(x, x_0; \Delta t)} \exp\left(\frac{i}{\hbar} \tilde{S}(x, x_0; \Delta t)\right) + O(\Delta t^2) \tag{21}$$

where

$$\rho(x, x_0; \Delta t) = \det\left(-\frac{\partial^2 \tilde{S}(x, x_0; \Delta t)}{\partial x_j \partial x_{0,k}}\right)_{1 \leq j, k \leq n}$$

It turns out that this formula can be somewhat improved. The Van Vleck determinant $\rho(x, x_0; \Delta t)$ is explicitly given, taking formula (20) into account, by

$$\rho(x, x_0; \Delta t) = \det\left(-\frac{1}{\Delta t} M - \tilde{U}''_{x, x_0}(x, x_0) \Delta t\right)$$

where M is the mass matrix (the diagonal matrix with positive entries the masses m_j) and

$$\tilde{U}''_{x, x_0} = \left(-\frac{\partial^2 \tilde{U}(x, x_0)}{\partial x_j \partial x_k}\right)_{1 \leq j, k \leq n}$$

Writing

$$\begin{aligned} &\left(-\frac{1}{\Delta t} M - \tilde{U}''_{x, x_0}(x, x_0) \Delta t\right) \\ &= -\frac{1}{\Delta t} M [I_{n \times n} - M^{-1} \tilde{U}''_{x, x_0}(x, x_0) \Delta t^2] \\ &= -\frac{1}{\Delta t} M [I_{n \times n} + O(\Delta t^2)], \end{aligned}$$

we have by taking the determinant of both sides

$$\rho(x, x_0; \Delta t) = \frac{m_1 \cdot \dots \cdot m_n}{(\Delta t)^n} \det(I_{n \times n} + O(\Delta t^2)).$$

Noting that $\det(I_{n \times n} + O(\Delta t^2)) = 1 + O(\Delta t^2)$, we thus have

$$\rho(x, x_0; \Delta t) = \frac{m_1 \cdot \dots \cdot m_n}{(\Delta t)^n} (1 + O(\Delta t^2)). \tag{22}$$

Writing

$$\tilde{\rho}(\Delta t) = \frac{m_1 \cdot \dots \cdot m_n}{(\Delta t)^n} \tag{23}$$

which is just the Van Vleck density for the free particle Hamiltonian. We thus have

$$\rho(x, x_0; \Delta t) = \tilde{\rho}(\Delta t) (1 + O(\Delta t^2)) \tag{24}$$

and hence we can rewrite formula (21) as

$$K(x, x_0; \Delta t) = \left(\frac{1}{2\pi i\hbar}\right)^{n/2} \sqrt{\tilde{\rho}(\Delta t)} \exp\left(\frac{i}{\hbar} \tilde{S}(x, x_0; \Delta t)\right) + O(\Delta t^2). \tag{25}$$

We will see below that this formula allows an easy study of the quantum potential for K .

4. Short-time Bohmian trajectories

Let us determine the quantum potential Q corresponding to the propagator $K = K(x, x_0; t)$ using the asymptotic formulas above. Recall that it describes an isotropic source of point-like particle emanating from the point x_0 at initial time $t_0 = 0$. We have, by definition,

$$Q = -\sum_{j=1}^n \frac{\hbar^2}{2m_j \sqrt{\rho}} \frac{\partial^2 \sqrt{\rho}}{\partial x_j^2}$$

which we can rewrite

$$Q = -\frac{\hbar^2}{2} \frac{M^{-1} \nabla_x \cdot \nabla_x \sqrt{\rho}}{\sqrt{\rho}}$$

where M is the mass matrix defined above. We have, using (24),

$$\sqrt{\rho} = \sqrt{\tilde{\rho}(\Delta t)} (1 + O(\Delta t^2))$$

and hence

$$\frac{\partial^2 \sqrt{\rho}}{\partial x_j^2} = O((\Delta t)^2).$$

From this it follows that the quantum potential associated with the propagator satisfies

$$Q(x, x_0; \Delta t) = O(\Delta t^2). \tag{26}$$

The discussion above suggests that the quantum trajectory of a sharply located particle should be identical with the classical (Hamiltonian) trajectory for short times. Let us show this is indeed the case. If we want to monitor the motion of such a particle, we have of course to specify its initial momentum which gives its direction of propagation at time $t_0 = 0$; we set

$$p(0) = p_0. \tag{27}$$

In view of formula (10), the trajectory in position space is obtained by solving the system of differential equations

$$\dot{x} = \hbar \operatorname{Im} \frac{M^{-1} \nabla_x K}{K}, \quad x(0) = x_0. \tag{28}$$

Replacing K with its approximation

$$\tilde{K}(x, x_0; \Delta t) = \left(\frac{1}{2\pi i\hbar}\right)^{n/2} \sqrt{\tilde{\rho}(\Delta t)} \exp\left(\frac{i}{\hbar} \tilde{S}(x, x_0; \Delta t)\right)$$

we have, since $K - \tilde{K} = O(\Delta t^2)$ in view of (25),

$$\dot{x} = \hbar \operatorname{Im} \frac{M^{-1} \nabla_x \tilde{K}}{\tilde{K}} + O(\Delta t^2).$$

A straightforward calculation, using the expression (20) for the approximate action $\tilde{S}(x, x_0; \Delta t)$, leads to the equation

$$\dot{x}(\Delta t) = \frac{x(\Delta t) - x_0}{\Delta t} - M^{-1} \nabla_x \tilde{U}(x(\Delta t), x_0) \Delta t + O(\Delta t^2) \tag{29}$$

(cf. the proof of Lemma 248 in [3]). This equation is singular at time $t = 0$ hence the initial condition $x(0) = x_0$ is not sufficient for finding a unique solution; this is of course consistent with the fact that (29) describes an arbitrary particle emanating from x_0 ; to single out one quantum trajectory we have to use the additional condition (27) giving the direction of the particle at time $t = 0$ (see the discussion in Holland (1993) [6, §6.9]). We thus have

$$x(\Delta t) = x_0 + M^{-1} p_0 \Delta t + O(\Delta t^2);$$

in particular $x(\Delta t) = x_0 + O(\Delta t)$ and hence, by continuity,

$$\nabla_x \tilde{U}(x(\Delta t), x_0) = \nabla_x \tilde{U}(x_0, x_0) + O(\Delta t).$$

Let us calculate $\nabla_x \tilde{U}(x_0, x_0)$. We have, taking definition (17) into account,

$$\nabla_x \tilde{U}(x, x_0) = \int_0^1 \lambda \nabla_x U(\lambda x + (1 - \lambda)x_0, 0) d\lambda$$

and hence

$$\nabla_x \tilde{U}(x_0, x_0) = \int_0^1 \lambda \nabla_x U(x_0, 0) d\lambda = \frac{1}{2} \nabla_x U(x_0, 0).$$

We can thus rewrite Eq. (29) as

$$\dot{x}(\Delta t) = \frac{x(\Delta t) - x_0}{\Delta t} - \frac{1}{2} M^{-1} \nabla_x U(x_0, 0) \Delta t + O(\Delta t^2).$$

Let us now differentiate both sides of this equation with respect to Δt :

$$\ddot{x}(t) = \frac{x(t) - x_0}{(\Delta t)^2} + \frac{\dot{x}(t)}{\Delta t} - \frac{1}{2} M^{-1} \nabla_x U(x_0, 0) + O(\Delta t) \quad (30)$$

that is, replacing $\dot{x}(\Delta t)$ by the value given by (29),

$$\dot{p}(t) = M \ddot{x}(t) = -\nabla_x U(x_0, 0) + O(\Delta t). \quad (31)$$

Solving this equation we get

$$p(t) = p_0 - \nabla_x U(x_0, 0) \Delta t + O(\Delta t^2). \quad (32)$$

Summarizing, the solutions of the Hamilton equations are given by

$$x(\Delta t) = x_0 + \frac{p_0}{m} \Delta t + O(\Delta t^2), \quad (33)$$

$$p(\Delta t) = p_0 - \nabla_x U(x_0, 0) \Delta t + O(\Delta t^2). \quad (34)$$

These equations are, up to the error terms $O(\Delta t^2)$ the equations of motion of a classical particle moving under the influence of the potential U ; there is no trace of the quantum potential, which is being absorbed by the terms $O(\Delta t^2)$. The motion is thus identical with the classical motion on time scales of order Δt^2 .

5. Conclusion

This result puts Bohm's original perception, which led him to the causal interpretation, on a firm mathematical footing. He writes [1]

Indeed it had long been known that when one makes a certain approximation (WKB) Schrödinger's equation becomes equivalent to the classical Hamilton–Jacobi theory. At a certain point I asked myself: What would happen, in the demonstration of this equivalence, if we did not make this approximation? I saw immediately that there would be an additional potential, representing a kind of force, that would be acting on the particle.

The source of this “force” was the quantum potential. In our approach we see that while any classical potential acts immediately,

the quantum potential does not. From this fact two consequences follow.

Firstly, it provides a rigorous treatment of the “watched pot” effect. If we keep observing a particle that, if unwatched would make a transition from one quantum state to another, will now no longer make that transition. The unwatched transition occurs when the quantum potential grows to produce the transition. Continuously observing the particle does not allow the quantum potential to develop so the transition does not take place. We will not discuss this effect further here as it has been reported in detail in [4]. For an other application see Section 6.9 in Bohm and Hiley [2].

Secondly, in the situation when the quantum potential decreases continuously with time, the quantum trajectory continuously deforms into a classical trajectory [5]. This means that there is no need to appeal to decoherence to reach the classical domain.

Acknowledgements

Maurice de Gosson has been supported by a research grant from the Austrian Research Agency FWF (Projektnummer P23902-N13).

The authors wish to thank the referees for useful remarks and suggestions, and for having pointed out misprints.

References

- [1] D. Bohm, *Hidden variables and the implicate order*, in: B.J. Hiley, D.F. Peat (Eds.), *Quantum Implications: Essays in Honour of David Bohm*, Routledge, London, 1987, p. 35.
- [2] D. Bohm, B.J. Hiley, *The Undivided Universe: An Ontological Interpretation of Quantum Theory*, Routledge, London, New York, 1993.
- [3] M. de Gosson, *The Principles of Newtonian and Quantum Mechanics. The Need for Planck's Constant \hbar* , Imperial College Press, London, 2001.
- [4] M. de Gosson, B.J. Hiley, *Zeno paradox for Bohmian trajectories: the unfolding of the metatron*, arXiv:1001.2622.
- [5] B.J. Hiley, A.H. Aziz Mufti, *The ontological interpretation of quantum field theory applied in a cosmological context*, in: M. Ferrero, A. van der Merwe (Eds.), *Fundamental Problems in Quantum Physics*, Kluwer, Dordrecht, 1995, pp. 141–156.
- [6] P.R. Holland, *The Quantum Theory of Motion: An Account of the de Broglie–Bohm Causal Interpretation of Quantum Mechanics*, Cambridge University Press, 1993.
- [7] P. Holland, *Hamiltonian theory of wave and particle in quantum mechanics I: Liouville's theorem and the interpretation of the de Broglie–Bohm theory*, *Nuovo Cimento B* 116 (2001) 1043–1070.
- [8] P. Holland, *Hamiltonian theory of wave and particle in quantum mechanics II: Hamilton–Jacobi theory and particle back-reaction*, *Nuovo Cimento B* 116 (2001) 1143–1172.
- [9] N. Makri, W.H. Miller, *Correct short time propagator for Feynman path integration by power series expansion in Δt* , *Chem. Phys. Lett.* 151 (1988) 1–8.
- [10] N. Makri, W.H. Miller, *Exponential power series expansion for the quantum time evolution operator*, *J. Chem. Phys.* 90 (1989) 904–911.
- [11] L.S. Schulman, *Techniques and Applications of Path Integration*, Wiley, New York, 1981.