

nind



Up Will. Research

> **104** December 1992

# SYMPTOM RESPONSE OF SORGHUM HYBRIDS INFECTED BY SUGARCANE MOSAIC VIRUS STRAIN MDMV-B D. L. Seifers and K. D. Kofoid Fort Hays Branch Experiment Station

Sorghum [Sorghum bicolor (L.) Moench] infected by sugarcane mosaic virus strain MDMV-B (SCMV-MDMV-B) (formerly maize dwarf mosaic virus strain B) expresses variable symptoms depending on plant genotype and temperature. Infection causes a yellow to dark green pattern (mosaic), and a necrotic reaction (red-leaf) develops on certain genotypes when the temperature drops below 21°C (69°F). Hybrids expressing the redleaf symptom show greater reduction in yield. Immunity to SCMV-MDMV-B is not present in commercially available hybrid sorghums.

## Procedure

Studies were conducted at the Fort Hays Branch Experiment Station to determine symptom response of sorghum hybrids infected by SCMV-MDMV-B both under field conditions (2 years of testing) and in growth chambers. Sorghum seed was planted in 6.1 m rows. When at the two- to three-leaf stage, plants in half of each row were inoculated, and plants in the remaining half of the

AGRICULTURAL EXPERIMENT STATION Kansas State University, Manhattan Marc A. Johnson, Director row served as healthy controls. Plants were inoculated with a Devilbiss EGA-502 spray gun attached to a 3.5 L (1 gal) inoculum reservoir. Compressor line pressure was maintained at 7.0 kg/cm<sup>2</sup> (100 lbs/in<sup>2</sup>). Symptom notes were recorded at biweekly intervals during the growing season beginning 2 weeks after inoculation. Planting was delayed until late June to assure cool temperatures for induction of red-leaf necrosis.

In growth chamber tests, sorghum hybrids were planted in metal flats with 15 seeds/entry and 33 entries/flat. Seedlings were inoculated at the two-leaf stage (on second leaf) with a Devilbiss atomizer at the same pressure used for field inoculations. Inoculated plants were maintained in growth chambers at  $27 \pm 2^{\circ}$ C ( $80 \pm 2^{\circ}$ F) with 14 hr of fluorescent light. Control plants, inoculated with buffer and abrasive only, were maintained in the same environment. The presence of mosaic symptoms was recorded 14 days following inoculation. The temperature of the growth chambers then was decreased to  $15^{\circ}$ C ( $59^{\circ}$ F). Four days later, plants were rated for the presence of red-leaf necrosis. Growth chamber tests were conducted once.

### Results

Reactions of hybrids are shown in Table 1. SS = inoculated leaf and two to three leaves above the inoculated lea have necrotic streaks and spots that often coalesce and cover 1-25 percent of a leaf. New leaves above these show few or no symptoms, so that at maturity, no symptoms of any type occur in the upper leaves. RL1 = seedling and juvenile plants have a necrotic red-leaf reaction consisting of various widths of streaks that often coalesce and cover 25-50 percent of a leaf. At maturity, the flag leaf and the subtending 3-4 leaves have a few necrotic streaks covering 1-10 percent of a leaf. RL2 = seedling and juvenile plants have a severe necrotic red-leaf reaction, not in streaks or spots, but continuous necrosis covering 80-100 percent of a leaf. In plants that survive to maturity, any new leaf will develop a like necrosis with onset of cool temperatures. M = plants having only a mosaic reaction. Current data indicate that plants with the M reaction suffer less yield loss than plants developing necrosis (RL tupes).

Brand	Hybrid	Symptom Response*
Agrigene	AG2470W	SS
Agrigene	AG2725	RL2
Agripro	AP 9250	SS
oripro	AP 9830	SS
arinro	AP 9850	M
arinro	ST 686	M
gripro	ST D701G	PI 2
Subio	A201	SS
egrow	Madora	
sgrow	Ospan	141
egrou	Songer	M
agrow	Tomas	
layofill	1000	
argiii \	1099	RLZ
argiii	5/5	55
argill	607E	55
argill	618Y	M
Cargill	630	SS
argill	837	SS
argill	847	М
asterline	SR 305	М
asterline	SR 314EC	RL2
asterline	SR 315E	RL2
asterline	SR 319E	RL2
asterline	SR 324E	RL1
ekalb	DK-39Y	RL2
)ekalb	DK-40Y	SS
ekalb	DK-41Y	SS
ekalb	DK-48	SS
ekalb	DK-56	М
Vekalb	DK-66	RL2
Vekalb	M-572	RL2
Velange	DSA 117	М
elange	DSA 121	SS
elange	DSA 131	RL2
)elange	DSA 141C	SS
elange	DSA 922	RL2
eltapine	1506	RL2
eltapine	G-1492	SS
eltapine	G-1616	SS
eltapine	G-550E	RL1
wNA	DN 31	RI 1
ontanelle	5550	RL2
ontanelle	5590	RI 2
ontanelle	W 5000	SS
arrison	SG-651	50
arrison	SG-821	55
arrison	SG_0/9	00 0 10
aret	5210	
laret	2200	M
aret	0072 5502	1v1
rui ai Iarrat	5503	M DI O
Jarst	2211	KLZ

Ĺ

		Symptom
Brand	Hybrid	Response
Garst	5522Y	SS
Garst	5681	SS
Golden Acres	T-E 76	М
Golden Acres	T-E 77-E	RL2
Golden Acres	T-E Dinero	RL2
Golden Acres	T-E Eden	М
Golden Acres	T-E Rio	RL2
Golden Acres	T-E Wahoo	SS
Golden Acres	T-E Y-60	SS
Golden Acres	T-E Y-75	М
Golden Acres	T-E Y-101G	RL1
Golden Harvest	H-388W	SS
Golden Harvest	H-444W	SS
Golden Harvest	H-514B	RL2
Golden Harvest	H-515E	SS
GRI	890111	RL2
Groagri	GSC 3146	М
Groagri	GSC 3148	М
Groagri	GSC 3150	RL2
Groagri	GSC 3605	RL1
Hoegemeyer	6636	RL1
Hoegemeyer	671	SS
Hoegemeyer	6744	SS
Hoegemeyer	6878	RL2
Hoegemeyer	688	RL2
Horizon	200Y	SS
Horizon	213Y	SS
Horizon	216G	M
Horizon	76G	M
Hyperformer	1255 DR	RL2
Hyperformer	Honcho	RL2
Hyperformer	HSC 1289C	M
Hyperformer	HSC Cherokee	M
Jacques	Z11 277 U	M
Jacques	377-W	55
Jacques		M DI 1
NC+	000E	
NC+	270	RLI CC
NC+	071 179	33
NC+	7R81	55
NC+	7001	55
NC+	V363	55
Northrun-King	2656	PI 2
Northrup-King	KS-383Y	SS
Northrup-King	KS-555Y	SS
Northrup-King	KS-710	RI 2
Northrup-King	KS-714Y	SS
Northrup-King	KS-737	ŝŝ
Northrup-King	S-9740Y	ŝŝ
Northrup-King	S 9750	RL2

# Table 1. Reactions of Sorghum Hybrids to Infection by Sugarcane Mosaic Virus Strain MDMV-B

		Symptom
Brand	Hybrid	Response*
Ohlde	120C	SS
Ohlde	134	SS
Ohlde	139A	RL2
Ohlde	140W	SS
Ohlde	246y	RL1
Oro	Alpha	М
Oro	Amigo	RL2
Oro	Baron	М
Oro	Edge	М
Oro	G XTRA	RL2
Oro	Hombre	М
Oro	Ivory	SS
Oro	Ultra	RL2
Pioneer	8231Y	RL1
Pioneer	8260	SS
Pioneer	8358	RL1
Pioneer	8379	SS
Pioneer	8500	SS
Pioneer	8536Y	SS
Pioneer	8557Y	М
Pioneer	8601	RL1
Pioneer	8696Y	М
Pioneer	8699	М
Pioneer	8771	М
SSI	SBP001	RL1
SSI	SBP005	RL1
SSI	SBP011	SS
Stine	SM 68 BR	RL2
Stine	SM 69	М
Stine	SM 74	RL2
Stine	SM 75R	RL2
Terra	TR 4580	RL1
Terra	TR 4620	RL2
Terra	TR 4637Y	SS
Triumph	TR 46	Μ
Triumph	TR 50yG	М
Triumph	TR 58Y	SS
Triumph	TR 60-G	М
Triumph	TR 65-G+	RL1
Triumph	TR 74CR	SS
Triumph	TWO 64yG	RL2
Triumph	TWO 80-D	RL2
Warner	W-624-Y	RL2
Warner	W-625-Y	RL2
Warner	W-902-W	М
Warner	W-917-E	М
Wilson	515W	SS
Wilson	522W	SS
Wilson	535Y	RL2
Wilson	622E	Μ

<sup>a</sup>Symptom responses are the same for both field and growth chamber tests.

## Acknowledgements

This research was partially funded by a grant from the Kansas Sorghum Commission.

Contribution no. 93-145-S from the KAES.



### Agricultural Experiment Station Kansas State University Manhattan 66506–4008

Keeping Up With Research 104

December 1992

Kansas State University is committed to a policy of nondiscrimination on the basis of race, sex, national origin, handicap, religion, age, sexual orientation, or other nonmetil reasons, in admissions, educational programs or activities, and employment, all as required by applicable laws and regulations. Responsibility for coordination of compliance efforts and receipt of inquiries, including those concerning Title IX of the Education Amendments of 1972 and Section 504 of the Rehabilition Ant of 1973, has been delegated to Jane D. Rowlett, Ph.D., Director, Aflirmative Action Office, 214 Anderson Hall, Kansas State University, Manhattan, Kansas 66506-0104, (013/532-6220). 1M