

# *Ab initio* rotation-vibration energy levels of triatomics to spectroscopic accuracy

Jonathan Tennyson,<sup>a,1</sup> Paolo Barletta,<sup>a</sup> Maxim A Kostin,<sup>a b</sup>  
Oleg L Polyansky<sup>a b</sup> and Nikolai F Zobov<sup>a b</sup>

<sup>a</sup>*Department of Physics and Astronomy, University College London, London, WC1E 6BT, UK*

<sup>b</sup>*Permanent address: Institute of Applied Physics, Russian Academy of Science, Uljanov Street 46, Nizhnii Novgorod, Russia 603024.*

The factors that need to be taken into account to achieve spectroscopic accuracy for triatomic molecules are considered focusing on  $\text{H}_3^+$  and water as examples. The magnitude of the adiabatic and non-adiabatic corrections to the Born-Oppenheimer approximation is illustrated for both molecules, and methods of including them *ab initio* are discussed. Electronic relativistic effects are not important for  $\text{H}_3^+$ , but are for water for which the magnitude of the various effects are discussed. For  $\text{H}_3^+$  inclusion of rotational non-adiabatic effects means that levels can be generated to an accuracy approaching  $0.01 \text{ cm}^{-1}$ ; for water the error is still dominated by the error in the correlation energy in the electronic structure calculation. Prospects for improving this aspect of the calculation are discussed.

---

<sup>1</sup> Author to whom correspondence should be addressed. email: [j.tennyson@ucl.ac.uk](mailto:j.tennyson@ucl.ac.uk)

## 1 Introduction

It has become standard in the quantum chemistry literature to describe calculations with an error of  $1\text{ cm}^{-1}$  as being of ‘spectroscopic accuracy’. Yet high resolution spectroscopy routinely measures rotation-vibration spectra to accuracies of  $0.001\text{ cm}^{-1}$  or even better. Indeed spectra of even light triatomics such as  $\text{H}_3^+$  [1] or water [2] can have tens or hundreds of transitions per  $\text{cm}^{-1}$ . An accuracy of  $1\text{ cm}^{-1}$  has practical convenience because, as discussed below, going beyond these limits requires the consideration of a number of effects which are neglected in standard *ab initio* formulations.

Spectroscopy not only provides a detailed probe of molecular structure and interactions, it also provides data for numerous applications. Some applications, particularly those which involve the spectra of hot ( $T > 1000\text{ K}$ ) species, require datasets which are too large for their laboratory determination to be possible. For example recent compilations of water rotation-vibration lines considered well in excess of  $10^8$  individual transitions [3–5]. Perforce these datasets have to be constructed computationally but at the same time high accuracy is required. Thus the cited linelists, despite being based on very sophisticated models, are still found to be inadequate for models of cool ( $T \sim 3000\text{ K}$ ) oxygen-rich stars [6,7]. The ability to compute reliable linelists thus relies not only on developing computational methods capable of yielding large amounts of data required but also on constructing high accuracy models. This has led us to explore in detail what is actually required to compute spectra of small molecules to real spectroscopic accuracy, ie at least better than  $0.01\text{ cm}^{-1}$ .

We have performed high accuracy *ab initio* studies for a number of systems including  $\text{H}_3^+$ ,  $\text{H}_2\text{O}$ ,  $\text{H}_2\text{S}$  [8] and  $\text{HCN}$  [9]. In this article we only consider results for the first two of these molecules, which are important prototypes. In particular  $\text{H}_3^+$  is the one triatomic system for which the Born-Oppenheimer electronic structure problem has been solved to spectroscopic accuracy [10] and water, besides being particularly important, has been found to be sensitive, perhaps unusually so, to effects often neglected in standard models, see for example [11].

## 2 $\text{H}_3^+$

$\text{H}_3^+$  is probably unique in that essentially all spectral assignments made have relied on first principles calculations based on the use of high accuracy potential energy surfaces and variational nuclear motion calculations [12]. The first assigned spectrum of  $\text{H}_3^+$ , the infrared emissions from its bending fundamen-

tal [13], relied heavily on the *ab initio* calculations of Carney and Porter [14]. Table 1 shows the improvement in electronic structure calculations since these pioneering calculations, concentrating only on those works that gave potential energy surfaces suitable for nuclear motion calculations.

For reliable calculations of vibration-rotation spectra the location of the minimum of the potential and the shape of the potential about the minimum is more important than the absolute error in the potential. Thus Carney and Porter [15] made excellent predictions despite an absolute error approaching the value of the quantities they were predicting. Similarly Meyer *et al.* (MBB) [18] used careful error control techniques to produce a potential which proved highly successful for spectral assignments [12]. Interestingly the potential of Frye *et al.* [19] actually gives results for vibrational band origins closer to the experimental ones than the near-exact potential of Cencek *et al.* [10]. This is due to fortuitous cancellation of errors: including corrections to the Born-Oppenheimer approximation with Frye *et al.*'s potential actually moves their results away from the observed ones.

Table 1 shows an improvement by at least a factor 20 for the potentials computed within the last decade. The common feature of these more recent potentials is the use of explicitly correlated wavefunctions *ie* ones which include the  $r_{12}$  electron-electron coordinate in the wavefunction in some form. The potential of Cencek *et al* [10] is so accurate that it makes a natural starting point for further investigations. Indeed these workers also calculated the relativistic correction to the electronic motion for  $\text{H}_3^+$ . They found this to be approximately  $3 \text{ cm}^{-1}$  but to vary little with the geometry of the molecule, see Table III of [10]. This correction therefore makes little significant contribution, less than  $0.01 \text{ cm}^{-1}$ , to the vibration-rotation spectrum of  $\text{H}_3^+$ .

The same cannot be said for corrections to the Born-Oppenheimer approximation. Table 2 shows the sensitivity of the vibrational band origins of  $\text{H}_3^+$  and its isotopomers to both adiabatic and non-adiabatic corrections to the Born-Oppenheimer calculation. It is relatively easy to calculate the adiabatic correction which is also known as the Born-Oppenheimer diagonal correction (BODC) as second derivatives of the electronic wavefunction with respect to displacements of the nuclear positions [21]. The original calculation of BODC surfaces for  $\text{H}_3^+$  [22] and its mixed isotopomers [23] used self-consistent field (SCF) wavefunctions and showed that the adiabatic correction was important at the  $1 \text{ cm}^{-1}$  level. More recently Cencek *et al.* [10] recalculated the BODC using their highly correlated wavefunctions. Their calculations give an effect  $13 \text{ cm}^{-1}$  larger than the SCF calculations, but found this shift to be nearly uniform with  $\text{H}_3^+$  geometry, see Table II of [10]. Use of the improved BODC changes the calculated vibrational band origins by less than  $0.1 \text{ cm}^{-1}$ .

Inclusion of non-adiabatic corrections to the Born-Oppenheimer approxima-

tion is far from simple. Recently Schwenke has developed a method which involves estimating the effect by including the coupling due to excited states by summing over a complete set of one-electron excitations [25]. This method has yet to be widely used and has not yet been applied to  $\text{H}_3^+$ . The most common approach for diatomic systems is not to attempt such complete summations but to model the non-adiabatic correction to the vibrational motion by using an effective vibrational reduced mass for the system,  $\mu^V$  [27], whose value should lie between that determined using atomic masses,  $\mu^A$ , and nuclear masses,  $\mu^N$ . Polyansky and Tennyson [28] adopted this approach and effective masses obtained by Moss [29] for  $\text{H}_2^+$  and its isotopomers. The success of this approach is shown by the excellent results obtained in the final column of Table 2.

Polyansky and Tennyson [28] also concluded that the non-adiabatic correction to the rotational motion was small for low values of the rotational quantum number,  $J$ , they considered. They therefore adopted a rotational mass,  $\mu^R$ , equal to the nuclear mass,  $\mu^N$ . However they found that use of  $\mu^V \neq \mu^R$  led to an extra term in the Hamiltonian, which, for  $\text{H}_3^+$ , is small but important for obtaining smooth results. It should be noted that the method adopted by Polyansky and Tennyson involves identifying vibrational and rotational motions, and that different ways of making this identification by, for example using different embedding of the body-fixed rotational axes, will lead to different results. It has long been known that the optimal separation between vibrational and rotational motions is obtained by using the Eckart conditions [30].

Use of the Eckart conditions is not standard in most internal coordinate based variational calculations since they lead to considerably more complicated Hamiltonians, see [31,32]. However for  $\text{H}_3^+$  it is possible to get an embedding close to the Eckart one by fixing the  $z$ -axis perpendicular to the plane of the molecules and noting that, for the equilibrium equilateral triangle structure of  $\text{H}_3^+$ , all orientations of axes within the plane of the molecule are equivalent. Recently Kostin *et al.* [33] have implemented a  $z$ -perpendicular embedding and used it to tackle the  $\text{H}_3^+$  problem. They obtained rotational energy levels up to  $J \leq 15$ , the full range for which there is experimental data [34].

These calculations, because of their higher accuracy and larger range of  $J$ 's considered, show the clear signature of a small contribution due to non-adiabatic effects. This contribution can be estimated approximately as  $0.003J(J+1) - 0.002K^2 \text{ cm}^{-1}$ , where  $K$  is the projection of  $J$  on the molecular  $z$ -axis.

Some time ago Bunker and Moss [35] made a rather complete formulation of the non-adiabatic contributions to the vibration-rotation problem for triatomic molecules. In this formulation they showed that the matrix elements which

determine the the non-adiabatic correction to the rotational motion can be directly related to the electronic contribution to the rotational g-factor,  $g^e$ . For  $\text{H}_3^+$  these factors are rather small but nevertheless have been calculated *ab initio* [36]. Use of these *ab initio*  $g^e$  in our calculations almost completely account for the observed rotational non-adiabatic effect. Full results of this work will be reported elsewhere.

### 3 Water

Until recently spectral analysis of the water molecule relied heavily on the use of effective Hamiltonians based on perturbation theory. However improvements in the theory of variational calculations led to the assignment of spectra which had defied traditional analysis [2] leading to something of a paradigm shift in the analysis of water spectra [37]. The use of variational methods has emphasized the need for accurate, *ab initio* procedures.

Water spectra has long been used as a testbed for the accuracy of *ab initio* methods [38]. Table 3 charts the progress in *ab initio* predictions of the vibrational band origins of water since the original, pioneering variational calculations of Bucknell and Handy [39]. Although the error in the fundamentals has improved by about a factor of about 100 since then, there is some way to go before spectroscopic accuracy can be achieved. We will return to this point in section 4.

The simple comparison of the fundamentals computed with a non-relativistic electronic structure methods with experiment is actually misleading. Calculations by Császár and co-workers [11,44] have shown that the electronic relativistic corrections are surprising large for water and therefore cannot be ignored. The result of including this effect is to significantly improve the predictions for the stretching fundamentals (and indeed overtones [11]), to within  $1 \text{ cm}^{-1}$ , but at the expense of worsening the agreement with the bending overtones. This finding is very much in line with error analysis of Partridge and Schwenke [3] who found their *ab initio* model to be significantly poorer for bending than stretching modes; however it suggests that the excellent agreement with experiment obtained in the non-relativistic, force field calculations of Martin *et al.* [42] is actually fortuitous.

The finding that the first-order relativistic correction to the electron kinetic energy, strictly the one-electron mass-velocity plus Darwin (MVD1) corrections, is significant has led to the investigation of other high-order relativistic effects. In particular Quiney *et al.* [46] found that the two-electron kinetic contribution via the two-electron Darwin term (D2) was fairly small, but the relativistic correction to the Coulomb potential, represented by the Breit in-

teraction or the simpler, approximate Gaunt term, often contributed more than  $1 \text{ cm}^{-1}$  to the higher vibrational band origins and therefore should not be neglected. Perhaps even more intriguingly an estimate of the quantum electrodynamic Lamb shift suggests that this too can contribute changes of the order  $1 \text{ cm}^{-1}$  to the vibrational band origins and a similar amount to the  $J = 20$  rotational levels [47]. Conversely recent calculations have shown that spin-orbit interactions, which might have been thought to be significant in the region of linear geometries, can safely be neglected [48].

Inclusion of non-Born-Oppenheimer effects can play an important contribution to the spectrum of water. Zobov *et al.* [49] computed an SCF adiabatic or BODC surface. In contrast to  $\text{H}_3^+$ , where use of such a surface resolved most of the problems of treating the vibrations of the isotopomers, Zobov *et al.* found that the BODC did not explain differences between the various isotopomers of water although its use did lead to significant shifts in the  $J = 20$  energy levels. More recently Schwenke [26] has recalculated the BODC using a correlated wavefunction calculated at the Complete Active Space Self Consistent Field (CASSCF) level. He found that this correction is on average  $8 \text{ cm}^{-1}$  larger than the SCF level calculation, and that the new surface showed significant differences to surface calculated at the SCF level.

Unfortunately Schwenke’s fit to his CASSCF BODC data extrapolates very poorly outside the relatively small region defined by his *ab initio* calculations. Use of this surface with our codes gave unstable results. We have therefore refitted his CASSCF data to the functional form

$$\Delta V^{\text{ad}}(r_1, r_2, \theta) = \sum_{i,j,k} C_{i,j,k} \left[ \frac{1}{2}(r_1 + r_2) - r_e \right]^i [\cos(\theta) - \cos(\theta_e)]^j \left[ \frac{1}{2}(r_1 - r_2) \right]^k (1)$$

where  $r_1$  and  $r_2$  are the two O–H bondlengths and  $\theta$  is the bondangle. Equilibrium is given by  $r_e = 1.80965034 \text{ a}_0$  and  $\theta_e = 1.82404493$  radians. The coefficients of this fit, which reproduces the original data with a standard deviation of  $0.01 \text{ cm}^{-1}$ , are given in table 4. This surface is no longer unstable but its behaviour for large values of the stretching coordinates are not well constrained by the *ab initio* data.

Table 5 shows the effect of the various BODC surfaces on the lower vibrational band origins. The BO results in this table used the *ab initio* surface of Partridge and Schwenke [3], which includes, separately, both valence correlation and core correlation effects. The calculations all used nuclear masses except for the penultimate column where  $\mu^V$  is defined, as previously [11,49], by the O atomic mass and an H mass midway between the atomic and nuclear mass. Use of our re-fitted BODC surface, while not giving the same results as Schwenke’s surface, does support his finding that there are significant difference between the BODC surface calculated at the SCF level and that

calculated using a CASSCF model. The two surfaces give similar behaviour for the bending modes but significant difference with respect to stretching excitations.

The main purpose of Schwenke’s work [26] was to develop a model for including non-adiabatic corrections in the nuclear motion problem. His method involved explicitly calculating full, three-dimensional coupling surfaces and is therefore a much more complete method than the use of mass scaling discussed for  $\text{H}_3^+$  above. Interestingly Schwenke found that use of the relatively simple, vibrationally averaged, diagonal components of his coupling surfaces gave results very similar to those of his full calculation. The final column of Table 4 presents results obtained by Schwenke using his full non-adiabatic correction. We have implemented a simplified, diagonal version of this correction within our vibration-rotation program DVR3D [50]. In this we have only considered two non-adiabatic parameters, the ones which scale the second derivative (or kinetic energy terms) in  $\theta$  and  $r_i$ . These results are compared to ones computed using the simple mass scaling procedure used for  $\text{H}_3^+$ .

Non-adiabatic effects, as calculated either by mass scaling or more explicitly are significantly larger for water than for  $\text{H}_3^+$ . This finding is in line with previous predictions [49,51]. The simplified two-term diagonal treatment of the non-adiabatic problem gives results in excellent agreement with Schwenke’s full calculation; however the use of mass scaling to include non-adiabatic effects predicts shifts of similar magnitude to the accurate treatment but differs considerably in detail. The mass scaling method systematically over-estimates non-adiabatic effects for the bending motion and under-estimates them for the stretches. Since the mass scaling results are insensitive to the choice of O mass used [49], this method essentially involves the change of a single parameter, the H mass. Schwenke’s diagonal approximation, while very greatly simplified compared to his full treatment, involves 4 parameters for the vibrational motion, plus a constant which does not affect band origins. We have used only two of these constants, with no noticeable further loss of accuracy. These are the parameters that scale the bending and stretching kinetic energy operators and are the same parameters that are scaled in the mass scaling method, where they are constrained to the same value. Table 6 presents results for the higher band origins of water. Schwenke does not present results for these states so no comparison is made with his work.

## 4 Discussion and conclusions

From the high accuracy study of only two molecules, which show markedly contrasting sensitivity to the inclusion of effects normally neglected in the *ab initio* treatment of vibration-rotation spectra, it would be unwise to draw

too many conclusions. However a number of the observations made below are supported by other studies, in particular our high accuracy work on H<sub>2</sub>S [8] and HCN [9]. Table 7 presents a summary of the influence of each effect. In each case the largest shift, or an estimate of the largest shift, is given. It should be noted that in compiling this table only band origins for which experimental data are available have been considered. Since many more vibrational bands are known for water than H<sub>3</sub><sup>+</sup>, this leads to some bias in the magnitude of the corrections towards the water case. In particular the adiabatic corrections for H<sub>3</sub><sup>+</sup> are actually larger for the low-lying states, the only ones for which a direct comparison can be made.

As a purely hydrogenic system, it is probably not surprising that relativistic effects are found to be largely unimportant for H<sub>3</sub><sup>+</sup>. Conversely, Császár *et al.* [44] found that the potential energy surface of water showed a particularly strong sensitivity to relativistic effects. This they ascribed to the re-hybridization of the O electrons as the molecule changes from a bent to linear geometries. Although other systems containing first row elements do not seem to show the same sensitivity to electronic relativistic effects as water [9,44], these effects are certainly not negligible. Methods for calculating these effects are becoming readily available. Unsurprisingly, relativistic effects in the heavier H<sub>2</sub>S system are larger than those in water [8]. Studies on whether higher order relativistic corrections are important in other systems, particularly H<sub>2</sub>S, are presently underway.

Corrections to the Born-Oppenheimer approximation are probably only important for high accuracy work on molecules containing H atoms. The adiabatic or Born-Oppenheimer diagonal correction is relatively easy to calculate *ab initio* [21]. For H<sub>3</sub><sup>+</sup> the adiabatic correction to the Born-Oppenheimer approximation is more important than the non-adiabatic correction. For water the two effects have similar magnitude. As yet there is no other triatomic system for which similar comparisons can be made but when these results are combined with studies of diatomics [29,52,53], it would appear to be a general property of only pure hydrogen systems that the adiabatic correction is the most important correction to the Born-Oppenheimer approximation.

We have compared a number of models for including non-adiabatic corrections. The full treatment by Schwenke [26] represents a significant advance, but it is to be hoped that it will not generally prove necessary to calculate the large number of coupling surfaces involved to model what is a fairly small effect. It is therefore encouraging that the two parameter diagonal reduction of Schwenke's full treatment gives such good results. This method gives significantly better results than the single parameter mass scaling which works well for H<sub>3</sub><sup>+</sup>. It is unclear at this stage whether the single parameter method works well for H<sub>3</sub><sup>+</sup> but not water because (a) the non-adiabatic effects are smaller in H<sub>3</sub><sup>+</sup> so a lower accuracy is acceptable, (b) the higher symmetry of H<sub>3</sub><sup>+</sup> means that only



one parameter is required, (c) the lack of available experimental data on the band origins  $H_3^+$  or (d) because of some other reason.

While there is a Born-Oppenheimer electronic potential for  $H_3^+$  [10] of unprecedented accuracy, the Born-Oppenheimer potential remains the major source of error for water and is likely to be the main error in a similar study for any other triatomic molecule. Partridge and Schwenke [3] expended considerable efforts in both trying to produce well converged results and to analyze the errors in their results. Their conclusion, which is supported by the study of Császár *et al.* [44], is that the major source of error is due to the lack of convergence of the basis set used to represent valence correlation in the bending coordinate. That the main error with the BO potential is in the bending coordinate is consistent with our error analysis of the vibrational energy levels.

It would seem unlikely that this problem will be resolved by simply using larger correlation consistent (CC) basis sets since such calculations are likely to suffer from problems with linear dependence. However there are two new approaches which offer hope that further improvements in the Born-Oppenheimer potential may well be achievable in the near future. One is the careful use of extrapolation techniques to accelerate the convergence of such studies with respect to basis set size [44] and level of electron correlation [54,55]; such methods has already been shown to yield satisfactory results for  $H_2S$  [8,55]. The other method is the use of wavefunctions which explicitly include the electron-electron coordinate [56,57]. Such methods are known to lead to accelerated convergence of the basis set required to represent correlation effects and have already been demonstrated to give high accuracy for HF [58] which is a ten electron system like water. It is therefore to be hoped that an *ab initio* solution to the vibration-rotation energy levels of water to spectroscopic accuracy, along the lines of the one that has already been achieved for  $H_3^+$ , will be achievable in the fairly near future. This will represent a major triumph for *ab initio* quantum chemistry.

Finally it should be noted that besides the obvious goal of achieving spectroscopic accuracy by *ab initio* procedures, there is another important use of improved *ab initio* procedures. Potential energy surfaces derived from fitting to spectroscopic data are now becoming a standard tool for the analysis of spectra of small molecules. Our experience shows that the quality of such potentials improves significantly with the quality of the *ab initio* starting point.

## Acknowledgements

We thank Attila G. Császár and David W. Schwenke for helpful discussions and for communicating results prior to publication. Funding from the Royal

Society, EPSRC, the British Council and the Russian Fund for Science is also gratefully acknowledged.

## References

- [1] A. Carrington and R.A. Kennedy, *J. Chem. Phys.* 81 (1984) 91.
- [2] O. L. Polyansky, N. F. Zobov, S. Viti, J. Tennyson, P. F. Bernath and L. Wallace, *Science*, 277 (1997) 346.
- [3] H. Partridge and D.W. Schwenke, *J. Chem. Phys.*, 106 (1997) 4618.
- [4] S. Viti, PhD Thesis, University of London (1997).
- [5] U.G. Jorgenssen, P. Jensen, G.O. Sorensen and B. Aringer, *Astron. Astrophys.*, 372 (2001) 249.
- [6] F. Allard, P.H. Hauschildt and D. Schwenke, *Astrophys. J.*, 540 (2000) 1005.
- [7] H.R.A. Jones, Y. Pavlenko, S. Viti and J. Tennyson, *Mon. Not. R. Astr. Soc.* (submitted).
- [8] G. Tarczay, A.G. Császár, O.L. Polyansky and J. Tennyson, *J. Chem. Phys.*, , in press.
- [9] T. van Mourik, G.J. Harris, O.L. Polyansky, J. Tennyson, A.G. Császár and P.J. Knowles, *J. Chem. Phys.*, in press.
- [10] W. Cencek, J. Rychlewski, R. Jaquet and W. Kutzelnigg, *J. Chem. Phys.*, 108 (1998) 2831.
- [11] A.G. Császár, J.S. Kain, O.L. Polyansky, N.F. Zobov and J. Tennyson, *Chem. Phys. Letts.*, 293 (1998) 317; 312 (1999) 613.
- [12] J. Tennyson, *Rep. Prog. Phys.*, 58 (1995) 421.
- [13] T. Oka, *Phys. Rev. Lett.*, 45 (1980) 531.
- [14] G.D. Carney and R.N. Porter, *Phys. Rev. Lett.*, 45 (1980) 537.
- [15] G.D. Carney and R.N. Porter, *J. Chem. Phys.*, 65 (1976) 3547.
- [16] R. Schinke, M. Dupius and W.A. Lester, *J. Chem. Phys.*, 72 (1980) 3909.
- [17] P.G. Burton, E. von-Nagy-Felsobuki, G. Doherty and M. Hamilton, *Mol. Phys.*, 55 (1985) 527.
- [18] W. Meyer, P. Botschwina and P.G. Burton, *J. Chem. Phys.*, 84 (1986) 981.
- [19] G.C. Lie and D. Frye, *J. Chem. Phys.*, 96 (1992) 6784.
- [20] R. Röhse, W. Kutzelnigg, R. Jaquet and W. Klopper, *J. Chem. Phys.*, 101 (1994) 2231 (1994)

- [21] N.C. Handy, Y. Yamaguchi and H.F. Schaefer III, *J. Chem. Phys.*, 84 (1986) 4481.
- [22] B.M. Dinelli, C.R. Le Sueur, J. Tennyson and R.D. Amos, *Chem. Phys. Letters* 232 (1995) 295.
- [23] O.L. Polyansky, B.M. Dinelli, C.R. Le Sueur and J. Tennyson, *J. Chem. Phys.*, 102 (1995) 9322.
- [24] B.M. Dinelli, L. Neale, O.L. Polyansky and J. Tennyson, *J. Mol. Spectrosc.*, 181 (1997) 142.
- [25] D.W. Schwenke, *J. Chem. Phys.*, 114 (2001) 1693.
- [26] D.W. Schwenke, *J. Phys. Chem. A*, 105 (2001) 2352.
- [27] P.R. Bunker and R.E. Moss, *Mol. Phys.*, 33 (1977) 417.
- [28] O.L. Polyansky and J. Tennyson, *J. Chem. Phys.*, 110 (1999) 5056.
- [29] R.E. Moss, *Mol. Phys.*, 89 (1996) 195.
- [30] C. Eckart, *Phys. Rev.*, 47 (1935) 552.
- [31] C.R. Le Sueur, S. Miller, J. Tennyson and B.T. Sutcliffe, *Mol. Phys.*, 76 (1992) 1147.
- [32] H. Wei and T. Carrington Jr., *Chem. Phys. Lett.*, 287 (1998) 289.
- [33] M.A. Kostin, O.L. Polyansky and J. Tennyson, *J. Chem. Phys.*, to be submitted.
- [34] C.M. Lindsay and B.J. McCall, *J. Mol. Spectrosc.*, in press.
- [35] P.R. Bunker and R.E. Moss, *J. Mol. Spectrosc.*, 80 (1980) 217.
- [36] J. Oddershede and J.R. Sabin, *Chem. Phys.*, 122 (1988) 291.
- [37] O.L. Polyansky, J. Tennyson and N.F. Zobov, *Spectrochimica Acta*, 55A (1999) 659.
- [38] B.T. Sutcliffe and J. Tennyson, *J. Chem. Soc., Faraday Trans. 2*, 83 (1987) 1663.
- [39] M.G. Bucknell and N.C. Handy, *Mol. Phys.*, 28 (1974) 759.
- [40] R.J. Bartlett, I. Shavitt and G.D. Purvis III, *J. Chem. Phys.*, 71 (1979) 281.
- [41] P.J. Knowles, S.J. Saxon and N.C. Handy, *Chem. Phys.*, 69 (1982) 305.
- [42] J.M.L. Martin, J.P. Francois and R. Gijbels, *J. Chem. Phys.*, 96 (1992) 7633.
- [43] G.S. Kedziora and I. Shavitt, *J. Chem. Phys.* 106 (1997) 8733.
- [44] A.G. Császár, W.D. Allen and H.F. Schaefer III, *J. Chem. Phys.*, 108 (1998) 9751.

- [45] J. Tennyson, N.F. Zobov, R. Williamson, O.L. Polyansky and P.F. Bernath, J. Phys. Chem. Ref. Data, (in press).
- [46] H.M. Quiney, P. Barletta, G. Tarczay, A.G. Császár, O.L. Polyansky and J. Tennyson, Chem. Phys. Letters, (in press).
- [47] P. Pyykkö, K.G. Dyall, A.G. Császár, G. Tarczay, O.L. Polyansky and J. Tennyson, Phys. Rev. A., 63 (2001) 024502.
- [48] G. Tarczay, A. G. Császár, W. Klopper, H. M. Quiney, Mol. Phys., submitted.
- [49] N.F. Zobov, O. L. Polyansky, C.R. Le Sueur and J. Tennyson, Chem. Phys. Letters, 260 (1996) 381.
- [50] J. Tennyson, J.R. Henderson and N.G. Fulton, Computer Phys. Comms., 86 (1995) 175.
- [51] O.L. Polyansky, P. Jensen, and J. Tennyson, J. Chem. Phys., 105 (1996) 6490.
- [52] N.C. Handy and A.M. Lee, Chem. Phys. Letts., 252 (1996) 425.
- [53] C. Schwartz and R.J. Le Roy, J. Mol. Spectrosc., 121 (1987) 420.
- [54] A.G. Császár and M.L. Leininger, J. Chem. Phys., 114 (2001) 5491.
- [55] A.G. Császár G. Tarczay, M.L. Leininger, O.L. Polyansky, J. Tennyson and W.D. Allen, in 'Molecules in Space', NATO ASI (2001).
- [56] W. Klopper, K. Bak, P. Jorgensen, J. Olsen and T. Helgaker, J. Phys. B: At. Mol. Phys., 32 (1999) R103.
- [57] E.F. Valeev, W.D. Allen, H.F. Schaefer III and A.G. Császár, J. Chem. Phys., 114 (2001) 2875.
- [58] H. Müller, R. Frank, S. Vogtner, R. Jaquet and W. Kutzelnigg, Theor. Chem. Acc. 100 (1998) 85.

Table 1

Minimum electronic energy,  $E_{\min}$ , as a function of time for  $\text{H}_3^+$  *ab initio* potential energy surfaces.  $\Delta E$  is the approximate error in the absolute electronic energy at the minimum.

Reference	Year	$E_{\min} / E_h$	$\Delta E / \text{cm}^{-1}$
Carney & Porter [15]	1976	-1.33519	1900
Schinke <i>et al.</i> [16]	1980	-1.34023	790
Burton <i>et al.</i> [17]	1985	-1.34188	430
Meyer <i>et al.</i> [18]	1986	-1.34309	160
Lie & Frye [19]	1992	-1.343828	9
Röhse <i>et al.</i> [20]	1994	-1.3438336	1
Cencek <i>et al.</i> [10]	1998	-1.3438355	0.04

Table 2

Vibrational band origins<sup>a</sup>, in  $\text{cm}^{-1}$ , for  $\text{H}_3^+$  and its observed isotopomers [28]. Results are given as observed – calculated for various models<sup>b</sup>.

	$E_{\text{obs}}$	BO	$+\Delta V_{ad}$	$\mu^V \neq \mu^N$
$\text{H}_3^+$				
$01^1$	2521.409	-0.11	-0.24	0.056
$10^0$	3178.290	-1.30	-0.40	0.025
$02^0$	4778.350	0.00	-0.50	0.020
$02^2$	4998.045	-0.30	-0.64	0.010
$11^1$	5554.155	-1.40	-0.50	0.000
$\text{H}_2\text{D}^+$				
$\nu_1$	2992.505	-1.46	-0.36	-0.020
$\nu_2$	2205.869	-0.47	-0.25	-0.050
$\nu_3$	2335.449	0.47	-0.14	0.090
$\text{D}_2\text{H}^+$				
$\nu_1$	2736.981	-1.04	-0.28	0.001
$\nu_2$	1968.169	0.58	-0.11	0.023
$\nu_3$	2078.430	-0.74	-0.18	-0.004

<sup>a</sup> Experimentally derived data [24]

<sup>b</sup> Models defined as follows:

BO: *Ab initio* BO potential + relativistic surface.

$+\Delta V_{ad}$ : as BO plus adiabatic correction surface,  $\Delta V_{ad}$ .

$\mu^V \neq \mu^N$ : as  $+\Delta V_{ad}$  with vibrational mass,  $\mu^V$ , greater than the nuclear mass,  $\mu^N$ .

Table 3

*Ab initio* predictions of the vibrational fundamentals of water, in  $\text{cm}^{-1}$ , as a function of year

Reference	Method <sup>a</sup>	Year	$\nu_2$	$\nu_1$	$\nu_3$
Bucknell & Handy [39]	SCF	1974	1728	4045	4139
Bartlett <i>et al.</i> [40]	MBBT	1979	1610	3702	3789
Knowles <i>et al.</i> [41]	CASSCF	1982	1645	3691	3794
Martin <i>et al.</i> [42]	QCISD(T)	1992	1595	3657	3756
Kedziora & Shavitt [43]	MRCISD	1997	1604.6	3650.5	3758.2
Partridge & Schwenke (PS) [3]	CCSD(T)	1997	1597.4	3660.5	3757.2
PS + relativistic correction [11]		1998	1598.2	3657.7	3755.3
Experiment [45]			1594.75	3657.05	3755.93

<sup>a</sup> For basis sets and explanations of methods see original publication. Note that all band origins are based on the use of variational nuclear motion calculations except those of Martin *et al.*, who used vibrational perturbation theory.

Table 4

Coefficients for the H<sub>2</sub>O Born-Oppenheimer Diagonal Correction (BODC) surface, see eq. (1). The coefficients are in cm<sup>-1</sup> for bondlengths in a<sub>0</sub> and angles in radians.

<i>i</i>	<i>j</i>	<i>k</i>	$C_{ijk}$	<i>i</i>	<i>j</i>	<i>k</i>	$C_{ijk}$
0	0	0	2745.1055606459	3	1	0	-26.4139825271
1	0	0	-59.4634051943	1	3	0	6.0253244059
0	1	0	52.4468059982	2	2	0	96.0354596696
2	0	0	209.0157347415	2	0	2	400.8671847238
0	2	0	-30.9569390720	0	2	2	64.3311129088
0	0	2	222.8220666155	1	1	2	-159.3984157719
1	1	0	-32.0954739957	5	0	0	26.5108357242
3	0	0	-84.5603232875	0	5	0	27.9528744878
0	3	0	28.3636764469	4	1	0	3.3262475087
2	1	0	-82.9317097732	1	4	0	23.3582782463
1	2	0	74.5702444206	1	0	4	-62.8395365075
1	0	2	-301.3739913256	0	1	4	33.1944328259
0	1	2	0.4721526758	3	2	0	31.8104538207
4	0	0	17.0778691448	3	0	2	-11.0876529850
0	4	0	9.6100487312	2	3	0	-151.9675922959
0	0	4	118.9501245531	0	3	2	-33.6314521634



Table 5

Band origins, in  $\text{cm}^{-1}$ , for water showing the effects of introducing non-Born-Oppenheimer effects. Comparison with results of Schwenke [26]

$(v_1, v_2, v_3)$	BO <sup>a</sup>	Adiabatic correction <sup>b</sup>			Non-adiabatic correction <sup>c</sup>		
		SCF	CASSCF		$\mu^V \neq \mu^N$	diag <sup>d</sup>	accurate
		[49]	[26]	refit			
(010)	1597.60	-0.50	-0.46	-0.46	-0.19	-0.06	-0.07
(020)	3157.14	-0.99	-0.94	-0.93	-0.38	-0.12	-0.15
(100)	3661.00	-0.06	0.39	0.55	-0.46	-0.72	-0.70
(030)	4674.88	-1.48	-1.46	-1.43	-0.55	-0.18	-0.23
(110)	5241.83	-0.51	-0.01	0.16	-0.65	-0.77	-0.76
(040)	6144.64	-2.02	-2.03	-2.00	-0.71	-0.23	-0.30
(120)	6784.56	-0.94	-0.42	-0.23	-0.83	-0.83	-0.84
(200)	7208.80	-0.08	0.96	1.25	-0.88	-1.39	-1.37
(002)	7450.86	0.19	1.22	1.47	-0.90	-1.47	-1.57
(050)	7555.62	-2.65	-2.71	-2.71	-0.84	-0.28	-0.37
(130)	8286.03	-1.37	-0.87	-0.67	-1.00	-0.89	-0.91
(210)	8771.71	-0.47	0.66	0.95	-1.07	-1.44	-1.43
(012)	9008.72	-0.20	0.90	1.17	-1.08	-1.52	-1.65

<sup>a</sup> BO: *ab initio* Born-Oppenheimer potential of Partridge and Schwenke [3].

<sup>b</sup> Adiabatic correction given relative to BO.

<sup>c</sup> Non-adiabatic correction given relative to the refitted CASSCF BODC.

<sup>d</sup> Simplified diagonal correction, see text, multiplied by 1.1 as recommended by Schwenke [26].

Table 6

Band origins for higher vibrational states, in  $\text{cm}^{-1}$ , for water showing the effects of introducing non-Born-Oppenheimer effects. Footnotes are as given in Table 5.

$(v_1, v_2, v_3)$	BO <sup>a</sup>	Adiabatic correction <sup>b</sup>		Non-adiabatic correction <sup>c</sup>	
		SCF [49]	CASSCF	$\mu^V \neq \mu^N$	diag <sup>d</sup>
(220)	10297.00	-0.83	0.64	-1.24	-1.49
(300)	10609.14	-0.02	2.23	-1.30	-1.99
(102)	10879.00	0.01	2.01	-1.32	-2.10
(230)	11782.64	-1.19	0.30	-1.41	-1.55
(032)	12021.78	-0.88	0.57	-1.42	-1.64
(310)	12151.51	-0.34	2.03	-1.47	-2.03
(112)	12400.72	-3.59	-3.36	-1.42	-0.78
(240)	13222.66	-1.63	-0.15	-1.56	-1.61
(042)	13470.04	-1.25	0.19	-1.58	-1.69
(320)	13655.39	-0.72	1.65	-1.64	-2.05
(170)	13682.51	-3.65	-3.15	-1.54	-1.05
(400)	13839.28	0.05	3.44	-1.73	-2.52
(122)	13926.37	-0.61	1.56	-1.67	-2.19
(202)	14234.86	-0.11	2.73	-1.74	-2.70
(004)	14548.39	0.37	3.22	-1.72	-2.79
(330)	15125.11	-0.88	1.63	-1.81	-2.15
(410)	15358.27	-0.22	3.27	-1.93	-2.54
(132)	15395.63	-0.85	1.40	-1.83	-2.26
(212)	15759.25	-0.36	2.60	-1.91	-2.73
(500)	16910.87	0.08	4.71	-1.79	-2.92
(302)	17473.72	-0.06	3.86	-2.17	-3.23
(430)	18284.98	-0.56	3.28	-2.20	-2.71
(510)	18407.30	-0.03	4.77	-2.11	-2.94
(600)	19793.59	0.05	6.27	-0.77	-3.31
(402)	20549.65	0.06	5.31	-2.22	-3.65
(610)	21237.93	-0.17	6.12	-1.15	-3.34
(700)	22540.31	0.18	8.87	-3.35	-3.75
(620)	22646.30	-0.44	5.63	-1.91	-3.32
(800)	25119.49	0.24	1.40	-9.20	-2.17

Table 7

Sensitivity of vibrational band origins to the various effects, see text for details

Effect	Contribution / $\text{cm}^{-1}$	
	H <sub>2</sub> O	H <sub>3</sub> <sup>+</sup>
BO convergence	+30	$\pm 0.003$
Relativistic correction (1e)	-19	$\pm 0.003$
Darwin term (2e)	-0.8	0 <sup>a</sup>
Gaunt correction	+5	0 <sup>a</sup>
Breit correction	+6	0 <sup>a</sup>
QED Lamb shift	+1.3	0 <sup>a</sup>
Adiabatic correction (BODC)	+5	$\pm 1.5$
Non-adiabatic correction	-4	-0.5

<sup>a</sup> Not known, assumed to be negligible.