

Psychological Medicine, 2003, **33**, 1369–1379. © 2003 Cambridge University Press
DOI: 10.1017/S0033291703008390 Printed in the United Kingdom

Examining the relationship between pubertal stage, adolescent health behaviours and stress

A. E. SIMON, J. WARDLE,¹ M. J. JARVIS, N. STEGGLES AND M. CARTWRIGHT

From the Cancer Research UK Health Behaviour Unit, Department of Epidemiology and Public Health, University College London

ABSTRACT

Background. This paper examines the associations between puberty and three important health behaviours (smoking, food intake and exercise) and explores whether these associations are mediated by puberty's relationship to stress and psychological difficulties.

Method. Data were taken from the first year of the ongoing, 5-year, Health and Behaviours in Teenagers Study (HABITS). This is a school-based study set in 36 schools in London. In the first year of the study, 4320 students (2578 boys, 1742 girls) in their first year of secondary education took part.

Results. Among girls, being more pubertally advanced was associated with a greater likelihood of having tried smoking. Among boys, being more pubertally advanced was associated with a greater likelihood of having tried smoking, a higher intake of high-fat food and higher levels of exercise. More pubertally advanced girls experienced more stress but not more psychological difficulties. There were no associations between puberty and either stress or psychological difficulties in boys. Stress and psychological difficulties were associated with health behaviours in girls and boys, but neither of these factors mediated the relationship between pubertal stage and health behaviours found in girls.

Conclusions. These results suggest that the onset of puberty has a marked effect on the development of health behaviours. Puberty was related to an acceleration of the development of unhealthy behaviours, except for exercise behaviour in boys, where advanced puberty was associated with more exercise. These changes were unrelated to adolescent issues of stress and a causal explanation for these associations must be sought elsewhere.

INTRODUCTION

Puberty is a period of rapid hormonal, physiological and physical changes. These dramatic changes might be expected to engender some psychological effects, insofar as they represent a developmental transition for which the individual is not always fully prepared. Onset of puberty has been associated with a range of social and emotional effects (Steinberg & Sheffield-Morris, 2001). Behaviour changes also appear,

including varied sleeping and eating patterns (Gillman *et al.* 2000; Laberge *et al.* 2001), new sexual behaviours (Goodson *et al.* 1997) and involvement in delinquency and substance use (Williams & Dunlop, 1999; Dick *et al.* 2000).

The hormonal and physical changes of puberty will have a direct influence on some aspects of emotion and behaviour. In addition, the perception of changes by the adolescents themselves, or the reactions of others to their more adult bodies, could play a part. There is evidence that the timing of puberty influences the individual's reactions. Being pubertally 'off-time', i.e. early or late compared to others in the social group, has been hypothesized to be stressful

¹ Address for correspondence: Professor Jane Wardle, Cancer Research UK Health Behaviour Unit, Department of Epidemiology and Public Health, University College London, 2–16 Torrington Place, London WC1E 6BT.

(Petersen & Taylor, 1980). This is partly because the 'off-time' adolescent is in the minority and may lack the social or institutional support that smooths the transitional path for the 'on-time' individuals (Alsaker, 1996). The stress of off-time puberty has been hypothesized to lead to adaptive problems, which display themselves in the form of emotional problems or delinquency (Williams & Dunlop, 1999). An alternative hypothesis is that only early maturers are at particular risk because their physical maturity means that both adults and peers attribute to them a psychological maturity that may outstrip their actual development (Stattin & Magnusson, 1990). Early maturing girls tend to develop friendship networks with other equally mature, and therefore older girls, where more 'adult-type' behaviour is acceptable (Silbereisen *et al.* 1989). Early maturers may also perceive themselves as more grown-up and emulate 'adult' behaviour, such as smoking and dieting. A conflict can arise because these behaviours, which are acceptable within the adult world, are perceived by adults to be 'norm-breaking' or rule transgressing in teenagers (Alsaker, 1996).

Behavioural research in this field has had a particular focus on behaviours such as substance abuse and delinquency (Tschann *et al.* 1994; Williams & Dunlop, 1999; Dick *et al.* 2000), which pose problems in the adolescent years, compromising family and social relationships, as well as educational progress. Delinquent behaviour peaks during mid-adolescence, then declines to pre-adolescent levels by the mid-twenties. This increase in delinquency may reflect a 'maturity gap' between biological maturity and access to adult roles in society (Moffitt, 1996). As regards substance use, studies have shown that early-maturing adolescents, but not all off-time adolescents, have higher substance use, including alcohol, marijuana and cigarette use (Tschann *et al.* 1994; Harrell *et al.* 1998). Early menarche has also been associated with earlier and more frequent smoking and drinking (Dick *et al.* 2000).

Smoking has most often been examined in this broader context of substance use or in the delinquency context where it is represented as a 'transgressing act'. However, adolescence is also an important time for establishing health behaviours, and it is possible that the onset of puberty could be linked with a higher risk not only of

smoking, but also of other health-compromising behaviours such as a sedentary lifestyle and an unhealthy diet. Early initiation of unhealthy behaviours might place the individual at higher risk for health problems in later life.

Research on exercise in adolescence shows that girls' participation in exercise is consistently lower than boys' participation (Lindquist *et al.* 1999; Lasheras *et al.* 2001). Boys' interest in school-based physical education (PE) remains relatively stable throughout secondary school, but girls' interest in PE, while higher than boys at younger ages, drops off, so that by the age of 14 the gender balance is reversed (Van Wersch *et al.* 1992). It has been suggested that PE loses its status as an important school subject for girls, because being 'good at sports' does not entail popularity for girls in the way that it does for boys (Van Wersch *et al.* 1992). Boys have also been shown to use exercise rather than dieting as their primary route for body shape change (Ricciaridelli *et al.* 1999). On the basis of these results, being more pubertally advanced might be expected to be associated with higher levels of exercise in boys and lower levels in girls. One study, which examined pubertal stage in relation to activity choices, found that higher pubertal stage was associated with an increase in sedentary activities in girls, although there was no such association in boys (Bradley *et al.* 2000).

Energy intake has been investigated in relation to pubertal stage in a number of studies, but there are few studies of food choices. Total energy intake increases across pubertal stage, peaking for girls at Tanner stage 3, and continually increasing with stage for boys (Clavien *et al.* 1996), although after controlling for body weight, the more pubertally advanced adolescents are not consuming more in relation to their body weight. Adolescence is a time when children take more responsibility for their own food choices, and are more likely to eat at different times from the family or away from the family home. Adolescents also appear to consume an excess of fat at the expense of carbohydrate and fibre intake (Clavien *et al.* 1996; Rolland-Cachera *et al.* 2000), and eat little fruit and vegetables (Neumark-Sztainer *et al.* 1996) although this has not been examined extensively in terms of pubertal development.

Some of the work relating pubertal stage to behavioural outcomes has considered the

possibility that emotional factors provide the link with pubertal stage (Tschann *et al.* 1994; McCabe *et al.* 2001). Puberty has been shown to be related to depression, antisocial behaviour, aggression, anxiety, withdrawal and attention problems (Caspi & Moffitt, 1991; Hayward *et al.* 1999). In addition, the increase in weight at puberty leads to body dissatisfaction and body change behaviour, such as dieting, that are also related to increased levels of anxiety (McCabe *et al.* 2001). However, few studies have directly tested the mediational model. One exception is a study by Tschann *et al.* (1994), which found that emotional distress did not appear to mediate the relationship between pubertal timing and substance use. This study's focus was an examination of substance use and it would be useful to both broaden the exploration of this topic to other kinds of health behaviours and to provide further evidence either to replicate or refute this finding.

The present study examines associations between puberty and three significant health behaviours: smoking, food choice and exercise and examines the hypothesis that emotional stress mediates the associations between puberty and behaviour. In this way, a fuller picture can be developed of the relationships between pubertal development, health behaviour and emotional distress. Data are taken from the baseline year of a 5-year cohort study, giving the advantage of a large, mixed-sex sample (Wardle *et al.* 2003).

METHOD

Design

The HABITS study is a 5-year longitudinal study of a cohort of adolescents attending 36 secondary schools in Greater London. The sample was stratified by school type (inner-city state; suburban state; privately funded) and gender mix (boys, girls, both). A list was compiled of all secondary schools in the South London boroughs taking students from the age of 11–16 years. Four schools were drawn at random from each of the nine cells formed by crossing the stratification factors. Each school was approached. If a school declined to take part a substitute was drawn from the list until all cells were complete. Twenty-eight schools

declined to take part. Thirty-six schools who agreed to participate were identified. The data for this paper were from the first year of data collection between January and December 1999.

All students in the first year (age 11–12 years) of the secondary schools were eligible for inclusion ($N=5153$). Parental approval was obtained prior to data collection in the form of a letter taken home from school by students. Parents returned forms to the school if they wished to exclude their child from the study. In the first year 4320 students (2578 boys and 1742 girls) in Year 7 took part. Eighty-four per cent of the total student body were present in the classes when data collection took place. Ten per cent were absent on the day of data collection, and 5.5% opted out of the study (either by parental or self-exclusion).

Participants completed a self-report questionnaire on smoking, diet, exercise, body image, pubertal development, stress, psychological health, and personality. Demographic information included ethnicity, gender, and socio-economic status (SES). Smoking status was confirmed with a saliva cotinine test. Body size was assessed with height, weight, and waist circumference. Ethical approval for the study was obtained from the Joint UCL/UCH Committees on the Ethics of Human Research.

Procedure

Data collection was carried out in school classes under the supervision of at least two researchers. Data were collected annually during the Spring and Autumn terms, with 18 schools visited each term.

A researcher introduced the session by explaining the purpose, the procedure, the confidentiality of answers, and the voluntary nature of the study. An opportunity for the students to ask questions was provided. Students completed a sheet (with a unique identifier) detailing their name, address, school name, date and confirming their consent to participate. This was stored separately for anonymity and used to track students across the 5 years. Students completed the questionnaire and were individually weighed out of the sight of the other students. Saliva samples for cotinine testing were obtained using a dental roll placed in the side of the mouth for 10 min (Benowitz, 1996).

Measures

Pubertal stage

Pubertal stage was assessed with items from the Pubertal Development Scale (PDS) (Petersen *et al.* 1988). The PDS was developed for use in schools, where other measures such as direct observation or explicit photographs (e.g. Sexual Maturation Scale (SMS); Tanner, 1962) are inappropriate. The PDS scale is correlated with the SMS ($r=0.72$ to 0.80) (Brooks-Gunn *et al.* 1987). Students rate themselves on five items: growth spurt, pubic hair, skin changes, and, for girls, menarche and breast development and, for boys, voice change and facial hair. Scores from each item were summed (range 5 to 19). Petersen *et al.* (1988) report that this measure can be used either continuously or with a categorical, 'staging' classification. In this sample, a 5-stage classification system has been used: pre-pubertal (score 5), beginning pubertal (scores 6–10), mid-pubertal (scores 11–15), advanced pubertal (scores 16–18), and post-pubertal (score 19).

Health behaviours

Smoking status was assessed using two questions from the UK National Smoking Surveys (Goddard & Higgins, 1999). The classification from these questions is: non-smokers (never smoked), tried smokers (tried only once), ex-smokers (used to smoke), occasional smokers (<1 a week), sometimes smokers (1–6 a week) and regular smokers (≥ 6 a week). For some analyses, because cell sizes of regular smokers at this age were small, students were categorized into 'never smokers' and 'ever smokers'. Susceptibility to future smoking among non-smokers was assessed using a scale developed by Pierce *et al.* (1996) that taps the pre-experimentation stage of the smoking onset process. The scale in this study is based on two items, referring to the possibility of smoking in the next 12 months and smoking a cigarette offered by a best friend. The response options are: 'yes definitely', 'yes probably', 'probably not' and 'definitely not'. If the respondent answers in any other way than 'definitely not' they are considered cognitively susceptible to experimentation with smoking.

Weekend exercise rates were measured by two questions developed for the study which asked if participants usually took part in sports or

physical activities on Saturday and Sunday. The response options were dichotomous (yes/no). This indicated voluntary participation in extra-curricular sports and provides a measure of exercise independent of compulsory school sports frequency, which shows considerable variation between schools. This variable is likely to reflect a social aspect of exercise behaviour.

Days per week exercised was assessed using a question developed in the Youth Risk Behaviour Surveillance System Questionnaire (YRBSSQ) (Heath *et al.* 1993). This asks how many days a person did exercise during the past week that made them 'sweat and breathe heavily'. The response ranged from no days to every day. This indicated overall exercise, including participation in sports at school.

Fruit and vegetable intake was assessed using two questions, originally based on items from the DINE (Roe *et al.* 1994) and used in a similar, modified format in other studies (e.g. Wardle *et al.* 2000). The questions asked: 'about how many servings of fruit/vegetables do you usually eat in a day?'. Response options ranged from less than one to five or more servings per day. The responses were summed to give a scale of 0–10, '0' represents intake of less than one serving per day and '10' represents consuming five or more servings of fruit and five or more servings of vegetables per day.

Intake of high-fat foods was based on six items using a scale adapted from the 'foods that you eat scale' (Sweeting *et al.* 1994). These items related to intake of crisps, cakes, biscuits, chips, fried food and sausages/burgers. These items provide proportionally high levels of fat and are often used as 'snack' foods outside mealtimes. Intake was assessed on a six-point scale ranging from 'never/less than once a week' to 'more than once a day'. Scores were summed, producing a range from 0–30, higher scores indicate higher intake. This scale had good reliability ($\alpha=0.78$).

Stress and psychological difficulties

Psychological difficulties were measured using the 'Difficulties' score from the Strengths and Difficulties Questionnaire (SDQ) (Goodman *et al.* 1998). This is a brief behavioural screening questionnaire for use with children from age 4–16 years. The SDQ measures hyperactivity, emotional symptoms, conduct problems, peer

problems and also has a pro-social scale. The pro-social scale is excluded when examining psychological problems. 'Difficulties' scores range from 0 to 34 with a higher score indicating more difficulties. In this sample it had good internal reliability ($\alpha=0.72$). The SDQ has good concurrent validity and is highly correlated with scores on the Rutter questionnaires (Goodman *et al.* 1998).

Stress was assessed with the 4-item version of the Perceived Stress Scale (PSS) (Cohen *et al.* 1983) a global measure of perceived stress. The scale ranges from 0 to 16, higher scores indicate more stress. The PSS has good concurrent and predictive validity and is correlated in the expected direction with a range of questionnaires and behaviours (Cohen *et al.* 1983). This shortened version has relatively low internal reliability ($\alpha=0.6$) (Cohen & Williamson, 1988), which was lower ($\alpha=0.50$) in the present sample, possibly because of the age of the respondents. Its advantage is that it is short and enables repeated measures of perceived stress in large samples.

Demographic measures

Demographic information included gender, age, self-reported ethnicity and SES. Students were asked their exact date of birth, and age in decimal years was calculated. Ethnicity was assessed with five response categories: white, black, Asian, mixed or other (specified). The categories 'mixed' and 'other' contained heterogeneous samples and were combined for the purpose of analyses. SES was assessed using indicators of material conditions in the family home. More traditional methods of assessing SES (e.g. parental occupation) are unreliable in lower SES adolescents (Wardle *et al.* 2002). Three material indicators were used: housing tenure, car ownership and possession of a dishwasher.

Body Mass Index (BMI)

BMI was derived from height and weight measurements.

Analytical strategy

Descriptive statistics are presented for the PDS scale and the demographic characteristics of the sample. Chi-square statistics and one-way analyses of variance (ANOVA) are then shown for the relationships between pubertal stage,

demographics, health behaviours and psychological variables.

The data were then analysed using logistic and ordinal regressions showing the relationship of pubertal stage to health behaviours for each gender, while also controlling for SES, ethnicity, age and BMI. The dependent variables in the logistic regressions were: smoking status (tried smoking or not), susceptibility to smoking (susceptible or not) and weekend exercise (1–2 days or none) and in the ordinal regressions were: days per week exercised (0–7 days), fruit and vegetable intake (0–10 servings) and high-fat food intake (separated into deciles). In order to assess the possibility that stress/psychological difficulties mediate the relationship between pubertal stage and health behaviours, further regression analyses were carried out which compared odds ratios from a model that included stress/psychological difficulties and a model without these variables. The regression analyses were carried out in STATA v.7 and took account of school clustering and a combined weighting based on a weight for design and missing data. The design weight was based on the total number of schools in South London within each stratification cell and the refusal rate of the schools within each cell. The missing data weight took account of the proportionately greater amount of missing data from the lower SES and ethnic minority groups.

RESULTS

The demographic characteristics of the participants are shown in Table 1. The average age of the students was 11.83 decimal years ($s.d.=0.35$). Approximately 40% of the participants considered themselves to be from non-white ethnic groups.

Table 1 also shows pubertal stage using the PDS scores. For both boys and girls, few are pre-pubertal, the majority are classified as 'beginning pubertal', with more boys than girls in this category, and more girls represented in the more advanced categories. This is to be expected at this age as girls mature earlier than boys. For most of the analyses, the sample was divided into three groups. This was developed because the cell sizes for boys who had reached advanced puberty and for girls who had still not begun puberty at this age were small. For girls the

Table 1. *Distribution of the sample by SES, ethnicity, pubertal stage, BMI and age*

	Girls (N=1742)	Boys (N=2578)
Age, mean (s.d.)	11.85 (0.32)	11.81 (0.37)
BMI, mean (s.d.)	19.98 (3.84)	19.02 (3.40)
SES quartiles, %		
Low	25.2	18.7
Medium	25.0	21.4
High	27.1	31.5
Highest	22.7	28.4
Ethnicity, %		
Black	21.6	15.3
White	58.1	64.3
Asian	7.8	8.9
Mixed/Other	12.6	9.8
Pubertal stage, %		
Pre-	2.1	8.1
Beginning	53.1	73.0
Mid-	36.0	18.1
Advanced	8.4	0.6
Post-	0.4	0.2

original 'pre' and 'beginning' categories were collapsed into a single category (pre/beginning puberty) and, for boys, the original 'mid' and 'advanced/post' categories were collapsed into a single category ('mid/advanced'). Therefore, the three groups in girls were 'pre/beginning puberty', 'mid-puberty' and 'advanced/post puberty', whereas the three groups in boys represented 'pre-puberty', 'beginning puberty' and 'mid/advanced puberty'. All analyses are carried out separately for boys and girls.

Table 2 shows the associations between demographic characteristics including ethnicity, SES, age and BMI and pubertal stage. There was a significant relationship between SES and pubertal stage in boys, $\chi^2(N=1715)=18.92$, $df=6$, $P<0.01$ and a similar trend in girls, $\chi^2(N=1316)=11.64$, $df=6$, $P=0.07$, with the students from lower SES backgrounds being more likely to be advanced maturers. Among both boys and girls, the black students were more likely to be in the more advanced pubertal stages, compared with children from white and Asian backgrounds: boys, $\chi^2(N=1770)=22.91$, $df=6$, $P<0.001$; girls, $\chi^2(N=1327)=76.54$, $df=6$, $P<0.001$.

The age range for Year 7 is 10.3 to 13.7 decimal years. The typical age of students in Year 7 is 11–12 years, the sample also included nine participants aged <11 and 10 participants aged >12. These represent children who have

either been accelerated or held back at school. As would be expected, the mean age increases across pubertal stages in girls, $F(2,1301)=26.20$, $P<0.001$; and boys, $F(2,1720)=13.00$, $P<0.001$. Students who were advanced, and thus more developmentally mature, also had a higher BMI: girls, $F(2,1327)=48.50$, $P<0.001$; boys, $F(2,1775)=8.55$, $P<0.001$.

Pubertal stage and health behaviours

Table 2 also shows the raw data for the relationships between pubertal stage and health behaviours. These indicated that, in boys, a relationship existed between pubertal stage and high-fat food intake, $F(2,1719)=8.22$, $P<0.001$, fruit and vegetable consumption, $F(2,1758)=3.64$, $P<0.05$, having tried smoking $\chi^2(N=1757)=26.61$, $df=2$, $P<0.001$ and weekend exercise, $\chi^2(N=1723)=15.94$, $df=4$, $P<0.01$. In girls a relationship was seen between pubertal stage and having tried smoking, $\chi^2(N=1313)=31.47$, $df=2$, $P<0.001$.

In order to explore these relationships more rigorously a series of logistic and ordinal regression analyses were carried out to assess the odds of reporting each behaviour in relation to pubertal stage, controlling for age, SES, ethnicity, and BMI and including clustering for schools and weighting for design and missing data, as previously discussed. The lowest pubertal stage was used as the reference group. In the first sets of analyses, the gender-by-pubertal stage interactions were included in the model. These were significant or close to significant, for weekend exercise, days of exercise, high-fat food intake, and fruit and vegetable intake, although not for either of the smoking variables. In the light of these results, and because of the different groupings of pubertal stage in boys and girls, the analyses of the associations between pubertal stage and health behaviours were carried out separately for each gender. These results are shown in Tables 3 and 4.

The odds of smoking were higher in the more pubertally advanced students in both boys and girls. The pattern of odds ratios supported a graded effect across the stages, indicating that relative to students in the earliest pubertal stage, those at successively later stages were increasingly likely to have smoked. There was also a relationship between pubertal stage and non-smokers' susceptibility to smoking in both boys

Table 2. Descriptive statistics showing the relationship between pubertal stage, demographics, health behaviours, stress and psychological difficulties

	Girls			Boys		
	Pubertal stage			Pubertal stage		
	Pre/Beginning (N=739)	Mid (N=482)	Advanced/Post (N=118)	Pre (N=145)	Beginning (N=1307)	Mid/Advanced (N=339)
SES						
Low, %	50.5	39.2	10.3	3.8	71.9	24.3
Middle, %	52.7	36.0	11.3	7.8	71.2	21.0
High, %	59.5	33.8	6.7	8.1	73.8	18.1
Highest, %	58.8	34.4	6.8	10.9	72.8	16.3
Ethnicity						
White, %	62.6	31.8	5.6	8.9	74.3	16.8
Black, %	35.5	46.6	17.9	5.8	66.8	27.4
Asian, %	54.3	40.0	5.7	7.8	77.1	15.1
Mixed, %	50.6	38.1	11.3	5.8	67.9	26.3
Tried smoking, %						
No	90.0	79.5	76.7	92.3	82.6	73.4
Yes	10.0	20.5	23.3	7.7	17.4	26.6
Susceptibility to smoking, %						
No	85.8	80.8	82.0	79.5	79.6	77.6
Yes	14.2	19.2	18.0	20.5	20.4	22.4
Weekend exercise, %						
No days	33.5	36.5	40.0	30.7	24.8	16.3
1 day	43.1	43.0	37.3	26.3	33.2	35.3
2 days	23.4	20.5	22.7	43.1	42.0	48.3
Age, mean (s.d.)	11.80 (0.32)	11.90 (0.32)	11.98 (0.31)	11.72 (0.34)	11.80 (0.36)	11.89 (0.38)
BMI, mean (s.d.)	19.15 (3.56)	20.57 (3.73)	22.25 (3.67)	18.24 (2.75)	18.78 (3.21)	19.45 (3.58)
Days/week exercised, mean (s.d.)	3.70 (1.78)	3.63 (1.76)	3.94 (2.11)	3.97 (2.07)	4.19 (2.00)	4.33 (2.14)
High-fat food intake, mean (s.d.)	13.77 (4.90)	14.47 (5.10)	14.17 (5.34)	14.22 (5.41)	15.37 (5.47)	16.38 (5.77)
Fruit/vegetable intake, mean (s.d.)	4.24 (2.26)	4.29 (2.41)	4.39 (2.66)	4.00 (2.30)	4.03 (2.32)	4.41 (2.64)
Perceived stress, mean (s.d.)	5.70 (2.82)	5.79 (2.91)	6.47 (3.30)	5.34 (2.93)	5.29 (2.98)	5.54 (3.16)
Psychological difficulties, mean (s.d.)	11.80 (5.04)	12.24 (5.07)	12.66 (4.85)	12.19 (4.87)	12.05 (5.14)	12.57 (5.36)

and girls. This was difficult to interpret for girls, but again suggested in boys that susceptibility to smoking increased with pubertal stage.

For exercise behaviour, a significant relationship was found between pubertal stage and weekend exercise in boys. The more pubertally advanced boys were more likely to take exercise at the weekend. Although non-significant the results from the girls' analyses suggested a reversal of this pattern with girls taking less weekend exercise as they become more developed. Effects for overall days of exercise were not significant in either girls or boys, although the results showed the same direction for boys.

For food intake, there was a significant relationship between pubertal stage and high-fat food intake in boys with the more pubertally advanced boys consuming larger amounts of high-fat food, but no significant relationship between pubertal stage and high-fat food intake

in girls. There was no association between pubertal stage and fruit and vegetable intake in either boys or girls.

Mediation by stress and psychological difficulties

For a variable to be a potential mediator it has to be associated with both the independent and the dependent variables (Baron & Kenny, 1986). Mean scores for the stress and psychological difficulties scales across the pubertal groups are shown in Table 2. There was a significant association between pubertal stage and stress scores in girls, $F(2, 1324) = 3.50, P < 0.05$, and a trend for psychological difficulties to be higher in those who were more pubertally advanced, $F(2, 1328) = 2.08, P = 0.13$. The pattern of scores for boys was neither linear nor statistically significant, although the highest level of stress and psychological difficulties was in the most pubertally advanced group.

Table 3. Summary of regression analyses showing the associations between health behaviours and pubertal stage in boys: Model 1 controlling for SES, age, ethnicity, BMI and Model 2 additionally controlling for perceived stress and psychological difficulties – odds ratios (95% CI)

	N	Pubertal stage		
		Pre	Beginning	Mid/Advanced
Model 1				
Tried smoking	1244	1.0	2.39 (1.28–4.45)**	4.41 (2.18–8.91)***
Susceptibility to smoking†	1057	1.0	2.37 (1.12–5.04)*	2.55 (1.04–6.28)*
Days/week exercised	1259	1.0	1.22 (0.90–1.63)	1.31 (0.78–2.18)
Weekend exercise	1195	1.0	1.21 (0.89–1.65)	2.11 (1.34–3.33)**
Fruit and vegetable intake	1254	1.0	1.08 (0.79–1.48)	1.30 (0.96–1.75)
High-fat food intake	1216	1.0	1.54 (1.08–2.20)*	2.39 (1.52–3.70)***
Model 2				
Tried smoking	1229	1.0	2.23 (1.17–4.26)*	4.17 (2.02–8.61)***
Weekend exercise	1184	1.0	1.24 (0.90–1.72)	2.19 (1.39–3.47)**
High-fat food intake	1203	1.0	1.68 (1.18–2.39)**	2.48 (1.63–3.86)***

† Among never smokers.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Table 4. Summary of regression analyses showing the associations between health behaviours and pubertal stage in girls: Model 1 controlling for SES, age, ethnicity, BMI and Model 2 additionally controlling for perceived stress and psychological difficulties – odds ratios (95% CI)

	N	Pubertal stage		
		Pre	Beginning	Mid/Advanced
Model 1				
Tried smoking	1603	1.0	2.36 (1.32–4.25)**	2.64 (1.09–6.41)*
Susceptibility to smoking†	1303	1.0	1.60 (1.08–2.34)*	1.37 (0.77–2.45)
Days/week exercised	1628	1.0	0.90 (0.73–1.13)	1.21 (0.71–2.03)
Weekend exercise	1576	1.0	0.79 (0.60–1.05)	0.60 (0.30–1.20)
Fruit and vegetable intake	1606	1.0	1.01 (0.71–1.43)	0.88 (0.52–1.49)
High-fat food intake	1574	1.0	1.22 (0.95–1.58)	1.23 (0.88–1.73)
Model 2				
Tried smoking	1573	1.0	2.56 (1.45–4.54)**	2.42 (1.07–5.49)*
Weekend exercise	1552	1.0	0.80 (0.60–1.07)	0.62 (0.31–1.24)
High-fat food intake	1547	1.0	1.17 (0.92–1.51)	1.13 (0.77–1.67)

† Among never smokers.

* $P < 0.05$; ** $P < 0.01$.

To check for associations between stress or psychological difficulties and health behaviours we used Pearson correlations, which showed that they were significantly associated with each of the health behaviours that were associated with pubertal stage, that is, having tried smoking, weekend exercise and high-fat intake. Stress was modestly but significantly associated with having tried smoking ($r = 0.09$, $P < 0.001$) doing less weekend exercise ($r = -0.06$, $P < 0.05$) and higher intake of fatty foods ($r = 0.10$, $P < 0.001$) in boys. The same patterns emerged for associations with psychological difficulties (smoking,

$r = 0.18$, $P < 0.001$; exercise, $r = -0.06$, $P < 0.01$; intake of fatty foods, $r = 0.13$, $P < 0.001$). The results were very similar for girls showing significant associations with stress (smoking, $r = 0.11$, $P < 0.001$; exercise, $r = -0.08$, $P < 0.01$; intake of fatty foods, $r = 0.10$, $P < 0.001$) and with psychological difficulties (smoking, $r = 0.16$, $P < 0.001$; exercise, $r = -0.07$, $P < 0.01$; intake of fatty foods, $r = 0.14$, $P < 0.001$).

Given that the criteria for mediation were met, logistic and ordinal regressions were carried out controlling for stress and psychological difficulties to see if this reduced the variance

associated with pubertal stage (see Tables 3 and 4, Model 2). In the interests of symmetry, these were done for both boys and girls, even though the association between pubertal stage and stress in boys was less clear cut. Looking at the odds ratios and confidence intervals for pubertal stage, it is clear that including these factors, although slightly changing the values, did not have a substantial impact. Therefore, differences in stress and psychological difficulties between pubertal stages did not seem to be the key to the variation in health behaviours linked with puberty.

DISCUSSION

The results of this study support the hypothesis that pubertal stage was related to health behaviour in a large sample of adolescents although the patterning of results varied both by behaviour and gender. Being more pubertally advanced in boys was associated with a greater likelihood of smoking and a less healthy diet, but higher levels of exercise. In girls, more advanced puberty was linked with higher smoking, but there was little association with either diet or exercise. The association between pubertal stage and health behaviours, where it was seen, generally showed a graded effect, such that the likelihood of engaging in a particular behaviour changed incrementally with increases in pubertal stage.

The results on smoking status and susceptibility to smoking replicated other findings in this area (Tschann *et al.* 1994; Gritz *et al.* 1998; Harrell *et al.* 1998; Dick *et al.* 2000). For exercise behaviour, advanced puberty was associated with more weekend exercise in boys but not in girls. In girls there was the possibility that weekend exercise may actually decrease with pubertal stage, although this was a non-significant result. The results reflect established gender differences in exercise participation among adolescents (Lasheras *et al.* 2001). The results also support the suggestion that higher pubertal stage is associated with an increase in sedentary activity among girls, but not boys (Bradley *et al.* 2000). The variable measuring 'days per week exercised' produced no significant results. This was probably because this variable is confounded with compulsory physical education at school, which varies between schools and does

not necessarily index voluntary participation in exercise. Exercise outside school context may be more important, as it indicates an adolescent's deliberate choice to exercise as part of their leisure/social time. Pubertal stage was associated with high-fat food intake only in boys in our sample. The more developed boys may simply need to eat more as they are older and taller. Other studies have suggested that this gradient in intake disappears after controlling for body weight (Clavien *et al.* 1996) but in our data, the effect persisted after controlling for BMI. These findings suggest that pubertal development in boys is associated with an increasingly poor diet. This increase in fat intake may be related to the greater independence of older children to make food choices as they decrease the amount of meals eaten at home (Gillman *et al.* 2000). There was no relationship between fruit and vegetable consumption and pubertal stage in either boys or girls, so although previous studies have shown decreased consumption associated with age (Vereecken & Maes, 2000), our data suggest that this is not related to the pubertal changes occurring at this time.

The main aim of the present study was to test whether the associations between pubertal stage and health behaviours were linked to puberty's associations with either perceived stress or emotional and behavioural difficulties as measured by the SDQ. A relationship was seen between stress and pubertal stage in girls, although no such relationship was found in boys. Smaller effects between puberty and stress in boys may have gone undetected due to the low reliability of the PSS in this sample. However, the SDQ also failed to detect a relationship, although for both measures the most stress and difficulties were detected among the most advanced male puberty group. The boys were overall less developed than the girls. The relationship with stress and psychological difficulties may only emerge among more advanced puberty groups than the boys represented at this age. Therefore, there may be value in exploring this relationship further using longitudinal work which tracks these groups over time. Having noted these limitations, it was still clear from the logistic regression analyses that these variables did not mediate the relationship between pubertal stage and health behaviours in either gender. On the basis of the results shown here,

an alternative explanation for the link between pubertal stage and health behaviours must be found.

Although not tested here, it may be that other social, rather than emotional, processes have a role to play in explaining the health behaviour differences. Early maturation has been associated with social processes, e.g. socialization with older peers, greater peer pressure, emulating 'adult' type behaviour (Silbereisen *et al.* 1989) that may result in behavioural changes. Behaviours such as smoking and eating outside of the home are associated with a more 'adult' or 'independent' lifestyle and the changes in behaviour seen in this study could demonstrate an attempt by the adolescent to assert their independence.

There were some limitations to the conclusions that can be drawn from this study. As discussed, there were problems with the measurement of perceived stress, which have meant that the mediational hypothesis was not tested as rigorously as it could have been. The results from this study derive from cross-sectional data which has meant that some relationships have remained ambiguous. For example, the results found here could reflect differences in pubertal timing rather than differences in levels of pubertal stage. That is, those who are the most pubertally advanced at this age, are also the group that have developed relatively early compared to their peer group. As the current pre/beginning/mid pubertal groups progress through the higher pubertal stages they may display similar behaviour to the current advanced/post-pubertal groups, supporting a maturational model. On the other hand those who are currently more developed than their peers might always stand out as a distinct group, suggesting an effect of pubertal timing. It will be possible to pursue this line of enquiry longitudinally with future data from the HABITS project.

This study was important for a number of reasons. First, comparatively little research has examined smoking uptake as part of a wider issue of adoption of health behaviours such as exercise and dietary choice. There is some debate as to whether smoking is better conceived as a health behaviour or is more closely allied with problem/delinquent behaviour established during this period (Turbin *et al.* 2000).

The analyses here have shown that dietary behaviours in boys and exercise in girls are related to pubertal status in a similar way to smoking behaviour, giving some support to the health behaviour view. However, there was no association with diet in girls and a positive association with exercise in boys, suggesting complex pathways best understood on a behaviour by behaviour basis. Secondly, this study used a large sample comprising both boys and girls. Much research to date which considers pubertal stage and behaviour has focused either on boys or girls but not both. Incorporating both genders into the analyses has enabled a comparison to be made between genders and there were a number of important differences. Thirdly, this study provided the opportunity to explore a possible causal explanation for the relationship between puberty and health behaviour, namely that puberty related increases in emotional stress or distress were responsible for the changes in behaviour. This has been explored in detail in only one previous study (i.e. Tschann *et al.* 1994). The results here have substantiated the conclusions of that study that emotional distress does not mediate the relationship between puberty and behaviour.

In conclusion, more advanced puberty was found to be associated with changes in health behaviours including smoking, exercise and diet. The results suggest that maturity is a largely unfavourable event for both boys and girls in terms of the health behaviours that are established during adolescence, with the notable exception of exercise behaviour in boys. The implications for those involved in health promotion with adolescents, including school staff, are that early-maturing adolescents need extra support. This may be to help them withstand exposure to new social pressures, or to come to terms with the physical changes. The initiation of these behaviours is probably unrelated to issues of psychological stress and further work could establish more clearly whether, and in what way, pubescent adolescents are exposed to unhelpful social influences.

REFERENCES

- Alsaker, F. (1996). Annotation: the impact of puberty. *Journal of Child Psychology and Psychiatry* 37, 249–258.
- Baron, R. M. & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: conceptual,

- strategic and statistical consideration. *Journal of Personality and Social Psychology* **51**, 1173–1182.
- Benowitz, N. L. (1996). Cotinine as a biomarker of environmental tobacco smoke exposure. *Epidemiology Review* **18**, 188–204.
- Bradley, C. B., McMurray, R. G., Harrell, J. S. & Deng, S. (2000). Changes in common activities of 3rd through 10th graders: the CHIC study. *Medicine and Science in Sports and Exercise* **32**, 2071–2078.
- Brooks-Gunn, J., Warren, M. P., Rosso, J. & Garigulo, J. (1987). Validity of self-report measures of girl's pubertal status. *Child Development* **58**, 829–841.
- Caspi, A. & Moffitt, T. E. (1991). Individual differences are accentuated during periods of social change: the sample case of girls at puberty. *Journal of Personality and Social Psychology* **1**, 157–168.
- Clavien, H., Theintz, G., Rizzoli, R. & Bonjour, J.-P. (1996). Does puberty alter dietary habits in adolescents living in a Western society? *Journal of Adolescent Health* **19**, 68–75.
- Cohen, S. & Williamson, G. (1988). Perceived stress in a probability sample of the United States. In *The Social Psychology of Health* (ed. S. Spacapan and S. Oskamp), pp. 31–67. Sage: California.
- Cohen, S., Kamarck, T. & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behaviour* **24**, 385–396.
- Dick, D. M., Rose, R. J., Viken, R. J. & Kaprio, J. (2000). Pubertal timing and substance use: Associations between and within families across late adolescence. *Developmental Psychology* **36**, 180–189.
- Gillman, M. W., Rifas-Shiman, S. L., Frazier, L., Rockett, H. R., Camargo, C. A., Field, A. E., Berkey, C. S. & Colditz, G. A. (2000). Family dinner and diet quality among older children and adolescents. *Archives of Family Medicine* **9**, 235–240.
- Goddard, E. & Higgins, V. (1999). *Smoking, Drinking and Drug use Among Young Teenagers in 1998. UK National Smoking Surveys Volume 1*. The Stationery Office: London.
- Goodman, R., Meltzer, H. & Bailey, V. (1998). The strengths and difficulties questionnaire: a pilot study on the validity of the self-report version. *European Child and Adolescent Psychiatry* **7**, 125–130.
- Goodson, P., Evans, A. & Edmundson, E. (1997). Female adolescents and onset of sexual intercourse: a theory-based review of research from 1984 to 1994. *Journal of Adolescent Health* **21**, 147–156.
- Gritz, E. R., Prokhorov, A. V., Suchanek Hudmon, K., Chamberlain, R., Taylor, W., Di Clemente, C., Johnston, D., Hu, S., Jones, L., Mullin Jones, M., Roseblum, C., Ayars, C. & Amos, C. (1998). Cigarette smoking in a multiethnic population of youth: Methods and baseline findings. *Preventive Medicine* **27**, 365–384.
- Harrell, J. S., Bangdiwala, S. I., Deng, S., Webb, J. & Bradley, C. (1998). Smoking initiation in youth: the role of gender, race, socioeconomic and developmental status. *Journal of Adolescent Health* **1998** **23**, 271–279.
- Hayward, C., Gotlib, I. H., Schraedley, P. K. & Litt, I. (1999). Ethnic differences in the association between pubertal status and symptoms of depression in adolescent girls. *Journal of Adolescent Health* **25**, 143–149.
- Heath, G. W., Pate, R. R. & Pratt, M. (1993). Measuring physical activity among adolescents. *Centres for Disease Control and Prevention Public Health Reports* **108**, 42S–46S.
- Laberge, L., Petit, D., Simard, C., Vitaro, F. & Tremblay, R. (2001). Development of sleep patterns in early adolescence. *Journal of Sleep Research* **10**, 59–67.
- Lasheras, L., Aznar, S. & Merion, B. (2001). Factors associated with physical activity among Spanish youth through the national health survey. *Preventive Medicine* **32**, 455–464.
- Lindquist, C. H., Reynolds, K. D. & Goran, M. I. (1999). Socio-cultural determinants of physical activity among children. *Preventive Medicine* **29**, 305–312.
- McCabe, M. P., Ricciardelli, L. A. & Banfield, S. (2001). Body image, strategies to change muscles and weight, and puberty. Do they impact on positive and negative affect among adolescent boys and girls? *Eating Behaviours* **2**, 129–149.
- Neumark-Sztainer, D., Story, M., Resnick, M. D. & Blum, R. (1996). Correlates of inadequate fruit and vegetable consumption among adolescents. *Preventive Medicine* **25**, 497–505.
- Petersen, A. C. & Taylor, B. (1980). The biological approach to adolescence: biological change and psychological adaptation. In *Handbook of Adolescent Psychology* (ed. J. Adelson), pp. 117–155. Wiley: New York.
- Petersen, A. C., Crockett, L., Richards, M. & Boxer, A. (1988). A self-report measure of pubertal status: reliability, validity and initial norms. *Journal of Youth and Adolescence* **17**, 117–132.
- Pierce, J. P., Choi, W. S., Gilpin, E. A., Farkas, A. & Merritt, K. (1996). Validation of susceptibility as a predictor of which adolescents take up smoking in the United States. *Health Psychology* **15**, 355–361.
- Ricciardelli, L. A., McCabe, M. P. & Banfield, S. (1999). Body image and body change methods in adolescent boys: role of parents, friends, and the media. *Journal of Psychosomatic Research* **49**, 189–197.
- Roe, L., Strong, C., Whiteside, C., Neil, A. & Mant, D. (1994). Dietary intervention in primary care: validity of the DINE method for diet assessment. *Family Practice* **11**, 375–381.
- Rolland-Cachera, M.-F., Bellisle, F. & Deheeger M. (2000). Nutritional status and food intake in adolescents living in Western Europe. *European Journal of Clinical Nutrition* **54**, 41S–46S.
- Silbereisen, R. K., Petersen, A. C., Albrecht, H. T. & Kracke, B. (1989). Maturation timing and the development of problem behaviour: longitudinal studies in adolescence. *Journal of Early Adolescence* **9**, 247–268.
- Stattin, H. & Magnusson, D. (1990). *Pubertal Maturation in Female Development*. Erlbaum: New Jersey.
- Steinberg, L. & Sheffield-Morris, A. (2001). Adolescent development. *Annual Review of Psychology* **52**, 83–110.
- Sweeting, H., Anderson, A. & West, P. (1994). Socio-demographic correlates of dietary habits in mid to late adolescence. *European Journal of Clinical Nutrition* **48**, 736–748.
- Tanner, J. M. (1962). *Growth at Adolescence: With a General Consideration of the Effects of Hereditary Environmental Factors upon Growth and Maturation from Birth to Maturity*, 2nd edn. Blackwell Scientific: Oxford.
- Tschann, J. M., Adler, N. E., Irwin, C. E., Millstein, S., Turner, R. & Kegeles, S. (1994). Initiation of substance use in early adolescence: the roles of pubertal timing and emotional distress. *Health Psychology* **13**, 326–333.
- Turbin, M. S., Jessor, R. & Costa, F. M. (2000). Adolescent cigarette smoking: health-related behaviour or normative transgression? *Prevention Science* **1**, 115–124.
- Van Wersch, A., Trew, K. & Turner, I. (1992). Post-Primary school pupils' interest in physical education: age and gender differences. *British Journal of Educational Psychology* **62**, 56–72.
- Vereecken, C. & Maes, L. (2000). Eating habits, dental care and dieting. In *Health and Health Behaviour among Young People, Health Behaviour in School-aged Children: A WHO Cross-National Study (HBSC) International Report* (ed. C. Currie, K. Hurrelmann and R. Smith), pp. 83–85. World Health Organization: Copenhagen.
- Wardle, J., Parmenter, K. & Waller, J. (2000). Nutrition knowledge and food intake. *Appetite* **34**, 269–275.
- Wardle, J., Robb, K. & Johnson, F. (2002). Assessing SES in adolescents: the validity of a home affluence scale. *Journal of Epidemiology and Community Health* **56**, 595–599.
- Wardle, J., Jarvis, M. J., Steggle, N., Sutton, S., Williamson, S., Farrimond, H., Cartwright, M. & Simon, A. E. (2003). Socio-economic disparities in cancer-risk behaviours in adolescence: baseline results from the Health and Behaviour in Teenagers Study (HABITS). *Preventive Medicine* **36**, 721–730.
- Williams, J. M. & Dunlop, L. C. (1999). Pubertal timing and self-reported delinquency among male adolescents. *Journal of Adolescence* **22**, 157–171.