

# Gap junction reduction in cardiomyocytes following transforming growth factor- $\beta$ treatment and *Trypanosoma cruzi* infection

Mariana C Waghbi<sup>1,2</sup>, Robson Coutinho-Silva<sup>2,3</sup>, Jean-Jacques Feige<sup>4</sup>, Maria de Lourdes Higuchi<sup>5</sup>, David Becker<sup>6</sup>, Geoffrey Burnstock<sup>2</sup>, Tania C de Araújo-Jorge<sup>1/+</sup>

<sup>1</sup>Laboratório de Inovações Terapêuticas, Instituto Oswaldo Cruz-Fiocruz, Rio de Janeiro, RJ, Brasil <sup>2</sup>Autonomic Neuroscience Centre, Royal Free and University College Medical School, Rowland Hill Street, London, UK <sup>3</sup>Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, Brasil <sup>4</sup>Institut National de la Santé et de la Recherche Médicale, Unité 878, and Commissariat à l'Énergie Atomique, Institut de Recherches en Technologies et Sciences pour le Vivant, Grenoble, France <sup>5</sup>Instituto do Coração, Faculdade de Medicina, Universidade de São Paulo, São Paulo, SP, Brasil <sup>6</sup>Department of Anatomy and Developmental Biology, University College London, Gower Street, London, UK

*Gap junction connexin-43 (Cx43) molecules are responsible for electrical impulse conduction in the heart and are affected by transforming growth factor- $\beta$  (TGF- $\beta$ ). This cytokine increases during *Trypanosoma cruzi* infection, modulating fibrosis and the parasite cell cycle. We studied Cx43 expression in cardiomyocytes exposed or not to TGF- $\beta$  T. cruzi, or SB-431542, an inhibitor of TGF- $\beta$  receptor type I (ALK-5). Cx43 expression was also examined in hearts with dilated cardiopathy from chronic Chagas disease patients, in which TGF- $\beta$  signalling had been shown previously to be highly activated. We demonstrated that TGF- $\beta$  treatment induced disorganised gap junctions in non-infected cardiomyocytes, leading to a punctate, diffuse and non-uniform Cx43 staining. A similar pattern was detected in T. cruzi-infected cardiomyocytes concomitant with high TGF- $\beta$  secretion. Both results were reversed if the cells were incubated with SB-431542. Similar tests were performed using human chronic chagasic patients and we confirmed a down-regulation of Cx43 expression, an altered distribution of plaques in the heart and a significant reduction in the number and length of Cx43 plaques, which correlated negatively with cardiomegaly. We conclude that elevated TGF- $\beta$  levels during T. cruzi infection promote heart fibrosis and disorganise gap junctions, possibly contributing to abnormal impulse conduction and arrhythmia that characterise severe cardiopathy in Chagas disease.*

Key words: gap junction - heart - Chagas disease - transforming growth factor- $\beta$  - connexin 43

Gap junctions are aggregates (plaques) of intercellular channels that connect the cytoplasm of adjacent cells. They mediate both electrical and metabolic coupling. The gap channels are composed of 12 connexin (Cx) molecules assembled from the hexameric hemichannels (connexons) in humans (Sohl & Willecke 2004). In the heart, intercellular gap junction channels constructed from Cx molecules are crucial for propagation of electrical impulses in excitable tissues, most notably including myocardium, where Cx43 is the most important isoform and has been studied extensively (Severs 2001, Rodriguez-Sinovas et al. 2007).

Some myocardial diseases show alterations in expression levels of Cxs and/or the anisotropic distribution of gap junction plaques (Severs et al. 2004, Chen & Zhang 2006). Decrease in expression levels of ventricular Cx43 has been described in the hypertrophic heart of hypertensive rats (Bastide et al. 1993) and in ischaemic

human and rabbit hearts (Peters et al. 1993, Tansey et al. 2006). A redistribution of Cx43 gap junctions has been reported in the borderzone of infarcts (Severs et al. 2004, Chen & Zhang 2006), in hypertrophied hearts (Uzzaman et al. 2000), ischemic hearts (Peters et al. 1993, Beardslee et al. 2000) and hypertrophic cardiomyopathy (Sepp et al. 1996). The redistribution is frequently accompanied by a decrease in the size of gap junction plaques (Kaprielian et al. 1998) or in the total amount of plaques (Uzzaman et al. 2000).

Some dysfunction in gap junction intercellular communication is regulated by growth factors [e.g., epithelial growth factor, transforming growth factor- $\beta$  (TGF- $\beta$ ) and bovine pituitary extract], which may act to isolate affected cells from one another by down-regulating the transfer of ions and small molecules through gap junctions (Chanson et al. 2005). TGF- $\beta$  is a cytokine that contributes to fibrosis by increasing the synthesis of collagen and other extracellular matrix proteins and decreasing their degradation. Physiological levels of TGF- $\beta$  maintain tissue homeostasis and aberrant over-expression of TGF- $\beta$  leads to tissue fibrosis (Tabibzadeh 2002). Through regulation of Cx43 expression, TGF- $\beta$  affects gap junction-mediated intercellular communication (van Zoelen & Tertoolen 1991, Hurst et al. 1999, Rudkin et al. 1996) and the outcome varies between positive and inverse relationships, depending on the cell type, the type of membrane receptors employed and the initial activation/phosphorylation state of the

Financial support: CNPq, FAPERJ, PRONEX, Wellcome Trust (062754/Z00Z to RCS)

+ Corresponding author: [taniaaj@ioc.fiocruz.br](mailto:taniaaj@ioc.fiocruz.br)

None of the authors have commercial or other associations that might pose a conflict of interest.

Received 30 April 2009

Accepted 19 October 2009

cells (Chanson et al. 2005). Acute down-regulation of Cx43 at wound sites leads to a reduced inflammatory response, enhanced keratinocyte proliferation and wound fibroblast migration (Mori et al. 2006).

Cardiomyopathy occurs in chronic Chagasic disease, which is an infection caused by *Trypanosoma cruzi* that is characterised by focal myocarditis following parasite invasion and inflammatory cell infiltration of heart myofibers (Palomino et al. 2000) that is likely to be triggered by parasite-host interaction during the initial acute phase. *T. cruzi* infection affects over 17 million people in Latin America, leading to 45,000 deaths/year (Schmunis 2007). However, a clear understanding of its pathophysiology remains a challenge (Marin-Neto et al. 2007). Most of the infected people remain asymptomatic in the so-called "indeterminate form"; however, about 25% of the patients develop the cardiac clinical form, wherein the heart increases due to exacerbated fibrosis, leading to dysfunctions such as arrhythmias, conduction disturbances and congestive heart failure (Higuchi et al. 1999). *T. cruzi* infection of cultured rat and mouse cardiac myocytes results in a decrease in electrical coupling between the infected cells, decreased passage of dyes into or away from infected cells and a decrease in the expression of organised plates of gap junction channels at appositional membrane areas (Campos de Carvalho et al. 1992, Adesse et al. 2008). We have demonstrated previously TGF- $\beta$  involvement in heart pathology in mice following acute infection and in human chronic heart fibrosis occurring in Chagas disease (Araújo-Jorge et al. 2008).

Herein we studied *in vitro* cultures and heart samples from chagasic patients, in which TGF- $\beta$  signalling has previously been shown to be highly activated, to test the hypothesis that the expression of Cx43 would decrease under high levels of TGF- $\beta$  (Araújo-Jorge et al. 2002).

## PATIENTS AND METHODS

***In vitro T. cruzi heart cell infection*** - Cardiomyocytes from mouse embryos were obtained as described previously (Meirelles et al. 1986). Cells were seeded in 24-well plates and maintained in Eagle's medium containing 0.1% fetal calf serum, gamicine, 1 mM glutamine and 2.5 mM CaCl<sub>2</sub>. To analyse the localisation of Cx43, some subconfluent monolayers were incubated with *T. cruzi* trypomastigotes from the Y strain, employing a parasite/host cell ratio of 10:1, along with TGF- $\beta$  1 (2 ng/mL) for 24 h at 37°C. After washing, some monolayers were treated with SB-431542 (10  $\mu$ M) for longer than 24 h, washed twice in phosphate buffered saline (PBS), fixed in 4% paraformaldehyde for 10 min at 4°C and processed for immunocytochemistry analysis. SB-431542 is an activin receptor-like kinase (ALK)-5 inhibitor that blocks the TGF- $\beta$  signalling pathway (Inman et al. 2002). All procedures were carried out in accordance with the guidelines established by the Fiocruz Committee of Ethical for the Use of Animals, resolution 242/99.

***Immunofluorescence and confocal microscopy of immunocytochemical staining*** - Cells were incubated once with 0.1% Triton X-100 in PBS for 10 min, followed by three 10 min washes in PBS/2% bovine se-

rum albumin and then reacted with rabbit anti-Cx43 (Zymed, San Francisco, CA, USA) diluted 1:50 in PBS overnight at 4°C. The monolayers were incubated for 1 h at RT with goat-anti-rabbit IgG-fluorescein isothiocyanate diluted 1:200 (Jackson ImmunoResearch Laboratories, West Grove, PA, USA). The cultures were then incubated with 4,6-diamidino-2-phenylindole (DAPI) (1:5000) to stain DNA, washed three times in PBS and mounted with CitiFluor AF1 (Agar Scientific Ltd, Stansted, Essex, UK). The sections were observed using a microscope Olympus Fluoview 3.3 (Olympus Corporation, Japan) coupled with an upright BX51 microscope image acquisition system.

***TGF- $\beta$  measurement*** - Measurement was performed using a TGF- $\beta$  1-specific commercial enzyme linked immunosorbent assay (ELISA) kit (TGF- $\beta$  1 Emax; Promega Corporation, Madison, Wisconsin, USA) according to the manufacturer's instructions (the minimum detectable dose of TGF- $\beta$  1 in this assay is less than 32 pg mL<sup>-1</sup>). Supernatants of control cardiomyocytes or those infected with *T. cruzi* at 48, 72 and 96 h post-infection were assayed after acidic pH activation of latent TGF- $\beta$  1 (total TGF- $\beta$ ) or without activation (active TGF- $\beta$ ) to calculate the ratio between active and latent TGF- $\beta$  1.

***Immunoblot analysis*** - Cells were treated or not with SB-431542 (10  $\mu$ M) for 1 h and then TGF- $\beta$  1 (2 ng/mL) or *T. cruzi* Y strain were added for 48 h. Cardiomyocytes were then washed twice with PBS and lysed in radioimmunoprecipitation assay buffer [50 mM Tris-HCl, NaCl 150 mM, 1% Triton X-100, 1 mM ethylene glycol-bis-(b-amino-ethyl ether) N,N,N',N'-tetra-acetic acid, 100  $\mu$ g/mL phenylmethylsulfonyl fluoride, 1  $\mu$ g/mL pepstatin, 10  $\mu$ g/mL leupeptin and 1  $\mu$ g aprotinin, pH 8.0]. Proteins in the lysates (20  $\mu$ g/lane) were separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (12%) and analysed by immunoblotting with polyclonal rabbit anti-Cx43 (Sigma Aldrich, St. Louis, IL, USA) diluted 1:8000. The same membrane was stripped and re-probed with a monoclonal antibody against glyceraldehyde 3-phosphate dehydrogenase (GAPDH) to confirm equal protein loading in each lane. Four independent experiments were performed and the densitometric histograms of the normalised levels of Cx43 as related to GAPDH are shown.

***Heart cryosections*** - Left ventricle fragments from the heart of patients chronic chagasic cardiopathy (CCC) (n = 7) as well as from patients with non-infectious dilated cardiopathy (IDCM) (n = 1) and a reference control from a non-infected person (Healthy control) (n = 1) were obtained as described elsewhere (Araújo-Jorge et al. 2002). Hearts were taken at autopsy from four patients (Table II, A1-4) who had died from cardiac failure, including two men aged 39 and 43 years (A1, 4) who died from ventricular fibrillation leading to severe arrhythmia, a man aged 46 years (A2) who died from cardiogenic shock by ventricular failure, a young man aged 22 years (A3) and biopsies from four patients (Table II, B1-4) that underwent heart transplant (heart explanted from the recipients). Three hearts were

obtained from a chagasic group (B1-3, 2 men aged 28 and 53 years, or class IV and a woman aged 24 years, NYHA class III) and one from a patient from the IDCM group (B4) with IDCM (NYHA class III, a man aged 41 years, presenting idiopathic dilated cardiopathy). The biopsy of one young woman who presented normal heart function, but was submitted to valve prosthesis, was used as a non-infected control (B5).

**Immunohistochemical staining** - Processing for immunofluorescence was performed as described previously (Coutinho-Silva et al. 2001). Frozen sections (10  $\mu\text{m}$ ) were fixed for 10 min in 4% paraformaldehyde (VWR International, Poole, UK) in PBS followed by blocking with 10% normal horse serum (Harlan Sera-Lab, Loughborough, UK) in PBS containing 0.05% thimerosal (Merthiolate; Sigma Chemical Co, Poole, UK) for 20 min. The slides were then incubated overnight with mouse monoclonal antibodies against Cx43 (1:200) (Becker & Davies 1995). All incubations were held at RT and separated by three 5 min washes in PBS. For confocal microscopy, the secondary antibody was anti-mouse Cy5 diluted 1:200 in PBS for 1 h at 37°C, washed and mounted with glycerol CitiFluor AF1 (Agar Scientific). The sections were observed using a Zeiss Axioplan microscope (Zeiss, Oberkochen, Germany) coupled with a Leica DC 200 image acquisition system (Leica, Heerbrugg, Switzerland).

**Cx43 plaque quantification** - Leica TCS NT confocal images were digitally processed using Photoshop 5.0 software (Adobe Systems, San Jose, CA, USA). Each test area was photographed at a final print magnification of 400X. At this magnification, a typical test area included approximately 34.600  $\mu\text{m}^2$  of tissue area. The images were projected on a square screen and adjusted for 1 cm (arbitrary value) of length of Cx43 from healthy heart. The minimum gap junction length measured was 0.1 cm that corresponded to five contiguous pixels. The mean number of Cx43 plaques as well as their sizes was evaluated in each case studied by quantification of three confocal field images/patient. The means obtained in measurements from the IDCM and healthy patients were compared to the mean calculated using data collected from the seven chagasic cases. Analyses and tests were carried out using GraphPad Prism version 3.0 for Windows (GraphPad Software, San Diego, CA, USA). Statistical differences between the three groups were determined by analysis of variance followed by Dunnett's post-test. Statistical significance was defined as  $p < 0.05$  and  $p < 0.001$ .

**Fibronectin and phosphorylated Smad 2 detection in heart cryosections** - Cryostat slides were obtained as described above. Histology and immune labelling of *T. cruzi* antigens and of inflammatory infiltrates were already studied and described (Higuchi et al. 1997, Reis et al. 1997). Sections were fixed in paraformaldehyde, incubated with 1%  $\text{H}_2\text{O}_2$ , washed and saturated with 5% normal goat serum before an overnight incubation with rabbit IgG anti-fibronectin (Sigma) or anti-phosphosmad2 (PS2). The anti-PS2 antibody was raised in rab-

bits against the peptide KKKSSpMSp (where Sp stands for phosphorylated serine residues) and was given to us by Dr P ten Dijke (The Netherlands Cancer Institute, Amsterdam, The Netherlands). Its specificity was assessed previously by western blotting and immunohistochemical methods (Rosendahl et al. 2001). A biotin-avidin-peroxidase system (Dako, Glostrup, Denmark) was used for immune detection.

The patients gave signed consent for their participation according to the rules of the official Fundação Oswaldo Cruz Ethical Committee (Brazilian Ministry of Health).

## RESULTS

**Expression of Cx43 in cardiomyocytes in vitro is reduced and disorganised after TGF- $\beta$  treatment and after *T. cruzi* infection** - To test for a biological effect of TGF- $\beta$  on Cx43 expression, we treated in vitro cardiomyocytes with 2 ng/mL TGF- $\beta$  for 48 h. In non-treated cultures, typical gap junction immunoreactivity was seen at virtually every interface between the cultured cardiomyocytes (Fig. 1A, arrow). A different, punctate and disorganised pattern in the Cx43 staining was observed following the addition of 2 ng/mL TGF- $\beta$  (Fig. 1B, open arrow). This pattern indicated a clear effect of TGF- $\beta$  on Cx43 expression and/or distribution. Western blot detection of Cx43 expression further confirmed the effect of TGF- $\beta$  (Fig. 2A, C).

Infection of cardiomyocytes with *T. cruzi* led to a decrease in Cx43 expression (Campos de Carvalho et al. 1992, Adesse et al. 2008) as detected by confocal microscopy and typical *T. cruzi* DNA staining by DAPI indicated the infected cells (Fig. 1D-F, small arrows). We confirmed the down-regulation of Cx43 staining in *T. cruzi*-infected cells (Fig. 1D, E, green labelling, open arrows) compared to non-infected cells (Fig. 1A, arrow). The cells containing parasites were negative for Cx43 staining (Fig. 1D, E, open arrow); however, non-infected cells from the same cultures sometimes retained the normal pattern of Cx43 expression (Fig. 1D, large arrow). Western blot detection of Cx43 confirmed its reduction after *T. cruzi* infection (Fig. 2A, Lane 4).

**SB-431542 treatment inhibits the effect of TGF- $\beta$  on Cx43 expression in normal and *T. cruzi*-infected cardiomyocytes** - Treatment of cardiomyocytes with SB-431542 strongly increased the Cx43 immunostaining (Fig. 1C), thus confirming that the TGF- $\beta$  signalling effect did indeed specifically disorganise Cx43 expression in cardiomyocytes (Fig. 1B). Western blot analysis also showed that the addition of SB-431542 (10  $\mu\text{M}$ ) to non-infected TGF- $\beta$ -treated cardiomyocytes up-regulated the expression of Cx43 (Fig. 2A, Lane 3).

We recently demonstrated that SB-431542 inhibited *T. cruzi* infection (Waghabi et al. 2007), but in the present work we aimed to verify if the effect of SB-431542 was due to the blocked expression of TGF- $\beta$ -responsive genes rather than a decreased infection rate. We then extended this approach by adding SB-431542 after 24 h of infection. We verified that treatment with SB-431542 of the 24 h-infected cultures (Fig. 1F) reversed the typical pattern of Cx43 reduction (Fig. 1D, E). Importantly, even






Fig. 1: confocal images of cardiomyocytes stained for connexin (Cx)-43 after treatment with TGF- $\beta$  at 2ng/mL (B, C, E) or infected with *Trypanosoma cruzi* (D, E, F) in the absence of (A, B, D, E) or the presence of SB-431542 (C, F). Changes in Cx43 expression in cells incubated with exogenous recombinant TGF- $\beta$  for 48 h (B) were observed as compared to non-treated control cells (A). SB-431542 addition (10  $\mu$ M) for 24 h on TGF- $\beta$  treated cells increased Cx43 reaction (C). Changes in Cx43 expression are also detected in 48 h *T. cruzi* infected cultures (D, E) and increased in 10  $\mu$ M SB-431542 treated cells (F). White large arrows indicate Cx43 staining, open arrows indicate altered punctate staining. Small arrows indicate *T. cruzi* amastigotes. Representative confocal images are shown, with DNA staining in blue and Cx43 in green.

infected cells expressed Cx43 (Fig. 1F, arrows). Western blot analysis confirmed that cardiomyocytes treated with SB-431542 (10  $\mu$ M) prior to *T. cruzi* infection and/or TGF- $\beta$  addition overexpressed Cx43 compared to the same procedure performed without SB-431542-mediated inhibition (Fig. 2A, Lanes 3, 5).

#### TGF- $\beta$ secretion by *T. cruzi*-infected cardiomyocytes

- Previous reports have shown that cardiomyocytes secrete TGF- $\beta$  in vitro (Flanders et al. 1995) and that Chagas disease is accompanied by increased TGF- $\beta$  levels (Silva et al. 1991, Araújo-Jorge et al. 2002, Waghabi et al. 2002). We compared TGF- $\beta$  levels measured in the supernatants of control cardiomyocytes and those infected by *T. cruzi* after 48, 72 and 96 h post-infection (Table I). Non-infected cells secreted regular levels of TGF- $\beta$  over




Fig. 2: SB-431542 treatment reverses *Trypanosoma cruzi*-induced connexin (Cx)-43 down expression in cardiomyocytes. Control cardiomyocytes or those treated with SB-431542 (10  $\mu$ M) for 1 h and then treated with TGF- $\beta$ 1 (2 ng/mL) or *T. cruzi* Y strain for 48 h. Cell lysates (20  $\mu$ g) were resolved on a 12% SDS-polyacrylamide gel and immunoblotted with anti-Cx43 (A) and anti-glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (B) antibodies. Densitometric histograms of the normalized levels of Cx43 as related to GAPDH are shown in (C). n = 4.

the period studied (70-250 pg/mL). Infected cardiomyocytes secreted higher TGF- $\beta$  levels, starting from 1,600 and reaching 2,200 pg/mL, which is seven times higher than total TGF- $\beta$  secretion from non-infected cells. Detection of active TGF- $\beta$  in culture supernatants was similar in infected and non-infected cells after 48 and 72 h, but was 2.6-fold higher in the infected cells 96 h after infection. To confirm these observations, we performed immunostaining of cardiomyocytes infected by *T. cruzi* using an anti-TGF- $\beta$  antibody. Amastigote-positive cells stained strongly with anti-TGF- $\beta$  antibody (Waghabi et al. 2005). Non-infected cardiomyocytes surrounding infected ones were strongly immunoreactive for TGF- $\beta$  by 24 h post-infection (Supplementary data A, arrow) as well as at 48 h (data not shown), 72 h (Supplementary data B) and 96 h (data not shown). These data confirmed TGF- $\beta$  secretion by non-infected cardiomyocytes in infected cultures, which contributed to the available total and active amounts of this cytokine as measured by ELISA (Table I). We showed, by RT-PCR assays, that the addition of TGF- $\beta$  to cardiomyocyte cultures increased by 3.6-fold the transcription of the TGF- $\beta$  gene, thus demonstrating an autocrine/paracrine self-regulation (Supplementary data C). In addition, western blot analysis showed that Smad-2 phosphorylation following TGF- $\beta$  addition to cardiomyocytes was decreased when incubated with SB-431542 in a concentration-dependent manner (Supplementary data D).

*Cx43, fibronectin and PS2 detection on human left ventricle of chagasic patients* - Circulating levels of

TGF-β and tissue expression of PS2 were associated with the progression of heart damage in Chagas disease (Araújo-Jorge et al. 2002). To compare and confirm the relevance of the in vitro data with human pathology, the expression and localisation of Cx43 was studied in the human left ventricle from healthy or cardiopathic patients. Confocal images showed staining for Cx43 (dark blue) in the different groups studied (Fig. 3A, D, G). In the healthy control sample, Cx43 labelling was localised at intercalated discs as a dense structure of plaques (Fig. 3A, arrows) as described previously (4). A change in Cx43 expression occurred in chagasic patients and we detected the disassembly of the gap junctions and a marked decreased in total staining (Fig. 3G). The mean number of Cx43 plaques was 20% lower in chagasic patients compared to healthy or IDCM groups (Table II, Fig. 4A). Cx43 plaque size was reduced by 2.2-fold (from 1 ± 0.08 to 0.45 ± 0.2) in chagasic patients, this being the main characteristic of the observed disassembly of this group (Fig. 4B), thus corroborating the observed loss of

Cx43 plaques (Fig. 3D, G). To compare the intensity of cardiac fibrosis and positive TGF-β-reactive cells with Cx43 expression, we analysed in human heart sections the presence of fibronectin and PS2, markers of fibrosis and of active TGF-β signalling, respectively. When compared to healthy controls (Fig. 3B, C), higher levels of pericellular fibronectin staining (Fig. 3H) and nuclear staining to PS2 (Fig. 3I) were observed on the sections of cardiac tissue from chagasic patients in parallel to the reduced Cx43 staining (Figs 3G, 4A, B).

The main clinical data and quantitative analysis of Cx43 size and expression from each patient are shown in Table II. The histology analysis and immunolabelling of *T. cruzi* antigens and of inflammatory infiltrates of these patients have already been studied and described elsewhere (Higuchi et al. 1997, 1999, Reis et al. 1997, Palomino et al. 2000). The normal pattern of Cx43 staining in human ventricular myocardium (healthy donors) has already been determined (Jiang et al. 2005), thus only one heart was used as a control due to ethi-

TABLE I  
Evidence for the presence of active TGF-β in *Trypanosoma cruzi*-infected cardiomyocyte cultures

Parameter	Hours after infection					
	48		72		96	
	NI	I	NI	I	NI	I
Total TGF-β (HCl treated) <sup>a</sup>	218. ± 77	1643.4 ± 6.3	239.9 ± 27	1841.9 ± 299	338.5 ± 162.9	2237.8 ± 78
Active TGF-β (non-HCl treated) <sup>a</sup>	134.2 ± 10	73.1 ± 3	87.1 ± 7.2	67.4 ± 17	273.02 ± 19	721.12 ± 16
Number of nuclei stained for PS2/ number of total non-infected cell nuclei - DAPI (%)	0	0	0	0	0	65/250 (26)
Number of nuclei stained for PS2 in infected cell/ number of total infected cell nuclei - DAPI (%)	0	0	0	0	0	83/126 (66)

a: pg/mL; DAPI: 4,6-diamidino-2-phenylindole; I: infected; NI: non infected; PS2: phosphorylated Smad 2.

TABLE II  
Clinical data in patients with normal, non-infectious dilated cardiopathy (IDCM) and chronic chagasic cardiopathy (CCC) hearts

Patient code	Group	Age (years)	Sex (M/F)	Heart weight (g)	Cx43 number	Cx43 size	CI (+/-)	Myocarditis (+/-)	Amastigote detection (+/-)
A1	CCC	39	M	350	74.33	0.73	-	+	nd
A2	CCC	43	M	600	29	0.33	+	+	+
A3	CCC	46	M	nd	56	0.26	+	+	nd
A4	CCC	22	M	nd	81.66	0.55	+	+	nd
B1	CCC	28	M	480	34	0.46	-	+	nd
B2	CCC	53	M	380	64.5	0.38	+	+	nd
B3	CCC	24	F	370	64.33	0.32	nd	nd	nd
B4	IDCM	41	M	nd	73.5	0.38	nd	nd	-
B5	Healthy control	nd	F	235	72	1	-	-	-

CI: cardiac insufficiency; Cx: connexin; nd: non-determinate; +: present; -: not present.




Fig. 3: fragments from human left ventricles from a healthy heart (A-C), a non-chagasic patient with idiopathic dilated cardiomyopathy (D-F) or a chagasic patient with cardiomyopathy (G-I). Blue staining corresponds to connexin (Cx)-43 labelling (arrows) in confocal images from hearts immunostained with anti-Cx43 antibody (A, D, G). The immune-reactivity for fibronectin and phosphorylated Smad 2 (PS2) was visualized as red-brown. Markedly elevated expression of PS2 and fibronectin concomitant to the decreased expression of Cx43 in heart biopsies from chagasic patients with cardiomyopathy are observed. Bar = 31  $\mu$ m for A, D, G; Bar = 63  $\mu$ m for B, C, E, F, H, I.




Fig. 4: quantitative analysis of the number of connexin (Cx)-43 plaques (A) and size (B) after measurement in confocal images for each group studied. Values express mean  $\pm$  standard deviation obtained with images from patients in each group. Cx43 size plaques were measured in centimetres in all images under the same magnification and expressed as relative arbitrary values. Correlation between Cx43 average plaque number and heart weight values (C) found for chagasic and control patients.  $r^2 = 0.8244$ . CCC: chagasic patients with severe cardiomyopathy; IDCM: patients with non-infectious dilated cardiomyopathy. \*  $p < 0.05$ ; \*\*  $p < 0.001$ .

cal constraints. Heart weight from the chagasic group was higher than the non-infected controls (mean weight, 457.5 g compared to 235 g, respectively). Data in Table II allowed a correlation study between Cx43 protein expression levels and heart weight (Fig. 4C), depicting a clear and significant negative correlation between these parameters ( $r^2 = 0.8244$ ). Fewer Cx43 plaques were observed in heavier hearts. We could not predict a correlation with the presence of cardiac insufficiency and/or myocarditis with the number of Cx43 plaques because all chagasic patients presented with myocarditis; however, the Cx43 level was not homogeneous (Table II).

## DISCUSSION

In the heart, gap junctions mediate electronic current flow, thereby coordinating the spread of excitation and subsequent contraction throughout the myocardium (Spray & Burt 1990). Intercalated discs join neighbouring cardiomyocytes and provide mechanical adhesion as well as electrical and metabolic coupling. Protein changes in these junctions, particularly in gap junctions, are known to be involved in pathological features (Severs 2001). The disturbance of gap junction signalling could lead to altered impulse propagation between cardiomyocytes and to ventricular arrhythmogenesis in the myopathic heart (Peters et al. 1993). In CCC, one of the main complications is heart failure in association with extensive fibrosis (Higuchi et al. 1999), high circulating TGF- $\beta$  levels (Araújo-Jorge et al. 2008) and arrhythmia (Casado et al. 1990). To examine the effect of TGF- $\beta$  on Cx43 protein expression, we employed an in vitro model of cardiomyocytes submitted to treatment with the cytokine, its inhibitor, or *T. cruzi* infection. Our findings demonstrated that TGF- $\beta$  addition altered the Cx43 protein expression in non-infected myocytes, resulting in reduced organisation of gap junctions, which led to a punctate, diffuse and non-uniform pattern of staining.

We observed a decrease in Cx43 staining in infected cells and their neighbours, thus confirming what has already been demonstrated (Campos de Carvalho et al. 1992, Adesse et al. 2008). These results predicted that direct *T. cruzi* infection of heart cells could slow cardiac conduction and alter the pattern of impulse propagation as a consequence of decreased strength of electrotonic coupling. Increased local production and secretion of TGF- $\beta$ -1, as we demonstrated, could also contribute to impaired gap junctional communication in cardiomyocytes. The involvement of TGF- $\beta$  signalling in the mechanism of Cx43 decrease was verified when non-infected cardiomyocytes also developed a punctate Cx43 labelling after exposure to exogenous TGF- $\beta$ , similar to the pattern observed in infected cells. It was further reinforced by the interesting results obtained with the ALK5 inhibitor, SB-431542, which completely reversed the effect of TGF- $\beta$  and of *T. cruzi* infection on Cx43 expression as detected by immunoreaction and by western blot analysis. These results indicate a possible role of endogenously secreted TGF- $\beta$  as an autocrine/paracrine agent. This result suggests that TGF- $\beta$  produced in infected cultures could affect both infected and non-infected cells and induce the diffuse pattern of Cx43



staining and could be implicated in the decrease of Cx43 protein expression in infected cells. In addition, this decrease may be related to a general instability of global cytoplasmic mRNA that is commonly associated with *T. cruzi* intracellular proliferation (Pereira et al. 2000). In macrophages, it has been demonstrated that the altered electrical conduction between cells was due to a different pattern of Cx43 localisation rather than altered Cx43 expression levels (Fortes et al. 2004).

Our present data agree well with data obtained in chagasic patients, who exhibited severe diffuse fibrosis (increased fibronectin expression associated with a strong nuclear stain for PS2), indicating higher levels of active TGF- $\beta$  (Araújo-Jorge et al. 2008). In this study, we observed a significant down-regulation of Cx43 protein expression, a reduction in the length of Cx43 plaques and an altered distribution of these plaques in the chagasic group. These observations may account for the abnormal impulse conduction commonly observed in electrocardiogram profiles of Chagas disease patients and for myocytolysis leading to intense heart fibrosis contributing to cardiomegaly, as shown in Fig. 4C. These data are consistent with previous studies of patients with heart failure of different origins (Kostin et al. 2003). Disruption of the normal pattern of gap junction distribution also occurs in myocytes bordering areas of fibrotic myocardial damage and in areas of myocardial inflammation, where impairment of the impulse conduction was observed (Spach & Boineau 1997). *T. cruzi* infection is a strong inducer of TGF- $\beta$  expression in several cell types (Waghabi et al. 2002). Higher amounts of TGF- $\beta$  could be implicated either in the gap junction disassembly or in the observed extensive fibrosis, but probably plays a role in both phenomena. Higher levels of TGF- $\beta$  and disorganised expression of Cx43 could both act in synergy to promote heart fibrosis, thus leading to a more severe cardiopathy.

In summary, our results indicate that an important component of the autocrine myocyte response to TGF- $\beta$  secretion is the change of Cx43 protein expression, which contributes to the disturbance of cardiac electrical conduction. Taken together, the results shown here introduce more elements to the complex picture of heart fibrogenesis and arrhythmia, thus connecting TGF- $\beta$  and Cx43 dysregulation as key events in Chagas disease cardiomyopathy.

#### ACKNOWLEDGEMENTS

To Mr. Daniel Ciantar and Dr. Carlos Alberto Bizzarro Rodrigues, for the confocal analysis, to Dr. Gillian E. Knight, for editorial assistance, and Dr. Joseli Lannes-Vieira, for her generous gift of anti-Cx43 antibodies.

#### REFERENCES

- Adesse D, Garzoni LR, Huang H, Tanowitz HB, Meirelles MN, Spray DC 2008. *Trypanosoma cruzi* induces changes in cardiac connexin43 expression. *Microbes Infect* 10: 21-28.
- Araújo-Jorge TC, Waghabi MC, Hasslocher-Moreno AM, Xavier SS, Higuchi ML, Keramidas M, Bailly S, Feige JJ 2002. Implication of transforming growth factor-beta1 in Chagas disease cardiomyopathy. *J Infect Dis* 186: 1823-1828.
- Araújo-Jorge TC, Waghabi MC, Soeiro MNC, Keramidas M, Bailly S, Feige JJ 2008. Pivotal role for TGF- $\beta$  in infectious heart disease: the case of *Trypanosoma cruzi* infection and its consequent chagasic cardiomyopathy. *Cytok Growth Factors Rev* 19: 405-413.
- Bastide B, Neyses L, Ganten D, Paul M, Willecke K, Traub O 1993. Gap junction protein connexin40 is preferentially expressed in vascular endothelium and conductive bundles of rat myocardium and is increased under hypertensive conditions. *Circ Res* 73: 1138-1149.
- Beardslee MA, Lerner DL, Tadros PN, Laing JG, Beyer EC, Yamada KA, Kleber AG, Schuessler RB, Saffitz JE 2000. Dephosphorylation and intracellular redistribution of ventricular connexin43 during electrical uncoupling induced by ischemia. *Circ Res* 87: 656-662.
- Becker DL, Davies CS 1995. Role of gap junctions in the development of the preimplantation mouse embryo. *Microsc Res Tech* 31: 364-374.
- Campos de Carvalho AC, Tanowitz HB, Wittner M, Dermietzel R, Roy C, Hertzberg EL, Spray DC 1992. Gap junction distribution is altered between cardiac myocytes infected with *Trypanosoma cruzi*. *Circ Res* 70: 733-742.
- Casado J, Davila DF, Donis JH, Torres A, Payares A, Colmenares R, Gottberg CF 1990. Electrocardiographic abnormalities and left ventricular systolic function in Chagas' heart disease. *Int J Cardiol* 27: 55-62.
- Chanson M, Derouette JP, Roth I, Foglia B, Scerri I, Dudez T, Kwak BR 2005. Gap junctional communication in tissue inflammation and repair. *Biochim Biophys Acta* 1711: 197-207.
- Chen X, Zhang Y 2006. Myocardial Cx43 expression in the cases of sudden death due to dilated cardiomyopathy. *Forensic Sci Int* 162: 170-173.
- Coutinho-Silva R, Parsons M, Robson T, Burnstock G 2001. Changes in expression of P2 receptor in rat and mouse pancreas during development and aging. *Cell Tissue Res* 306: 373-383.
- Flanders KC, Holder MG, Winokur TS 1995. Autoinduction of mRNA and protein expression for transforming growth factor-beta S in cultured cardiac cells. *J Mol Cell Cardiol* 27: 805-812.
- Fortes FSA, Pecora IL, Costa V, Coutinho-Silva R, Bisaggio RC, Pires-de-Farias F, Persechini PM, Campos-de-Carvalho AC, Goldenberg RC 2004. Modulation of intercellular communication in macrophages: possible interactions between gap junctions and P2 receptors. *J Cell Sci* 117: 4717-4726.
- Higuchi ML, Fukasawa S, De Brito T, Parzianello LC, Bellotti G, Ramirez JA 1999. Different microcirculatory and interstitial matrix patterns in idiopathic dilated cardiomyopathy and Chagas' disease: a three dimensional confocal microscopy study. *Heart* 82: 279-285.
- Higuchi ML, Reis MM, Aiello VD, Benvenuti LA, Gutierrez PS, Bellotti G, Pileggi F 1997. Association of an increase in CD8<sup>+</sup> T cells with the presence of *Trypanosoma cruzi* antigens in chronic, human, chagasic myocarditis. *Am J Trop Med Hyg* 56: 485-489.
- Hurst VIV, Goldberg PL, Minnear FL, Heimark RL, Vincent PA 1999. Rearrangement of adherens junctions by transforming growth factor-beta1: role of contraction. *Am J Physiol* 276: 582-595.
- Inman GJ, Nicolas FJ, Callahan JF, Harling JD, Gaster LM, Reith AD, Laping NJ, Hill CS 2002. SB-431542 is a potent and specific inhibitor of transforming growth factor-beta superfamily type I activin receptor-like kinase (ALK) receptors ALK4, ALK5 and ALK7. *Mol Pharmacol* 62: 65-74.
- Jiang L, Bardini M, Keogh A, Remedios CG, Burnstock G 2005. P2X1 receptors are closely associated with connexin43 in human ventricular myocardium. *Int J of Cardiol* 98: 291-297.

- Kaprielian RR, Gunning M, Dupont E, Sheppard MN, Rothery SM, Underwood R, Pennell DJ, Fox K, Pepper J, Poole-Wilson PA, Severs NJ 1998. Downregulation of immunodetectable connexin43 and decreased gap junction size in the pathogenesis of chronic hibernation in the human left ventricle. *Circulation* 97: 651-660.
- Kostin S, Rieger M, Dammer S, Hein S, Richter M, Klovekorn WP, Bauer EP, Schaper J 2003. Gap junction remodeling and altered connexin43 expression in the failing human heart. *Mol Cell Biochem* 242: 135-144.
- Marin-Neto JA, Cunha-Neto E, Maciel BC, Simões MV 2007. Pathogenesis of chronic Chagas heart disease. *Circulation* 115: 1109-1123.
- Meirelles MNL, Araújo-Jorge TC, Miranda CF, De Souza W, Barbosa HS 1986. Interaction of *Trypanosoma cruzi* with heart muscle cells: ultrastructural and cytochemical analysis of endocytic vacuole formation and effect upon myogenesis in vitro. *Eur J Cell Biol* 41: 198-206.
- Mori R, Power KT, Wang CM, Martin P, Becker DL 2006. Acute downregulation of connexin43 at wound sites leads to a reduced inflammatory response, enhanced keratinocyte proliferation and wound fibroblast migration. *J Cell Sci* 119: 5193-5203.
- Palomino SA, Aiello VD, Higuchi ML 2000. Systematic mapping of hearts from chronic chagasic patients: the association between the occurrence of histopathological lesions and *Trypanosoma cruzi* antigens. *Ann Trop Med Parasitol* 94: 571-579.
- Pereira MC, Singer RH, Meirelles MN 2000. Ultrastructural distribution of poly (A)+ RNA during *Trypanosoma cruzi*-cardiomyocyte interaction *in vitro*: a quantitative analysis of the total mRNA content by in situ hybridization. *J Eukaryot Microbiol* 47: 264-270.
- Peters NS, Green CR, Poole-Wilson PA, Severs NJ 1993. Reduced content of connexin43 gap junctions in ventricular myocardium from hypertrophied and ischemic human hearts. *Circulation* 88: 864-875.
- Reis MM, Higuchi ML, Benvenuti LA, Aiello VD, Gutierrez PS, Bellotti G, Pileggi F 1997. An *in situ* quantitative immunohistochemical study of cytokines and IL-2R+ in chronic human chagasic myocarditis: correlation with the presence of myocardial *Trypanosoma cruzi* antigens. *Clin Immunol Immunopathol* 83: 165-172.
- Rodriguez-Sinovas A, Cabestrero A, Lopez D, Torre I, Morente M, Abellan A, Miro E, Ruiz-Meana M, Garcia-Dorado D 2007. The modulatory effects of connexin43 on cell death/survival beyond cell coupling. *Prog Biophys Mol Biol* 94: 219-232.
- Rosendahl A, Checchin D, Fehniger TE, ten Dijke P, Heldin CH, Sideras P 2001. Activation of the TGF- $\beta$ /activin-Smad2 pathway during allergic airway inflammation. *Am J Respir Cell Mol Biol* 25: 60-68.
- Rudkin GH, Yamaguchi DT, Ishida K, Peterson WJ, Bahadosingh F, Thye D, Miller TA 1996. Transforming growth factor-beta, osteogenin and bone morphogenetic protein-2 inhibit intercellular communication and alter cell proliferation in MC3T3-E1 cells. *J Cell Physiol* 168: 433-441.
- Schmunis G A 2007. Epidemiology of Chagas disease in non-endemic countries: the role of international migration. *Mem Inst Oswaldo Cruz* 102: 75-85.
- Sepp R, Severs NJ, Gourdie RG 1996. Altered patterns of cardiac intercellular junction distribution in hypertrophic cardiomyopathy. *Heart* 76: 412-417.
- Severs NJ 2001. Gap junction remodeling and cardiac arrhythmogenesis: cause or coincidence? *J Cell Mol Med* 5: 355-366.
- Severs NJ, Coppin SR, Dupont E, Yeh HI, Ko YS, Matsushita T 2004. Gap junction alterations in human cardiac disease. *Cardiovasc Res* 62: 368-377.
- Severs NJ, Dupont E, Coppin SR, Halliday D, Inett E, Baylis D, Rothery S 2004. Remodelling of gap junctions and connexin expression in heart disease. *Biochim Biophys Acta* 1662: 138-148.
- Silva JS, Twardzik DR, Reed SG 1991. Regulation of *Trypanosoma cruzi* infections in vitro and *in vivo* by transforming growth factor beta (TGF-beta). *J Exp Med* 174: 539-545.
- Sohl G, Willecke K 2004. Gap junctions and the connexin protein family. *Cardiovasc Res* 62: 228-232.
- Spach MS, Boineau JP 1997. Microfibrosis produces electrical load variations due to loss of side-to-side cell connections: a major mechanism of structural heart disease arrhythmias. *Pacing Clin Electrophysiol* 20: 397-413.
- Spray DC, Burt JM 1990. Structure-activity relations of the cardiac gap junction channel. *Am J Physiol* 258: 195-205.
- Tabibzadeh S 2002. Homeostasis of extracellular matrix by TGF-beta and lefty. *Front Biosci* 7: 1231-1246.
- Tansey EE, Kwaku KF, Hammer PE, Cowan DB, Federman M, Levitsky S, McCully JD 2006. Reduction and redistribution of gap and adherens junction proteins after ischemia and reperfusion. *Ann Thorac Surg* 82: 1472-1479.
- Uzzaman M, Honjo H, Takagishi Y, Emdad L, Magee AI, Severs NJ, Kodama I 2000. Remodeling of gap junctional coupling in hypertrophied right ventricles of rats with monocrotaline-induced pulmonary hypertension. *Circ Res* 86: 871-878.
- van Zoelen EJ, Tertoolen LG 1991. Transforming growth factor-beta enhances the extent of intercellular communication between normal rat kidney cells. *J Biol Chem* 266: 12075-12081.
- Waghbi MC, Keramidas M, Bailly S, Degraeve W, Mendonça-Lima L, Soeiro MNC, Meirelles MN, Paciornik S, Araújo-Jorge TC, Feige JJ 2005. Uptake of host cell transforming growth factor-beta by *Trypanosoma cruzi* amastigotes in cardiomyocytes: potential role in parasite cycle completion. *Am J Pathol* 167: 993-1003.
- Waghbi MC, Keramidas M, Calvet CM, Meuser M, Soeiro MNC, Mendonça-Lima L, Araújo-Jorge TC, Feige JJ, Bailly S 2002. Increased *Trypanosoma cruzi* invasion and heart fibrosis associated with high transforming growth factor beta levels in mice deficient in alpha(2)-macroglobulin. *Infect Immun* 70: 5115-5123.
- Waghbi MC, Keramidas M, Calvet CM, Meuser M, Soeiro MNC, Mendonça-Lima L, Araújo-Jorge TC, Feige JJ, Bailly S 2007. SB-431542, a transforming growth factor beta inhibitor, impairs *Trypanosoma cruzi* infection in cardiomyocytes and parasite cycle completion. *Antimicrob Agents Chemother* 51: 2905-2910.