

Search for New Physics in High Mass Electron-Positron Events in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

- T. Aaltonen,²³ A. Abulencia,²⁴ J. Adelman,¹³ T. Affolder,¹⁰ T. Akimoto,⁵⁵ M.G. Albrow,¹⁷ S. Amerio,⁴³ D. Amidei,³⁵ A. Anastassov,⁵² K. Anikeev,¹⁷ A. Annovi,¹⁹ J. Antos,¹⁴ M. Aoki,⁵⁵ G. Apollinari,¹⁷ T. Arisawa,⁵⁷ A. Artikov,¹⁵ W. Ashmanskas,¹⁷ A. Attal,³ A. Aurisano,⁵³ F. Azfar,⁴² P. Azzi-Bacchetta,⁴³ P. Azzurri,⁴⁶ N. Bacchetta,⁴³ W. Badgett,¹⁷ A. Barbaro-Galtieri,²⁹ V.E. Barnes,⁴⁸ B.A. Barnett,²⁵ S. Baroiant,⁷ V. Bartsch,³¹ G. Bauer,³³ P.-H. Beauchemin,³⁴ F. Bedeschi,⁴⁶ S. Behari,²⁵ G. Bellettini,⁴⁶ J. Bellinger,⁵⁹ A. Belloni,³³ D. Benjamin,¹⁶ A. Beretvas,¹⁷ J. Beringer,²⁹ T. Berry,³⁰ A. Bhatti,⁵⁰ M. Binkley,¹⁷ D. Bisello,⁴³ I. Bizjak,³¹ R.E. Blair,² C. Blocker,⁶ B. Blumenfeld,²⁵ A. Bocci,¹⁶ A. Bodek,⁴⁹ V. Boisvert,⁴⁹ G. Bolla,⁴⁸ A. Bolshov,³³ D. Bortoletto,⁴⁸ J. Boudreau,⁴⁷ A. Boveia,¹⁰ B. Brau,¹⁰ L. Brigliadori,⁵ C. Bromberg,³⁶ E. Brubaker,¹³ J. Budagov,¹⁵ H.S. Budd,⁴⁹ S. Budd,²⁴ K. Burkett,¹⁷ G. Busetto,⁴³ P. Bussey,²¹ A. Buzatu,³⁴ K. L. Byrum,² S. Cabrera^q,¹⁶ M. Campanelli,²⁰ M. Campbell,³⁵ F. Canelli,¹⁷ A. Canepa,⁴⁵ S. Carrilloⁱ,¹⁸ D. Carlsmith,⁵⁹ R. Carosi,⁴⁶ S. Carron,³⁴ B. Casal,¹¹ M. Casarsa,⁵⁴ A. Castro,⁵ P. Catastini,⁴⁶ D. Cauz,⁵⁴ M. Cavalli-Sforza,³ A. Cerri,²⁹ L. Cerrito^m,³¹ S.H. Chang,²⁸ Y.C. Chen,¹ M. Chertok,⁷ G. Chiarelli,⁴⁶ G. Chlachidze,¹⁷ F. Chlebana,¹⁷ I. Cho,²⁸ K. Cho,²⁸ D. Chokheli,¹⁵ J.P. Chou,²² G. Choudalakis,³³ S.H. Chuang,⁵² K. Chung,¹² W.H. Chung,⁵⁹ Y.S. Chung,⁴⁹ M. Cilijak,⁴⁶ C.I. Ciobanu,²⁴ M.A. Ciocci,⁴⁶ A. Clark,²⁰ D. Clark,⁶ M. Coca,¹⁶ G. Compostella,⁴³ M.E. Convery,⁵⁰ J. Conway,⁷ B. Cooper,³¹ K. Copic,³⁵ M. Cordelli,¹⁹ G. Cortiana,⁴³ F. Crescioli,⁴⁶ C. Cuenca Almenar^q,⁷ J. Cuevas^l,¹¹ R. Culbertson,¹⁷ J.C. Cully,³⁵ S. DaRonco,⁴³ M. Datta,¹⁷ S. D'Auria,²¹ T. Davies,²¹ D. Dagenhart,¹⁷ P. de Barbaro,⁴⁹ S. De Cecco,⁵¹ A. Deisher,²⁹ G. De Lentdecker^c,⁴⁹ G. De Lorenzo,³ M. Dell'Orso,⁴⁶ F. Delli Paoli,⁴³ L. Demortier,⁵⁰ J. Deng,¹⁶ M. Deninno,⁵ D. De Pedis,⁵¹ P.F. Derwent,¹⁷ G.P. Di Giovanni,⁴⁴ C. Dionisi,⁵¹ B. Di Ruzza,⁵⁴ J.R. Dittmann,⁴ M. D'Onofrio,³ C. Dörr,²⁶ S. Donati,⁴⁶ P. Dong,⁸ J. Donini,⁴³ T. Dorigo,⁴³ S. Dube,⁵² J. Efron,³⁹ R. Erbacher,⁷ D. Errede,²⁴ S. Errede,²⁴ R. Eusebi,¹⁷ H.C. Fang,²⁹ S. Farrington,³⁰ I. Fedorko,⁴⁶ W.T. Fedorko,¹³ R.G. Feild,⁶⁰ M. Feindt,²⁶ J.P. Fernandez,³² R. Field,¹⁸ G. Flanagan,⁴⁸ R. Forrest,⁷ S. Forrester,⁷ M. Franklin,²² J.C. Freeman,²⁹ I. Furic,¹³ M. Gallinaro,⁵⁰ J. Galyardt,¹² J.E. Garcia,⁴⁶ F. Garberson,¹⁰ A.F. Garfinkel,⁴⁸ C. Gay,⁶⁰ H. Gerberich,²⁴ D. Gerdes,³⁵ S. Giagu,⁵¹ P. Giannetti,⁴⁶ K. Gibson,⁴⁷ J.L. Gimmell,⁴⁹ C. Ginsburg,¹⁷ N. Giokaris^a,¹⁵ M. Giordani,⁵⁴ P. Giromini,¹⁹ M. Giunta,⁴⁶ G. Giurgiu,²⁵ V. Glagolev,¹⁵ D. Glenzinski,¹⁷ M. Gold,³⁷ N. Goldschmidt,¹⁸ J. Goldstein^b,⁴² A. Golossanov,¹⁷ G. Gomez,¹¹ G. Gomez-Ceballos,³³ M. Goncharov,⁵³ O. González,³² I. Gorelov,³⁷ A.T. Goshaw,¹⁶ K. Goulianatos,⁵⁰ A. Gresele,⁴³ S. Grinstein,²² C. Gross-Pilcher,¹³ R.C. Group,¹⁷ U. Grundler,²⁴ J. Guimaraes da Costa,²² Z. Gunay-Unalan,³⁶ C. Haber,²⁹ K. Hahn,³³ S.R. Hahn,¹⁷ E. Halkiadakis,⁵² A. Hamilton,²⁰ B.-Y. Han,⁴⁹ J.Y. Han,⁴⁹ R. Handler,⁵⁹ F. Happacher,¹⁹ K. Hara,⁵⁵ D. Hare,⁵² M. Hare,⁵⁶ S. Harper,⁴² R.F. Harr,⁵⁸ R.M. Harris,¹⁷ M. Hartz,⁴⁷ K. Hatakeyama,⁵⁰ J. Hauser,⁸ C. Hays,⁴² M. Heck,²⁶ A. Heijboer,⁴⁵ B. Heinemann,²⁹ J. Heinrich,⁴⁵ C. Henderson,³³ M. Herndon,⁵⁹ J. Heuser,²⁶ D. Hidas,¹⁶ C.S. Hill^b,¹⁰ D. Hirschbuehl,²⁶ A. Hocker,¹⁷ A. Holloway,²² S. Hou,¹ M. Houlden,³⁰ S.-C. Hsu,⁹ B.T. Huffman,⁴² R.E. Hughes,³⁹ U. Husemann,⁶⁰ J. Huston,³⁶ J. Incandela,¹⁰ G. Introzzi,⁴⁶ M. Iori,⁵¹ A. Ivanov,⁷ B. Iyutin,³³ E. James,¹⁷ D. Jang,⁵² B. Jayatilaka,¹⁶ D. Jeans,⁵¹ E.J. Jeon,²⁸ S. Jindariani,¹⁸ W. Johnson,⁷ M. Jones,⁴⁸ K.K. Joo,²⁸ S.Y. Jun,¹² J.E. Jung,²⁸ T.R. Junk,²⁴ T. Kamon,⁵³ P.E. Karchin,⁵⁸ Y. Kato,⁴¹ Y. Kemp,²⁶ R. Kephart,¹⁷ U. Kerzel,²⁶ V. Khotilovich,⁵³ B. Kilminster,³⁹ D.H. Kim,²⁸ H.S. Kim,²⁸ J.E. Kim,²⁸ M.J. Kim,¹⁷ S.B. Kim,²⁸ S.H. Kim,⁵⁵ Y.K. Kim,¹³ N. Kimura,⁵⁵ L. Kirsch,⁶ S. Klimenko,¹⁸ M. Klute,³³ B. Knuteson,³³ B.R. Ko,¹⁶ K. Kondo,⁵⁷ D.J. Kong,²⁸ J. Konigsberg,¹⁸ A. Korytov,¹⁸ A.V. Kotwal,¹⁶ A.C. Kraan,⁴⁵ J. Kraus,²⁴ M. Kreps,²⁶ J. Kroll,⁴⁵ N. Krumnack,⁴ M. Kruse,¹⁶ V. Krutelyov,¹⁰ T. Kubo,⁵⁵ S. E. Kuhlmann,² T. Kuhr,²⁶ N.P. Kulkarni,⁵⁸ Y. Kusakabe,⁵⁷ S. Kwang,¹³ A.T. Laasanen,⁴⁸ S. Lai,³⁴ S. Lami,⁴⁶ S. Lammel,¹⁷ M. Lancaster,³¹ R.L. Lander,⁷ K. Lannon,³⁹ A. Lath,⁵² G. Latino,⁴⁶ I. Lazzizzera,⁴³ T. LeCompte,² J. Lee,⁴⁹ J. Lee,²⁸ Y.J. Lee,²⁸ S.W. Lee^o,⁵³ R. Lefèvre,²⁰ N. Leonardo,³³ S. Leone,⁴⁶ S. Levy,¹³ J.D. Lewis,¹⁷ C. Lin,⁶⁰ C.S. Lin,¹⁷ M. Lindgren,¹⁷ E. Lipeles,⁹ A. Lister,⁷ D.O. Litvintsev,¹⁷ T. Liu,¹⁷ N.S. Lockyer,⁴⁵ A. Loginov,⁶⁰ M. Loretta,⁴³ R.-S. Lu,¹ D. Lucchesi,⁴³ P. Lujan,²⁹ P. Lukens,¹⁷ G. Lungu,¹⁸ L. Lyons,⁴² J. Lys,²⁹ R. Lysak,¹⁴ E. Lytken,⁴⁸ P. Mack,²⁶ D. MacQueen,³⁴ R. Madrak,¹⁷ K. Maeshima,¹⁷ K. Makhoul,³³ T. Maki,²³ P. Maksimovic,²⁵ S. Malde,⁴² S. Malik,³¹ G. Manca,³⁰ A. Manousakis^a,¹⁵ F. Margaroli,⁵ R. Marginean,¹⁷ C. Marino,²⁶ C.P. Marino,²⁴ A. Martin,⁶⁰ M. Martin,²⁵ V. Martin^g,²¹ M. Martínez,³ R. Martínez-Ballarín,³² T. Maruyama,⁵⁵ P. Mastrandrea,⁵¹ T. Masubuchi,⁵⁵ H. Matsunaga,⁵⁵ M.E. Mattson,⁵⁸ R. Mazini,³⁴ P. Mazzanti,⁵ K.S. McFarland,⁴⁹ P. McIntyre,⁵³ R. McNulty^f,³⁰ A. Mehta,³⁰ P. Mehtala,²³ S. Menzemer^h,¹¹ A. Menzione,⁴⁶

P. Merkel,⁴⁸ C. Mesropian,⁵⁰ A. Messina,³⁶ T. Miao,¹⁷ N. Miladinovic,⁶ J. Miles,³³ R. Miller,³⁶ C. Mills,¹⁰ M. Milnik,²⁶ A. Mitra,¹ G. Mitselmakher,¹⁸ A. Miyamoto,²⁷ S. Moed,²⁰ N. Moggi,⁵ B. Mohr,⁸ C.S. Moon,²⁸ R. Moore,¹⁷ M. Morello,⁴⁶ P. Movilla Fernandez,²⁹ J. Mülmenstädt,²⁹ A. Mukherjee,¹⁷ Th. Muller,²⁶ R. Mumford,²⁵ P. Murat,¹⁷ M. Mussini,⁵ J. Nachtman,¹⁷ A. Nagano,⁵⁵ J. Naganoma,⁵⁷ K. Nakamura,⁵⁵ I. Nakano,⁴⁰ A. Napier,⁵⁶ V. Necula,¹⁶ C. Neu,⁴⁵ M.S. Neubauer,⁹ J. Nielsenⁿ,²⁹ L. Nodulman,² O. Norniella,³ E. Nurse,³¹ S.H. Oh,¹⁶ Y.D. Oh,²⁸ I. Oksuzian,¹⁸ T. Okusawa,⁴¹ R. Oldeman,³⁰ R. Orava,²³ K. Osterberg,²³ C. Pagliarone,⁴⁶ E. Palencia,¹¹ V. Papadimitriou,¹⁷ A. Papaikonomou,²⁶ A.A. Paramonov,¹³ B. Parks,³⁹ S. Pashapour,³⁴ J. Patrick,¹⁷ G. Pauletta,⁵⁴ M. Paulini,¹² C. Paus,³³ D.E. Pellett,⁷ A. Penzo,⁵⁴ T.J. Phillips,¹⁶ G. Piacentino,⁴⁶ J. Piedra,⁴⁴ L. Pinera,¹⁸ K. Pitts,²⁴ C. Plager,⁸ L. Pondrom,⁵⁹ X. Portell,³ O. Poukhov,¹⁵ N. Pounder,⁴² F. Prakoshyn,¹⁵ A. Pronko,¹⁷ J. Proudfoot,² F. Ptohos^e,¹⁹ G. Punzi,⁴⁶ J. Pursley,²⁵ J. Rademacker^b,⁴² A. Rahaman,⁴⁷ V. Ramakrishnan,⁵⁹ N. Ranjan,⁴⁸ I. Redondo,³² B. Reisert,¹⁷ V. Rekovic,³⁷ P. Renton,⁴² M. Rescigno,⁵¹ S. Richter,²⁶ F. Rimondi,⁵ L. Ristori,⁴⁶ A. Robson,²¹ T. Rodrigo,¹¹ E. Rogers,²⁴ S. Rolli,⁵⁶ R. Roser,¹⁷ M. Rossi,⁵⁴ R. Rossin,¹⁰ P. Roy,³⁴ A. Ruiz,¹¹ J. Russ,¹² V. Rusu,¹³ H. Saarikko,²³ A. Safonov,⁵³ W.K. Sakumoto,⁴⁹ G. Salamanna,⁵¹ O. Saltó,³ L. Santi,⁵⁴ S. Sarkar,⁵¹ L. Sartori,⁴⁶ K. Sato,¹⁷ P. Savard,³⁴ A. Savoy-Navarro,⁴⁴ T. Scheidle,²⁶ P. Schlabach,¹⁷ E.E. Schmidt,¹⁷ M.P. Schmidt,⁶⁰ M. Schmitt,³⁸ T. Schwarz,⁷ L. Scodellaro,¹¹ A.L. Scott,¹⁰ A. Scribano,⁴⁶ F. Scuri,⁴⁶ A. Sedov,⁴⁸ S. Seidel,³⁷ Y. Seiya,⁴¹ A. Semenov,¹⁵ L. Sexton-Kennedy,¹⁷ A. Sfyrla,²⁰ S.Z. Shalhout,⁵⁸ M.D. Shapiro,²⁹ T. Shears,³⁰ P.F. Shepard,⁴⁷ D. Sherman,²² M. Shimojima^k,⁵⁵ M. Shochet,¹³ Y. Shon,⁵⁹ I. Shreyber,²⁰ A. Sidoti,⁴⁶ P. Sinervo,³⁴ A. Sisakyan,¹⁵ A.J. Slaughter,¹⁷ J. Slaunwhite,³⁹ K. Sliwa,⁵⁶ J.R. Smith,⁷ F.D. Snider,¹⁷ R. Snihur,³⁴ M. Soderberg,³⁵ A. Soha,⁷ S. Somalwar,⁵² V. Sorin,³⁶ J. Spalding,¹⁷ F. Spinella,⁴⁶ T. Spreitzer,³⁴ P. Squillacioti,⁴⁶ M. Stanitzki,⁶⁰ A. Staveris-Polykalas,⁴⁶ R. St. Denis,²¹ B. Stelzer,⁸ O. Stelzer-Chilton,⁴² D. Stentz,³⁸ J. Strologas,³⁷ D. Stuart,¹⁰ J.S. Suh,²⁸ A. Sukhanov,¹⁸ H. Sun,⁵⁶ I. Suslov,¹⁵ T. Suzuki,⁵⁵ A. Taffard^p,²⁴ R. Takashima,⁴⁰ Y. Takeuchi,⁵⁵ R. Tanaka,⁴⁰ M. Tecchio,³⁵ P.K. Teng,¹ K. Terashi,⁵⁰ J. Thom^d,¹⁷ A.S. Thompson,²¹ E. Thomson,⁴⁵ P. Tipton,⁶⁰ V. Tiwari,¹² S. Tkaczyk,¹⁷ D. Toback,⁵³ S. Tokar,¹⁴ K. Tollefson,³⁶ T. Tomura,⁵⁵ D. Tonelli,⁴⁶ S. Torre,¹⁹ D. Torretta,¹⁷ S. Tourneur,⁴⁴ W. Trischuk,³⁴ S. Tsuno,⁴⁰ Y. Tu,⁴⁵ N. Turini,⁴⁶ F. Ukegawa,⁵⁵ S. Uozumi,⁵⁵ S. Vallecorsa,²⁰ N. van Remortel,²³ A. Varganov,³⁵ E. Vataga,³⁷ F. Vazquezⁱ,¹⁸ G. Velev,¹⁷ C. Vellidis^a,⁴⁶ G. Veramendi,²⁴ V. Veszpreni,⁴⁸ M. Vidal,³² R. Vidal,¹⁷ I. Vila,¹¹ R. Vilar,¹¹ T. Vine,³¹ M. Vogel,³⁷ I. Vollrath,³⁴ I. Volobouev^o,²⁹ G. Volpi,⁴⁶ F. Würthwein,⁹ P. Wagner,⁵³ R.G. Wagner,² R.L. Wagner,¹⁷ J. Wagner,²⁶ W. Wagner,²⁶ R. Wallny,⁸ S.M. Wang,¹ A. Warburton,³⁴ D. Waters,³¹ M. Weinberger,⁵³ W.C. Wester III,¹⁷ B. Whitehouse,⁵⁶ D. Whiteson^p,⁴⁵ A.B. Wicklund,² E. Wicklund,¹⁷ G. Williams,³⁴ H.H. Williams,⁴⁵ P. Wilson,¹⁷ B.L. Winer,³⁹ P. Wittich^d,¹⁷ S. Wolbers,¹⁷ C. Wolfe,¹³ T. Wright,³⁵ X. Wu,²⁰ S.M. Wynne,³⁰ A. Yagil,⁹ K. Yamamoto,⁴¹ J. Yamaoka,⁵² T. Yamashita,⁴⁰ C. Yang,⁶⁰ U.K. Yang^j,¹³ Y.C. Yang,²⁸ W.M. Yao,²⁹ G.P. Yeh,¹⁷ J. Yoh,¹⁷ K. Yorita,¹³ T. Yoshida,⁴¹ G.B. Yu,⁴⁹ I. Yu,²⁸ S.S. Yu,¹⁷ J.C. Yun,¹⁷ L. Zanello,⁵¹ A. Zanetti,⁵⁴ I. Zaw,²² X. Zhang,²⁴ J. Zhou,⁵² and S. Zucchelli⁵

(CDF Collaboration*)

¹Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China

²Argonne National Laboratory, Argonne, Illinois 60439

³Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain

⁴Baylor University, Waco, Texas 76798

⁵Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy

⁶Brandeis University, Waltham, Massachusetts 02254

⁷University of California, Davis, Davis, California 95616

⁸University of California, Los Angeles, Los Angeles, California 90024

⁹University of California, San Diego, La Jolla, California 92093

¹⁰University of California, Santa Barbara, Santa Barbara, California 93106

¹¹Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain

¹²Carnegie Mellon University, Pittsburgh, PA 15213

¹³Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637

¹⁴Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia

¹⁵Joint Institute for Nuclear Research, RU-141980 Dubna, Russia

¹⁶Duke University, Durham, North Carolina 27708

¹⁷Fermi National Accelerator Laboratory, Batavia, Illinois 60510

¹⁸University of Florida, Gainesville, Florida 32611

¹⁹Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy

²⁰University of Geneva, CH-1211 Geneva 4, Switzerland

²¹Glasgow University, Glasgow G12 8QQ, United Kingdom

- ²²Harvard University, Cambridge, Massachusetts 02138
²³Division of High Energy Physics, Department of Physics,
 University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland
²⁴University of Illinois, Urbana, Illinois 61801
²⁵The Johns Hopkins University, Baltimore, Maryland 21218
²⁶Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany
²⁷High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan
²⁸Center for High Energy Physics: Kyungpook National University,
 Taegu 702-701, Korea; Seoul National University, Seoul 151-742,
 Korea; SungKyunKwan University, Suwon 440-746, Korea
²⁹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720
³⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom
³¹University College London, London WC1E 6BT, United Kingdom
³²Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
³³Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
³⁴Institute of Particle Physics: McGill University, Montréal,
 Canada H3A 2T8; and University of Toronto, Toronto, Canada M5S 1A7
³⁵University of Michigan, Ann Arbor, Michigan 48109
³⁶Michigan State University, East Lansing, Michigan 48824
³⁷University of New Mexico, Albuquerque, New Mexico 87131
³⁸Northwestern University, Evanston, Illinois 60208
³⁹The Ohio State University, Columbus, Ohio 43210
⁴⁰Okayama University, Okayama 700-8530, Japan
⁴¹Osaka City University, Osaka 588, Japan
⁴²University of Oxford, Oxford OX1 3RH, United Kingdom
⁴³University of Padova, Istituto Nazionale di Fisica Nucleare,
 Sezione di Padova-Trento, I-35131 Padova, Italy
⁴⁴LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
⁴⁵University of Pennsylvania, Philadelphia, Pennsylvania 19104
⁴⁶Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa,
 Siena and Scuola Normale Superiore, I-56127 Pisa, Italy
⁴⁷University of Pittsburgh, Pittsburgh, Pennsylvania 15260
⁴⁸Purdue University, West Lafayette, Indiana 47907
⁴⁹University of Rochester, Rochester, New York 14627
⁵⁰The Rockefeller University, New York, New York 10021
⁵¹Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,
 University of Rome "La Sapienza," I-00185 Roma, Italy
⁵²Rutgers University, Piscataway, New Jersey 08855
⁵³Texas A&M University, College Station, Texas 77843
⁵⁴Istituto Nazionale di Fisica Nucleare, University of Trieste/ Udine, Italy
⁵⁵University of Tsukuba, Tsukuba, Ibaraki 305, Japan
⁵⁶Tufts University, Medford, Massachusetts 02155
⁵⁷Waseda University, Tokyo 169, Japan
⁵⁸Wayne State University, Detroit, Michigan 48201
⁵⁹University of Wisconsin, Madison, Wisconsin 53706
⁶⁰Yale University, New Haven, Connecticut 06520

(Dated: February 1, 2008)

We report the results of a search for a narrow resonance in electron-positron events in the invariant mass range of $150 - 950 \text{ GeV}/c^2$ using 1.3 fb^{-1} of $p\bar{p}$ collision data at $\sqrt{s} = 1.96 \text{ TeV}$ collected by the CDF II detector at Fermilab. No significant evidence of such a resonance is observed and we interpret the results to exclude the standard model-like Z' with a mass below $923 \text{ GeV}/c^2$ and the Randall-Sundrum graviton with a mass below $807 \text{ GeV}/c^2$ for $k/\bar{M}_{pl}=0.1$, both at the 95% confidence level. Combining with di-photon data excludes the Randall-Sundrum graviton for masses below $889 \text{ GeV}/c^2$ for $k/\bar{M}_{pl}=0.1$.

PACS numbers: 13.85.Rm, 13.85.Qk, 14.70.Pw, 04.50.+h

*With visitors from ^aUniversity of Athens, 15784 Athens, Greece,
^bUniversity of Bristol, Bristol BS8 1TL, United Kingdom,

^cUniversity Libre de Bruxelles, B-1050 Brussels, Belgium, ^dCornell

At hadron colliders, electron-positron pairs (ee) are a distinct experimental signature with a low background rate. Since many models introducing new physics beyond the standard model of particle physics (SM) predict an excess in ee production at a hadron collider, this channel has a strong discovery potential. This Letter describes a search for a new high mass state in ee events from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. The data used in this analysis were collected by the CDF II detector at the Fermilab Tevatron and correspond to an integrated luminosity of 1.3 fb^{-1} . The analysis also uses results from the $\gamma\gamma$ channel described in [1] to increase the analysis's sensitivity to the Randall-Sundrum (RS) graviton [2].

The search is optimized for new physics processes which produce narrow ee resonances [3], but is otherwise model-independent. In addition to the above search, cross section times branching ratio ($\sigma \cdot \mathcal{B}$) upper limits [4] are set for generic neutral spin-1 and spin-2 bosons. These $\sigma \cdot \mathcal{B}$ limits are then used to set lower bounds on the masses of specific particles predicted by new physics models. These particles are the E_6 Z 's [5] and the RS graviton. The E_6 model unifies the forces of the SM into the E_6 gauge group and in doing so predicts the presence of two additional neutral massive spin-1 bosons, referred to as Z 's, which can mix with some arbitrary mixing angle. The Z'_η , Z'_χ , Z'_ψ , and Z'_I Z 's correspond to specific values of the mixing angle and are used to benchmark the model. The RS graviton is predicted by the RS model of warped extra dimensions which solves the hierarchy between the weak and Planck scales by introducing an extra spatial dimension with negative curvature k . The model predicts a series of narrow neutral spin-2 resonances which couple to all SM particles, with the lowest mass resonance referred to here as the RS graviton. The properties of this model are determined by the mass of the RS graviton and the ratio k/\bar{M}_{pl} , where \bar{M}_{pl} is the reduced effective Planck scale. This ratio governs the couplings of the graviton to SM particles and it has a favored range of 0.01 to 0.1 [6].

The most recent similar search by the DØ collaboration used an integrated luminosity of 260 pb^{-1} and treated the ee and the di-photon ($\gamma\gamma$) channels as a single channel to perform the first dedicated RS graviton search [7]. The

most recent search for new physics in the ee channel by the CDF collaboration was a dedicated Z' search and used an integrated luminosity of 448 pb^{-1} [8].

This analysis is based on an integrated luminosity of 1.3 fb^{-1} collected with the CDF II detector. The CDF II detector is a general purpose detector which is azimuthally and forward-backward symmetric and is described in detail elsewhere [9]. The relevant components for this analysis are the central tracking chamber (COT) and the central and plug calorimeters. The COT is a 96-layer drift chamber placed within a 1.4 Tesla magnetic field and is used to measure the momenta of charged particles within the pseudorapidity range $|\eta| \leq 1.1$ [10]. The COT is complemented by a silicon microstrip detector which directly surrounds the beampipe and has a tracking coverage of $|\eta| \leq 2.0$ [11]. The central and plug calorimeters are sampling calorimeters that surround the COT; they consist of electromagnetic (EM) and hadronic sections that measure the energy of particles in the range $|\eta| \leq 1.1$ and $1.2 \leq |\eta| \leq 3.6$ respectively.

The trigger used in this analysis requires two separate deposits of EM energy in the calorimeter and is effectively 100% efficient for selecting ee events within the acceptance of the analysis. Events are selected by requiring two electron candidates with $E_T \geq 25 \text{ GeV}$. Events are separated into two channels: central-central (CC) where both electrons are in the central EM calorimeter (CEM) ($|\eta| \leq 1.1$) and central-plug (CP) where one electron is in the plug EM calorimeter (PEM) ($1.2 \leq |\eta| \leq 3.0$). Electrons in the CEM are required to have a well-measured track, whereas there is no tracking requirement for electron candidates in the PEM. Due to the lack of tracking in the plug region there is no opposite sign requirement for any of the ee pairs. In the CC channel, 5% of the electron pairs in the signal region ($M_{ee} \geq 150 \text{ GeV}/c^2$) are same sign, which is compatible with the fraction of misidentified opposite sign pairs predicted by simulation. Electrons are identified in an identical way to the previously published analysis [8], with the exception that a photon conversion veto is applied to central electrons in CP events. This selection cut improves the sensitivity of the analysis by reducing the $\gamma\gamma$ background in this channel. The event selection and the search method defined later were chosen without regard to events observed in the signal region to ensure a statistical robust result.

The geometric and kinematic acceptance as a function of resonance mass is estimated using event samples generated by Monte Carlo (MC) simulation. The PYTHIA event generator [12], with the CTEQ5L parton distribution functions (PDF) [13] and the CDF II detector simulation based on GEANT 3 [14] are used to generate all simulation samples unless otherwise stated. A Z' with the couplings of the SM Z (SM-like Z') is used for the simulated spin-1 signal sample and a RS graviton with $k/\bar{M}_{pl} = 0.1$ is used for the spin-2 sample. Both the Z' and RS graviton bosons are constrained to be within

University, Ithaca, NY 14853, ^eUniversity of Cyprus, Nicosia CY-1678, Cyprus, ^fUniversity College Dublin, Dublin 4, Ireland, ^gUniversity of Edinburgh, Edinburgh EH9 3JZ, United Kingdom, ^hUniversity of Heidelberg, D-69120 Heidelberg, Germany, ⁱUniversidad Iberoamericana, Mexico D.F., Mexico, ^jUniversity of Manchester, Manchester M13 9PL, England, ^kNagasaki Institute of Applied Science, Nagasaki, Japan, ^lUniversity de Oviedo, E-33007 Oviedo, Spain, ^mUniversity of London, Queen Mary College, London, E1 4NS, England, ⁿUniversity of California Santa Cruz, Santa Cruz, CA 95064, ^oTexas Tech University, Lubbock, TX 79409, ^pUniversity of California, Irvine, Irvine, CA 92697, ^qIFIC(CSIC-Universitat de Valencia), 46071 Valencia, Spain,

$\pm 10\%$ of their on-shell mass at generator level. The uncertainty on the acceptance resulting from the PDF parameterization is estimated to be 2-4% using the procedure recommended by the CTEQ Collaboration [15]. The uncertainty on the acceptance due to initial state radiation (ISR) is estimated to be 4% by varying the parameters governing ISR in PYTHIA. The electron identification efficiency ranges from 90% to 95% per electron and is estimated using the simulated signal samples. These estimates are corrected for imperfections in the simulation by comparing the simulation with the data at the Z pole, resulting in a 2% systematic uncertainty. The total selection efficiency \times acceptance over the entire signal region is in the range of 40% to 45% for spin-1 and spin-2 bosons.

The most significant source of background to new physics in the ee channel is the SM Drell-Yan process via Z/γ^* which is an irreducible background. Jet events, such as di-jet or W +jet events where the jets are misidentified as electrons, represent the most significant reducible background. Other less significant backgrounds result from $t\bar{t}$, $\tau^+\tau^-$, WW , WZ , $W\gamma$, and $\gamma\gamma$ events. Figure 1 shows the invariant mass distribution for all the background components properly normalized, together with the observed data for the CC and CP channels combined.




FIG. 1: The measured ee mass spectrum with the expected background for the CC and CP channels combined. The backgrounds are displayed cumulatively. There are no observed events above $550 \text{ GeV}/c^2$.

The SM Drell-Yan contribution is estimated using MC simulated events normalized to the data at the Z pole. By investigating the stability of the normalization factor a 4% systematic uncertainty is obtained on the SM Drell-Yan normalization. An uncertainty on the SM Drell-Yan shape due to PDF uncertainties is determined using the same method as used for the acceptance. The di-jet and W +jet backgrounds are treated as a single background,

referred to as the jet background. The size of this background is estimated from a sample of jet events constructed from the data identically to the signal sample except that at least one electron candidate is not isolated [16] and therefore likely to be a jet. From the distribution of the isolation of these ‘electron’ candidates, the number of jet events in the signal sample and its uncertainty is extracted. The shape of the jet background is estimated from a jet+ Y sample, where Y is either an electron or a jet misidentified as an electron. In the region where the jet background is significant, the normalization uncertainty is the dominant uncertainty on the jet background. Using these methods, the jet background is estimated to account for $0.8 \pm 0.7\%$ and $25 \pm 8\%$ of the total background above $150 \text{ GeV}/c^2$ in the CC and CP channels respectively. The remaining backgrounds are all estimated using MC simulation normalized to the theoretical next-to-leading order (NLO) or higher order cross section. The uncertainties in these background estimates are dominated by the 6% uncertainty on the luminosity [17].

A model-independent search for an excess over SM predictions is performed in an invariant mass range of $150 - 950 \text{ GeV}/c^2$. The search is optimized for a narrow resonance, but still retains sensitivity to other signals which would produce an excess over SM predictions. Using $1 \text{ GeV}/c^2$ steps from $M_{ee} = 150$ to $950 \text{ GeV}/c^2$, the probability, referred to as the p-value, of observing at least as many events as recorded in the real experiment given the expected background rate is calculated using Poisson statistics in a mass window of $4.8 + 0.044 \times M_{ee} \text{ GeV}/c^2$. This mass window is approximately the width a narrow resonance would have if observed in the CDF detector, and this choice of mass window maximizes the sensitivity to discovering such a resonance for this particular analysis, as verified from studies of simulated events. The uncertainty on the background estimate is treated as a nuisance parameter with a Gaussian distribution. The p-values for the CC and CP channels are combined multiplicatively and the results are shown in Fig. 2. The minimum p-value expected in the absence of new physics depends on the size of the search range, with the expected minimum p-value decreasing as the search range increases. The range in which minimum p-value is expected to occur is shown in Fig. 2 and is defined to include 68.3% of the minimum p-values centered on the median value, using one million simulated mass spectra. Similarly, the 3σ evidence line is the p-value above which the minimum p-value in 99.85% of the simulated mass spectra fall; any p-value lower than this would be taken as evidence for the presence of new physics. The lowest p-value observed is at $367 \text{ GeV}/c^2$ and is within the expected range. It is therefore concluded that the results of this analysis are consistent with the SM.

To complement the above search, a Bayesian binned




FIG. 2: The probability that the background alone could give rise to the observed number of events in a mass window equal to the width of a narrow resonance in the CDF detector. The expected range and the 3σ evidence line are defined in the text. The region $550 - 950 \text{ GeV}/c^2$ is not shown as no events are observed in data and therefore the p-value is always 1 in this region.

likelihood method is used to extract limits on $\sigma \cdot \mathcal{B}(X \rightarrow ee)$, where the mass of X is within $\pm 10\%$ of its on-shell mass. As the acceptance of the final-state ee system is required to extract a cross section, it is necessary to specify the spin of the particle. Both spin-1 and spin-2 particles are considered here. The likelihood is 1-dimensional with the signal cross section as the free parameter and the bin contents treated using Poisson statistics. The likelihood is then convoluted with a Gaussian to allow for the uncertainty on the cross section from the acceptance, background, and luminosity estimates. The probability density function is formed by taking a flat prior for the signal cross section and is numerically integrated to obtain the limit on $\sigma \cdot \mathcal{B}(X \rightarrow ee)$. The observed limits are shown in Fig. 3 for the spin-1 case. The Z' model lines are obtained using the leading-order PYTHIA event generator using the couplings in [18], with a factor of 1.3 applied multiplicatively to account for NLO corrections [19].

For the specific case of the RS graviton, which has a branching ratio to $\gamma\gamma$ twice that of ee , the analysis sensitivity can be improved by combining with the $\gamma\gamma$ channel described in [1]. The $\gamma\gamma$ events are required to have two photons with $E_T \geq 15 \text{ GeV}$, with one in the CEM ($|\eta| \leq 1.04$) and the other either in the CEM or PEM ($1.2 \leq |\eta| \leq 2.8$). The CC and CP channels use integrated luminosities of 1.2 fb^{-1} and 1.1 fb^{-1} respectively, with the difference arising from the requirement that the silicon system be operational for track veto purposes for photons in the PEM. Selected photons have similar isolation and shower shape requirements to electrons; however photons are required not to have an associated track,

which ensures there is no overlap between the $\gamma\gamma$ and ee sample.

The $\gamma\gamma$ channel is combined with the ee channel by multiplying their individual likelihoods together. The uncertainties on the background estimates are considered to be uncorrelated. The uncertainties on the acceptance and luminosity are taken to be 100% correlated. The combined limits, together with the k/\bar{M}_{pl} vs graviton mass exclusion region, are shown in Fig. 4. The RS graviton model lines are obtained using the HERWIG event generator [20] with a factor of 1.3 applied multiplicatively to account for NLO corrections.




FIG. 3: The observed and expected limits on the $\sigma \cdot \mathcal{B}(X \rightarrow ee)$ of a spin-1 particle.

In summary a search has been made for new physics in the ee channel, and no significant excess over the standard model prediction is observed. Limits are placed on new spin-1 and spin-2 bosons. The SM-like Z' is found to be excluded for masses below $923 \text{ GeV}/c^2$ and the E_6 Z' bosons; the Z'_I , the Z'_ψ , the Z'_χ , and the Z'_η bosons are excluded with masses below 729 , 822 , 822 , and $891 \text{ GeV}/c^2$ respectively. The direct limits presented here on all the E_6 Z' bosons surpass the corresponding indirect limits from LEP [21]. The RS graviton with $k/\bar{M}_{pl} = 0.1$ is excluded for masses below $807 \text{ GeV}/c^2$. When combined with the $\gamma\gamma$ channel, masses less than $889 \text{ GeV}/c^2$ are excluded for $k/\bar{M}_{pl} = 0.1$. The above limits on Z' 's and the RS graviton represent the best single-experiment direct limits to date.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the




FIG. 4: The limits on $\sigma \cdot \mathcal{B}(G \rightarrow ee)$ (top) and k/\bar{M}_{pl} (bottom) for a RS graviton in the ee , $\gamma\gamma$ channels separately and combined.

Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucléaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Comisión Interministerial de Ciencia y Tecnología, Spain; the European Community's Human Potential Programme; the

Slovak R&D Agency; and the Academy of Finland.

-
- [1] T. Aaltonen *et al.* (CDF Collaboration), to be submitted to *Phys. Rev. Lett.*
 - [2] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 3370 (1999).
 - [3] A narrow resonance is defined as a resonance where the resolution of the CDF detector solely determines the observed width.
 - [4] All limits presented in this Letter are at the 95% confidence level.
 - [5] F. del Aguila, M. Quiros, and F. Zwirner, *Nucl. Phys.* **B287**, 419 (1987); J. L. Hewett and T. G. Rizzo, *Phys. Report.* **183**, 193 (1989).
 - [6] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, *Phys. Rev. Lett.* **84**, 2080 (2000);
B. C. Allanach *et al.* *J. High Energy Phys.* **0212**, 039 (2002).
 - [7] V. M. Abazov *et al.* (DØ Collaboration), *Phys. Rev. Lett.* **95**, 091801 (2005).
 - [8] A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **96**, 211801 (2006).
 - [9] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005).
 - [10] CDF uses a cylindrical coordinate system with the z -axis along the proton direction, where $z=0$ is the center of the detector. Pseudorapidity defined as $\eta = -\ln(\tan(\theta/2))$, where θ is the polar angle while $E_T = E \sin(\theta)$, where E is the energy deposited in the cluster of calorimeter towers.
 - [11] A. Sill *et al.*, *Nucl. Instrum. Methods A* **447**, 1 (2000).
 - [12] T. Sjostrand *et al.*, *Comput. Phys. Commun.* **135**, 238 (2001). We use PYTHIA version 6.216.
 - [13] H. L. Lai *et al.*, *Eur. Phys. J.* **C12** 375 (2000).
 - [14] S. Agostinelli *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **506**, 250 (2003).
 - [15] J. Pumplin *et al.*, *J. High Energy Phys.* **0207**, 012 (2002).
 - [16] Electrons are isolated if in a surrounding cone of 0.4 radius the E_T deposited minus $1.02 \times$ the electron E_T is less than 3 GeV (1.6 GeV) in the CEM (PEM).
 - [17] S. Klimenko, J. Konigsberg, and T.M. Liss, FERMILAB-FN-0741 (2003).
 - [18] C. Ciobanu *et. al.*, FERMILAB-FN-0773-E (2005).
 - [19] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **94**, 091803 (2005).
 - [20] G. Corcella *et al.*, *J. High Energy Phys.* **0101**, 10 (2001).
 - [21] W.-M. Yao *et al.*, *J. Phys. G* **33**, 1 (2006).