

Lepton flavor changing Higgs Boson decays in a Two Higgs Doublet Model with a fourth generation of fermions

S. Chamorro-Solano*, A. Moyotl and M. A. Pérez

*Departamento de Ciencias Naturales y Exactas, Universidad de la Costa, Calle 58 num. 55-66, Barranquilla, Colombia.

Departamento de Física, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apdo. Postal 14-740, 07000 México D.F., México.

E-mail: schamorr1@cuc.edu.co
amoyotl@fis.cinvestav.mx
mperez@fis.cinvestav.mx

Abstract. We analyze the flavor changing decay $h \rightarrow \mu\tau$ in the framework of a Two Higgs Doublet Model with a fourth generation of fermions (4G2HDM) which couples only to the heavy scalar doublet. We obtain that the respective branching ratio at one-loop level can reach values as high as $10^{-4} - 10^{-6}$ for masses of 300 GeV to 1 TeV for the heavy leptons in the fourth family and the new heavy Higgs bosons. These radiative corrections are of the same order of magnitude as the tree level prediction of the 4G2HDM.

1. Introduction

In the Standard Model (SM) there are no Flavor Changing Neutral Currents (FCNC) transitions at tree level. This type of transitions may be induced at one-loop level due to the virtual exchange of scalar or gauge bosons but they are highly suppressed due to the Glashow-Iliopolous-Maiami mechanism [1]. However, they can be relaxed by extended flavor structures in extensions of the SM [2, 3]. For this reason, the excess reported recently by the CMS [4] and ATLAS [5] Collaborations for the branching ratio $\text{Br}(h \rightarrow \tau\mu)$ has been widely addressed in the literature [6–10] and, in particular in some versions of the 2HDM [11–13] the respective branching ratio could be at the level of few percent. The CMS and ATLAS Collaborations have reported upper limits on the branching ratios of the $h \rightarrow \tau\mu$ decay mode of 1.51×10^{-2} and 1.85×10^{-2} at 95% C.L., respectively [4, 5]. CMS has also reported a slight excess, with a significance of 2.4 standard deviations at $m_h = 125$ GeV, with a best fit $\text{BR}(h \rightarrow \tau\mu) = 18.4 + 3.9 / - 3.7 \times 10^{-3}$ [4]. Even though a new CMS report does not throw a clear conclusion on the evidence of this decay mode [14], any evidence of Higgs lepton flavor violating decays (LFVD) would point towards a

neat signal of physics beyond the SM since this type of transitions are highly suppressed in the SM [2, 3].

Lepton flavor violating Higgs decays (LFVHD) were considered first in the context of the SM with three heavy Majorana neutrinos [6, 7] and later in models beyond the SM [7]. In particular, it was found that within the effective Lagrangian approach, the LFV decays of the Higgs boson involving the tau lepton could have branching ratios at the percentage level [8]. The study of LFVHD has received an increased interest recently in different models [9, 10, 15–17]. In some cases, the enhancement due to loops involving new degrees of freedom could be sufficiently large as to explain the CMS and ATLAS signals [4, 5].

LFVHD have been analyzed also in Two Higgs Doublet Models (2HDM) where all flavor-changing neutral processes are determined by the weak mixing matrix [18, 19]. Although some electroweak precision observables allow the existence of a fourth family of quarks [20], the most severe bounds come from the invisible width of the Z^0 at the LEP [21] and the Higgs boson production at the LHC [22, 23]. The main mechanism of Higgs boson production at LHC is the gluon fusion induced by top and bottom quarks loops. If extra heavy quarks exist, then the respective cross section may be enhanced by a factor of 9 compared to the SM with three families [24]. However, the actual Higgs boson production is in agreement with the top and bottom contributions [22, 23]. This experimental circumstance could exclude a fourth family within of the SM, but in the 4G2HDM [25] the fourth family interacts only with the extra heavy scalar bosons, and thus the gluon fusion mechanism remains unchanged by the presence of the extra heavy quarks. Previously we presented only the one-loop contribution of the heavy neutral scalar bosons in 4G2HDM [17]. In the present paper we include the contributions of all heavy scalar bosons and heavy leptons. In this context, we consider the possibility that a fourth family of leptons could induce a large contribution to the branching ratio $\text{Br}(h \rightarrow \tau\mu)$ through one-loop effects. We will take a 2HDM with a fourth family of quarks and leptons which has been found not to be in conflict with electroweak precision data [25–27]. Even more, it has been pointed out that Higgs data at the LHC does not rule out a sequential fourth family [28–31]. We will find that the heavy lepton of the fourth family and the heavy neutral Higgs boson induce contributions of the same order of magnitude as the tree level prediction of the 4G2HDM. We would like to stress that our calculation is not just a direct extension of some old calculations in the SM since in our case is not possible to neglect the virtual contributions associated to new heavy fermions and scalar bosons with masses heavier than the Higgs boson mass.

The plan of the paper is as follows. In section 2 we introduce the way to include a fourth family in the 2HDM. In section 3 we present details of the one-loop calculations for the $h^0 \rightarrow \mu\tau$ decay, as well as the tree level prediction of this model. In section 4 we include our analysis and results; finally, we present the conclusions in section 5.

2. The 2HDM with a fourth generation

The 2HDM have been introduced in order to explain the large top-quark mass through the inequality between the two vacuum expectation values (VEVs) $v_1 \gg v_2$ [14]. In these models, the Higgs sector is composed of two isospin doublet scalar fields Φ_1 and Φ_2 with VEVs $v_{1,2}$, respectively. We shall use the definitions $v \equiv \sqrt{v_1^2 + v_2^2}$ and $\tan\beta \equiv v_2/v_1$. The fourth family can be incorporated in three different 2HDM scenarios [25]. In scenario I, Φ_1 gives masses only to fermions in the fourth family while Φ_2 generates masses to the rest of the fermions; in this case, $\tan\beta \sim m_{t_4}/m_t \simeq \mathcal{O}(1)$. In scenario II, Φ_1 is responsible for the mass generation of the heavy fermion states of both the third and fourth generations, whereas Φ_2 induces masses to

the light fermions of the first and second generations. In scenario III, we have $m_{t_4} \propto v_1$ and only the fermions with masses at the electroweak scale are coupled to Φ_1 . In the latter two cases $\tan\beta \gg 1$ is a natural choice. In the present paper we shall consider only scenario I whose phenomenology has been studied recently by Bar-Shalom and Soni [31]; the Z_2 charge assignments in this scenario are show in the Table 1. Physically we have chosen this scenario because the heavy fermions and quarks (of the SM) have the same order of mass, then the production of Higgs boson by the fusion of gluons is not altered. Scenario I of the 2HDM will be appropriate to the present analysis since the heavy fermions of the fourth family will couple only to the Φ_1 doublet and we should expect that $\tan\beta \sim \mathcal{O}(1)$.

Φ_1	Φ_2	u_R	d_R	ℓ_R	Q_L	f_L
+	-	-	-	-	+	+

Table 1. Z_2 charge assignments in the model we are considering.

We consider an extension of the 2HDM with the heavy scalar fermions of the fourth family coupled to scalar and pseudoscalar fields that are remanant from the softly broken ¹ Z_2 symmetry: the 4G2HDM. In this model, there is a FCNC at tree level between scalar bosons and fermions. Of course, FCNC may arise also through charged and neutral currents. These FCNC Yukawa interactions for the leptons of the fourth family are given by

$$\mathcal{L}_{\text{FV}} = \frac{g}{4m_W} f_\beta^\phi \bar{\ell}_i \left[\left(g_s^\phi \right)_{ij} + \left(g_p^\phi \right)_{ij} \gamma_5 \right] \ell_j \phi, \quad (1)$$

where the indices i, j run through the four generations of families of leptons ℓ_i , $\phi = h^0, H^0, A^0$, while for $\phi = H^+$ we have $\ell_i \rightarrow \nu_i$ and the coupling constant f_β^ϕ acquires values according to Table 2. On other hand, the scalar (g_s^ϕ) and pseudoscalar (g_p^ϕ) couplings are depicted in Table 3, where the subscripts i, j run through the e, μ, τ and a lepton (neutrino) of the fourth family $\ell_4(\nu_4)$ only for the heavy scalar bosons.

ϕ	f_β^ϕ
h^0	$c_\alpha/s_\beta + s_\alpha/c_\beta$
H^0	$c_\alpha/c_\beta - s_\alpha/s_\beta$
A^0	$2iI_\ell(t_\beta + 1/t_\beta)$
H^+	$2/\sqrt{2}$

Table 2. The f_β^ϕ values for the scalar boson ϕ , with I_ℓ the weak isospin and we will use the short notation $s_\theta \equiv \sin \theta$, $c_\theta \equiv \cos \theta$ and $t_\beta \equiv \tan \beta$.

On the other hand, we shall assume CP invariance in the Higgs sector and thus the triple scalar couplings $\lambda_{\phi\phi h^0}$ are obtained from the usual Higgs potential of any 2HDM; these couplings are given by [32]

$$\lambda_{H^+H^-h} = \frac{1}{v} \left[(2M^2 - 2m_{H^\pm}^2 - m_h^2) s_{\beta-\alpha} + 2(M^2 - m_h^2) \cot 2\beta c_{\beta-\alpha} \right], \quad (2)$$

¹ Z_2 symmetry is softly broken to achieve a CP conserving potential as this is phenomenologically more interesting.

ϕ	$(g_s^\phi)_{ij}$	$(g_p^\phi)_{ij}$
h^0	$m_{\ell_i} \Sigma_{ij}^\ell + m_{\ell_j} \Sigma_{ji}^{\ell*}$	$m_{\ell_i} \Sigma_{ij}^\ell - m_{\ell_j} \Sigma_{ji}^{\ell*}$
H^0	$m_{\ell_i} \Sigma_{ij}^\ell + m_{\ell_j} \Sigma_{ji}^{\ell*}$	$m_{\ell_i} \Sigma_{ij}^\ell - m_{\ell_j} \Sigma_{ji}^{\ell*}$
A^0	$m_{\ell_i} \Sigma_{ij}^\ell - m_{\ell_j} \Sigma_{ji}^{\ell*}$	$m_{\ell_i} \Sigma_{ij}^\ell + m_{\ell_j} \Sigma_{ji}^{\ell*}$
H^+	$t_\beta U_{ki}^* (m_{\ell_k} - m_{\nu_i}) - (t_\beta + 1/t_\beta) (m_{\nu_i} \Sigma_{ii}^\nu U_{ki}^* - m_{\ell_i} U_{ii}^* \Sigma_{ik}^{\ell*})$	$t_\beta U_{ki}^* (m_{\ell_k} + m_{\nu_i}) + (t_\beta + 1/t_\beta) (m_{\nu_i} \Sigma_{ii}^\nu U_{ki}^* + m_{\ell_i} U_{ii}^* \Sigma_{ik}^{\ell*})$

Table 3. Scalar (g_s^ϕ) and pseudoscalar (g_p^ϕ) couplings, the subscripts i and j run through the e, μ, τ or ℓ_4 (ν_4) only for heavy scalar bosons. Further, $\Sigma_{ij}^{\ell, \nu}$ is a complex element of a mixing 4×4 matrix in the 4G2HDM, while U_{ij} is a complex element of the 4×4 PMNS matrix.

$$\lambda_{AAh} = \frac{1}{2v} \left[(2M^2 - 2m_A^2 - m_h^2) s_{\beta-\alpha} + 2(M^2 - m_h^2) \cot 2\beta c_{\beta-\alpha} \right], \quad (3)$$

$$\lambda_{HHh} = \frac{s_{\beta-\alpha}}{2v} \left[(2M^2 - 2m_H^2 - m_h^2) s_{\beta-\alpha}^2 + 2(3M^2 - 2m_H^2 - m_h^2) \cot 2\beta s_{\beta-\alpha} c_{\beta-\alpha} - (4M^2 - 2m_H^2 - m_h^2) c_{\beta-\alpha}^2 \right], \quad (4)$$

with α and β the usual mixing angles, M describes the soft breaking scale of the Z_2 symmetry and it is fixed by the relation $\sqrt{\lambda v^2} = \sqrt{m_\phi^2 - M^2}$. Finally, the scaling factors that describe the deviations in the Higgs boson couplings from the SM prediction are given in Ref. [32]. At tree level, these scaling factors become unit if $\sin(\beta - \alpha) = 1$ and further, since LHC data suggest that the observed Higgs boson coincide with the SM-like, then $\sin(\beta - \alpha) \simeq 1$ can be allowed. Thus, for convenience we introduce the tiny parameter

$$\chi = \frac{\pi}{2} - (\beta - \alpha), \quad (5)$$

with $\chi \rightarrow 0$ the SM-like limit.

3. The $h^0 \rightarrow \mu\tau$ decay in the 4G2HDM

The 4G2HDM has some free parameters such as the masses of the scalar bosons and the fourth family leptons, as well as $\tan \beta$ and the mixing matrix elements $\Sigma_{4,4}^{\ell, \nu}$, $\Sigma_{4k}^{\ell, \nu}$, $U_{4,4}$ and U_{4k} . In this model the FCNC may occur at tree level but we will be interested in computing the one-loop contributions coming from the virtual exchange of heavy scalars and fourth family fermions, as it is shown in Figure 1. Our results will show that the tree and one-loop contributions are of the same order of magnitude. In the case of the virtual exchange of heavy neutral scalar bosons, there is also a heavy lepton τ_4 exchange, while for the heavy charged scalar boson exchange, there is a heavy neutrino ν_4 . There are thus four diagrams that contribute to one-loop order in the $h^0 \rightarrow \mu\tau$ decay. The respective transition amplitude is given by

$$\mathcal{M} = i\bar{u}(p_2, m_j)(A + iB\gamma_5)u(p_1, m_i), \quad (6)$$

where m_k and p_k correspond to the mass and 4-momenta of the final fermion ℓ_k , respectively, A and B are form factors. At tree level, for the Higgs boson (h^0) contribution, we have from the Lagrangian in the Eq. (1):




Figure 1. One-loop diagrams for $h^0 \rightarrow \mu\tau$ in the 4G2HDM. Here ϕ is any neutral scalar A^0 , H^0 or charged scalar H^\pm ; while f_4 can be a charged lepton τ_4 for the heavy neutral scalar bosons, or a ν_4 for the scalars charged scalar boson.

$$A_{h^0} = \frac{gf_\beta^{h^0}}{4m_W} (g_s^{h^0})_{ij}, \quad (7)$$

and the B_{h^0} form factor is obtained by the exchange $(g_s^{h^0})_{ij} \rightarrow -i(g_p^{h^0})_{ij}$ in A_{h^0} . If we neglect the muon mass in $(g_s^{h^0})_{\mu\tau}$ of the Table 3, the tree level contribution to the $BR(h^0 \rightarrow \tau\mu)$ depends only of the χ parameter and the $\Sigma_{2,3}$ matrix element, if we assume that $|\Sigma_{2,3}^\ell|^2 \simeq |\Sigma_{4,3}^\ell \Sigma_{4,2}^\ell|$. Under these conditions, the branching ratio to tree level decreases as we diminish the χ parameter, where we have values between 1.2×10^{-6} and 1.2×10^{-8} , for $\chi = 0.1$ and $\chi = 0.01$ respectively.

The one-loop level contributions are constructed from the explicit interactions given in Eq. (1) and the A_ϕ form factor, where the exact form is given by

$$A_\phi = \frac{g^2 (f_\beta^\phi)^2 \lambda_{\phi\phi h^0}}{256\pi^2 m_h m_W^2} \int_{x=0}^1 \int_{y=0}^{1-x} \Xi(x, y) dx dy, \quad (8)$$

$\Xi(x, y)$ is a dimensionless function that corresponds to the Feynman parametrization

$$\Xi(x, y) = \frac{(g_p^\phi)_{i4} (g_p^\phi)_{j4} (-r_{f_4} + r_i x + r_j y) + (g_s^\phi)_{i4} (g_s^\phi)_{j4} (r_{f_4} + r_i x + r_j y)}{r_\phi^2 (x+y) + (r_i^2 x + r_j^2 y - r_{f_4}^2)(x+y-1) - xy}, \quad (9)$$

and we have defined $r_k = m_k/m_h$ ($k = i, j, f_4, \phi$). In order to get the B_ϕ form factor, we use the following products of scalar and pseudoscalar coupling constants:

$$\begin{aligned} (g_p^\phi)_{i4} (g_p^\phi)_{j4} &\rightarrow (g_p^\phi)_{j4} (g_s^\phi)_{i4}, \\ (g_s^\phi)_{i4} (g_s^\phi)_{j4} &\rightarrow (g_p^\phi)_{i4} (g_s^\phi)_{j4}. \end{aligned}$$

It is important to emphasize that our results are free from UV divergences in a natural way, i.e., all UV divergences cancel among themselves. In the approximation that the masses of the outgoing fermions are neglected in (9), we obtain the simple expression:

$$\Xi(x, y) = \frac{r_{f_4}[(g_s^\phi)_{i4}(g_s^\phi)_{j4} - (g_p^\phi)_{i4}(g_p^\phi)_{j4}]}{r_\phi^2(x+y) - r_{f_4}^2(x+y-1) - xy}. \quad (10)$$

In this approximation the respective decay width can be written as:

$$\Gamma(h^0 \rightarrow \tau\mu) = \frac{m_h}{8\pi}(|A|^2 + |B|^2). \quad (11)$$

4. Results and analysis

In the 4G2HDM the new fermions are expected to be very heavy and the PDG has included, at 95 % CL, the following limits: $m_{\tau_4} > 102.6$ GeV and $m_{\nu_4} > 90.3$ GeV [21]; these limits were obtained through the search of the τ_4 decays into a $W\nu$ pair and an analysis of the three neutrino mixing scheme. However, in the context of other two Higgs doublet models with a Higgs boson mass of 124.5 GeV, the most favored value for m_{τ_4} is 110.8 GeV [33]. In this framework, the respective fourth leptonic mass splitting is given by,

$$\Delta_\ell = m_{\nu_4} - m_{\tau_4} \lesssim -m_W, \quad (12)$$

which is the value used by the LHC Higgs Cross Section Working Group for the study of the Higgs boson decays to fourth family fermions [22, 34]. This mass splitting is also valid for all values of the Higgs boson mass from constraints of unitarity and oblique parameters [35].

The search of heavy scalar bosons, in some 2HDMs and model independent frameworks, has been performed by the ATLAS [36–38] and CMS [39–43] collaborations. In the case of the heavy neutral boson H^0 , the search was done in channels like ZZ [38], h^0h^0 [39, 43] and $\gamma\gamma$ [41], while for the heavy pseudoscalar boson A^0 , the respective channels consist of Zh [37, 39, 40, 43] and $\gamma\gamma$ [41]. Additionally, the heavy charged scalar boson H^\pm was searched in the decays $t \rightarrow H^\pm b$ [36, 42, 44] and $H^\pm \rightarrow \tau^\pm \nu_\tau$ [42]. These searches covered the mass range from 150 GeV to 1000 GeV. On other hand, $\tan\beta$ is assumed to be in the range around $\tan\beta \sim 1$, where the 4G2HDM is consistent with both electroweak precision data [25] and the observed 125 GeV Higgs boson [45].

For the present analysis we consider the mass range for the fourth family lepton $100 \text{ GeV} < m_{f_4} < 350 \text{ GeV}$, and for the mass of the heavy neutral scalar $200 \text{ GeV} < m_\phi < 700 \text{ GeV}$. We shall take values of $\tan\beta$ close to unity. The Higgs total decay width is taken by the SM prediction of $\Gamma_{h^0} \simeq 4.07 \text{ MeV}$. For simplicity, we assume that the mixing matrices are symmetrical $\Sigma_{3,4}^{\nu,\ell} \sim U_{3,4}$. Finally, the branching ratio for the $\tau \rightarrow \gamma\mu$ decay in the 4G2HDM is not too far below the current bound [46] if we take $|U_{4,4}| = |\Sigma_{4,4}|^{\nu,\ell} = 1$ and $U_{2,4}\Sigma_{4,3}, U_{3,4}\Sigma_{4,2} \sim \mathcal{O}(10^{-3})$. Similar constraints have been derived for the values of the mixing matrices of the known three generations from the analysis of $g-2$ of the muon and the decay $\mu \rightarrow e\gamma$ [46, 47].

At one loop level and for suppressed values of χ , it is possible to demonstrate from the Table 2 and the Eq. (5), that the A_{H^0, A^0} and B_{H^0, A^0} form factors are basically independent of the χ parameter. Additionally, we found that the respective form factors of H^0 and A^0 have different sign, they have basically the same magnitude and therefore the contribution of these heavy neutral scalar bosons is very small [17]. On other hand, the A_{H^\pm} and B_{H^\pm} form factors do not depend of the χ parameter and further, the respective magnitudes are different to the previous case. Thus, the main contributions of the $h^0 \rightarrow \tau\mu$ decay come from the heavy charged scalar boson. We present in Figure 2 the respective values for the branching ratio as a function of the mass of the heavy scalar bosons, where we have considered the degenerate case for the scalars bosons and we used $m_{\tau_4} = 150 \text{ GeV}$, $m_{\nu_4} = m_{\tau_4} - m_W$ for the splitting and different values of $\sqrt{\lambda v^2}$. We observe that the branching ratio decreases for high mass values of the heavy

scalar bosons. But as expected from the triple scalar couplings $\lambda_{\phi\phi h^0}$, we observe also that the branching ratio increases as the value of $\sqrt{\lambda v^2}$ increases. Thus, there are approximately two orders of magnitude between the contributions for $\sqrt{\lambda v^2} = 150$ GeV (black dashed line) and $\sqrt{\lambda v^2} = 450$ GeV (blue line).




Figure 2. One loop contribution to the branching ratio for the $h^0 \rightarrow \tau\mu$ decay as a function of m_ϕ . We consider that all the scalar bosons are degenerate, $m_{\tau_4} = 150$ GeV, $\tan\beta = 1$ and different values of $\sqrt{\lambda v^2}$. The tree level prediction is above these results and of order 1.2×10^{-6} for $\chi = 0.1$.

In Figures 3, 4, 5 and 6 we depict the dependence of $\text{BR}(h^0 \rightarrow \mu\tau)$ as a function of scale parameter $\sqrt{\lambda v^2}$ and the mass of the fourth family neutrino m_{ν_4} . In all cases we fixed the masses of the heavy neutral scalar bosons and the horizontal line corresponds to the tree-level contribution for $\chi = 0.1$, which amounts to about 1.2×10^{-6} .




Figure 3. One loop contribution to the branching ratio for the $h^0 \rightarrow \tau\mu$ decay as a function of $\sqrt{\lambda v^2}$. We consider $m_\phi = 300$ GeV, $\tan\beta = 1$ and different values of m_{τ_4} . The horizontal line corresponds to the contribution at tree level with $\chi = 0.1$

In Figures 3 - 5 we depict the dependence of the branching ratio as a function of $\sqrt{\lambda v^2}$ and the mass of the fourth family neutrino m_{ν_4} , respectively. In both cases we fixed the masses




Figure 4. One-loop contribution to $\text{BR}(h^0 \rightarrow \mu\tau)$ as a function of the scale parameter $\sqrt{\lambda v^2}$. We considered $m_\phi = 600$ GeV, $\tan\beta = 2$, and two values for the mass of the heavy lepton $m_{\tau_4} = 500$ GeV (black line) and 1 TeV (blue line).




Figure 5. One loop contribution to the branching ratio for the $h^0 \rightarrow \tau\mu$ decay as a function of the fourth family neutrino mass m_{ν_4} . We consider $m_{H^+} = 300$ GeV, $\tan\beta = 1$ and different values of $\sqrt{\lambda v^2}$. The horizontal line corresponds to the contribution at tree level with $\chi = 0.1$.

of the heavy scalar bosons to 300 GeV and the horizontal line corresponds to the tree level branching ratio with $\chi = 0.1$, which amounts approximately to 1.2×10^{-6} . In the Figure 3 we have used three different values of the mass of the fourth family lepton m_{τ_4} . In this case we observe that the branching ratio increases as the value of $\sqrt{\lambda v^2}$ increases, where from 100 GeV to 500 GeV there is a difference approximately of more than two orders of magnitude. But more important, for very heavy masses of τ_4 , the branching ratio at one-loop level can be greater than the respective prediction at tree level. We observe explicitly this behaviour for $m_{\tau_4} = 350$ GeV (blue line) and $\sqrt{\lambda v^2} \simeq 210$ GeV, as well as for $m_{\tau_4} = 250$ GeV (blue dashed line) and $\sqrt{\lambda v^2} \simeq 400$ GeV, while for $m_{\tau_4} = 150$ GeV (black dashed line) the one-loop contribution to the branching ratio remains below the tree level value. In the Figure 5 we have used the same values of $\sqrt{\lambda v^2}$ than in Figure 2, and we observed also the same situation as in the Figure 3. In these figures the branching ratio at one-loop level exceeds the prediction at tree level for $\sqrt{\lambda v^2} = 450$ GeV (blue line) and $m_{\tau_4} \simeq 230$ GeV, as well as for $\sqrt{\lambda v^2} = 300$ GeV (blue dashed line) and $m_{\tau_4} \simeq 290$ GeV. Particular for $\sqrt{\lambda v^2} = 450$ GeV, $m_{\tau_4} = 350$ GeV and masses of the heavy




Figure 6. One-loop contribution to $BR(h^0 \rightarrow \mu\tau)$ as a function of the mass of the heavy neutrino m_{ν_4} . We have taken $m_{H^+} = 1$ TeV, $\tan\beta = 1$, and three values of $\sqrt{\lambda v^2} = 150$ GeV (dotted black line), 300 GeV (dotted blue line) and 450 GeV (blue line). The tree-level contribution is given by the horizontal line.

scalar bosons between 200 GeV and 300 GeV, we can have up to $BR(h^0 \rightarrow \mu\tau) \simeq 7.2 \times 10^{-5}$ and $BR(h^0 \rightarrow \mu\tau) \simeq 2.4 \times 10^{-5}$ respectively. This significantly improves the respective tree level contribution with $\chi = 0.1$, and it would be better if $\chi = 0.01$ or for more suppressed values of χ . Thus, we can appreciate that the one-loop result for this branching ratio can be as high as the tree level contribution in a wide range of values of the mass of the fourth family lepton and the $\sqrt{\lambda v^2}$ parameter.

In Figures 4 and 6 we have taken larger values of the different heavy masses involved and $\tan\beta = 2$ (see Figure 4). We can appreciate that our results are sensitive to these variations, in particular to a heavy mass of the charged Higgs boson. The increase in the branching ratio could be of order one or two orders of magnitude.

Finally, from the Eq. (8) we can obtain also the branching ratio for the $h^0 \rightarrow \mu e$ and $h^0 \rightarrow \tau e$ decays. But in this case the respective mixing matrix elements are smaller than for the $\tau\mu$ element; therefore, the respective branching ratios are more suppressed. At one-loop level, if we use the current bound $U_{1,4}\Sigma_{4,2}, U_{2,4}\Sigma_{4,1} \sim \mathcal{O}(10^{-6})$ [46] and the same methodology for $h^0 \rightarrow \tau\mu$, we obtain $BR(h^0 \rightarrow \mu e) \sim 10^{-11}$ for $m_\phi = 200$ GeV and large values of m_{τ_4} and $\sqrt{\lambda v^2}$.

5. Concluding remarks

We have calculated the one-loop corrections to the flavor changing Higgs boson decay $h^0 \rightarrow \mu\tau$ in the framework of the 4G2HDM, and we showed that the loop contributions exceed the tree level prediction of the branching ratio in certain regions of the parameter space. Since the tree level contribution depends directly on the χ parameter, the respective predictions can be very suppressed. On other hand, the contributions of heavy neutral scalar bosons are basically independent of small values of the χ parameter, while the contribution of the heavy charged scalar boson does not depend on the χ parameter. However, our results show that the dominant contributions at one-loop level arise from the virtual exchange of the heavy charged scalar boson H^\pm , in a wide range of the parameters associated to this model. We showed that for very heavy fourth family leptons and high values of the $\sqrt{\lambda v^2}$ parameter, the branching ratio at one-loop

level exceeds the prediction at tree level. In particular for values of $\sqrt{\lambda v^2} = 450$ GeV, $m_{\tau_4} = 350$ GeV and $m_\phi = 200$ (300) GeV, we have obtained $\text{BR}(h^0 \rightarrow \mu\tau) \simeq 7.2(2.4) \times 10^{-5}$. While at tree level we have $\text{BR}(h^0 \rightarrow \mu\tau) \simeq 1.2 \times 10^{-6}$ for $\chi = 0.1$ and $\text{BR}(h^0 \rightarrow \mu\tau) \simeq 1.2 \times 10^{-8}$ for $\chi = 0.01$.

We have shown also that these results are sensitivity to larger masses of the charged Higgs boson and the heavy neutrino. The respective branching ratio could increase by one or two orders of magnitude. However, even in this case it will be a challenge to test these results in the forthcoming run of the LHC.

It is also important to notice that our results differ with respect to several models that predict LFV Higgs decays. Some of them obtain branching ratios for the $h^0 \rightarrow \tau\mu$ decay mode of order few percent in the framework of flavor symmetry models: with a continuous Abelian or a discrete non-Abelian symmetries [48], as well as with a discrete S_4 symmetry [49]. A large prediction for the $\text{BR}(h^0 \rightarrow \tau\mu)$ has been also advanced as a strong probe of neutrino mass models [50] or for the existence of a singlet dark matter candidate [51]. Finally, there are some versions of 2HDM that do not allow the $h\tau\mu$ vertex at tree level [52]. In the latter case, the exact alignment limit of 2HDM induces a one-loop prediction of $\text{BR}(h^0 \rightarrow \tau\mu)$ of the order of few percent.

Acknowledgements

We acknowledge support from CONACYT (Mexico) and useful suggestions made by Pablo Roig.

References

- [1] S. L. Glashow, J. Iliopoulos, and L. Maiani, “Weak interactions with lepton-hadron symmetry,” *Phys. Rev. D*, vol. **2**, pp. 1285–1292, (1970).
- [2] M. A. Perez, G. Tavares-Velasco, and J. J. Toscano, “New physics effects in rare Z decays,” *Int. J. Mod. Phys*, vol. **A19**, pp. 159–178, (2004). [arXiv:hep-ph/0305227].
- [3] F. Larios, R. Martinez, and M. A. Perez, “New physics effects in the flavor-changing neutral couplings of the top quark,” *Int. J. Mod. Phys*, vol. **A21**, pp. 3473–3494, (2006). [arXiv:hep-ph/0605003].
- [4] V. Khachatryan *et al.*, “Search for Lepton-Flavour-Violating Decays of the Higgs Boson,” *Phys. Lett*, vol. **B749**, pp. 337–362, (2015). [arXiv:hep-ex/1502.07400].
- [5] G. Aad *et al.*, “Search for lepton-flavour-violating $H \rightarrow \mu\tau$ decays of the Higgs boson with the ATLAS detector,” *JHEP*, vol. **11**, p. 211, (2015). [arXiv:hep-ex/1508.03372].
- [6] A. Pilaftsis, “Lepton flavour nonconservation in h^0 decays,” *Physics Letters B*, vol. **285**, no. 1, pp. 68–74, (1992).
- [7] L. Diaz-Cruz and J. J. Toscano, “Lepton flavor violating decays of Higgs bosons beyond the standard model,” *Phys. Rev*, vol. **D62**, p. 116005, (2000). [arXiv:hep-ph/9910233].
- [8] C. Alvarado *et al.*, “Minimal models of loop-induced lepton flavor violation in Higgs boson decays,” *Phys. Rev*, vol. **D94**, no. 7, p. 075010, (2016). [arXiv:hep-ph/1602.08506].
- [9] J. Lee and K. Lee, “ $B_s \rightarrow \mu\tau$ and $h \rightarrow \mu\tau$ decays in the general two Higgs doublet model,” 2016. [arXiv:hep-ph/1612.04057].
- [10] A. Lami and P. Roig, “ $H \rightarrow \ell\ell'$ in the simplest little Higgs model,” *Phys. Rev*, vol. **D94**, no. 5, p. 056001, (2016). [arXiv:hep-ph/1603.09663].

- [11] J. Herrero-Garcia *et al.*, “Full parameter scan of the Zee model: exploring Higgs lepton flavor violation,” *Journal of High Energy Physics*, vol. **2017**, p. 130, Apr 2017. [arXiv:hep-ph/1701.05345v2].
- [12] D. Aristizabal Sierra and A. Vicente, “Explaining the CMS Higgs flavor-violating decay excess,” *Phys. Rev. D*, vol. **90**, p. 115004, Dec 2014. [arXiv:hep-ph/1409.7690v2].
- [13] D. Das and A. Kundu, “Two hidden scalars around 125 GeV and $h \rightarrow \mu\tau$,” *Phys. Rev. D*, vol. **92**, p. 015009, Jul 2015. [arXiv:hep-ph/1504.01125v2].
- [14] “Observation of the SM scalar boson decaying to a pair of τ leptons with the CMS experiment at the LHC,” Tech. Rep. CMS-PAS-HIG-16-043, CERN, Geneva, (2017).
- [15] F. del Aguila *et al.*, “Lepton Flavor Changing Higgs decays in the Littlest Higgs Model with T-parity,” (2017). [arXiv:hep-ph/1705.08827].
- [16] S. Chamorro-Solano, A. Moyotl, and M. A. Perez, “The decay $h \rightarrow \mu\tau$ in the Littlest Higgs Model with T-parity,” *J. Phys. Conf. Ser.*, vol. **761**, no. 1, p. 012051, (2016).
- [17] A. Moyotl, S. Chamorro-Solano, and M. Perez, “The $h \rightarrow \mu\tau$ decay in a two higgs doublet model with a fourth generation of fermions,” *Nucl. Particle Phys. Proce.*, vol. **287-288**, pp. 205–207, (2017). The 14th International Workshop on Tau Lepton Physics.
- [18] F. J. Botella, G. C. Branco, M. Nebot, and M. N. Rebelo, “Flavour Changing Higgs Couplings in a Class of Two Higgs Doublet Models,” *Eur. Phys. J.*, vol. **C76**, no. 3, p. 161, (2016). [arXiv:hep-ph/1508.05101].
- [19] M. Sher and K. Thrasher, “Flavor Changing Leptonic Decays of Heavy Higgs Bosons,” *Phys. Rev.*, vol. **D93**, no. 5, p. 055021, (2016). [arXiv:hep-ph/1601.03973].
- [20] O. Eberhardt, A. Lenz, and J. Rohrwild, “Less space for a new family of fermions,” *Phys. Rev. D*, vol. **82**, no. 20, p. 095006, 2010. [arXiv:hep-ph/1005.3505v31].
- [21] C. Patrignani *et al.*, “Review of Particle Physics,” *Chin. Phys.*, vol. **C40**, no. 10, p. 100001, (2016).
- [22] O. Eberhardt *et al.*, “Status of the fourth fermion generation before ICHEP2012: Higgs data and electroweak precision observables,” *Phys. Rev.*, vol. **D86**, p. 074014, (2012). [arXiv:hep-ph/1207.0438].
- [23] O. Eberhardt *et al.*, “Impact of a Higgs Boson at a Mass of 126 GeV on the Standard Model with Three and Four Fermion Generations,” *Phys. Rev. Lett.*, vol. **109**, p. 241802, Dec 2012.
- [24] Q. Li *et al.*, “Higgs Boson Production via Gluon Fusion in the Standard Model with four Generations,” *Phys. Rev. D*, vol. **83**, p. 094018, May 2011. [arXiv:hep-ph/1011.4484].
- [25] S. Bar-Shalom, S. Nandi, and A. Soni, “Two Higgs doublets with 4th generation fermions - models for TeV-scale compositeness,” *Phys. Rev.*, vol. **D84**, p. 053009, (2011). [arXiv:hep-ph/1105.6095].
- [26] M. Hashimoto, “Constraints on the mass spectrum of fourth generation fermions and higgs bosons,” *Phys. Rev. D*, vol. **81**, p. 075023, Apr (2010).
- [27] M. Baak *et al.*, “Updated status of the global electroweak fit and constraints on new physics,” *Eur. Phys. J. C*, vol. **72**, no. 5, p. 2003, (2012).
- [28] S. Banerjee, M. Frank, and S. K. Rai, “Higgs data confronts Sequential Fourth Generation Fermions in the Higgs Triplet Model,” *Phys. Rev.*, vol. **D89**, no. 7, p. 075005, (2014). [arXiv/hep-ph:1312.4249].
- [29] N. Chen and H.-J. He, “LHC Signatures of Two-Higgs-Doublets with Fourth Family,” *JHEP*, vol. **04**, p. 062, (2012). [arXiv/hep-ph:1202.3072].

- [30] D. Das, A. Kundu, and I. Saha, “Higgs data does not rule out a sequential fourth generation,” (2017). [arXiv/hep-ph:1707.03000].
- [31] S. Bar-Shalom and A. Soni, “Chiral heavy fermions in a two Higgs doublet model: 750 GeV resonance or not,” *Physics Letters B*, vol. 766, pp. 1 – 10, (2017). [arXiv:1607.04643].
- [32] S. Kanemura, M. Kikuchi, and K. Yagyu, “Fingerprinting the extended Higgs sector using one-loop corrected Higgs boson couplings and future precision measurements,” *Nucl. Phys.*, vol. **B896**, pp. 80–137, (2015). [arXiv:hep-ph/1502.07716].
- [33] L. Bellantoni *et al.*, “Masses of a Fourth Generation with Two Higgs Doublets,” *Phys. Rev.*, vol. **D86**, p. 034022, (2012). [arXiv:hep-ph/1205.5580].
- [34] A. Denner *et al.*, “Higgs Production and Decay with a Fourth Standard-Model-Like Fermion Generation,” *Eur. Phys. J.*, vol. **C72**, p. 1992, (2012). [arXiv:hep-ph/1111.6395].
- [35] A. Dighe *et al.*, “Large mass splittings for fourth generation fermions allowed by LHC Higgs exclusion,” *Phys. Rev.*, vol. **D85**, p. 114035, (2012). [arXiv:hep-ph/1204.3550].
- [36] G. Aad *et al.*, “Search for charged Higgs bosons through the violation of lepton universality in $t\bar{t}$ events using pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS experiment,” *JHEP*, vol. **03**, p. 076, (2013). [arXiv:hep-ex/1212.3572].
- [37] G. Aad *et al.*, “Search for a CP-odd Higgs boson decaying to Zh in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector,” *Phys. Lett.*, vol. **B744**, pp. 163–183, (2015). [arXiv:hep-ex/1502.04478].
- [38] G. Aad *et al.*, “Search for an additional, heavy Higgs boson in the $H \rightarrow ZZ$ decay channel at $\sqrt{s} = 8$ TeV in pp collision data with the ATLAS detector,” *Eur. Phys. J.*, vol. **C76**, no. 1, p. 45, (2016). [arXiv:hep-ex/1507.05930].
- [39] V. Khachatryan *et al.*, “Searches for heavy Higgs bosons in two-Higgs-doublet models and for $t \rightarrow ch$ decay using multilepton and diphoton final states in pp collisions at 8 TeV,” *Phys. Rev.*, vol. **D90**, p. 112013, (2014). [arXiv:hep-ex/1410.2751].
- [40] V. Khachatryan *et al.*, “Search for a pseudoscalar boson decaying into a Z boson and the 125 GeV Higgs boson in $\ell^+\ell^-b\bar{b}$ final states,” *Phys. Lett.*, vol. **B748**, pp. 221–243, (2015). [arXiv:hep-ex/1504.04710].
- [41] V. Khachatryan *et al.*, “Search for diphoton resonances in the mass range from 150 to 850 GeV in pp collisions at $\sqrt{s} = 8$ TeV,” *Phys. Lett.*, vol. **B750**, pp. 494–519, (2015). [arXiv:hep-ex/1506.02301].
- [42] V. Khachatryan *et al.*, “Search for a charged Higgs boson in pp collisions at $\sqrt{s} = 8$ TeV,” *JHEP*, vol. **11**, p. 018, (2015). [arXiv:hep-ex/1508.07774].
- [43] V. Khachatryan *et al.*, “Searches for a heavy scalar boson H decaying to a pair of 125 GeV Higgs bosons hh or for a heavy pseudoscalar boson A decaying to Zh , in the final states with $h \rightarrow \tau\tau$,” *Phys. Lett.*, vol. **B755**, pp. 217–244, (2016). [arXiv:hep-ex/1510.01181].
- [44] A. G. Akeroyd *et al.*, “Prospects for charged higgs searches at the lhc,” *Eur. Phys. J. C*, vol. **77**, no. 5, p. 276, (2017).
- [45] M. Geller *et al.*, “The 125 GeV Higgs in the context of four generations with 2 Higgs doublets,” *Phys. Rev.*, vol. **D86**, p. 115008, (2012). [arXiv:hep-ph/1209.4081].
- [46] S. Bar-Shalom, S. Nandi, and A. Soni, “Muon $g - 2$ and lepton flavor violation in a two Higgs doublets model for the fourth generation,” *Phys. Lett.*, vol. **B709**, pp. 207–217, (2012). [arXiv:hep-ph/1112.3661].
- [47] M. Lindner, M. Platscher, and F. S. Queiroz, “A Call for New Physics : The Muon Anomalous Magnetic Moment and Lepton Flavor Violation,” (2016). [arXiv/hep-ph:1610.06587].

- [48] J. Heeck, M. Holthausen, W. Rodejohann, and Y. Shimizu, “ $h \rightarrow \mu\tau$ in Abelian and non-Abelian flavor symmetry models,” *Nuclear Physics B*, vol. **896**, pp. 281–310, (2015). [arXiv:hep-ph/1412.3671].
- [49] M. Campos, A. Cárcamo Hernández, H. Päs, and E. Schumacher, “Higgs $\rightarrow \mu\tau$ as an indication for S_4 flavor symmetry,” *Phys. Rev. D*, vol. **91**, p. 116011, Jun (2015). [arXiv:hep-ph/1408.1652].
- [50] M. Aoki, S. Kanemura, K. Sakurai, and H. Sugiyama, “Testing neutrino mass generation mechanisms from the lepton flavor violating decay of the Higgs boson,” *Physics Letters B*, vol. **763**, pp. 352 – 357, (2016). [arXiv:hep-ph/1607.08548].
- [51] S. Baek, T. Nomura, and H. Okada, “An explanation of one-loop induced $h \rightarrow \mu\tau$ decay,” *Physics Letters B*, vol. **759**, pp. 91 – 98, (2016). [arXiv:hep-ph/1604.03738].
- [52] L. Wang, S. Yang, and X. Han, “ $h \rightarrow \mu\tau$ and muon $g - 2$ in the alignment limit of two-Higgs-doublet model,” *Nuclear Physics B*, vol. **919**, pp. 123 – 141, (2017). [arXiv:hep-ph/1606.04408].