



Open Research Online

The Open University's repository of research publications and other research outputs

Draft Genome Sequences of Obligate Methylophs *Methylovorus* sp. Strain MM2 and *Methylobacillus* sp. Strain MM3, Isolated from Grassland Soil

Journal Item

How to cite:

Macey, Michael Christopher; Pratscher, Jennifer; Crombie, Andrew and Murrell, J. Colin (2018). Draft Genome Sequences of Obligate Methylophs *Methylovorus* sp. Strain MM2 and *Methylobacillus* sp. Strain MM3, Isolated from Grassland Soil. *Microbiology Resource Announcements*, 7(8), article no. e00824-18.

For guidance on citations see [FAQs](#).

© 2018 The Authors

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1128/MRA.00824-18>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk



Draft Genome Sequences of Obligate Methylophils *Methylovorus* sp. Strain MM2 and *Methylobacillus* sp. Strain MM3, Isolated from Grassland Soil

Michael Christopher Macey,^{a*} Jennifer Pratscher,^{a*} Andrew Crombie,^b J. Colin Murrell^a

^aSchool of Environmental Sciences, University of East Anglia, Norwich, United Kingdom

^bSchool of Biological Sciences, University of East Anglia, Norwich, United Kingdom

ABSTRACT Methylophils of the family *Methylophilaceae* were isolated from grassland soil. Here, we report the draft genome sequences of two obligate methylophils, *Methylovorus* sp. strain MM2 and *Methylobacillus* sp. strain MM3. These genome sequences provide further insights into the genetic and metabolic diversity of the *Methylophilaceae*.

The family *Methylophilaceae* is composed of four genera containing facultative and obligate methanol-utilizing methylophils (1–4). Members of the *Methylophilaceae* have been isolated from a range of locations, including terrestrial and marine environments (5–10). Here, we report the draft genome sequences of two obligate methylophils, *Methylovorus* sp. strain MM2 and *Methylobacillus* sp. strain MM3. The obligate nature of these methylophils was confirmed through growth experiments. The strains were isolated from soil collected at a 5-cm depth from grassland in Bawburgh, Norfolk, United Kingdom (52.6276 N, 1.1784 E).

Genome sequencing was performed by MicrobesNG using the Illumina MiSeq platform, producing 2 × 250-bp paired-end reads. Trimmed sequences were assembled using SPAdes version 3.7.1, and genome annotation was performed using the RAST annotation server (<http://rast.nmpdr.org>) (11, 12). Coverage of the genomes was calculated using BWA, SAMtools, and BEDTools genomecov (13–15). The *Methylovorus* sp. MM2 genome is composed of 27 contigs and includes 2,291 coding sequences (CDSs), 1 16S rRNA gene copy, and 46 tRNAs. The genome size is 2.42 Mb, with 46% G+C content. The genome of *Methylobacillus* sp. MM3, with 2.95 Mb and 57% G+C content, is composed of 64 contigs and includes 2,897 CDSs and 3 copies of 16S rRNA genes. Both genomes had 30-fold coverage.

Both genomes contain pyrroloquinoline quinone methanol dehydrogenases. *Methylobacillus* sp. MM3 possesses three separate gene clusters for the alternative methanol dehydrogenase XoxF (16, 17) and no copies of the canonical methanol dehydrogenase-encoding genes *mxoA*. *Methylovorus* sp. MM2 possesses three copies of *xoxF* and one set of the genes *mxoA*. All genes encoding the *N*-methylglutamate pathway for methylamine utilization (*mgdABC*, *gmaS*, and *mgsABC*) (18, 19) are present only in the genome of *Methylobacillus* sp. MM3, in addition to genes that encode dimethylamine dehydrogenase and trimethylamine dehydrogenase enzymes (*dmd* and *tmd*) (20, 21). The genes for an assimilatory nitrate reductase (*nasAB*) and the complete denitrification pathway (*narGHI*, *nirK*, *nirS*, *norB*, and *nosZ*) are present in the genome of *Methylobacillus* sp. MM3, while *Methylovorus* sp. MM2 possesses only an assimilatory nitrate reductase (*nasAB*) and a dissimilatory nitrite reductase (*nirBD*).

Data availability. These whole-genome shotgun projects have been deposited at DDBJ/ENA/GenBank under accession numbers [LXTQ00000000](https://www.ncbi.nlm.nih.gov/nuccore/LXTQ00000000) for *Methylobacillus* sp.

Received 5 June 2018 Accepted 31 July 2018 Published 30 August 2018

Citation Macey MC, Pratscher J, Crombie A, Murrell JC. 2018. Draft genome sequences of obligate methylophils *Methylovorus* sp. strain MM2 and *Methylobacillus* sp. strain MM3, isolated from grassland soil. *Microbiol Resour Announc* 7:e00824-18. <https://doi.org/10.1128/MRA.00824-18>.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2018 Macey et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to J. Colin Murrell, j.c.murrell@uea.ac.uk.

* Present address: Michael Christopher Macey, School of Environment, Earth and Ecosystem Sciences, The Open University, Milton Keynes, United Kingdom; Jennifer Pratscher, Lyell Centre, Heriot-Watt University, Edinburgh, United Kingdom.

MM3 and [LXUF0000000](#) for *Methylovorus* sp. MM2. The versions described in this paper are the first versions. The strains are available from the authors upon request.

ACKNOWLEDGMENTS

This work was supported by the Norwich Research Park Doctoral Training Program, the Earth and Life Systems Alliance (ELSA) at the University of East Anglia, NERC Independent Research Fellowship NE/L010771/1, and Leverhulme Early Career Fellowship ECF-2016-626. Genome sequencing was provided by MicrobesNG (<http://www.microbesng.uk>), which is supported by the BBSRC (grant number BB/Lo24209/1).

REFERENCES

- Kaparullina EN, Trotsenko YA, Doronina NV. 2017. *Methylobacillus methanolivorans* sp. nov., a novel non-pigmented obligately methylotrophic bacterium. *Int J Syst Evol Microbiol* 67:425–431. <https://doi.org/10.1099/ijsem.0.001646>.
- Doronina NV, Kaparullina EN, Trotsenko YA. 2016. Emended description of *Methylovorus glucosotrophus* Govorukhina and Trotsenko 1991. *Microbiology* 85:548–552. <https://doi.org/10.1134/S0026261716050040>.
- Kalyuzhnaya MG, Beck DAC, Vorobev A, Smalley N, Kunkel DD, Lidstrom ME, Chistoserdova L. 2012. Novel methylotrophic isolates from lake sediment, description of *Methylotenera versatilis* sp. nov. and emended description of the genus *Methylotenera*. *Int J Syst Evol Microbiol* 62: 106–111. <https://doi.org/10.1099/ijs.0.029165-0>.
- Jenkins O, Byrom D, Jones D. 1987. *Methylophilus*: a new genus of methanol-utilizing bacteria. *Int J Syst Evol Microbiol* 37:446–448. <https://doi.org/10.1099/00207713-37-4-446>.
- Doronina NV, Gogleva AA, Trotsenko YA. 2012. *Methylophilus glucosoxydans* sp. nov., a restricted facultative methylotroph from rice rhizosphere. *Int J Syst Evol Microbiol* 62:196–201. <https://doi.org/10.1099/ijs.0.024620-0>.
- Gogleva AA, Kaparullina EN, Doronina NV, Trotsenko YA. 2011. *Methylobacillus arboreus* sp. nov., and *Methylobacillus gramineus* sp. nov., novel non-pigmented obligately methylotrophic bacteria associated with plants. *Syst Appl Microbiol* 34:477–481. <https://doi.org/10.1016/j.syapm.2011.03.005>.
- Doronina NV, Trotsenko YA, Kolganova TV, Tourova TP, Salkinoja-Salonen MS. 2004. *Methylobacillus pratensis* sp. nov., a novel non-pigmented, aerobic, obligately methylotrophic bacterium isolated from meadow grass. *Int J Syst Evol Microbiol* 54:1453–1457. <https://doi.org/10.1099/ijs.0.02956-0>.
- Madhaiyan M, Poonguzhali S, Kwon S-W, Sa T-M. 2009. *Methylophilus rhizosphaerae* sp. nov., a restricted facultative methylotroph isolated from rice rhizosphere soil. *Int J Syst Evol Microbiol* 59:2904–2908. <https://doi.org/10.1099/ijs.0.009811-0>.
- Doronina NV, Kaparullina EN, Trotsenko YA. 2011. *Methylovorus mentalis*, a novel species of aerobic obligate methylotrophic bacteria associated with plants. *Microbiology* 80:713–719. <https://doi.org/10.1134/S0026261711050043>.
- Xia F, Zou B, Shen C, Zhu T, Gao X-H, Quan Z-X. 2015. Complete genome sequence of *Methylophilus* sp. TWE2 isolated from methane oxidation enrichment culture of tap-water. *J Biotechnol* 211:121–122. <https://doi.org/10.1016/j.jbiotec.2015.07.023>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Aziz RK, Bartels D, Best AA, Dejongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, Mcneil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *Nucleic Acids Res* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map (SAM) format and SAMtools. *Bioinformatics* 25:2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>.
- Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26:841–842. <https://doi.org/10.1093/bioinformatics/btq033>.
- Chistoserdova L. 2011. Modularity of methylotrophy, revisited. *Environ Microbiol* 13:2603–2622. <https://doi.org/10.1111/j.1462-2920.2011.02464.x>.
- Keltjens JT, Pol A, Reimann J, Op den Camp HJM. 2014. PQQ-dependent methanol dehydrogenases: rare-earth elements make a difference. *Appl Microbiol Biotechnol* 98:6163–6183. <https://doi.org/10.1007/s00253-014-5766-8>.
- Chen Y, Scanlan J, Song L, Crombie A, Rahman MT, Schafer H, Murrell JC. 2010. γ -Glutamylmethylamide is an essential intermediate in the metabolism of methylamine by *Methylocella silvestris*. *Appl Environ Microbiol* 76:4530–4537. <https://doi.org/10.1128/AEM.00739-10>.
- Latypova E, Yang S, Wang Y-S, Wang T, Chavkin TA, Hackett M, Schäfer H, Kalyuzhnaya MG. 2010. Genetics of the glutamate-mediated methylamine utilization pathway in the facultative methylotrophic beta-proteobacterium *Methyloversatilis universalis* FAM5. *Mol Microbiol* 75: 426–439. <https://doi.org/10.1111/j.1365-2958.2009.06989.x>.
- Boyd G, Math FS, Leonard C, Scrutton NS. 1992. Trimethylamine dehydrogenase of bacterium W₃A₁ molecular cloning, sequence determination and over-expression of the gene. *FEBS Lett* 308:271–276. [https://doi.org/10.1016/0014-5793\(92\)81291-5](https://doi.org/10.1016/0014-5793(92)81291-5).
- Yang C-C, Packman LC, Scrutton NS. 1995. The primary structure of *Hyphomicrobium* X dimethylamine dehydrogenase. *Eur J Biochem* 232: 264–271. <https://doi.org/10.1111/j.1432-1033.1995.tb20808.x>.