

Measurement of the neutrino component of an antineutrino beam observed by a nonmagnetized detector

A. A. Aguilar-Arevalo,¹² C. E. Anderson,¹⁵ S. J. Brice,⁶ B. C. Brown,⁶ L. Bugel,¹¹ J. M. Conrad,¹¹ R. Dharmapalan,¹ Z. Djurcic,² B. T. Fleming,¹⁵ R. Ford,⁶ F. G. Garcia,⁶ G. T. Garvey,⁹ J. Grange,⁷ J. A. Green,^{8,9} R. Imlay,¹⁰ R. A. Johnson,³ G. Karagiorgi,¹¹ T. Katori,^{8,11} T. Kobilarcik,⁶ S. K. Linden,¹⁵ W. C. Louis,⁹ K. B. M. Mahn,⁵ W. Marsh,⁶ C. Mauger,⁹ W. Metcalf,¹⁰ G. B. Mills,⁹ J. Mirabal,⁹ C. D. Moore,⁶ J. Mousseau,⁷ R. H. Nelson,⁴ V. Nguyen,¹¹ P. Nienaber,¹⁴ J. A. Nowak,¹⁰ B. Osmanov,⁷ A. Patch,^{9,11} Z. Pavlovic,⁹ D. Perevalov,¹ C. C. Polly,⁶ H. Ray,⁷ B. P. Roe,¹³ A. D. Russell,⁶ M. H. Shaevitz,⁵ M. Sorel,^{5,*} J. Spitz,¹⁵ I. Stancu,¹ R. J. Stefanski,⁶ R. Tayloe,⁸ M. Tzanov,⁴ R. G. Van de Water,⁹ M. O. Wascko,^{10,†} D. H. White,⁹ M. J. Wilking,⁴ G. P. Zeller,⁶ and E. D. Zimmerman⁴

(The MiniBooNE Collaboration)

¹University of Alabama, Tuscaloosa, Alabama 35487, USA²Argonne National Laboratory, Argonne, Illinois 60439, USA³University of Cincinnati, Cincinnati, Ohio 45221, USA⁴University of Colorado, Boulder, Colorado 80309, USA⁵Columbia University, New York, New York 10027, USA⁶Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA⁷University of Florida, Gainesville, Florida 32611, USA⁸Indiana University, Bloomington, Indiana 47405, USA⁹Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA¹⁰Louisiana State University, Baton Rouge, Louisiana 70803, USA¹¹Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA¹²Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, D.F. 04510, México¹³University of Michigan, Ann Arbor, Michigan 48109, USA¹⁴Saint Mary's University of Minnesota, Winona, Minnesota 55987, USA¹⁵Yale University, New Haven, Connecticut 06520, USA

(Received 20 March 2011; published 26 October 2011)

Two methods are employed to measure the neutrino flux of the antineutrino-mode beam observed by the MiniBooNE detector. The first method compares data to simulated event rates in a high-purity ν_μ -induced charged-current single π^+ (CC1 π^+) sample while the second exploits the difference between the angular distributions of muons created in ν_μ and $\bar{\nu}_\mu$ charged-current quasielastic (CCQE) interactions. The results from both analyses indicate the prediction of the neutrino flux component of the predominately antineutrino beam is overestimated—the CC1 π^+ analysis indicates the predicted ν_μ flux should be scaled by 0.76 ± 0.11 , while the CCQE angular fit yields 0.65 ± 0.23 . The energy spectrum of the flux prediction is checked by repeating the analyses in bins of reconstructed neutrino energy, and the results show that the spectral shape is well-modeled. These analyses are a demonstration of techniques for measuring the neutrino contamination of antineutrino beams observed by future nonmagnetized detectors.

DOI: [10.1103/PhysRevD.84.072005](https://doi.org/10.1103/PhysRevD.84.072005)

PACS numbers: 14.60.Lm, 14.60.Pq, 14.60.St

I. INTRODUCTION

If θ_{13} is nonzero, next-generation neutrino oscillation experiments will embark on a program to measure the neutrino mass ordering and look for evidence of CP violation in the neutrino sector. This effort will require precise oscillation measurements with both neutrino and antineutrino beams in order to isolate these effects. Since beams

produced in an accelerator environment are never purely neutrino nor antineutrino in content, detectors must be able to separate the two contributions. Most commonly, this is achieved by employing a magnetic field to identify the final-state μ^- (or μ^+) produced in charged-current ν_μ (or $\bar{\nu}_\mu$) interactions. A handle on the overall level and energy dependence of ν_μ versus $\bar{\nu}_\mu$ induced events, however, is also possible in unmagnetized detectors with a suitable choice of reaction channels.

Accelerator-based neutrino beams are typically created by colliding proton beams with thick nuclear targets. Mesons produced at a variety of energies and angles are focused by a magnetic horn before entering a decay tunnel.

*Present address: IFIC, Universidad de Valencia and CSIC, Valencia 46071, Spain.

†Present address: Imperial College; London SW7 2AZ, United Kingdom., USA

Meson decays can be calculated sufficiently well for a given beam geometry that the neutrino flux uncertainties arise mainly from uncertainties in the meson production cross sections. In particular, to avoid extrapolating data taken with diverse nuclear target materials or proton energies, neutrino experiments require dedicated hadron production cross section measurements taken with the same beam energy and target to obtain a reliable flux prediction. If an accelerator-based neutrino experiment lacks such hadron production data, it may be able to meet its oscillation analysis goals using calibrations from a near detector; however, the secondary physics goal of measuring neutrino-nucleon absolute cross sections will still be limited by flux uncertainties.

To avoid ambiguity, in this paper references to “neutrinos” are not meant to also refer to antineutrinos, and “mode” refers to the polarity of the magnetic focusing horn used in the beamline. In this way, for example, “antineutrino events” refers to antineutrino-induced events exclusively while “antineutrino-mode events” refers to data obtained when the horn polarity focuses negatively charged particles, which is a mix of neutrino and antineutrino-induced events.

The Mini Booster Neutrino Experiment (MiniBooNE) is located at Fermilab in Batavia, Illinois, and has made many oscillation [1–5] and cross section [6–13] measurements. For MiniBooNE, the pion production data crucial to the flux model come from proton-beryllium cross sections on a 5% interaction length target reported by the HARP experiment [14]. However, even with dedicated data appropriate to the experimental setup of MiniBooNE, there remain small regions of phase space relevant to MiniBooNE not covered by these HARP measurements. Of particular importance is the production of very forward pions with respect to the direction of the incoming proton beam. This is the dominant production region of parent particles contributing neutrinos to the antineutrino-mode beam, or vice versa. MiniBooNE uses a magnetic horn to defocus the majority of these background parent particles, but as Fig. 1 suggests, the very forward pions can escape magnetic deflection. This same angular region suffers from a sizable beam-related proton background and would also require a model-dependent acceptance correction [15]. For these reasons, pion cross sections in the $\theta_\pi < 30$ mrad region, where θ_π is the angle the outgoing pion makes with respect to the incoming proton beam, are not reported by HARP and the majority of the MiniBooNE flux prediction arising from π^+ (π^-) decay while focusing π^- (π^+) is extrapolated from the available hadron production data. The hadron production data cover $\sim 90\%$ of sign-selected pions, while less than 25% of oppositely charged pions in the same beam are constrained. Some of these acceptance limitations could be reduced by use of the long-target data taken by HARP, which are actively being analyzed.




FIG. 1 (color online). Predicted angular distributions of pions with respect to the incident proton beam (θ_π) producing ν_μ and $\bar{\nu}_\mu$ in (a) neutrino mode and (b) antineutrino mode. Only pions leading to ν_μ and $\bar{\nu}_\mu$ events in the detector are shown, and all distributions are normalized to 5.66×10^{20} protons on target. Arrows indicate the region where HARP data [14] are available.

The overall contamination rate is more significant in antineutrino mode due to effects from both flux and cross section: the leading-particle effect at the target preferentially produces about twice as many π^+ as π^- , and the neutrino cross section is about 3 times higher than the antineutrino cross section in the MiniBooNE energy range (~ 1 GeV) [16]. For these reasons antineutrino-induced events are not a serious complication for neutrino-mode running, as these flux and cross section effects conspire to suppress their contribution, while the same effects amplify the neutrino contamination in antineutrino-mode data. Simulation predicts antineutrino events account for $\sim 1\%$ of neutrino-mode data while neutrinos are responsible for $\sim 30\%$ of antineutrino-mode data. This motivates a dedicated study of the neutrino flux contribution to the antineutrino-mode data. A data set corresponding to 5.66×10^{20} protons on target is analyzed here and is important for both the ongoing MiniBooNE antineutrino oscillation search [4,5] and antineutrino cross section measurements.

Two approaches for measuring the neutrino flux in antineutrino mode are taken. In the first method, a high-purity sample of charged-current single π^+ ($\text{CC1}\pi^+$) events isolate the ν_μ contribution in the beam. The second method exploits the interference term in the charged-current quasielastic (CCQE) cross section, where the angular distribution of final-state muons are predicted to be distinct for ν_μ compared to $\bar{\nu}_\mu$ interactions. Both techniques were introduced in the MiniBooNE antineutrino-mode run proposal [17]. These two approaches offer complementary means of measuring the neutrino flux component in antineutrino-mode data, with the CCQE sample providing a constraint at lower neutrino energies while the $\text{CC1}\pi^+$ measurement covers higher energies. They provide both a check of the MiniBooNE beam simulation in a region not covered by external data and demonstrate a set of techniques for measuring the ν_μ contamination in an antineutrino-mode beam in the absence of a magnetized detector. It has been argued elsewhere that even modest statistical separation of charged-current neutrino and antineutrino events, afforded by the kind of analyses presented here, may be sufficient to meet the physics goals in proposed future experiments such as neutrino factories [18].

This paper is organized as follows: the MiniBooNE experiment is described in Sec. II while Sec. III details the neutrino and antineutrino scattering models. Two techniques to measure the neutrino contribution to the antineutrino flux are presented in Secs. IV and V. The results are compared in Sec. VI, implications for other neutrino experiments are discussed in Sec. VII and this work is summarized in Sec. VIII.

II. THE MINIBOONE EXPERIMENT

A. Beamline and flux

The Booster Neutrino Beamline (BNB) provides the neutrino and antineutrino flux to MiniBooNE. A beam of 8 GeV-kinetic-energy protons is extracted from the Booster synchrotron in “spills” of 5×10^{12} protons over 1.6 μs at a maximum rate of 5 Hz. A lattice of alternatively focusing and defocusing quadrupole magnets steer the proton bunches to a beryllium target 71 cm (1.75 interaction lengths) long. The protons collide with the target to create a spray of secondary particles. An aluminum electromagnetic horn surrounding the target is pulsed to coincide with the p-Be collisions, creating a toroidal magnetic field to focus mesons of the desired charge. The horn pulses are such that the magnetic field is constant for the duration of the proton spill. In neutrino mode, the magnetic horn focuses positively charged secondary particles while defocusing those with negative charge; the horn effects are reversed in antineutrino mode. The focused mesons are allowed to decay in a 50 m air-filled decay region which terminates at a steel beam dump. The dominant decay

modes of the mesons, mostly pions, produce muon neutrinos and antineutrinos.

A GEANT4-based model [19] is used to predict the neutrino and antineutrino flux at the detector. The simulation considers proton travel to the target, p-Be interactions in the target including meson production, magnetic horn focusing, particle propagation, meson decay, and finally neutrino and antineutrino transport to the detector. As mentioned earlier, measurement of pion cross sections from p-Be interactions are obtained from the HARP experiment. The HARP double differential cross section error matrix is used to set pion production uncertainties [20]. Even with valuable data constraints, meson production at the target contributes the largest systematic error to the flux prediction. The fractional uncertainty on pion production is $\sim 8\%$ around the flux peak, while the uncertainty grows significantly in regions dominated by pions unconstrained by HARP data. The flux prediction in neutrino and antineutrino modes is presented in Fig. 2. Details of the beamline and flux prediction are given in Ref. [20].




FIG. 2 (color online). The MiniBooNE flux prediction for (a) neutrino mode and (b) antineutrino mode. Because of the leading-particle effect, the neutrino contribution to the antineutrino-mode flux is more significant compared to the antineutrino component of the neutrino-mode beam. Plots taken from Ref. [20].

B. Detector

The MiniBooNE detector is a 6.1 m-radius sphere filled with 818 tons of pure Marcol7 mineral oil. It houses 1520 8-inch Hamamatsu photomultiplier tubes (PMTs) segregated into two optically isolated regions: an inner signal region of 575 cm radius and an outer veto shell of thickness 35 cm. The former contains 1280 PMTs (11.3% coverage) while the latter holds 240 PMTs. The veto region is used to enforce containment of charged particles produced by neutrinos and antineutrinos from the beam and reject charged particles entering the tank.

The mineral oil has a density of 0.845 g/cm^3 with an index of refraction of 1.47 at 20°C . Charged particles with velocity $\beta > 0.68$ produce Cherenkov radiation. Particle identification and reconstruction is principally obtained through the pattern and timing of this prompt Cherenkov light; however, delayed scintillation light present due to fluorescent components in the oil has also been used effectively to provide energy information for charged particles produced below Cherenkov threshold [10].

MiniBooNE electronics record PMT charge and time information beginning about $5 \mu\text{s}$ before the $1.6 \mu\text{s}$ BNB proton delivery. Data are recorded for a total of $19.2 \mu\text{s}$. The $5 \mu\text{s}$ interval before the beam spill is primarily present to minimize data contamination caused by cosmic ray muons stopping in the signal region prior to the start of the DAQ window. PMT activity is recorded for more than $10 \mu\text{s}$ after beam delivery to observe electrons from the at-rest decay of muons (hereafter referred to as “Michel” electrons) subsequent to the initial neutrino or antineutrino-induced interaction.

The detector response to muons is measured using a dedicated muon tagging system that independently measures the energy and direction of cosmic ray muons up to 800 MeV. MiniBooNE employs a scintillator hodoscope directly above the detector and seven internal scintillator cubes at different depths, each connected to a dedicated 1-inch PMT for readout. The measured ranges and directions of muons traversing the hodoscope and stopping in cubes are used to verify muon reconstruction algorithms. The energy (angle) resolution improves from 12% (5.4 deg) at 100 MeV to 3.4% (1.0 deg) at 800 MeV. Full detector details and calibrations are available in Ref. [21].

C. Detector simulation

The detector response to particle interactions and propagation is simulated using GEANT3 [22]. The entire detector geometry is considered, including the steel tank, external supports and main inner components. In addition, the surrounding environment composed of dirt external to the MiniBooNE enclosure, the concrete cylindrical housing and the air-filled gap between the detector and walls is treated. Of critical importance is the treatment of

particle transport in the detector medium. The GEANT3 program takes as input the final-state particles emerging from the nucleus and simulates their propagation in the detector.

With a few exceptions, MiniBooNE uses the standard GEANT3 settings to simulate physics processes. Deviations include a custom model for light propagation in the detector oil and a substitution of the hadronic interaction model. The default GFLUKA hadron model is replaced by the GCALOR [23] package, which better models pion absorption ($\pi^\pm + X \rightarrow X'$) and charge-exchange ($\pi^\pm + X \leftrightarrow \pi^0 + X'$) processes. This is particularly relevant for the present analysis, where the predicted event composition of the two interaction samples studied is dependent on the pion survival model. Based on comparisons with external data [24] and the GCALOR prediction, an uncertainty of 35% (50%) is assigned to the pion absorption (charge exchange) interaction *in the detector medium*. The uncertainty for the same processes *inside the nucleus* is discussed in Sec. III C.

The model for light propagation in the oil is formed using a combination of external measurements and calibration data. Photon emission through Cherenkov and scintillation processes is simulated and propagated until the photon either is absorbed or hits a PMT photocathode, possibly leading to photoelectron production. Light emission, attenuation and scattering are included. The optical model of the detector describes the wavelength, time, and angular dependence of these processes [25].

III. PREDICTED NEUTRINO AND ANTINEUTRINO INTERACTIONS

MiniBooNE uses the NUANCE [16] event generator to simulate neutrino and antineutrino interactions in the detector. NUANCE includes a comprehensive neutrino and antineutrino cross section model which considers known interactions in the neutrino and antineutrino energy range from $\sim 100 \text{ MeV}$ to 1 TeV. Ninety-nine reactions are modeled separately and combined with nuclear models describing bound nucleon states and final-state interactions to predict event rates and kinematics.

Bound nucleons in the detector medium are described by the Relativistic Fermi Gas model [26]. This assumes the nucleons to be independent and quasifree. Also specified is a hard cutoff in available struck nucleon energies as dictated by the exclusion principle.

The neutrino and antineutrino interaction types relevant to the analysis presented here are charged-current quasielastic (Sec. III A) and pion production (Sec. III B). The neutrino-induced absolute cross sections for both processes have been measured at MiniBooNE using a flux prediction well determined by HARP data. These cross section measurements are utilized in the antineutrino-mode simulation.

A. Charged-current quasielastic scattering

To model CCQE interactions, this analysis uses measured cross sections from the MiniBooNE neutrino-mode CCQE data [9] and a model which has been found to reproduce well the kinematics of such events. Specifically, MiniBooNE adopts the CCQE scattering formalism of Smith-Moniz [26]. The vector component of the interaction is measured by electron scattering experiments and is assumed to have a nondipole form [27]. The axial-vector form factor employs a dipole construction, containing an “axial mass,” M_A , taken either from MiniBooNE or external data, depending on the neutrino target.

The MiniBooNE mineral oil is composed of C_nH_{2n+2} , $n \sim 20$, and the prediction for CCQE scattering is different for the two flavors of target. In the present analysis, $M_A^{\text{eff}} = 1.35 \pm 0.17$ GeV, together with a Pauli blocking adjustment, $\kappa = 1.007 \pm 0.012$, are assumed for bound nucleon scattering. These values come from a high-statistics analysis of MiniBooNE ν_μ CCQE events on carbon [9] and are consistent with values recently determined from an independent MiniBooNE neutral-current elastic scattering sample [10]. A previous shape-only study has shown that these CCQE model parameters reproduce the MiniBooNE antineutrino-mode data shape [28], and therefore the same M_A^{eff} and κ values are applied to both ν_μ and $\bar{\nu}_\mu$ CCQE scattering events on carbon.

For free scattering of hydrogen, a process accessible to antineutrino and not neutrino CCQE events, a value of $M_A = 1.03 \pm 0.02$ GeV is used based on a global fit to previous light target data [29].

In the case of carbon scattering, the superscript “eff”, short for “effective,” on M_A is introduced to allow for the possibility that nuclear effects are responsible for the apparent discrepancy between the MiniBooNE carbon-based measurements and light target results. This is also theoretically motivated by a possible reconciliation between the measurements through a mechanism resulting in intranuclear correlations of greater importance than previously thought [30–34]. Such a mechanism would indicate a larger CCQE cross section for nuclear targets than for free scattering, which in this case, is reflected in the higher M_A choice for carbon versus hydrogen scattering.

B. Pion production

Baryonic resonances are the dominant source of single pion production at MiniBooNE. The formalism to describe these events is taken from the Rein-Sehgal model [35], where the relativistic harmonic oscillator quark model is assumed [36]. Eighteen resonances are considered, however the $\Delta(1232)$ is dominant in the energy range spanned by MiniBooNE. Multipion production mechanisms are also considered, though their contribution is predicted to be small.

The axial masses in the resonance channels are set simultaneously to reproduce inclusive non-MiniBooNE charged-current data [37]. The extracted values are $M_A^{1\pi} = 1.10 \pm 0.27$ GeV (single pion production) and $M_A^{\text{multi-}\pi} = 1.30 \pm 0.52$ GeV (multipion production).

In the present analysis the charged-current single π^+ (CC1 π^+) prediction with these assumptions is adjusted to reproduce the kinematic distributions measured in MiniBooNE neutrino-mode data [9,12].

C. Final-state interactions

For a neutrino or antineutrino interaction with a nucleon bound in carbon, NUANCE propagates the outgoing hadrons including nucleons, mesons and baryonic resonances, and simulates their reinteraction as they exit the nucleus. The initial interaction model employs the impulse approximation which assumes an instantaneous exchange with independent nucleons. Subsequent to the initial neutrino or antineutrino interaction, particles produced inside the nucleus are propagated stepwise in 0.3 fm increments until they emerge from the ~ 2.5 fm radius sphere. Intermittently, the probability for hadronic reinteraction is calculated using a radially-dependent nucleon density distribution [38] along with external $\pi - N$, $N - N$ cross section measurements [39]. For Δ reinteractions ($\Delta + N \rightarrow N + N$), an energy-independent probability of 20% (10%) is taken for $\Delta^+ + N$, $\Delta^0 + N$ ($\Delta^{++} + N$, $\Delta^- + N$) based on K2K data [37] and is assigned 100% uncertainty.

As mentioned earlier, out of all hadronic reinteraction processes, pion absorption and charge exchange ($\pi^\pm + X \leftrightarrow \pi^0 + X'$) are the most relevant in predicting the composition of the CC1 π^+ (Sec. IVA) and CCQE (Sec. VA) samples studied in this analysis. *Intranuclear* fractional uncertainties on pion absorption (charge exchange) are set to 25% (30%) based on comparisons between external data [24] and NUANCE. The simulation of these two processes in the detector medium is addressed separately in the detector simulation (Sec. IIC).

IV. MEASURING THE NEUTRINO FLUX COMPONENT IN THE CC1 π^+ SAMPLE

A. The CC1 π^+ sample

The events in the CC1 π^+ sample in antineutrino mode originate almost exclusively from ν_μ interactions, making it an excellent candidate for measuring the ν_μ content of the antineutrino-mode beam. In the few-GeV energy range, the dominant charged-current single pion production channels contain a final-state π^+ (π^-) in the case of ν_μ ($\bar{\nu}_\mu$) scattering. MiniBooNE cleanly identifies CC1 π^+ events by selecting 3 “subevents”, attributed to the muon from the primary ν_μ interaction and two subsequent decay electrons, one each from the μ^- and π^+ decay chain:

- 1: $\nu_\mu + p(n) \rightarrow \mu^- + p(n) + \pi^+ \hookrightarrow \mu^+ + \nu_\mu$
- 2: $\hookrightarrow e^- + \bar{\nu}_e + \nu_\mu$
- 3: $\hookrightarrow e^+ + \nu_e + \bar{\nu}_\mu$. (1)

The monoenergetic μ^+ from the decay of stopped π^+ does not lead to a separate subevent due to the short lifetime of the π^+ . Subevents are defined as clusters in time of PMT activity (or PMT “hits”). A hit is any PMT pulse passing the discriminator threshold of ~ 0.1 photoelectrons. A temporal cluster of PMT activity with at least 10 hits within a 200 ns window and individual hit times less than 10 ns apart, while allowing for at most two spacings of 10–20 ns, defines a subevent. Apart from detection efficiencies, some neutrino-induced CC1 π^+ events do not enter the three-subevent sample as $\sim 8\%$ of μ^- are captured in carbon [40] and therefore do not lead to the production of a Michel electron. Other selection cuts made to enhance sample purity and improve reconstruction are given with efficiencies in Table I. Cut 1 is the three-subevent criterion previously detailed. Cut 2 requires that the first subevent occur during a 3 μ s time window centered on the BNB proton spill. Cut 3 rejects events close to the detector edge that are likely to be poorly reconstructed. Selection cuts on the number of tank hits are based primarily on the observation that Michel electrons produce fewer than 200 tank hits. Cut 4 ensures the first subevent is not a Michel electron and rejects low-energy muons that might be reconstructed poorly. Cut 5 requires that the number of hits for the second and third subevents is consistent with a Michel electron. Veto PMT activity is monitored simultaneously with the main tank PMTs, thus Cut 6 ensures no subevent is due to charged particles entering the tank and that all charged particles produced inside the detector are contained. Cut 7 enforces spatial correlation between the end of the muon track and the closest Michel

TABLE I. Summary of selection cuts in the CC1 π^+ sample. Purity and efficiency numbers are sequential and are calculated for the “observable CC1 π^+ ” event signature— $1\mu^-, 1\pi^+$.

Cut #	Description	Efficiency (%)	Purity (%)
0	No cuts	100	10
1	Three subevents	30	29
2	1st subevent in event time window $4000 < T(\text{ns}) < 7000$	28	34
3	All subevents: reconstructed vertex <500 cm from tank center	23	36
4	1st subevent: tank hits >200	22	39
5	2nd, 3rd subevents: tank hits <200	19	65
6	All subevents: veto hits <6	16	78
7	Distance between reconstructed end of 1st subevent and nearest Michel electron vertex <150 cm	12	82

electron vertex. This reduces a class of backgrounds where neither the second nor the third subevent arise from the decay of the muon to a Michel electron. This cut is applied only to the Michel closest to the end of the reconstructed primary muon track as the pion lifetime compared to the muon is short enough that either Michel can occur temporally first.

Charged-current single π^- events induced by $\bar{\nu}_\mu$ are largely rejected by the primary requirement of three subevents because most π^- come to rest and are captured by carbon nuclei [41], yielding no decay electron. The predicted event composition after this selection is presented in Table II. The sample is 82% observable CC1 π^+ events (i.e., events with a single muon, a single π^+ , and any number of nucleons exiting the initial target nucleus). Some $\bar{\nu}_\mu$ CC1 π^- events do make it into the sample, primarily due to decay-in-flight π^- . Starting from an event population that is $\sim 70\%$ $\bar{\nu}_\mu$, this simple two-decay electron requirement remarkably yields a sample that is $\sim 80\%$ pure ν_μ .

B. CC1 π^+ event reconstruction

In this analysis, charged-current single π^+ event reconstruction relies exclusively on the observation of the outgoing muon. Muon kinematics are obtained by the pattern, timing, and total charge of prompt Cherenkov radiation collected by PMTs in the first subevent of the interaction. The topology and timing of the observed PMT hits are compared to a likelihood function operating under a muon hypothesis. This likelihood function predicts hit patterns and timing based on the interaction vertex and the momentum four-vector of the muon. The likelihood function simultaneously varies these seven parameters while comparing to the observed PMT hits. The parameters from the maximized likelihood function yield the reconstructed muon kinematics.

Under the assumption of $\Delta(1232)$ production by a neutrino scattering off a stationary nucleon target in carbon, the neutrino energy is given by

$$E_\nu^\Delta = \frac{2(M_p - E_B)E_\mu - (E_B^2 - 2M_p E_B + m_\mu^2 + \Delta M^2)}{2[(M_p - E_B) - E_\mu + p_\mu \cos\theta_\mu]}, \quad (2)$$

TABLE II. Predicted event composition of the CC1 π^+ sample in antineutrino mode.

Interaction channel	Contribution (%)
$\nu_\mu N \rightarrow \mu^- \pi^+ N$ (resonant)	64
$\nu_\mu A \rightarrow \mu^- \pi^+ A$ (coherent)	7
$\bar{\nu}_\mu N \rightarrow \mu^+ \pi^- N$ (resonant)	6
$\nu_\mu n \rightarrow \mu^- p$	6
$\nu_\mu n \rightarrow \mu^- \pi^0 p$	2
$\bar{\nu}_\mu p \rightarrow \mu^+ \pi^0 n$	1
Other (mostly DIS)	14
“Observable CC1 π^+ ” ($1\mu^-, 1\pi^+$)	82




FIG. 3 (color online). The reconstructed energy spectrum for simulation versus data in the antineutrino-mode $CC1\pi^+$ sample. Simulation is normalized to data, and only statistical errors are shown.

where $E_B = 34$ MeV is the binding energy, m_μ is the muon mass, $\Delta M^2 = M_p^2 - M_\Delta^2$, where M_Δ (M_p) is the $\Delta(1232)$ (proton) mass, p_μ is the muon momentum, and θ_μ is the outgoing muon angle relative to the incoming neutrino beam. Effects not accounted for in the reconstruction include nonresonant pion production, contributions from higher mass Δ resonances, and scattering off the quasifree protons in hydrogen instead of carbon. A shape comparison of reconstructed E_ν^Δ in data and simulation is presented in Fig. 3.

C. Measuring the neutrino flux component in the antineutrino-mode $CC1\pi^+$ sample

The simulation sample is separated into two components: observable $CC1\pi^+$ events and background. All observable $CC1\pi^+$ events in the simulation are modeled using the $CC1\pi^+$ cross section that has been measured in MiniBooNE neutrino-mode data [12]. Given that the majority of the $CC1\pi^+$ sample in antineutrino mode is induced by neutrinos, with this cross section measurement applied any remaining normalization difference between data and simulation is interpreted as a neutrino flux measurement. Results are presented in Table III. Events in the antineutrino-mode $CC1\pi^+$ sample indicate the neutrino flux in antineutrino mode is lower than the simulation predicts. The extracted calibration is 0.76 ± 0.11 of the nominal prediction over all reconstructed energies, while the analysis applied to individual energy ranges does not indicate any significant energy dependence.

D. Systematic errors

The systematic error on the neutrino flux measurement using the antineutrino-mode $CC1\pi^+$ sample comes from two sources that are treated as uncorrelated with each other: the uncertainty on the $CC1\pi^+$ cross section obtained

TABLE III. Antineutrino-mode $CC1\pi^+$ sample details and ν_μ flux component measurement. The measured cross section has been applied to simulation, and the ν_μ flux scale is found by calculating (observed events—expected $\bar{\nu}_\mu$ events)/expected ν_μ events. The reported error is discussed in more detail in Sec. IV D. The Monte Carlo sample is generated so that the associated statistical error is negligible compared to the other sources of uncertainty.

E_ν^Δ Range (MeV)	Mean gen. E_ν (MeV)	Events in data	Expected ν_μ	Expected $\bar{\nu}_\mu$	ν_μ Flux scale
600–700	961	465	556	104	0.65 ± 0.10
700–800	1072	643	666	118	0.79 ± 0.10
800–900	1181	573	586	97	0.81 ± 0.10
900–1000	1285	495	474	78	0.88 ± 0.11
1000–1200	1426	571	646	92	0.74 ± 0.10
1200–2400	1685	521	614	74	0.73 ± 0.15
Inclusive	1266	3268	3542	563	0.76 ± 0.11

from [12] and the uncertainty in the background prediction. The largest contribution to the uncertainty on the $CC1\pi^+$ cross section comes from the neutrino-mode flux uncertainty, which is the only systematic error associated with the cross section measurement that is also independent of the measurement made here. Because the other $CC1\pi^+$ uncertainties are treated as uncorrelated, a partial cancellation of errors is ignored in the present neutrino flux measurement. This results in a slight overestimate of the neutrino flux uncertainty. Both ν_μ and $\bar{\nu}_\mu$ background events in the sample are assigned 30% uncertainties to conservatively recognize the model dependence of the sample composition. The fractional uncertainty contributions to the flux measurement are presented in Table IV.

V. MEASURING THE NEUTRINO FLUX THROUGH MUON ANGULAR DISTRIBUTIONS IN THE CCQE SAMPLE

A. The CCQE sample

The CCQE interaction is the dominant channel in MiniBooNE's energy range. CCQE events typically have

TABLE IV. Fractional uncertainty (%) contributions to the neutrino flux measurement in the $CC1\pi^+$ sample. The ν_μ uncertainty is dominated by the $CC1\pi^+$ cross section error.

E_ν^{QE} Range (MeV)	Statistical	$\bar{\nu}_\mu$	ν_μ	Total fractional error
600–700	6	9	11	15
700–800	5	7	10	13
800–900	5	6	10	13
900–1000	5	6	10	13
1000–1200	5	6	11	13
1200–2400	5	5	19	20
Inclusive	2	6	13	14

two subevents, attributed to the primary muon and the associated decay positron:

$$\begin{aligned} 1: & \bar{\nu}_\mu + p \rightarrow \mu^+ + n \\ 2: & \quad \hookrightarrow e^+ + \nu_e + \bar{\nu}_\mu. \end{aligned} \quad (3)$$

The CCQE sample is therefore similar in formation to the CC1 π^+ sample with one major divergence: a requirement of two subevents instead of three. As shown in Table V, the CCQE selection cuts closely follow those motivated in Sec. IVA, with a few exceptions appropriate to the inclusion of a single Michel electron. The Michel tank hit and veto PMT hit cuts apply to the second subevent only now (Cuts 5 and 6, respectively), and the muon endpoint-electron vertex cut in Cut 7 is tightened to 100 cm in light of larger backgrounds. The selection cuts outlined here are identical to those employed in a previous shape-only extraction of CCQE model parameters [6] and closely follow those used in the absolute measurement of the ν_μ induced CCQE cross section [9], with only minor differences that result in approximately the same sample efficiency and purity.

Despite the selection cuts, there are formidable backgrounds to the antineutrino-mode CCQE sample. Prior to this analysis, simulation estimates the antineutrino-mode CCQE sample has a purity just above 50% as shown in Table VI. The major backgrounds include CC1 π^+ and CC1 π^- events, which account for a total of $\sim 20\%$ of the sample, and the ν_μ processes, predicted to be responsible for $\sim 30\%$ of the sample. The 30% predicted ν_μ contamination is investigated and ultimately constrained in this analysis.

A few additional modifications to the simulation are made to accommodate the backgrounds. The largest non-CCQE background in the sample is single pion production which enters the sample due to nuclear effects, including

TABLE V. Summary of selection cuts with efficiencies in the CCQE sample. ‘‘Purity’’ refers to $\bar{\nu}_\mu$ CCQE only, and purity and efficiency numbers are sequential.

Cut #	Description	Efficiency (%)	Purity (%)
0	No cuts	100	32
1	Two subevents	49	41
2	1st subevent in event time window $4000 < T(\text{ns}) < 7000$	47	42
3	1st subevent: reconstructed vertex <500 cm from tank center	38	43
4	1st subevent: tank hits >200	35	45
5	2nd subevent: tank hits <200	33	45
6	Both subevents: veto hits <6	29	49
7	Distance between reconstructed end of 1st subevent and 2nd subevent vertex <100 cm	25	54

TABLE VI. Predicted composition of the antineutrino-mode CCQE sample.

Channel	Contribution (%)
$\bar{\nu}_\mu p \rightarrow \mu^+ n$	54
$\nu_\mu n \rightarrow \mu^- p$	20
$\bar{\nu}_\mu N \rightarrow \mu^+ \pi^- N$ (resonant)	8
$\nu_\mu N \rightarrow \mu^- \pi^+ N$ (resonant)	6
$\bar{\nu}_\mu A \rightarrow \mu^+ \pi^- A$ (coherent)	4
$\bar{\nu}_\mu N \rightarrow \mu^+ \Lambda, \Sigma$	3
$\bar{\nu}_\mu p \rightarrow \mu^+ \pi^0 n$	2
Other	3
All $\bar{\nu}_\mu$	71
All ν_μ	29

μ^- , π^- capture and final-state interactions; however, in the case of antineutrino-induced CC1 π^- scattering, due to π^- nuclear capture almost 100% of CC1 π^- events have only two subevents and are experimentally indistinguishable from CCQE. This implies a direct background measurement of CC1 π^- events (analogous to what was done in Ref. [9]) is impossible. Therefore, though the CC1 π^+ yield constraint made in Ref. [9] is strictly appropriate to neutrino-induced CC1 π^+ events only, it is applied to both predicted CC1 π^+ and CC1 π^- background events in the CCQE sample.

Many backgrounds to the CCQE sample peak in the most forward scattering region of the muon angular distribution with respect to the incoming neutrino beam. This includes pion production and hydrogen CCQE scattering—while the latter is technically not a background, the proper handling of the difference in nuclear effects between bound and free targets is not straightforward. Additionally, the forward scattering region is strongly correlated with low Q^2 events, a problematic region both experimentally and theoretically [42]. Such low Q^2 data are dominated by $\bar{\nu}_\mu$ interactions, while the present analysis is principally interested in backwards scattering muons which is dominated by ν_μ . For these reasons, events with $\cos\theta_\mu > 0.91$ are not included in the fit to data, where θ_μ is the outgoing muon angle relative to the incoming neutrino beam.

B. CCQE event reconstruction

Event reconstruction in the antineutrino-mode CCQE sample proceeds similarly as in the CC1 π^+ sample, described in Sec. IV B. As in the CC1 π^+ reconstruction, the measurement of muon kinematics from the primary interaction is solely responsible for recreating the incident neutrino energy. No requirement is made on the ejected nucleon; this is an important distinction from the CCQE definitions used by other experiments [43,44], where a single proton track may be required in the case of neutrino-induced CCQE. A similar energy reconstruction as described in Sec. IV B is implemented, but in this sample a $\bar{\nu}_\mu$ probe is assumed:




FIG. 4 (color online). The reconstructed energy spectrum for simulation versus data in the antineutrino-mode CCQE sample. Simulation is normalized to data, and only statistical errors are shown.

$$E_{\bar{\nu}}^{\text{QE}} = \frac{2(M_p - E_B)E_{\mu} - (E_B^2 - 2M_p E_B + m_{\mu}^2 + \Delta M^2)}{2[(M_p - E_B) - E_{\mu} + p_{\mu} \cos\theta_{\mu}]}, \quad (4)$$

where the same definitions from Eq. (2) apply and $\Delta M^2 = M_p^2 - M_n^2$, where M_n is the neutron mass. Figure 4 presents the reconstructed energy distributions in simulation and data in the CCQE sample. CCQE scattering with free protons in hydrogen are indistinguishable from those on bound protons in carbon, so all events in data and simulation are reconstructed using the carbon scattering assumption implicit in Eq. (4).

C. Neutrino flux measurement using CCQE

Neutrino and antineutrino CCQE cross sections differ exclusively by an axial-vector interference term that amplifies ν scattering while suppressing $\bar{\nu}$ events. A particularly clean way to exploit this cross section difference is to fit the angular distribution of the primary muon. The contribution from $\bar{\nu}_{\mu}$ is suppressed in the backward scattering region. Figure 5 shows the predicted ν_{μ} and $\bar{\nu}_{\mu}$ contributions to the cosine of the outgoing muon angle.

To measure the neutrino content in the antineutrino-mode beam, the Monte Carlo sample is separated into two $\cos\theta_{\mu}$ templates, one arising from all ν_{μ} interactions and the other from $\bar{\nu}_{\mu}$, regardless of interaction channel and nuclear target. A linear combination of these two templates is then formed

$$T_{\text{MC}}(\alpha_{\nu}, \alpha_{\bar{\nu}}) \equiv \alpha_{\nu} \nu^{\text{MC}} + \alpha_{\bar{\nu}} \bar{\nu}^{\text{MC}}, \quad (5)$$

where T_{MC} is the total predicted $\cos\theta_{\mu}$ distribution to be compared to data, α_{ν} and $\alpha_{\bar{\nu}}$ are neutrino and antineutrino rate scales, and ν^{MC} and $\bar{\nu}^{\text{MC}}$ are the Monte Carlo neutrino and antineutrino scattering angular predictions, respectively. The modified simulation sample is compared to




FIG. 5 (color online). The $\cos\theta_{\mu}$ distribution of the CCQE sample by neutrino type before fitting. As printed on the figure, 29% of the sample is predicted to be induced by neutrinos. The Monte Carlo sample has been normalized to 5.66×10^{20} protons on target.

data by forming a goodness-of-fit χ^2 test as a function of the rate scales

$$\chi^2 = \sum_{i,j} (T_{\text{MC}}(\alpha_{\nu}, \alpha_{\bar{\nu}})_i - d_i) M_{ij}^{-1} (T_{\text{MC}}(\alpha_{\nu}, \alpha_{\bar{\nu}})_j - d_j), \quad (6)$$

where i and j label bins of $\cos\theta_{\mu}$, d is data and M is the symmetric error matrix given in Eq. (7). The error matrix is used to propagate correlated uncertainties on parameters and processes to the quantities reported in the analysis. It is made by first forming weights corresponding to simulation excursions set by Gaussian variations of parameters within their associated error. The difference of these weighted events from the simulated central value forms the error matrix

$$M_{ij} = \frac{1}{K} \sum_{s=1}^K (N_i^s - N_i^{\text{CV}}) \times (N_j^s - N_j^{\text{CV}}). \quad (7)$$

Here K simulation excursions are used ($K = 100$ in this analysis), N^s is the reweighted number of entries corresponding to the s th simulation set, and N^{CV} represents the MiniBooNE simulation central value. This technique is further described in Ref. [45]. Bin-by-bin $\cos\theta_{\mu}$ correlations between ν_{μ} and $\bar{\nu}_{\mu}$ are also treated. The specific systematic errors are discussed in the next section.

The fit is performed analytically in three bins of reconstructed energy and also in an inclusive energy sample. Results including statistical and systematic uncertainties are presented in Table VII. The fits to data are shown in the Appendix, where Fig. 7 contains both the fitted distributions and the fractional differences between the simulation and data before and after the fits. The adjusted contributions of ν_{μ} and $\bar{\nu}_{\mu}$ to the CCQE sample are compared to the prediction in Table VIII.

The χ^2 value for the angular fit in the reconstructed energy range $E_{\bar{\nu}}^{\text{QE}} > 900$ MeV is unusually low at

TABLE VII. Fit results in three energy bins and an inclusive sample. The results are consistent with an overprediction of the ν_μ contamination of the MiniBooNE antineutrino-mode CCQE sample.

$E_{\bar{\nu}}$ Range (MeV)	Mean generated E_ν (MeV)	Events in data	α_ν fit	$\alpha_{\bar{\nu}}$ fit	$\rho_{\alpha_\nu - \alpha_{\bar{\nu}}}$ fit correlation	χ^2 (DOF = 21)
<600	675	15242	0.65 ± 0.22	0.98 ± 0.18	0.33	13
600–900	897	16598	0.61 ± 0.20	1.05 ± 0.19	0.49	21
>900	1277	15626	0.64 ± 0.20	1.18 ± 0.21	0.45	7
Inclusive	950	47466	0.65 ± 0.23	1.00 ± 0.22	0.25	16

$\chi^2 = 7$ for 21 degrees of freedom. This is believed due simply to chance, as the statistical error only fit agrees with the data exceptionally well within the error, returning $\chi^2 = 13$ for 21 degrees of freedom.

As the ν_μ angular template has been corrected for the observed cross section per Ref. [9], α_ν may be interpreted as a flux scale factor, and significant deviations from unity would imply a flux mismodeling. Consistent with the results reported in Sec. IV C, fits in the antineutrino-mode CCQE sample indicate the true neutrino flux to be somewhat lower than the simulation predicts. Over all reconstructed energies, the neutrino flux component of the antineutrino-mode beam should be scaled by 0.65 to match the observed data. Fits in individual reconstructed energy bins show that the neutrino flux component shape is well-modeled. Finding the calibration on the neutrino flux component inconsistent with unity is not surprising, as the neutrino parent pions originate primarily in a poorly constrained production region (cf. Fig. 1). The rate scale $\alpha_{\bar{\nu}}$ is ambiguous in interpretation, as the cross section is yet unmeasured.

Care must be taken when comparing these results to the μ^+/μ^- yield numbers reported in the MiniBooNE $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillation analysis [4,5], since the interaction prediction is different. In the oscillation analysis the cross section parameters measured in Ref. [6] are employed, which includes $M_A^{\text{eff}} = 1.23(1.13)$ GeV for bound (free) nucleon CCQE scattering and $\kappa = 1.022$. When the muon angular fit technique described in this section is repeated with this prediction, yield rates of $\alpha_\nu = 0.99 \pm 0.23$ and $\alpha_{\bar{\nu}} = 1.20 \pm 0.23$ are found, as reported in Ref. [5]. With this alternate CCQE scattering model, the angular fit over all reconstructed energies reports a neutrino contamination in the sample of $23 \pm 6\%$,

TABLE VIII. Fractional composition of the antineutrino-mode CCQE sample before and after angular fits.

$E_{\bar{\nu}}$ Range (MeV)	Before fit (%)		After fit (%)	
	ν_μ	$\bar{\nu}_\mu$	ν_μ	$\bar{\nu}_\mu$
<600	25	75	18 ± 6	82 ± 16
600–900	26	74	17 ± 6	83 ± 15
>900	35	65	23 ± 7	77 ± 15
Inclusive	29	71	21 ± 8	79 ± 18

consistent with the $21 \pm 8\%$ contamination found with the scattering assumptions described in Sec. III A.

The results from this technique depend on knowing the angular distributions of neutrino and antineutrino CCQE interactions in the detector. While the procedure relies on exploiting the effect of the interference term in the CCQE cross section, the angular distributions may be somewhat altered by nuclear effects. In this analysis the measured angular distribution of neutrino interactions on carbon [9] is employed, but the measurement relies on the scattering model described in Sec. III A to predict antineutrino interactions. This model does not include two-body current effects which may be larger than previously expected [30] and may introduce additional neutrino and antineutrino angular differences. Despite this inherent model dependence, the results present a demonstration of a technique aimed at informing future experiments looking to separately constrain neutrino and antineutrino events in an unmagnetized environment. By that time, the effect of additional nuclear processes on the angular dependence of antineutrino CCQE scattering should be better known.

D. Systematic errors

As the present analysis directly measures the neutrino component in the antineutrino-mode beam, systematic errors relating to beam geometry and meson production




FIG. 6 (color online). Summary of the neutrino flux constraint in the antineutrino-mode beam from the CC1 π^+ (Sec. IV) and CCQE (Sec. V) measurements.

TABLE IX. Summary of systematic error contribution to the scale parameter α_ν in the inclusive energy fit. Individual error contributions are found for the i th systematic error by first repeating the fits with only independent systematics considered. The fractional error contributions are then found by $\sqrt{(\Delta\alpha_\nu/\alpha_\nu)_{\text{syst}_i+\text{stat}}^2 - (\Delta\alpha_\nu/\alpha_\nu)_{\text{stat}}^2}$, where $\Delta\alpha_\nu$ is the one-sigma error reported in Table VII. The statistical error is found by considering the second term only. This method does not account for small changes in the α_ν best fit parameter between the fits considering various errors, and so the individual fractional errors do not add in quadrature to produce the total fractional error reported in Table VII and in the final column.

Source of error	Fractional uncertainty (%)
Statistical	8
Detector modeling	11
CC1 π^+ Constraint	4
Cross section	26
Total fractional error	35

at the target are not considered. The remaining systematic errors include those arising from detector modeling, the single pion production background, and the cross section parameters in the underlying model. Contributions propagated from these errors to the uncertainty on the parameter α_ν in the inclusive energy sample are given in Table IX.

Apart from final-state interaction uncertainties leading to errors on the cross section, the error on the CC1 π^+ background contributes to the systematic error through the error labeled ‘‘CC1 π^+ Constraint’’ in Table IX. This measurement uncertainty is based on a Q^2 -dependent shape-only scale factor to improve data-simulation agreement in the neutrino-mode CC1 π^+ sample [3]. The cross section (both CCQE and CC1 π^+) uncertainty is dominant in these fits and warrants further discussion. Table X offers a breakdown of cross section parameters and associated errors. The error on carbon M_A^{eff} has been reduced from that reported in Ref. [9] to avoid double-counting MiniBooNE systematic errors applicable to both the measurement of M_A^{eff} and the measurement reported here. The 26% uncertainty due to cross section errors reported in Table IX can be expanded as the quadrature sum of 16% from the 10% normalization errors on $\bar{\nu}_\mu$ and CCQE processes, 14% from the error on M_A and κ , and 15% from the remaining.

As the main contributions to the dominant cross section systematic error apply to both ν_μ and $\bar{\nu}_\mu$ scattering, α_ν and $\alpha_{\bar{\nu}}$ are positively correlated as reported in Table VII.

VI. RESULT COMPARISON

Including all reconstructed energies in the CC1 π^+ sample, a neutrino flux component scale of 0.76 ± 0.11 is found, while the CCQE analysis yields 0.65 ± 0.23 . The measurements are compatible and complementary as each analysis includes energy regions not covered by the other

TABLE X. Summary of cross section errors used in this analysis. The bottom portion presents fractional uncertainties assigned to processes *in addition* to parameter errors. Errors given on pion absorption and charge exchange are relevant to pion propagation in the detector medium.

Parameter	Value with error
M_A^{eff} carbon target	1.35 ± 0.07 GeV
M_A^{eff} hydrogen target	1.03 ± 0.02 GeV
κ	1.007 ± 0.005
E_B	34 ± 9 MeV
Δs	0.0 ± 0.1
$M_A^{1\pi}$	1.10 ± 0.28 GeV
$M_A^{\text{multi-}\pi}$	1.30 ± 0.52 GeV
pF	220 ± 30 MeV
Process	Fractional uncertainty (%)
π^+ Charge exchange	50
π^+ Absorption	35
CCQE σ Normalization	10
All $\bar{\nu}_\mu$ σ Normalization	10
$\Delta + N \rightarrow N + N$	100

as shown in Fig. 6. The results indicate the simulated neutrino component of the antineutrino-mode flux is over-estimated by $\sim 30\%$. These flux measurements constrain the very forward π^+ created at the target, where an external data constraint is not available. Results from both methods are summarized in Fig. 6, where measurements are placed at the mean of the generated energy distribution for each reconstructed energy sample.

VII. IMPLICATIONS FOR OTHER EXPERIMENTS

The techniques applied here could also aid future neutrino experiments that will test for CP violation in the lepton sector using large unmagnetized detectors. This includes experiments such as NO ν A [46], T2K [47], LBNE [48], LAGUNA [49], and Hyper-K [50]. A magnetized near detector can provide a powerful constraint on the neutrino flux and provide precise cross sections. However, a measurement of the neutrino rate at the far detector can still be very useful given that the the beam spreads from an extended source and oscillates while traveling between the detectors.

Additional techniques could offer potentially helpful constraints on the neutrino component in an antineutrino-mode beam. This includes taking advantage of the effective lifetime difference between μ^-/μ^+ due to μ^- capture in a nuclear environment. Fitting the lifetime distributions or measuring how often a decay electron is produced could supply constraints that are especially useful as they are independent of the underlying neutrino interaction cross sections. Also, selection of CCQE interactions with and without a proton in the final state may afford additional neutrino versus antineutrino tagging capabilities [51,52].

VIII. CONCLUSIONS

Two analyses are presented to measure the neutrino flux in the MiniBooNE antineutrino-mode beam. The two measurements have a common dependence on the neutrino flux in the neutrino-mode beam [20] that MiniBooNE obtained from HARP hadroproduction data. At present the CCQE angular distribution method is largely limited by

uncertainties in the cross sections, especially the antineutrino cross section which MiniBooNE is in the process of measuring, while the uncertainty of the CC1 π method is dominated by the neutrino-mode flux uncertainty.

Using two event samples dominated by independent physics processes, compatible and complementary results are found. The results from both analyses indicate the




FIG. 7 (color online). Results of the muon angular fits to the CCQE data described in Sec. V. Shown are (a) the fits and (b) fractional differences (data—simulation)/simulation for both the unmodified prediction and the best fit. Along with an inclusive sample, three reconstructed energy bins are considered. The before-fit simulation is absolutely normalized to 5.66×10^{20} protons on target. Only events with $\cos \theta_\mu < 0.91$ are considered.

prediction of the neutrino flux component of the antineutrino beam is overestimated—the CC1 π^+ analysis (Sec. IV) indicate the predicted ν_μ flux should be scaled by 0.76 ± 0.11 , while the CCQE angular fits (Sec. V) yield 0.65 ± 0.23 . Results from repeating the analyses in bins of reconstructed neutrino and antineutrino energy indicate that the predicted flux spectrum shape is well-modeled. The results from fitting the muon angular distributions in the CCQE sample has already been employed in the MiniBooNE oscillation analysis [4,5], while the CC1 π^+ -based measurement will likely be more valuable to MiniBooNE antineutrino cross section extractions, as it is much less model-dependent and carries comparatively smaller uncertainty. These types of analyses, along with others discussed in Sec. VII may be of use to present and

future precision neutrino experiments testing CP violation with neutrino and antineutrino beams.

ACKNOWLEDGMENTS

We wish to acknowledge the support of Fermilab, the National Science Foundation, and the Department of Energy in the construction, operation, and data analysis of the MiniBooNE experiment.

APPENDIX A: CCQE ANGULAR FIT DETAILS

This appendix presents details on the CCQE angular fit results described in Sec. V. The fits to data are plotted in Fig. 7.

-
- [1] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **98**, 231801 (2007).
- [2] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **102**, 101802 (2009).
- [3] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **103**, 061802 (2009).
- [4] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **103**, 111801 (2009).
- [5] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **105**, 181801 (2010).
- [6] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **100**, 032301 (2008).
- [7] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **103**, 081801 (2009).
- [8] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **81**, 013005 (2010).
- [9] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **81**, 092005 (2010).
- [10] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **82**, 092005 (2010).
- [11] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), [arXiv:1010.3264](https://arxiv.org/abs/1010.3264).
- [12] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **83**, 052007 (2011).
- [13] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Lett. B* **664**, 41 (2008).
- [14] M. G. Catanesi *et al.* (HARP Collaboration), *Eur. Phys. J. C* **52**, 29 (2007).
- [15] M. G. Catanesi *et al.* (HARP Collaboration), *Nucl. Phys. B* **732**, 1 (2006).
- [16] D. Casper, *Nucl. Phys. B, Proc. Suppl.* **112**, 161 (2002).
- [17] M. O. Wascko (MiniBooNE Collaboration), [arXiv:0602051](https://arxiv.org/abs/0602051).
- [18] P. Huber and T. Schwetz, [arXiv:0805.2019](https://arxiv.org/abs/0805.2019).
- [19] S. Agostinelli *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [20] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **79**, 072002 (2009).
- [21] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **599**, 28 (2009).
- [22] CERN Program Library Long Writeup W5013, 1993, <http://wwwasdoc.web.cern.ch/wwwasdoc/geant/geantall.html>.
- [23] C. Zeitnitz and T. A. Gabriel, *Nucl. Instrum. Methods Phys. Res., Sect. A* **349**, 106 (1994).
- [24] D. Ashery *et al.*, *Phys. Rev. C* **23**, 2173 (1981); M. K. Jones *et al.*, *Phys. Rev. C* **48**, 2800 (1993); R. D. Ransome *et al.*, *Phys. Rev. C* **45**, R509 (1992).
- [25] B. C. Brown *et al.*, in *2004 IEEE Nuclear Science Symposium Conference Record*, Vol. 1, p. 652 (2006).
- [26] R. A. Smith and E. J. Moniz, *Nucl. Phys.* **B43**, 605 (1972); **B101**, 547(E) (1975).
- [27] H. Budd, A. Bodek, and J. Arrington, [arXiv:hep-ex/0308005](https://arxiv.org/abs/hep-ex/0308005) [*Nucl. Phys. B, Proc. Suppl.* (to be published)].
- [28] J. Grange, in *6th International Workshop on Neutrino-Nucleus Interactions in the Few-GeV Region (NUI NT 2009)*, edited by F. Sanchez *et al.*, AIP Conf. Proc. No. 1189, (AIP, New York, 2009), 331.
- [29] V. Bernard *et al.*, *J. Phys. G* **28**, R1 (2002).
- [30] M. Martini *et al.*, *Phys. Rev. C* **80**, 065501 (2009).
- [31] M. Martini *et al.*, *Phys. Rev. C* **81**, 045502 (2010).
- [32] J. Nieves *et al.*, *Phys. Rev. C* **83**, 045501 (2011).
- [33] J. Nieves *et al.*, [arXiv:1106.5374](https://arxiv.org/abs/1106.5374).
- [34] J. E. Amaro *et al.*, *Phys. Rev. D* **84**, 033004 (2011).
- [35] D. Rein and L. M. Sehgal, *Ann. Phys. (N.Y.)* **133**, 79 (1981).
- [36] R. P. Feynman, M. Kislinger, and F. Ravndal, *Phys. Rev. D* **3**, 2706 (1971).
- [37] D. Casper (private communication).
- [38] W. Reuter *et al.*, *Phys. Rev. C* **26**, 806 (1982).
- [39] V. Flaminio *et al.*, CERN-HERA-83-01, 1983; CERN-HERA-83-02, 1983; CERN-HERA-84-01, 1984.
- [40] T. Suzuki *et al.*, *Phys. Rev. C* **35**, 2212 (1987).
- [41] A. Shinohara *et al.*, *Phys. Rev. A* **53**, 130 (1996).
- [42] A. M. Ankowski *et al.*, [arXiv:1001.0481](https://arxiv.org/abs/1001.0481).

- [43] J.L. Alcaraz-Aunión and J. Walding, *AIP Conf. Proc.* **1189**, 145 (2009).
- [44] K. S. Kuzmin, V. V. Lyubushkin, and V. A. Naumov, *Eur. Phys. J. C* **54**, 517 (2008); V. V. Lyubushkin *et al.* (NOMAD Collaboration), *Eur. Phys. J. C* **63**, 355 (2009).
- [45] B. P. Roe, *Nucl. Instrum. Methods Phys. Res., Sect. A* **570**, 159 (2007).
- [46] D. S. Ayres *et al.* (NOvA Collaboration), FERMILAB-DESIGN-2007-01.
- [47] Y. Itow *et al.*, [arXiv:hep-ex/0106019](https://arxiv.org/abs/hep-ex/0106019).
- [48] V. Barger *et al.*, Report of the US long baseline neutrino experiment study, <http://nwg.phy.bnl.gov/fnal-bnl>.
- [49] A. de Bellefon *et al.*, [arXiv:hep-ex/0607026](https://arxiv.org/abs/hep-ex/0607026).
- [50] K. Nakamura, *Int. J. Mod. Phys. A* **18**, 4053 (2003).
- [51] A. A. Aguilar-Arevalo *et al.*, [arXiv:hep-ex/0601022](https://arxiv.org/abs/hep-ex/0601022).
- [52] M. Fechner and C. W. Walter, *J. High Energy Phys.* **11** (2009) 040.