

RKKY interaction between adsorbed magnetic impurities in graphene: Symmetry and strain effects - DTU Orbit (09/11/2017)

RKKY interaction between adsorbed magnetic impurities in graphene: Symmetry and strain effects

The growing interest in carbon-based spintronics has stimulated a number of recent theoretical studies on the Ruderman-Kittel-Kasuya-Yosida (RKKY) interaction in graphene, with the aim of determining the most energetically favorable alignments between embedded magnetic moments. The RKKY interaction in undoped graphene decays faster than expected for conventional two-dimensional materials, and recent studies suggest that the adsorption configurations favored by many transition-metal impurities may lead to even shorter-ranged decays and possible sign-changing oscillations. Here, we show that these features emerge in a mathematically transparent manner when the symmetry of the configurations is included in the calculation. Furthermore, we show that by breaking the symmetry of the graphene lattice, via uniaxial strain, the decay rate, and hence the range, of the RKKY interaction can be significantly altered. Our results suggest that magnetic interactions between adsorbed impurities in graphene can be manipulated by careful strain engineering of such systems.

General information

State: Published

Organisations: Department of Micro- and Nanotechnology, Theoretical Nanotechnology, Center for Nanostructured

Graphene, Trinity College Dublin

Authors: Gorman, P. D. (Ekstern), Duffy, J. M. (Ekstern), Ferreira, M. (Forskerdatabase), Power, S. (Intern)

Number of pages: 11 Publication date: 2013

Main Research Area: Technical/natural sciences

Publication information

Journal: Physical Review B

Volume: 88 Issue number: 8 ISSN (Print): 1098-0121

Ratings:

BFI (2017): BFI-level 2

Web of Science (2017): Indexed yes

BFI (2016): BFI-level 2

Scopus rating (2016): CiteScore 3.16 Web of Science (2016): Indexed yes

BFI (2015): BFI-level 2

Scopus rating (2015): SJR 1.933 SNIP 0.94 CiteScore 2.8

Web of Science (2015): Indexed yes

BFI (2014): BFI-level 2

Scopus rating (2014): SJR 2.667 SNIP 1.262 CiteScore 3.3

Web of Science (2014): Indexed yes

BFI (2013): BFI-level 2

Scopus rating (2013): SJR 2.785 SNIP 1.339 CiteScore 3.55

ISI indexed (2013): ISI indexed yes Web of Science (2013): Indexed yes

BFI (2012): BFI-level 2

Scopus rating (2012): SJR 3.206 SNIP 1.394 CiteScore 3.57

ISI indexed (2012): ISI indexed yes Web of Science (2012): Indexed yes

BFI (2011): BFI-level 2

Scopus rating (2011): SJR 3.382 SNIP 1.438 CiteScore 3.61

ISI indexed (2011): ISI indexed yes Web of Science (2011): Indexed yes

BFI (2010): BFI-level 2

Scopus rating (2010): SJR 3.417 SNIP 1.451

Web of Science (2010): Indexed yes

BFI (2009): BFI-level 2

Scopus rating (2009): SJR 3.109 SNIP 1.474

Web of Science (2009): Indexed yes

BFI (2008): BFI-level 1

Scopus rating (2008): SJR 2.982 SNIP 1.524

Web of Science (2008): Indexed yes

Scopus rating (2007): SJR 2.923 SNIP 1.546

Web of Science (2007): Indexed yes

Scopus rating (2006): SJR 2.796 SNIP 1.56

Web of Science (2006): Indexed yes

Scopus rating (2005): SJR 2.763 SNIP 1.607

Web of Science (2005): Indexed yes

Scopus rating (2004): SJR 2.742 SNIP 1.606

Web of Science (2004): Indexed yes

Scopus rating (2003): SJR 2.75 SNIP 1.536

Web of Science (2003): Indexed yes

Scopus rating (2002): SJR 2.788 SNIP 1.706

Web of Science (2002): Indexed yes

Scopus rating (2001): SJR 2.946 SNIP 1.635

Web of Science (2001): Indexed yes

Scopus rating (2000): SJR 2.986 SNIP 1.631

Web of Science (2000): Indexed yes

Scopus rating (1999): SJR 3.115 SNIP 1.58

Original language: English

Electronic versions:

 $RKKY_interaction_between_adsorbed_magnetic_impurities_in_graphene.pdf$

DOIs:

10.1103/PhysRevB.88.085405

Source: dtu

Source-ID: n::oai:DTIC-ART:inspec/391079532::31433

Publication: Research - peer-review > Journal article - Annual report year: 2012