
OCCURENCE OF THE INTRODUCED BLACK RAT
(*RATTUS RATTUS*) AND ITS POTENTIAL EFFECTS ON
ENDEMIC RODENTS IN SOUTHEASTERN MADAGASCAR

JUKKA T. LEHTONEN

Department of Biosciences
Faculty of Biological and Environmental Sciences
University of Helsinki
Finland

Academic dissertation

To be presented for public examination with the permission of the Faculty of Biological and Environmental Sciences of the University of Helsinki, in the Lecture Room 1 in the C-building (Latokartanonkaari 5), on October 18th 2013, at 12 o'clock noon

Supervised by: Prof. Jari Niemelä,
Department of Environmental Sciences
University of Helsinki, Finland

Reviewed by: Prof. Jörg Ganzhorn
Institute of Zoology
University of Hamburg, Germany

Doc. Otso Huitu
Finnish Forest Research Institute, Finland

Examined by: Prof. Herwig Leirs
Department of Biology
University of Antwerp, Belgium

Custos: Prof. Liselotte Sundström
Department of Biosciences
University of Helsinki, Finland

© Jukka T. Lehtonen (Summary)
© Kluwer Academic Publishers (Chapter I)
© Authors (Chapters II, III)
© Estonian Academy Publishers (Chapter IV)

Layout © Petri Clusius

ISBN 978-952-10-9307-4 (paperback)

ISBN 978-952-10-9308-1 (PDF)

<http://ethesis.helsinki.fi>

Unigrafia, Helsinki 2013

CONTENTS

ABSTRACT.	5
SUMMARY.	7
1. Introduction	8
2. Aims of thesis	13
3. Material and methods.	14
4. Main results and discussion	20
5. Final remark and future perspectives	26
6. Acknowledgements	27
7. References	29
CHAPTER I	37
CHAPTER II.	59
CHAPTER III	81
CHAPTER IV	103

LIST OF ORIGINAL PUBLICATIONS

The thesis is based on the following articles which are referred to in the text by their Roman numerals:

- I Lehtonen, J.T., Mustonen, O., Ramiarinjanahary, H., Niemelä, J & Rita, H. 2001: Habitat use by endemic and introduced rodents along gradient of forest disturbance in Madagascar – *Biodiversity and Conservation* 10 (7):1185–1202
- II Lehtonen, J.T., Rasataharilala, V., Dehgan, A. & Niemelä, J.: Effects of rainforest fragmentation on rodents in southeastern Madagascar
- III Rita, H. & Lehtonen, J.T. Explanatory framework (EFRA): a non-statistical statistical tool for ecologists – Case: habitat use of black rat in Madagascar
- IV Miljutin, A. & Lehtonen, J.T. 2008: Probability of competition between the introduced and endemic rodents in Madagascar: an estimation based on morphological traits. – *Estonian Journal of Ecology* 57 (2): 133.52.

Contributions

The following table shows the major contributions of authors to the original articles or manuscripts (referred to as I-IV)

	I	II	III	IV
Original idea	OM, JN, JTL	AD, JTL, JN	HRI, JTL	JTL, AM
Study design	JTL, OM	JTL, AD	JTL, HRI	AM
Data collecting	JTL, HRA, OM	JTL, VR	JTL, VR	AM, JTL
Analysis	JTL, HRI	JTL	JTL, HRI	AM
Manuscript preparation	JTL, JN, HRI	JTL, JN, AD	HRI, JTL	AM, JTL

AD: Alexander Dehgan, JTL: Jukka T. Lehtonen, AM: Andrei Miljutin, OM: Olli Mustonen, JN: Jari Niemelä, VR: Volatiana Rasataharilala, Hra: Haingotiana Ramiarinjanahary, HRI: Hannu Rita

ABSTRACT

Native faunas and floras are especially susceptible to negative effects of invasive alien species in islands. The world's fourth largest island, Madagascar, has very unique biota with high level of endemism. The black rat, *Rattus rattus*, is claimed to cause more extinctions of insular vertebrates than any other introduced rodent in the world. On Madagascar, *R. rattus* is suggested to compete with native rodents belonging to the endemic subfamily Nesomyinae. Extensive deforestation and fragmentation also threatens Malagasy forest-dwelling species.

The aims of this thesis were to 1) study the occurrence of native and introduced species and the different kind of factors determining occurrence in southeastern Madagascar and 2) help to target future studies by estimating which native rodents are the most potential competitors with introduced rodents on Madagascar.

Habitat use data of black rat and endemic rodents were collected both in fragmented and in unfragmented forest in Ranomafana National Park in southeastern Madagascar. A total of 698 rodent individuals were captured in 6204 trap nights. Logistic and Poisson regression models were used to determine the factors that influence the presence and abundance of rodent species and to investigate how sensitive one variable can be to other variables in regression models. This led to the introduction of a new approach called 'explanatory framework based regression analysis'(EFRA) which rests on the social science –based elaboration technique. EFRA enables systematization of the link between ecological knowledge and statistical analysis. From the point of it multicollinearity is more source of information than a problem for data analysis.

The abundance of *R. rattus* increased with increasing forest disturbance. The spread of *R. rattus* was suggested to be associated with deforestation but not directly with fragmentation. It is not surprising when remembering that *R. rattus* utilizes open areas too. The measured value of the size of the fragment can be viewed as an index capturing certain features of the forest fragment.

For Malagasy native rodents, clear-cutting is disastrous and forest fragmentation may have a diminishing effect on populations of *Nesomys audeberti*. However, none of the native species was more abundant in un-logged than in selectively logged forest. There was no evidence that *Eliurus webbi* suffers directly from forest fragmentation but it may be more susceptible to interactions with *R. rattus* in fragments than in continuous forest.

In the last paper, the comparison of morphological measurements leads to the conclusion that there is a high probability of competition between introduced *R. rattus* and the following native taxa: all species of *Nesomys*, larger semiarboreal *Eliurus* species (e.g. *E. tanala*, *E. webbi*), and especially *Gymnuromys roberti*.

SUMMARY

SUMMARY

1. Introduction

1.1 Invasive alien species

It is widely reported that invasive alien species are one of the most important threats to species extinctions worldwide (Vitousek et al. 1996, 1997; Wilcove et al. 1998, Mack et al. 2000, Clavero & Garcia-Berthou 2005) and particularly on islands (Sax & Gaines 2008, Fordman & Brook 2010). Although it has been criticized that many researches supporting invasion as a cause of extinctions are speculative and based upon limited data (Gurevitch & Padilla 2004), there are several examples where invasive species have driven local species to extinctions or in other ways outstandingly altered ecosystems. One of the best known examples is Guam's island where the accidental introduction of the brown treesnake around 1950 has caused the extinction or dramatic population decline of ten of 13 native bird species, nine of twelve reptile species, and all three mammal species (Fritts & Rodda 1998).

Invasive alien animal species can drive other species to extinction in three ways: 1) by direct competition, 2) by predation, and 3) by spreading parasites or pathogens. There are surprisingly few instances in which indigenous species have been driven to extinction by direct competition with invasive species (Davis 2003). Rather, numerous extinctions or replacement of long-term resident species have been suggested to be caused by the introduction of new parasites and pathogens (McCallum and Dobson 1995, Combes 1996, Daszak et al. 2000, Davis 2003, Tompkins et al. 2003) or predators (e.g. Savidge 1987, Fritts & Rodda 1998, Thibault et al. 2002). Gurevitch & Padilla (2004) examined the IUCN Red List database and noticed that 37 animal species were threatened by direct effects of competition, 58 species by pathogens or parasites, and 113 species by predation.

According to the enemy release hypothesis, invaders can get an advantage over their competitors when they lose a considerable part their enemies such as parasites, pathogens, predators, and herbivores in their new location (Keaney & Crawley 2002, Mitchell & Power 2003, Torchin et al. 2003, Prenter et al. 2004, Drake 2005). Torchin et al. (2003) analysed parasitological studies of 26 invasive animal species from seven taxa and demonstrated that the number of parasite species found in exotic populations is roughly half that found in their native populations. However, introduced species may also get harmful parasites from native species.

Most extinctions facilitated by invasive alien species have been on islands as opposed to mainland (Sax & Gaines 2008). Island native faunas and floras are especially susceptible to negative effects of exotic species because of low species richness, isolation, evolution under relatively stable conditions, and decreased competitive ability and predator awareness (Williamson 1989). Forest disturbance may also facilitate the invasion of undesirable non-native species (e.g. Diamond et al. 1987, Kesavaraju et al. 2008, Corlett 2010, Rickart et al. 2011). Logging may cause very long-lasting effects. In Madagascar logged rainforest areas never recovered native plant species diversity because of the dominance of nonnative plant species (Brown and Gurevitch 2004).

1.2. Introduced rodents on islands

Several rodent species have been introduced on islands, many of which have negative effects on other species (Simberloff 2008). The four species that are most commonly implicated to cause negative ecosystems impacts include the house mouse (*Mus musculus*), the black rat (*Rattus rattus*), Norway rat (*R. norvegicus*), and the Pacific rat (*R. exulans*) (Townsend et al. 2006, Harris 2009). All of these except *R. exulans* is present on Madagascar (Duplantier & Duchemin 2003). Three species of rats have occupied 90 % of the world's major islands (Oppel et al. 2011). Townsend et al. (2006) reviewed the global literature on the effects of introduced rats on island flora and fauna and found that black rats and Norway rats affected mostly seabird and landbird species and were the only species which also affected mammals. However, the fact that introduced rodents are known to cause more negative effects on insular birds than on rodents is partly because insular birds have been studied wider than rodents (Harris 2009).

Black rats are claimed to cause the largest number of declines or extinctions of insular vertebrate species (60), including five species of rodents (Townsend et al. 2006). Harris (2009) found three rodent species more and suggested the reason that *Rattus rattus* is considered in the majority of conservation problems is due to its extensive distribution and ecological plasticity. As most of the eight extinction cases of rodents happened more than hundred years ago, they are not properly studied and an extinction mechanism is not proven.

According to Harris (2009) the *Rattus rattus* –caused extinction mechanism is best supported in Christmas Island where two rodent species disappeared within ten years after the introduction of a blood parasite trypanosome infected *R. rattus*; heavy trypanosome infection was also recorded in native rodents (Pickering & Norris 1996). In Madagascar, only few trypanosomes were found in native *Nesomys rufus* rodents and they differed morphologically from those found in *Rattus rattus* (Laakkonen et al. 2003a, 2003b). However, it was unclear if the few trypanosomes found in Malagasy rodents were spread by *Rattus* and whether a lethality effect was the reason for the lack of *Trypanosoma lewisi* –like trypanomastigotes in Malagasy rodents.

In most other extinction cases, the loss of native species has quickly followed the introduction of *R. rattus* but an extinction mechanism is more or less speculative. The

disappearance of two rodent species, *Rhagamys orthodon* in Corsica and Sardinia islands (Wilson 1991) and *Oryzomys nelson* in Mexico (Vigne and Valladas 1996), has been suggested to be caused by competition with *R. rattus*. However, it is not very probably that the extinction can be explained by a single factor. Vigne and Valladas (1996) discussed that *R. rattus* may increase sensitivity of native rodent species to environmental change.

1.3. Effects of human disturbance on Malagasy biota

Madagascar is the fourth largest island in the world and separated from the African continent by the 400 km wide Mozambique Channel. It isolated from the African continent about 160 million years ago and from India about 80 million years ago (de Wit 2003).

Biota of Madagascar is very unique as species-level endemism reaches 82 % in vascular plants (Callmander et al. 2011). For animals the proportion is usually even higher, for example 96 % in dung beetles (Wirta 2009). All non-introduced land mammals are endemic (Goodman et al. 2003). Madagascar is one of the world's hotspot areas with 3.2 % and 2.8 % of the total global endemic plants and vertebrates, respectively (Myers et al. 2000, de Wit 2003). Almost 50 % of the world's chameleon species are seen only on Madagascar. Typical to Malagasy fauna is also a lack of many taxa. There are no native canids, felids, and ungulates except for three extinct pygmy hippopotami species.

Human colonization history on Madagascar is a disputable topic. Gommery et al. (2011) suggested that the cut marks of hippopotamus bones dated to approximately 2000 BC is the oldest evidence of human presence on Madagascar. However, according to geographical screening of Malagasy and Indonesian human genetic variation, Madagascar was settled approximately 1200 years ago (Cox et al. 2013). Regardless of whether the moment of human colonization was before or after the Common Era, it has been followed by several species extinction: 17 species of large-bodied lemurs, 3 species of hippopotami, the elephant bird, and giant tortoises (Burney et al. 2004). However, the causes of these extinctions are complex, and the role of humans in the extinction process is probable but not proven.

The forest decline is unambiguously human-mediated on Madagascar. The most important reason for deforestation on the island is charcoal production and the needs of agriculture, which usually implies a slash and burn -culture (Green & Sussman 1990). In Madagascar, slash and burn culture is called 'tavy' and is illegal. Estimates of deforestation indicate that 50–80 % of primary vegetation has already been lost on Madagascar (Green & Sussman 1990). The highest estimates are opposed with the argument that originally the highland areas were covered by wooded grassland instead of continuous forest (Burney 1997). Puhakka (2012) studied the tropical rainforest corridor between two national parks, Ranomafana NP and Andringitra NP, in south-

eastern Madagascar and found that the deforestation rate in the area varied between 1.2 % (1990–2000) and 1.5 % (2000–2005).

Deforestation can assist the spreading of invasive species, too. In Ranomafana NP, the colonization of invasive plant species in forests clear-cut or selectively logged in the past prevented return of native plant species even after 150 years (Brown & Gurevitch 2004).

Forest fragmentation is also an acute problem in Madagascar. In 2000, almost half of Malagasy forest existed in fragments less than 500 km² (Harper et al. 2007). This means that more and more animals and plants are living close to forest edge as over 80 % of forested area was less than one kilometre from the edge of forest (Harper et al. 2007). After reviewing the response of several different plant and animal taxa to anthropogenic disturbance, Irwin et al. (2010) concluded that forest disturbance usually decreases endemic species diversity in Madagascar, but different species and other taxa have very divergent reactions.

1.4. Native rodents of Madagascar

All 27 species (9 genera) of Malagasy native rodents belong to the same endemic subfamily Nesomyinae (Soarimalala & Goodman 2011). Based on a phylogenetic approach, Jansa et al. (1999) suggested that Madagascar was colonized by nesomyines via a single invasion; nesomyines later dispersed from Madagascar to Africa. However, the placement of nesomyines in murid rodent evolutionary history has afterward been questioned by other molecular studies (Steppan et al. 2004, Jansa & Weksler 2004), and their monophyletic origin is not clearly confirmed or rejected.

There is a great variety in size among the species of nesomyine rodents. The body weight varies in Nesomyine from 20 to 1300 g, and the head and body length vary from 80 to 345 mm (Carleton, Goodman, 1998; Garbutt, 2007). The contemporary fauna of Nesomyinae contains one very large species (*Hypogeomys antimena*), four small species with the head and body length less or around 100 mm (*Eliurus minor*, *Macrotarsomys bastardi*, *Monticolomys koopmani*, *Voalavo gymnocaudatus*), and 19 species of medium size with head and body length in adults between 120–250 mm. The size of introduced rodents are comparable with the two last groups. *Mus musculus* is about the size of small nesomyines, while two *Rattus* species are comparable with larger medium sized nesomyines. Therefore, most of the Malagasy rodent species are about the same size as introduced ones.

The most species-rich genus among Malagasy rodents is tuft-tailed rats, *Eliurus*, including twelve nocturnal and arboreal species (Goodman et al. 2009). Representatives of *Eliurus* have been found in all natural forest types in Madagascar (Goodman et al. 2003). Red forest rats, *Nesomys*, are the only diurnal rodents in Madagascar. All three species of *Nesomys* are terrestrial. Two of these species occur in the eastern rainforest of Madagascar. *N. audeberti* prefers lower elevations than *N. rufus* (Ryan et al. 1993; Goodman & Carleton 1999). Altitude is known to affect the presence and abundance

of Malagasy rodent species (Goodman et. al. 1996; Goodman and Carleton 1996, 1998; Carleton & Goodman 2000), but other factors influencing their habitat use have been analysed in more detail by a few studies only (Stephenson 1995, Ramanamanjato & Ganzhorn 2001, Ganzhorn 2003).

The subfossils of Malagasy small mammal fauna is very poorly studied and only three nesomyine rodent species are known to be extinct so far: *Brachytarsomys mahajambaensis*, *Hypogeomys australis*, and *Nesomys narindaensis* (Goodman & Rakotondravony 1996, Mein et al. 2010). They all have been recovered in upper pleistocene or Holocene deposits. Therefore, the fossil data does not give reason for speculation of possible nesomyine extinction caused by *Rattus rattus*.

Only a few studies have addressed the effects of fragmentation and deforestation on the eastern small mammal fauna in Madagascar. Research conducted in the montane humid forests of the Central High Plateau demonstrated that endemic rodents were rare at same time when the introduced *R. rattus* was abundant in forest fragments (Ganzhorn et al. 2003). In littoral forests of southeastern Madagascar, Ramanamanjato and Ganzhorn (2001) found that the endemic rodent *Eliurus webbi* was sensitive to fragmentation, but they found no effects in its association with *R. rattus*.

1.5. Introduced rodents on Madagascar

Contrary to many other islands, not many invasive alien animal species have occupied Madagascar. If domestic animals are not included, eight introduced mammal species are living in Madagascar (Garbutt 2007). Three of these are rodents: *Mus musculus*, *Rattus norvegicus*, and *Rattus rattus*.

The first archaeological record for *Rattus* on Madagascar was found in the north-west part of the island from the 11th-14th century (Rakotozafy 1996, Radimihilahy 1997). However, according to mitochondrial DNA analyses, the colonization of *Rattus rattus* probably happened much earlier but not prior to human arrival (Tollenaere et al. 2010). The sequences of Malagasy *Rattus rattus* were most closely related to the sequences found in Oman (Tollenaere et al. 2010), suggesting that the Arabian Peninsula is the strongest candidate for the origin of *R. rattus* colonization on Madagascar. Tollenaere's et al. (2010) observations support the hypothesis of a single colonization of Madagascar. The population of *R. rattus* seems to have expanded rapidly first in coastal areas and later in the central highlands.

Now *Rattus rattus* has invaded most of Madagascar (Goodman 1995). In many study sites in Madagascar (e.g. in Andringitra Strict Nature Reserve), *R. rattus* has not only been found close to human habitation but also deep within undisturbed forests (Goodman & Carleton 1996). Goodman (1995) has suggested that the invasion of *R. rattus* may threaten endemic rodent species on Madagascar. This is supported by two observations: 1) there is an overlap in food composition of *R. rattus* and some nesomyine rodents (Goodman and Sterling 1996), and 2) the abundance of *R. rattus* at certain sites is much higher than that of the native rodents (Stephenson 1993,

Goodman et al. 1996, 1997). The biggest proportion of *R. rattus* has been observed in the humid forest of Montagne d'Ambre at 1350 meters in elevation where all 104 captured rodents, with the exception of one individual, were *Rattus rattus* during 750 trap-nights (Goodman et al. 1996) even though this altitude is where native rodent species richness is generally high in Madagascar. However, Ganzhorn (2003) couldn't find evidence for competition between *R. rattus* and native small mammal fauna in dry deciduous forest in Western Madagascar. No experimental studies testing the competition hypothesis have been done in Madagascar.

The origin of Norway rat, *Rattus norvegicus*, is in southeastern Siberia and northern China and therefore it is not very common in tropical countries (Musser & Carleton 1993). In Madagascar, *R. norvegicus* occurs only in towns of central highlands and harbours but some single individuals have also been captured elsewhere (Duplantier & Duchemin 2003). A mitochondrial study showed that the house mouse in Madagascar originated from the Arabian Peninsula in a single colonization (Duplantier et al. 2002). The house mouse is found in almost all kinds of habitats except for closed forests (Langrand & Goodman 1997; Rakotondravony & Randrianjafy 1998; I).

2. Aims of thesis

It was necessary to obtain more information about the ecology of *Rattus rattus* on Madagascar because: 1) *Rattus rattus* is claimed to have caused more extinctions of insular vertebrate species than any other introduced rodent in the world (Townsend et al. 2006), 2) black rats are suggested to compete with native rodents on Madagascar even though there is little empirical evidence to support this claim (Goodman 1995), and 3) *R. rattus* is not only a problem for conservation, but it is one of the most damaging agricultural pests and an important transmitter of human diseases (e.g. plague, hantaviruses and intestinal bilharziasis) in Madagascar (Duplantier & Rakotondravony 1999)

Unlike many other tropical areas in the world, *R. rattus* has not only been found close by human habitation, but also deep in undisturbed forest (Goodman & Carleton 1996). However, during our first studies (I) in Ranomafana National Park in 1996 and 1998, we caught *R. rattus* and *Mus musculus* only close by human habitation. It was possible that in Ranomafana *R. rattus* was just in the beginning of its expansion to primary forest or it had not succeeded in doing so. Thus, the surrounding of Ranomafana National Park proved to be an interesting area to study the occurrence of *R. rattus*. All types of ecological data about rodents in the area was needed because the national park was established in 1991 and a new research station, Centre Valbio, was inaugurated in 2001. Only two papers were published about the rodents of Ranomafana before our study (Carleton and Schmidt 1990, Ryan et al. 1993).

Because the knowledge of Malagasy rodent ecology was scarce, it was good to start to study the presence and abundance of rodent species and the different factors de-

termining them (I, II and III). As *Rattus rattus* is a human-mediated species, special attention was paid to the effects of human disturbance on the occurrence of species. In chapters I, II, and III the aims were 1) which rodent species co-exist (I, II)?, (2) what are their species-specific habitat utilisation patterns (I, II)?, and (3) how does anthropogenic habitat disturbance and fragmentation affect the rodent community (I, II and III) ?

When analyzing the occurrence data we noticed how sensitive one variable can be to other variables in regression models. In the methodological chapter (III), the goal was to introduce the social science -based elaboration technique into ecological research and to show how elaboration enables better utilization of our ecological knowledge in statistical data analysis. As the habitat use of *R. rattus* on the fragmented forest areas was used as an example, the second aim was to get a more in-depth interpretation about the effect of forest fragmentation on the presence of *R. rattus* at the same time.

The competition between native and introduced rodents on Madagascar is very difficult to study without a good idea about target species. The last goal of this thesis was to help to target future studies by estimating which native rodents are the most potential competitors with introduced rodents on Madagascar (IV). This estimation would hardly be done on the basis of scarce ecological data. Rather, the degree of ecomorphological similarity between introduced and nesomyine rodents was used.


The original idea of the thesis was also to examine whether rats, mice, and native rodents share any endoparasites. We collected parasitological data and some parts of it have already published (Laakkonen et al. 2003a, 2003b; Haukisalmi et al. 2010). Because we found altogether at least twelve new endoparasite species and only five of them (Durette-Desset et al. 2002a, 2002b, 2007) have been described so far, we passed the parasitological study in this dissertation.

3. Material and methods

3.1. Study area

The study was conducted in the Ranomafana National Park (RNP: 21 × 16', 47 × 20') in southeastern Madagascar, in its 3 km wide surrounding peripheral zone (I, II and III), and in the fragmented area (II, III) on the eastern and southeastern side of the RNP (Map 1). The protected core area of RNP, created in 1991, covers 43 500 ha of relatively undisturbed lowland rain forest, cloud forest, and high plateau forest (Grenfell & Robinson 1995). The altitudinal gradient ranges from 500 meters to 1500 meters. Over 25 000 residents are living in more than 100 villages in the peripheral zone of the RNP.

Six monitoring study sites were visited between 15 September and 29 November 1996, 1998, 1999, and 2000. Three of these sites were located in primary forests



Map 1. In Chapter I trapping grids were situated in six annual monitoring sites (stars, M= Miaranony, AL = Ambatolahy, TK = Talatakely, VT = Vatoharanana, VL = Valohoaka) and three additional lines (close by monitoring sites but not shown in the map). In Chapter II and III trapping grid were situated in 12 forest fragments (arrows) and four control sites. The dark grey areas represent the remaining forest cover according to the satellite photos taken in 1997.

(i.e., unlogged or slightly selectively logged forest): Vatoharanana (VT), Valohoaka (VL), and Miaranony II (M2). Another three study sites were in secondary forests (i.e., heavily logged forest): Talatakely (TK), Ambatolahy (AL), and Miaranony I (M1) (Map 1). For further information about the monitoring study sites see Chapter I, Wright (1997). Each of the six study sites included 10 (M1 and M2) or 12 (other sites) trapping plots. In this thesis the monitoring data of years 1996 and 1998 is used in Chapter I. The monitoring data of all other years are shown in Table 1 but are not used in other parts of the thesis. In addition to the monitoring plots, 44 extra trapping plots were studied in 1998. These trapping plots were placed along three lines between the primary forest areas and areas where *R. rattus* was known to occur. In total, the data of 112 trapping plots was used in Chapter I.

The study fragments (II, III) were situated in a 40 km long and 5 km wide zone between 600 meters and 1000 meters in elevation (Map 1). Field studies were carried

out in 2000, and satellite photos taken in 1997 were used to measure fragments. There were approximately 240 forest fragments ranging in size from 0.5 to 200 ha in area in 1997. The habitat matrix between the forest fragments consists of agricultural fields, grasslands, and villages. The loss rate of forest patches has been alarming in the corridor area between the RNP and the Andringitra NP: the number of patches decreased 54 % in 1990–2005 (Puhakka 2012).

3.2. Trapping methods

We used trapping grids in all studies but trapping grids were different in Chapter I than in Chapters II and III. Each trapping plot was a square measuring 20 x 20 m in Chapter I and 15 x 15 m in Chapters II and III. Live trap types were also different. Only one big live trap was in every corner in Chapter I whereas two live traps were utilized in every corner in Chapters II and III (see more details in I and II). Traps were in the field three nights in Chapter I and five nights in Chapter II and III. In the fragmentation study (II, III), a total of 95 trapping grids were established in 20 study sites (16 forest fragments ranging from 0.5 to 170 ha and four control sites in unfragmented forests). All study grids were situated at an elevation of 650–910 m.

In all trapping for this thesis (I,II, III), live traps were baited daily, on afternoons, with fresh banana slices and peanut butter. The traps were checked in the morning beginning at 07:00 h. Captured animals were sexed, examined for reproductive condition, weighed, and measured. All captured individuals of *Rattus rattus* were sacrificed, but most of the endemic species were released after measuring and ear-marking. In order to study endoparasites or preserve voucher specimens, some representatives of nesomyine rodent species were collected.

3.3. Habitat variables

Twelve habitat variables of forest and landscape structure were the same at all trapping grids (I, II, III): canopy cover; fallen log density; herbaceous cover; liana stem density; overstory tree height; altitude; and slope and distances from a building, field, human habitation (distance from the grid to nearest house, permanent satellite campsite, or to a recently cultivated field), forest edge, and stream. All trapping plots were also divided into four categories of human disturbance: (1) secondary bush, (2) heavily logged forest, (3) selectively logged forest, and (4) pristine forest. If the exact logging history of the trapping plot was not known, it was estimated by the occurrence of the following plant indicator species of forest disturbance: *Clidemia hirta*, *Dombeya* sp., *Harunga madagascariensis*, *Psidium cattleianum*, *Ravenala madagascariensis* (Lowry et al. 1997). In addition to these variables, the area of the fragments and the distance of a fragment from continuous forest and from the three nearest neighboring patches were also measured in fragmented area (II, III). All habitat variables were measured by same person.

3.4. Data analyses of habitat variables

Logistic regression was used to examine the effect of various habitat variables on the presence of the rodent species in trapping grids (I,II, III). Explanatory variables affecting the abundance of rodent species were modeled by Poisson regression (I). Only those zero observations that according to the logistic model had at least 30 % probability of species presence (Lindenmayer et al. 1991) were included in the abundance analysis. Without the elimination of some of the empty trapping grids extreme variable values which only indicate presence but not abundance of the studied species would have been included. Finally, the best-fit models were tracked by the forward stepwise procedure (I).

The logistic models developed in Chapter I were tested for predicting the presence of rodent species in the fragments and in continuous forest in Chapter II. Error rate (an estimate of the number of incorrect predictions), sensitivity (a measure of the ability of the model to predict the presence of studied species at a grid), and specificity (the ability of the model to correctly predict that studied species will not occur on a given square) were calculated as provided by Lindenmayer et al. (1991).

Although best-fit models were used in Chapter I, problems with using them were later realised. Mac Nally (2000) highlighted that best-fit models can be appropriate when the purpose of the modelling is to achieve a predictive model, but because of multicollinearity problems you can not use it for explaining causality. Predictive models can occasionally be useful in conservation biology, for example, if you have to predict distribution of endangered species for the purpose of protected area planning. However, drawing inferences about the causality of variables is often of more interest in classical ecology as well as in applied ecology (Mac Nally 2002). For example, if you have to prevent introduced species from invading more, you have to know not only where it is going to invade but also why it is going to invade to a certain site. Macnally's (2002) solution to multicollinearity problems was hierarchical partitioning with using randomization of the data matrix. However, hierarchical partitioning is mostly a technical tool. Austin (2002) claimed that neglect of ecological theory is a limiting factor in the application of statistical modelling in ecology. We relied on ecological knowledge with utilizing the idea of elaboration (see Box 1, III).

In Chapter III, the idea of elaboration is applied to logistic regression models of habitat use by black rat in fragmented areas. We call our approach 'explanatory framework based regression analysis'(EFRA), which enables systematization of the link between ecological knowledge and statistical analysis. It is important to take into consideration how the explanatory variables are related to each other which is often reflected as multicollinearity. The starting point for EFRA is that multicollinearity is more a source of information than a nuisance to data analysis as often seen in ecological research.

When using the EFRA approach, the interplay between the different explanatory variables and the ecological importance of the coefficient of fragment size were explored. In addition to fragment size, other variables were included in the logistic re-

gression model and the resulting changes in the estimated values of the coefficient of fragment size were recorded. Based on these changes, the role of the fragment size in the habitat use of *R. rattus* were interpreted. The order of different variables included in the model was founded on a spatial scale: the variables affecting the presence of *R. rattus* on the largest spatial scale were included first. Thus, the variables determining the habitat use of *R. rattus* on a metapopulation or population level were included in the model earlier than those operating more in an individual level. As the interpretation of EFRA rested on an existing ecological knowledge and the hypotheses under study, it guided the data analysis using the best of the author's ecological understanding and its limitations.

3.5. Morphological measurements and analysis

A total of 87 introduced and 62 nesomyine rodent specimens were measured for estimating probability for competition between introduced and nesomyine rodents based on morphological traits (IV). The data included representatives of all genera of the Nesomyinae except *Hypogeomys*, *Monticolomys*, and *Voalavo*. In total, data on 11 species of native Malagasy rodents was used.

However, data on the length of the head and body for most of these excluded species was obtained from the literature and used for comparison of size. Specimens of *R. rattus* were collected by J. T. Lehtonen in Ranomafana National Park between September and November 2000. The specimens of native Malagasy rodent were from the collections of the Museum of Natural History, London; Muséum National d'Histoire Naturelle, Paris; and the Finnish Museum of Natural History, Helsinki (collected during the field work of this thesis). Data on *Rattus norvegicus* and *Mus musculus* belongs to the Estonian specimens from the Zoological Museum of the University of Tartu, Estonia. The external and cranial proportions were assumed not differ significantly between European and Malagasy populations of the same species.

A total of 19 measurements were used in this study: 8 external, 4 skeletal, and 7 craniodental. They were selected from a larger number of characters based on a previous ecomorphological study (Miljutin 1997 and 1999). Cluster analysis (Sneath & Sokal 1973) was used for the estimation of similarity between species. Two procedures were made: 1) ranging of character values and 2) calculation of the Manhattan distances between species using unweighted pair-group average linkage.


Box 1. Elaboration

From the viewpoint of complexity, ecology is very close to the social sciences (Pigliucci 2002, Hansson 2003). Interestingly, many social scientists and ecologists seem to rely on logistical regression models, but they use them in different ways. In social sciences, regression analysis is often supported with the analysis method called elaboration. Elaboration, which is more a way of thinking than a statistical method, studies the relationship between two variables in a context of several other variables. It allows causal inference in non-experimental research (Blalock 1961).


If it is observed that the variable X precedes Y and hence it would likely be concluded that X causes Y, it has to be proven that Y is not caused by some variable other than X. In experimental studies it is possible to use a control without X and treatment with X. If Y is observed only with X, the causality theorem is supported. However, it still has to be proven that Y can't occur without X.

Elaboration recognizes three situations which may occur to the original bivariate relationship (X,Y) with the introduction of a test variable (Z, Figure 1). Other variables (Z) can be excluded by standardizing them by means of models or by means of thinking experiments. If the standardizing of Z does not exclude a relation between X and Y, causality is supported. If the inclusion of Z removes the effect of X, it means that Z affects both X and Y and causes relation between them (explanation) or X affects Z which affects onward Y (interpretation). Also, Z may change a way in which X affects Y (specification). Of course, there are often more than three variables when the whole situation is much more complicated.

a) explanation



b) interpretation



c) specification




Figure 1. Three analysis tools of elaboration

4. Main results and discussion

4.1. Rodent species in Ranomafana area

A total 698 rodent individuals were captured in 6204 trap nights in 1996 and 1998–2000 (I, II, Table 1). Five endemic species (*Nesomys rufus*, *N. audeberti*, *Eliurus tanala*, *E. minor*, and *E. webbi*) and introduced *R. rattus* were found in both the secondary and the primary forests. The house mouse *M. musculus* was only found in the secondary forests. In addition to our regular trapping *Gymnuromys roberti* was captured twice in the secondary forest close to Miaranony village (Map 1) in 1999. We didn't capture *Brachytarsomys albicauda* with traps but we observed it close to both Miaranony and Talatakely villages. *Brachyuromys betsileonsis* was absent from our data probably because our traps were not located above 1120 m a.s.l. In other studies, *B. betsileonsis* has been captured in RNP above 1200 m a.s.l. (Carleton and Schmidt 1990, W. Rakotonirina, per. comm.). Also *Monticolomys koopmani* has been observed in RNP (Soarimalala & Goodman 2011) but we could not capture it.

4.2. Effects of human disturbance on endemic rodent occurrence

Although the total number of endemic rodents was higher in the primary forests than in the secondary forests, selective logging or other types of light forest disturbance doesn't seem to be a big threat to endemic rodent species in Ranomafana (I). None of the three most common native species (*Nesomys audeberti*, *N. rufus*, and *Eliurus tanala*) were more abundant in un-logged than in selectively logged forest (I, Figure 2). However, all other native species except *Eliurus tanala* were absent in secondary bushes (I). Thus, there is no doubt that deforestation contributes to the disappearance of most of the native rodent species. The same kind of result was found in the spiny forest in southern Madagascar where small mammal species diversity declined by 40 percent in response to clearing (Scott et al. 2005) and from Central China where small mammal diversity was higher in deforested than in afforested habitats (Raoul et al. 2008).

When we were studying the effects of forest fragmentation on rodents, the study fragments were at relatively low elevations (650–910 m) and presented only two common native species: *Eliurus webbi* and *Nesomys audeberti* (II). In our study area, *E. webbi* does not seem to be sensitive to forest fragmentation as its mean abundance per study grid was almost the same in every size class of fragments and in control sites (II, Figure 3). *N. audeberti* was virtually absent from the fragments although two or three individuals were trapped in each control site (II, Figure 3). In addition to fragmentation, *N. audeberti* probably suffers from rapid deforestation below 800 m in elevation where it used to be the most common. Dehgan (2003) studied the lemurs partly at the same fragments in RNP as us and found that lemur species richness could not be predicted by the size of forest fragments.

Table 1. Number of individuals trapped for each species of rodent in six monitoring sites in secondary forest (SF) and in primary forest (PF) in 1996 and 1998–2000 (total trap-days accrued is shown in parentheses). Data of years 1996 and 1998 was used in Chapter I but data of years 1999 and 2000 are not shown in other chapters of the thesis.

Species	SF (272)				PF (272)			
	-96	-98	-99	-00	-96	-98	-99	-00
<i>Mus musculus</i>	2	3	5	1	-	-	-	-
<i>Rattus rattus</i>	30	30	40	17	-	1	6	1
<i>Eliurus minor</i>	1	1	-	-	2	-	-	-
<i>Eliurus tanala</i>	1	6	3	6	8	7	8	3
<i>Eliurus webbi</i>	-	1	3	1	1	1	1	1
<i>Nesomys audeberti</i>	-	10	11	4	2	2	2	4
<i>Nesomys rufus</i>	7	3	1	1	8	24	19	12
Total rodents	41	54	63	30	21	34	36	21
Total nesomyines	9	21	18	12	21	33	30	20

Contrary to our results, *E. webbi* was found only in fragments larger than 28 hectares in the littoral forest of southeastern Madagascar (Ramanamanjato & Ganzhorn 2001). The fragments in Ranomafana were much younger than in the littoral forest. It is possible that all negative effects of isolation are not yet seen in Ranomafana, and the native rodent populations suffer from extinction debt (Tilman et al. 1994, Hanski & Ovaskainen 2002). In the dry deciduous forest of western Madagascar, the abundance of endemic rodents decreased as forest fragments became smaller (Ganzhorn 2003). In the Amazonian forest, small mammal abundance and species diversity were not affected by forest fragment size but rather by the quality of the open-habitat matrix (Santos-Vilho et al. 2012). It is not surprising that the response to forest fragmentation varies from species to species and from region to region. As shown in our methodological study (III), the effect of forest fragmentation is a very complicated phenomenon. There can be many other habitat variables which may effect via the fragment size. However, the situation is different with endemic rodents than with *R. rattus* because the forest fragments are really habitat fragments for most of the endemic rodent species but not for *R. rattus* (II, III).

4.3. Occurrence of black rat in Ranomafana area

The abundance of *R. rattus* increased with increasing forest disturbance and it was most common in heavily logged secondary forest (Figure 2). Therefore, *R. rattus* benefits from deforestation unlike endemic rodents. However, *R. rattus* can survive also in primary forests in eastern Madagascar (Goodman and Carleton 1996, 1998). We observed *R. rattus* the first time in selectively logged primary forest areas in Ranomafana in 1998 (I). Later it has been captured also in un-logged forest areas in Valohoaka in 1999, 2000 (Table 1), and 2012 by the students of a tropical field ecology course (L. Sandholm. pers. comm.). In Chapter I we said that *R. rattus* was never captured farther than 500 meters away from human habitation or permanent satellite camp-sites. The distance from Valohoaka to the nearest field is about 1800 meters but there is also a small satellite campsite which was used temporarily in 1999–2000 and in 2012. It is difficult to say if the temporarily used campsite and its exposed garbage pit are the only things attracting *R. rattus* there.

By means of the best-fit logistic regression model probability, of *R. rattus*'s presence increased with decreasing canopy cover, distance from forest edge, and from human habitation, and with increasing herbaceous cover, altitude, and overstory tree height (I). All these variables are in line with the suggestion that the occurrence of species is associated with forest disturbance. However, in the third chapter (III) we express our concern on the suitability of best-fit models to explain ecological phenomena. Mac Nally (2000) highlighted that the best-fit procedure can be appropriate when the




Figure 2. The mean abundance of different rodent species per trapping square in different categories of natural state in RNP (I).

purpose of the modelling is to achieve a ‘predictive’ model, but it shouldn’t be used it for explaining causality.

Because our first results suggested that the spread of *R. rattus* is associated with deforestation it was interesting to also study the effects of fragmentation on the occurrence of *R. rattus*. The abundance of *R. rattus* was higher in the fragments than in continuous forest, but the difference was not statistically significant (Figure 3, II). The size of fragment was also not significantly correlated with the abundance of *R. rattus* (II). Similar results were found in the littoral forest of southeastern Madagascar (Ramanamanjato and Ganzhorn 2001), but in the dry deciduous forest, the capture rate of *R. rattus* was negatively correlated with the fragment size (Ganzhorn 2003).

The above-mentioned best-fit logistic regression model could not predict the presence of *R. rattus* in fragmented areas well (II). When we were analysing the habitat data in fragmented areas, we tried to better understand what ecological factors determine the presence of *R. rattus* by using the EFRA approach (explanatory framework based regression analysis, III). We demonstrated the sensitivity of the size of fragment to other variables in the regression models (Figure 4). The dramatic variation of the coefficient of fragment size was not surprising when remembering that *R. rattus* utilizes open areas, too. A forest fragment does not necessarily mark the limits of a habitat fragment for this species and that is why its role in the habitat models of rat is fragile. It may also explain the difference in results about the correlation between the fragment size and the trapping rate of *R. rattus* in different areas in Madagascar.




Figure 3. Mean number of rodents captured per square and standard deviations in different size categories of fragments. The values were calculated from mean number of rodents per square of each fragment (II).

The association between the size of the fragment and other variables varies in different areas. The measured value of the size of the fragment can rather be viewed as an index capturing certain features of the forest fragment (III).

In our data especially two habitat variables had a dominant role: 1) distance from a field which may be connected to the food resources of the rat and 2) liana stem density which can be seen as an indicator of the virginity of forest (III). Food resources provided by fields and all kind of forest structure changes reflecting human presence are probably more important to *R. rattus* than the size or shape of the forest fragment.

4.4. Interaction between black rat and endemic rodents

Based on a comparison of body size, body construction, and craniodental characters it is probable that nesomyine rodents compete with *Rattus rattus* but not with *Rattus norvegicus* (IV). The probability of interspecific competition was estimated as:

1) very high between *R. rattus* and *Gymnuromys roberti*;




Figure 1. The coefficients of FRAGSIZE in models that are organized according to the operational level explanatory framework (see text). Horizontal axis shows the inclusion order of the variables into the logit-model, the vertical – coefficient of FRAGSIZE in the corresponding model. The effects of separate inclusion of variables in the distance and forest structure packages are represented as well. The exact values of the coefficients and their p-values are given in Table 2. The straight lines connecting the different models are just to make comparison of changes easier. It is by no means meant to indicate linearity of change between the models.

- 2) high between *R. rattus* and species of *Nesomys* and larger semiarboreal *Eliurus* (*E. tanala*, *E. webbi*);
- 3) moderately low between *R. rattus* and *Brachytarsomys albicauda* and larger species of arboreal *Eliurus* (e.g. *E. myoxinus*); and
- 4) low between *R. rattus* and species of *Brachyuromys* and *Macrotarsomys ingens*.

Other native Malagasy rodents than those mentioned above and the introduced rodents likely do not compete with each other, except maybe for *M. musculus* and the poorly studied *Monticolomys koopmani* and *Voalavo gymnocaudatus*.

Although Goodman and Sterling (1996) demonstrated that food preference of *R. rattus* overlaps with *Brachyuromys ramirohitra*, *Eliurus tanala*, *E. webbi*, *Gymnuromys roberti*, and *Nesomys rufus*, there are no other experimental studies testing the competition hypothesis between the black rat and Nesomyine rodents. In Madagascar, *R. rattus* is largely terrestrial and uses the same kind of ground burrow types as *G. roberti* and *N. rufus* (Laakkonen et al. 2003). In future studies, particular attention should be paid to species-pairs *R. rattus* – *Gymnuromys roberti* and *R. rattus* – *Nesomys rufus*. It is possible that the populations of *G. roberti* have already diminished in RNP. During four years of trapping it was observed only twice close by Miaranony village in RNP. Previously *G. roberti* has been reported in RNP close to Talatakeley village, which is the most intensively studied area in RNP (Carleton & Schmidt 1990). *G. roberti* is very abundant in the Andringitra National Park which is connected with RNP by a forest corridor (Goodman & Carleton 1996).

Interactions between *Eliurus webbi* and *Rattus rattus* could be worth studying in the future as well. Based on the fragmentation study in RNP (II), *E. webbi* may be more susceptible to competitive effects from *R. rattus* in fragments than in continuous forest although no negative correlations between capture rates of *R. rattus* and *Eliurus webbi* have been found in the littoral forest or in the dry deciduous forest (Ramanamajato & Ganzhorn 2001, and Ganzhorn 2003). In contrast, hardly any Nesomyine rodents were captured in fragments in the humid forests of Andranomay/ Ambohitantely where *R. rattus* was abundant (Ganzhorn et al. 2003). However, drawing conclusions about interspecific competition from correlation analyses should be treated with caution (Strong et al. 1984).

Among potential competitors with *R. rattus*, four species – *Brachyuromys albicauda*, *Brachyuromys villosa*, *Nesomys audeberti*, and *Nesomys lambertoni* – are on average larger than the black rat (IV). This larger size may help them to predominate over *R. rattus*. All other potential competitors with *R. rattus* are smaller or equal in size and may be threatened by the black rat.

When considering *R. rattus* in a global perspective, it has occupied a wide variety of habitats on islands without an extensive native fauna of small mammals (New Zealand, Hawaii, Puerto Rico, Galápagos islands, Timor, and numerous smaller islands), but where a diverse small-mammal fauna exists, black rats are usually absent or only small in numbers in undisturbed habitats (see Clark 1980, Aplin et al. 2003). How-

ever, Madagascar is an exceptional case as *R. rattus* has invaded many different kinds of habitats with a rich native rodent fauna.

We recommend the ecomorphological approach (IV) for preliminary studies of recently described, extinct, and rare mammal species. Usually it is very difficult to get other kind of data about species which should be the focus of conservation studies. Dayan and Simberloff (1998) concluded that mammals have determinate growth, which makes them favorable for morphometric studies and that while many studies of invertebrates fail to confirm the hypothesis of interspecific competition, many studies of vertebrates do suggest its operation.

5. Final remark and future perspectives

This thesis suggests that deforestation and heavy logging promotes the spreading of introduced *Rattus rattus* in the humid forest of Madagascar (I). The forest fragmentation does not directly affect the occurrence of *Rattus rattus* but it may intensify negative association between *R. rattus* and endemic rodents (II, III). Therefore, forest fragmentation can be problematic for the Malagasy rodent community.

For native Malagasy rodents, clear-cutting is disastrous and forest fragmentation may have a diminishing effect on populations of some species, e.g. *Nesomys audeberti* (I,II). The forest below 800 m in elevation where *N. audeberti* is most frequently documented is rapidly disappearing in the Ranomafana-Andringitra corridor area. Many other forest-dwelling animal and plant species may suffer from the same problem. Accordingly, it is important to ensure forest protection at different elevations.

The interspecific competition between *Rattus rattus* and morphological similar native rodent species (e.g. *Gymnuromys roberti*) will be an important topic to study in the future. One possibility to study it is via removal experiments. In response to *R. rattus* removal, the populations of the Australian native rodents increased due to immigration, juvenile recruitment, and increases in residency of females (Stokes et al. 2009). Other fruitful ways to study interspecific association could be a research on the effects of competition on individuals' fitness and physiological costs (Eccard and Ylönen 2003).

In the beginning of the thesis, it was discussed that invasive species drive native animal species to extinction more often by spreading parasites or pathogens or by predation than by competition. So far, there is evidence that some parasites are shared between *R. rattus* and nesomyine rodents, but there is no evidence that populations of native rodents suffer from those parasites (see Ganzhorn 2003, Laakkonen et al. 2003). We have also studied helminths of Malagasy rodents, but we found only one parasite species from endemic rodents (*Mastophorus muris* in *Eliurus tanala*) which is an endoparasite of *Rattus rattus* as well (Lehtonen et al. unpubl.).

One of the most potential pathogens that could spread from *Rattus rattus* to Nesomyine rodents is the plague. Duplantier and Duchemin (2003) mentioned the fol-

lowing case from Moramanga in 1932. Several specimens of *Brachytarsomys albicauda* invaded villages, later their bodies were found around houses and finally villagers came down with the plague. More parasitological and epidemiological studies are needed in Madagascar.

Moreover, it is important to start to study predator effects of introduced rodents on birds and other animals in Madagascar. It is an almost totally neglected research topic in Madagascar so far.

6. Acknowledgements

'Tsara tsara kokoa miandry.' This sentence, meaning 'It is better to wait,' was the first Malagasy sentence that I learnt. I am very sorry to everybody that I assimilated the sentence so completely in my dissertation process. First, I would like thank you all for your patience!

My sincerest gratitude is due to my supervisor, Jari Niemelä, who – despite his hectic schedule – has always found time for comments, recommendation letters, and advice. Over the years, he has remained very patient about my unfinished dissertation and reminded me about it very kindly. It was risky to begin to support young mammalogists who dreamed of conducting research on an island that they have never visited. Jari, thank you so much for taking a risk!

I am deeply grateful to Hannu Rita who has been like my second supervisor. It was always inspirational to visit your office where I learnt a lot. I would select you if I had to award the best university teacher prize to someone. Thank you for guiding me in the jungle of non-statistical statistics!

I have dreamt of being a biologist in the rainforest since I was a schoolboy, and I can never thank Olli Mustonen enough for making my dream come true! Olli was the initiator of the preliminary project and the expedition in 1996. I have good memories from the first expedition which changed my life. Misaotra betsaka Olli! I also thank Salla Hannunen who cancelled her expedition to Madagascar just one month before leaving and gave me the possibility to go in her place.

When we started the preliminary studies in Madagascar we knew almost nothing about Malagasy rodents. We were lucky that Haingotiana Ramiarinjanahary had the possibility to start working with us. Haingo has been the best teacher to help me become familiar with Malagasy rodents and culture. Haingo, I could never survive without your help! It has been great to cooperate in many other projects after that as well. During the fragmentation study I enjoyed the help of Volatiana Rasataharilala who was excellent both with working in the field and writing her Master's thesis. Misaotra tompoko Haingo sy Vola!

I met Andrei Miljutin in the African Small Mammal Symposium in Paris where we got the idea for the ecomorphological study. Andrei can understand the mental and

physical life of rodents better than anybody else I have met. Andrei, thank you very much for guiding me in ecomorphology and warm home hospitality in Tartu!

My fragmentation study couldn't be possible without Alexander Dehgan studying lemurs in the same area. It was easy to work with Alex's good groundwork. Alex, thank you for all your help and hospitality in Chicago!

I was also studying parasites; however, I left the parasitological study out of this dissertation. Voitto Haukisalmi and Juha Laakkonen were always ready to help with parasitological studies. It was almost as unforgettable to find new parasite species from the intestines of rodents as it was to trek in the rainforest. Voitto was helping me a lot with non-parasitological studies, as well. Thank you so much Voitto and Juha!

I want to thank François Zakamanana for being such a great help in the field and for being like my brother in Madagascar. It was always nice to work with you. Jean Claude Rakotoniaina, François Ratalata, Voahangy Soarimalala, Justin Solo, Desire Ramarison, Albert (from Ranovao), Helena Wirta, and many short time assistants helped with fieldwork as well. Misaotra betsaka Zaka et al.!

There were plenty of people sharing breakfasts, lunches, dinners, and many memorable parties with me in the old cabin in Talatakely in Ranomafana National Park. I would like to acknowledge you all for the good atmosphere! Unfortunately I can't mention all people by name. However, I would like to mention Aimée Razafiarimalala who was like the mother of the cabin and always ready to help. Misaotra tompoko Aimée!

Without Prof. Patricia Wright there would be no national park or any kind of research station in Ranomafana. Patricia, thank you very much for establishing the national park and for all the help and many good tips! I thank Prof. Jukka Jernvall for advice too!

Benjamin Andriamihaja and the entire staff of the Madagascar Institut pour la Conservation des Environnements Tropicaux (MICET) provided invaluable logistic support. It was always nice to visit in the office of MICET in Antananarivo. You solved many problems. A very important contact for me was Prof. Daniel Rakotondravony at the University of Antananarivo who was always ready for cooperation and sharing his knowledge of Malagasy small mammals. The Madagascar National Parks (before ANGAP) organization always gave permission to work without problems. Misaotra betsaka!

I want to express my gratitude to Steve Goodman, for advising me with identifying Malagasy small mammals and commenting on one of my papers. I am also grateful to my reviewers Prof. Jörg Ganzhorn and Doc. Otso Huitu for all of their comments. I would like to thank Prof. Veijo Kaitala for support and help in bureaucracy matters during the final stages, Victoria Veach for the language revision of the summary, Petri Clusius for layout, and Prof. Lotta Sundström for taking on the role of the custos for my defence! The head of the department Kurt Fagerstedt deserves a special mention: his 'kick' finally brought about the completion of the dissertation. This work would not have been possible without grants by the Academy of Finland, the Finnish

Cultural Foundation, the Oscar Öflunds stiftelse and the University of Helsinki. Halti company donated camping equipment.

I also acknowledge all members of the Finland-Madagascar Association, Tany Maitso project and RESPECT project for sharing interest in Madagascar. Especially I would like mention Chantal Andriaharimidy, Mar Cabeza, Ilkka Hanski, Kaisa Korhonen, Markku Lahtinen, Johanna Rainio, Tanjona Ramiadantsoa, Njara Rasolomanana, Essi Syrjäkari, Tarmo Virtanen, and Helena Wirta.

I also want to thank all my colleagues at the university and friends outside work and the Madagascar projects. During the last months the one-year bird spotting – jogging competition has kept me in good condition. Thank you Mikko, Olli A, and Paa! Although I can't mention all my friends, I would like acknowledge Jan, Ilona, Jyrki, and Panu for friendships lasting even longer than my PhD project!

Lopuksi haluan kiittää sukulaisiani ja perhettäni kaikesta avusta pitkän väitöskirjahankkeen aikana ja ennen sitä. Siskoni Sirpa on aina tukenut ja kannustanut minua parhaansa mukaan. Isäni on sytyttänyt minussa kiinnostuksen luontoon ja äidiltäni olen toivon mukaan oppinut jotain sellaista ihmisten kanssa toimimisesta, josta on voinut olla apua myös vieraan kulttuurin keskellä. Kaikki poikamme isovanhemmat, kummit Sirpa ja Jukka sekä Pasi-eno ja Maria ovat pyyteettömästi rientäneet lastenhoitoavuksi kun sitä on tarvittu. Ovaskaisen perhe on myös tarjonnut Teiskossa mitä mainiota täysihoitoa etätyöskentelyni aikana ja Paavo on auttanut Madagaskar-hankkeiden kirjanpidossa. Kiitoksia teille kaikille avusta! Viimeisimpänä kiitän minulle kaikkein tärkeimpiä ihmisiä Outia ja Okkoa. Ilman teidän kärsivällisyyttä, rakkautta ja antamaa Eliurus-T-paitaa tämä työ ei olisi koskaan valmistunut! Kirjoitan tätä viimeistä kappaletta kaksi päivää sen jälkeen kun Okko (3v) kysyi: ”Mikä on väitöskirja?” Tämä se on.

7. References

- Aplin KP, Chesser T and ten Have J. 2003. Evolutionary biology of the genus *Rattus*: profile of an archetypal pest. In: Singleton GR, Hinds LA, Krebs CJ and Spratt, DM (eds). *Rats, Mice and People: Rodent Biology and Management*. ACIAR Monograph No. 96. Canberra, Australian Centre for International Agricultural Research. 487–498.
- Austin MP. 2002. Spatial prediction of species distribution: an interface between ecological theory and statistical modelling. *Ecological Modelling* 157:101–118.
- Blalock Jr, HM 1961. *Causal inferences in nonexperimental research*. Chapel Hill
- Brown KA, Gurevitch J. 2004. Long-term impacts of logging on forest diversity in Madagascar. *Proceedings of the National Academy of Sciences*. 101: 6045–6049.
- Burney, D:A. 1997. Theories and facts regarding Holocene environmental change before and after human colonization. – In Goodman, S.M. and Patterson, B.D. (eds.), *Natural change and human impact in Madagascar*, Smithsonian Institution Press.
- Burney DA, Burbey LP, Godfrey LR, Jungers WL, Goodman SM, Wright HT and Jull AJT. 2004. A chronology for late prehistoric Madagascar. *Journal of Human Evolution* 47: 25–63.

- Callmander M.W., Phillipson P.B., Schatz G.E., Andriambololona S., Rabarimanarivo M., Rakotonirina N., Raharimampionona J., Chatelain C., Gautier L., Lowry P.P. 2011. The endemic and non-endemic vascular flora of Madagascar updated. *Plant Ecology and Evolution* 144:121–125
- Carleton MD and Goodman SM. 1998. New taxa of Nesomyinae rodents (Muroidea: Muridae) from Madagascar's northern highlands, with taxonomic comments on previously described forms. *Fieldiana: Zoology, New Series* 90: 163–200
- Carleton MD and Schmidt DF. 1990. Systematic studies of Madagascar's endemic rodents (Muroidea: Nesomyinae): An annotated gazetteer of collecting localities of known forms. *American Museum Novitates* 2987: 1–36.
- Clark DB 1980. Population ecology of *Rattus rattus* across a desert-montane forest gradient in the Galápagos islands. *Ecology* 61: 1422–1433. Davis M.A. 2003. Biotic globalization: does competition from introduced species threaten biodiversity? *BioScience* 53: 481–489.
- Clavero M, Garcia-Berthou E. 2005. Invasive species are a leading cause of animal extinctions. *Trends in Ecology Evolution* 20:110.
- Combes C. 1996. Parasites, biodiversity and ecosystem stability. *Biodiversity and Conservation* 5: 953–962.
- Corlett RT. 2010. Invasive aliens on tropical East Asian islands. *Biodiversity and Conservation* 19:411–423.
- Cox MP, Nelson MG, Tumonggor MK, Ricaut FX, Sudoyo H. 2012. A small cohort of island Southeast Asian women founded Madagascar. *Proceedings of the Royal Society B: Biological Sciences* 279:2761–2768.
- Davis MA. 2003. Biotic globalization: does competition from introduced species threaten biodiversity? *Bioscience* 53:481–489.
- Dayan T, Simberloff D. 1998. Size patterns among competitors: ecological character displacement and character release in mammals, with special reference to island populations. *Mammal Review* 28:99–124
- Daszak P, Cunningham AA and Hyatt AD. 2000. Emerging infectious diseases of wildlife threats to biodiversity and human health. *Science* 287: 443–449.
- de Wit, MJ 2003. Madagascar: Heads it's a continent, tails it's an island. *Annual review of Earth Planet Sciences* 31: 213–248.
- Dehgan A. 2003. The Behavior of Extinction: Predicting the Incidence and Local Extinction of Lemurs in Fragmented Habitats of Southeastern Madagascar. PhD Dissertation, University of Chicago, Chicago, IL, 248 p.
- Diamond JM, Bishop KD, Van Balen S. 1987. Bird survival in an isolated Javan woodland: Island or mirror? *Conservation Biology* 1:132–42.
- Drake, JM. 2003. The paradox of the parasites: implications for biological invasions. *Proceedings of the Royal Society of London, B (Suppl.)* 270: S133–S135.
- Duplantier JM, Duchemin JB. 2003. Introduced small mammals and their ectoparasites: a description of their colonisation and its consequences. In: Goodman SM, Benstead JP (eds) *The natural history of Madagascar*. University of Chicago Press, Chicago
- Duplantier JM, Orth A, Catalan J & Bonhomme F. 2002. Evidence for a mitochondrial lineage originating from the Arabian Peninsula in the Madagascar house mouse (*Mus musculus*). *Heredity*, 89, 154–158.
- Duplantier J, Rakotondravony D. 1999. The rodent problem in Madagascar : agricultural pest and threat to human health. In: H. L. Singleton G, Leirs H and Zhang Z, editors. *Ecologically-based rodent management*. ACIAR publ., 441–459 pp.

- Durette-Desset MC, Lehtonen JT & Haukisalmi V. 2007: Trichostrongylina (Nematoda) from Malagasy Muridae. III – Description of a new species of *Heligmonoides Baylis*, 1928 (Heligmonellidae) parasitic in *Mus musculus*. – Parasite 14: 53–60
- Durette-Desset MC, Lehtonen JT & Haukisalmi V. 2002a: Trichostrongylina (Nematoda) from Malagasy Muridae. I – Description of two new species of Heligmonina (Heligmonellidae) in *Nesomys* spp. – Parasite 9: 29–35
- Durette-Desset MC, Lehtonen JT, Haukisalmi V. 2002b: Trichostrongylina (Nematoda) from Malagasy Muridae. II – Description of two new species of Heligmonina (Heligmonellidae) in *Nesomys rufus* and *Eliurus tanala* – Parasite 9: 127–133
- Eccard JA, Ylönen H. 2003. Interspecific competition in small rodents: from populations to individuals. *Evolutionary Ecology*, 17, 423–440.
- Fritts TH, Rodda GH. 1998. The role of introduced species in the degradation of island ecosystems: A case history of Guam. *Annual Review of Ecology and Systematics* 29: 113–140.
- Fordham DA & Brook BW. 2010. Why tropical island endemics are acutely susceptible to global change. *Biodiversity and Conservation* 19:329–342
- Ganzhorn, JU. 2003. Effects of introduced *Rattus rattus* on endemic small mammals in dry deciduous forest fragments of western Madagascar. *Animal Conserv.* 6, 147–157.
- Ganzhorn JU, Goodman SM, Dehgan A. 2003. Effects of Forest Fragmentation on Small Mammals and Lemurs in *The natural history of Madagascar* (ed. by S.M. Goodman and J.P. Benstead), pp. 1228–1234. University of Chicago Press, Chicago.
- Garbutt N 2007. *Mammals of Madagascar: A Complete Guide*. A & C Black, 304 pp.
- Gommery D, Ramanivosoa B, Faure M, Gue'rin C, Kerloc'h P, Se'ne'gas F, Randrianantenaina, H. 2011. Les plus anciennes traces d'activités anthropiques de Madagascar sur des ossements d'hippopotames subfossiles d'Anjohibe (Province de Mahajanga) *Comptes Rendus Palevol.* 10: 271–278.
- Goodman SM. 1995. *Rattus* on Madagascar and the dilemma of protecting the endemic fauna. *Conservation Biology* 9: 450–453
- Goodman SM & Carleton MD. 1996. The rodents of the Réserve Naturelle Intégrale d'Andringitra, Madagascar. *Fieldiana: Zoology* 85: 257–283.
- Goodman SM and Carleton MD. 1998. The Rodents of the Réserve Spéciale d'Anjanaharibe-Sud, Madagascar. *Fieldiana: Zoology, New Series* 90: 201–221
- Goodman SM, Sterling EJ. 1996. The utilization of *Canarium* (Burseraceae) seeds by vertebrates in the RNI d'Andringitra, Madagascar. *Fieldiana: Zoology, New Series* 85: 83–89
- Goodman SM, Rakotondravony D. 1996. The Holocene distribution of Hypogeomys (Rodentia:Muridae: Nesomyinae) on Madagascar. In *Biogéographie de Madagascar*: 283–293. Lourenço, W.R. (Ed.). Paris: Editions de l'Ors.
- Goodman SM, Andrianarimisa A, Olson LE, Soarimalala V. 1996. Patterns of elevational distribution of birds and small mammals in the humid forests of Montagne d'Ambre, Madagascar. *Ecotropica* 2: 87–98.
- Goodman SM, Ganzhorn JU, Olson LE, Pidgeon M, Soarimalala V. 1997. Annual variation in species diversity and relative density of rodents and insectivores in the Parc National de la Montagne d'Ambre, Madagascar. *Ecotropica* 3: 109–118
- Goodman SM, Ganzhorn JU, Rakotondravony D. 2003. Introduction to mammals in *The natural history of Madagascar* (ed. by S.M. Goodman and J.P. Benstead), pp. 1159–1186. University of Chicago Press, Chicago.
- Goodman SM, Raheriarisena M, Jansa SA. 2009. A new species of *Eliurus* Milne Edwards, 1885 (Rodentia: Nesomyinae) from the Réserve Spéciale d'Ankarana, northern Madagascar. *Bonner Zoologisches Beiträge*, 56: 133–159.

- Green GM and Sussman RW. 1990. Deforestation history of the eastern rainforest of Madagascar. *Science* 248: 212–215.
- Grenfell S, Robinson L. 1995. The Ranomafana National Park Management Plan (1996–2000). Ranomafana National Park Project report
- Gurevitch J, Padilla DK. 2004. Are invasive species a major cause of extinctions? *Trends in Ecology and Evolution* 19:470–474.
- Hanski I, Ovaskainen O. 2002. Extinction Debt at Extinction Threshold. *Conservation Biology* 16: 666–673.
- Hansson L. 2003. Why ecology fails at application: should we consider variability more than regularity? – *Oikos* 100: 624–627.
- Harper GJ, Steininger MK, Tucker, CJ, Juhn D, Hawkins F. 2007. Fifty years of deforestation and forest fragmentation in Madagascar. *Environmental Conservation* 34: 325–333.
- Harris D. 2009. Review of negative effects of introduced rodents on small mammals on islands. *Biological Invasions* 11:1611–1630.
- Haukisalmi V, Hardman LM, Foronda P, Feliu C, Laakkonen J, Niemimaa J, Lehtonen JT, Henttonen H. 2010. Systematic relationships of hymenolepidid cestodes of rodents and shrews inferred from sequences of 28S ribosomal RNA. *Zoologica Scripta*. 39: 631–641
- Irwin MT, Wright PC, Birkinshaw C, Fisher BL, Gardner CJ, Glos J, Goodman SM, Loiselle P, Rabeson P, Raharison J-L, Raheirilalao MJ, Rakotondravony D, Raselimanana A, Ratsimbazafy J, Sparks JS, Wilmé L, Ganzhorn JU. 2010. Patterns of species change in anthropogenically disturbed forests of Madagascar. *Biological Conservation* 143:2351–2362
- Jansa SA, Goodman SM, Tucker PK. 1999. Molecular phylogeny and biogeography of the native rodents of Madagascar (Muridae: Nesomyinae): A test of the single-origin hypothesis. *Cladistics* 15: 253–270.
- Jansa SA, Weksler M. 2004. Phylogeny of muroid rodents: Relationships within and among major lineages as determined by IRBP gene sequences. *Molecular Phylogenetics and Evolution* 31: 256–76.
- Keane RM, Crawley MJ. 2002. Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology and Evolution* 17: 164–170.
- Kesavaraju B, Damal K, Juliano SA. 2008. Do natural container habitats impede invader dominance? Predator-mediated coexistence of invasive and native container-dwelling mosquitoes. *Oecologia* 155:631–639
- Laakkonen J, Goodman SM, Duchemin J-B, Duplantier J-M. 2003. Trypomastigotes and potential flea vectors of the endemic rodents and the introduced *Rattus rattus* in rainforests of Madagascar. *Biodiversity and Conservation* 12: 1775–1783.
- Laakkonen J, Lehtonen JT, Ramiarinjanahary H, Wright P. 2003. Trypanosome infections in the invading *Rattus rattus* and endemic rodents in rainforests in Madagascar. In: Singleton GR, Hinds LA, Krebs CJ, Spratt DM (eds). *Rats, Mice and People: Rodent Biology and Management*. ACIAR Monograph No. 96. Canberra, Australian Centre for International Agricultural Research. 37–39.
- Langrand O, Goodman SM. 1997. Inventaire des oiseaux et des micromammifères des zones sommitales de la Réserve Naturelle Intégrale d'Andringitra. *Akon'ny Ala* 20: 39–54
- Lindenmayer DB, Cunningham MT, Tanton MT, Nix HA and Smith AP. 1991. The conservation of arboreal marsupials in the montane ash forests of the central highlands of Victoria, south-east Australia: III. The habitat requirements of Leadbeaters Possum *Gymnobelideus leadbeateri* and models of the diversity and abundance of arboreal marsupials. *Biological Conservation* 56: 295–315
- Lowry II PP, Schatz GE and Phillipson PB. 1997. The classification of natural and anthropogenic vegetation in Madagascar. In: Goodman SM and Patterson BD (eds) *Natural Change and Human Impact in Madagascar*, pp 93–123. Smithsonian Institution Press, Washington, DC

- Mack RN, Simberloff D, Lonsdale WM, Evans H, Clout M, Bazzas FA. 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* 10: 689–710.
- Mac Nally R. 2000. Regression and model-building in conservation biology, biogeography and ecology: The distinction between – and reconciliation of – ‘predictive’ and ‘explanatory’ models. *Biodiversity and Conservation* 9: 655–671.
- Mac Nally R. 2002. Multiple regression and inference in ecology and conservation biology: further comments on identifying important predictor variables. *Biodiversity and Conservation* 11: 1397–1401.
- McCallum H, Dobson A. 1995. Detecting disease and parasite threats to endangered species and ecosystems. *Trends in Ecology and Evolution* 10: 190–194.
- Mein P, S en egas F, Gommery D, Ramanivosoa B, Randrianantenaina H, Kerloc’h P. 2010. Nouvelles esp eces subfossiles de rongeurs du Nord-Ouest de Madagascar. *Comptes Rendus Palevol* 9: 101–112.
- Miljutin A. 1997. Ecomorphology of the Baltic rodents: Body form, ecological strategies and adaptive evolution. *Folia Theriologica Estonica* 3: 1–111.
- Miljutin A. 1999. Trends of specialization in rodents: The birch mice, genus *Sicista* (Dipodoidea, Rodentia). *Folia Theriologica Estonica* 4: 91–128.
- Mitchell CE, Power AG. 2003. Release of invasive plants from fungal and viral pathogens. *Nature* 421: 625–627.
- Musser GG, Carleton MD. 1993. Family Muridae. In *Mammal species of the world*, eds. Wilson, DE, Reeder DM. pp. 501–753. Smithsonian Institution Press.
- Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB, Kents J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- Oppel S, Beaven BM, Bolton M, Vickery J & Bodey TW. 2011. Eradication of Invasive Mammals on Islands Inhabited by Humans and Domestic Animals. *Conservation Biology* 25: 232–240.
- Pickering J, Norris CA. 1996. New evidence concerning the extinction of the endemic murid *Rattus macleari* from Christmas Island, Indian Ocean. *Australian Mammalogy* 19: 19–25.
- Pigliucci M. 2002. Are ecology and evolutionary biology “soft” sciences? – *Annales Zoologici Fennici* 39: 87–98.
- Prenter J, MacNeil C, Dick JTA, Dinn AM. 2004. Roles of parasites in animal invasions. *Trends in Ecology and Evolution* 19: 385–390.
- Puhakka E. 2012. Tropical rainforest corridor in Madagascar: satellite image –based land cover classification, landscape analysis and forest quality measurements. Master’s Thesis. University of Helsinki. 102 pp.
- Radimilahy C. 1997. Mahilaka, an eleventh to fourteenth century Islamic port: The first impact of urbanism in Madagascar. In *Natural Change and Human Impact in Madagascar* (Goodman, S. M. & Patterson, B. D., eds), pp. 342–377. Smithsonian Institution Press, Washington, DC.
- Rakotozafy LMA. 1996. Etude de la constitution du r egime alimentaire des habitants du site de Mahilika du XI au XIV si ecle  a partir des produits de fouilles arch eologiques. PhD thesis, Universit e d’Antananarivo.
- Ramanamanjato J-B and Ganzhorn JU. 2001. Effects of forest fragmentation, introduced *Rattus rattus* and the role of exotic tree plantations and secondary vegetation for the conservation of an endemic rodent and a small lemur in littoral forest of southeastern Madagascar. *Animal Conservation* 4: 175–183.
- Raoul F, Pleydell D, Qu er e JP, Vaniscotte A, Rieffel D, Takahashi K, Bernard N, Wang J, Dobigny T, Galbreath KE, Giraudoux P. 2008. Small-mammal assemblage response to deforestation and afforestation in central China. *Mammalia* 72: 320–332

- Rickart EA, Baleta DS, Rowe RJ, Heaney LR. 2011. Mammals of the northern Philippines: tolerance for habitat disturbance and resistance to invasive species in an endemic insular fauna. *Diversity and distributions* 17: 530–541.
- Ryan JM, Creighton GK and Emmons LH. 1993. Activity patterns of two species of *Nesomys* (Muridae: Nesomyinae) in a Madagascar rain forest. *Journal of Tropical Ecology* 9: 101–107.
- Santos-Filho M, Peres CA, da Silva DJ, Sanaiotti TM. 2012. Habitat patch and matrix effects on small-mammal persistence in Amazonian forest fragments. *Biodiversity and Conservation* 21: 1127–1147.
- Savidge JE. 1987. Extinction of an island forest avifauna by an introduced snake. *Ecology* 68: 660–668.
- Sax DE, Gaines SD. 2008. Species invasions and extinction: the future of native biodiversity on islands. *Proceedings of the National Academy of Sciences of the United States of America* 105:11490–11497.
- Scott DM, Brown D, Mahood S, Denton B, Silburn A, Rakotondraparany F. 2005. The impacts of forest clearance on lizard, small mammals and bird communities in the arid spiny forest, southern Madagascar. *Biological Conservation* 127:72–87.
- Sneath PHA, Sokal RR. 1973. *Numerical Taxonomy. The Principles and Practice of Numerical Classification*. W.H. Freeman, San Francisco.
- Simberloff D. 2009. Rats aren't the only introduced rodents producing ecosystem impacts on islands. *Biological Invasions* 11: 1735–1742.
- Soarimalala V, Goodman SM. 2011. *Les petits mammifères de Madagascar*. Association Vahatra. Antananarivo, Madagascar. 176 pp.
- Stephenson PJ. 1993. The small mammal fauna of Réserve Spéciale d'Analamazaotra, Madagascar: the effects of human disturbance on endemic species diversity. *Biodiversity and Conservation*. 2: 603–615.
- Stephenson PJ. 1995. Small mammal microhabitat use in lowland rain forest of north-east Madagascar. *Acta Theriologica* 40: 425–438.
- Steppan SJ, Adkins RM and Anderson J. 2004. Phylogeny and divergence-date estimates of rapid radiations in muroid rodents based on multiple nuclear genes. *Systematic Biology*. 53: 533–553.
- Stokes VL, Banks PB, Pech RP, Spratt DM. 2009. Competition in an invaded rodent community reveals black rats as a threat to native bush rats in littoral rainforest of south-eastern Australia. *Journal of Applied Ecology* 46:1239–1247.
- Strong DRJ, Simberloff D, Abele LG and Thistle AB, Eds. 1984. *Ecological communities: conceptual issues and the evidence*. Princeton University Press, Princeton, NJ.
- Thibault J-C, Martin J-L, Penloup A, Meyer J-Y. 2002. Understanding the decline and extinction of monarchs (Aves) in Polynesian Islands. *Biological Conservation* 108: 161–174.
- Tilman D, May RM, Lehman CL, Nowak MA. 1994. Habitat destruction and the extinction debt. *Nature* 371: 65–66.
- Tollenaere C, Brouat C, Duplantier J-M, Rahalison L, Rahelinirina S, Pascal M, Moné H, Mouahid G, Leirs H, Cosson J-F. 2010. Phylogeography of the introduced species *Rattus rattus* in the western Indian Ocean, with special emphasis on the colonization history of Madagascar. *Journal of Biogeography* 37: 398–410.
- Tompkins DM, White AR, Boots M. 2003. Ecological replacement of native red squirrels by invasive greys driven by disease. *Ecology Letters* 6: 189–196.
- Torchin ME, Lafferty KD, Dobson AP, McKenzie VJ, Kuris AM. 2003. Introduced species and their missing parasites. *Nature* 421: 628–630.
- Towns DR, Atkinson IAE, Daugherty CH. 2006. Have the harmful effects of introduced rats on islands been exaggerated? *Biological Invasions* 8: 863–891.

- Vigne JD, Valladas H. 1996. Small mammal fossil assemblages as indicators of environmental change in northern Corsica during the last 2500 years. *Journal of Archaeological Science* 23: 199–215.
- Vitousek P, DAntonio C, Loope L, Westbrooks R. 1996. Biological invasions as global environmental change. *American Scientist* 84: 468–478.
- Vitousek PM, Mooney HA, Lubchenco J, Melillo JM. 1997. Human domination of Earth's ecosystems. *Science* 277: 494–499.
- Wilcove DS, Rothstein D, Dubow J, Phillips A, Losos E. 1998. Quantifying threats to imperilled species in the United States. *BioScience* 48: 607–615.
- Williamson M. 1989. Natural extinction on islands. *Philosophical Transactions of the Royal Society B: Biological Sciences* 325: 457–468.
- Wilson DE. 1991. Mammals of the Tres Mari'as Islands. *The Bulletin of the American Museum of Natural History* 206: 214–250.
- Wirta H. 2009. Dung beetle radiations in Madagascar. PhD Dissertation, University of Helsinki, 124 p.
- Wright PC. 1997. The future of biodiversity in Madagascar: a view from Ranomafana National Park. In: Goodman SM and Patterson BD (eds) *Natural Change and Human Impact in Madagascar*, pp 381–405. Smithsonian Institution Press, Washington, DC