

Technical University of Denmark



A rational approach for -transaminase-catalyzed process design Synthesis of p-Br-1-phenylethylamine

Deslauriers, Maria Gundersen; Lloyd, Richard; Tufvesson, Pär; Woodley, John

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

T. Gundersen, M., Lloyd, R., Tufvesson, P., & Woodley, J. (2013). A rational approach for -transaminase-catalyzed process design: Synthesis of p-Br-1-phenylethylamine. Poster session presented at 1st International Symposium on Transaminase Biocatalysis, Stockholm, Sweden.

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A rational approach for ω -transaminase-catalyzed process design: synthesis of p-Br-1-phenylethylamine






Maria T. Gundersen¹, Richard Lloyd², Pär Tufvesson¹, and John M. Woodley¹

1) Department of Chemical and Biochemical Engineering, DTU, Denmark,

2) Dr. Reddy's Chiretech Technology Centre, Cambridge, UK



INTRODUCTION

Herein we describe a novel rational approach to the design of a ω -transaminase process such that it will fulfill criteria necessary for industrial use. By first determining the fundamental properties of the reaction system, it is possible to suggest appropriate process strategies that may be used to overcome any unfavorable parameters. The ω -transaminase is used as a model system because it is an important enzyme class and developing a systematic methodology would have significant value.



$$K_{eq} = \frac{[\text{product}][\text{co-product}]}{[\text{Acceptor}][\text{donor}]} = \frac{[1,89][1,89]}{[0,11][998,11]} = 1/30$$

Parameter	Acceptor	Product
Water solubility	2mM	15mM
Boiling point	255 °C	?



DISCUSSION

It is clear that the reaction studied exhibits many of the challenges often found with the use of transaminases for the synthesis of pharmaceutical intermediates such as product inhibition, unfavorable thermodynamics and poor solubility. We here propose a number of process strategies which, if adopted additively, may make the process economically viable.

Acknowledgements

Work financed through the European Union 7th Framework Programme (Grant agreement no.: 245144, AMBIOCAS).