

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Biochemistry -- Faculty Publications

Biochemistry, Department of

2008

Plant-like Substitutions in the Large-Subunit Carboxy Terminus of *Chlamydomonas* Rubisco Increase CO₂/O₂ Specificity

Sriram Satagopan

Ohio State University, Columbus, OH

Robert Spreitzer

University of Nebraska - Lincoln, rspreitzer1@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/biochemfacpub>



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#)

Satagopan, Sriram and Spreitzer, Robert, "Plant-like Substitutions in the Large-Subunit Carboxy Terminus of *Chlamydomonas* Rubisco Increase CO₂/O₂ Specificity" (2008). *Biochemistry -- Faculty Publications*. 26. <https://digitalcommons.unl.edu/biochemfacpub/26>

This Article is brought to you for free and open access by the Biochemistry, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Biochemistry -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Research article

Open Access

Plant-like substitutions in the large-subunit carboxy terminus of *Chlamydomonas* Rubisco increase CO₂/O₂ Specificity

Sriram Satagopan^{*1,3} and Robert J Spreitzer²

Address: ¹Department of Microbiology, Ohio State University, Columbus, OH 43210, USA, ²Department of Biochemistry, University of Nebraska, Lincoln, NE 68588-0664, USA and ³DuPont Knowledge Center, ICICI Knowledge Park, Hyderabad 500078, India

Email: Sriram Satagopan^{*} - satagopan.1@osu.edu; Robert J Spreitzer - rspreitzer1@unl.edu

^{*} Corresponding author

Published: 30 July 2008

Received: 28 April 2008

BMC Plant Biology 2008, 8:85 doi:10.1186/1471-2229-8-85

Accepted: 30 July 2008

This article is available from: <http://www.biomedcentral.com/1471-2229/8/85>

© 2008 Satagopan and Spreitzer; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Ribulose-1,5-bisphosphate is the rate-limiting enzyme in photosynthesis. The catalytic large subunit of the green-algal enzyme from *Chlamydomonas reinhardtii* is ~90% identical to the flowering-plant sequences, although they confer diverse kinetic properties. To identify the regions that may account for species variation in kinetic properties, directed mutagenesis and chloroplast transformation were used to create four amino-acid substitutions in the carboxy terminus of the *Chlamydomonas* large subunit to mimic the sequence of higher-specificity plant enzymes.

Results: The quadruple-mutant enzyme has a 10% increase in CO₂/O₂ specificity and a lower carboxylation catalytic efficiency. The mutations do not seem to influence the protein expression, structural stability or the function in vivo.

Conclusion: Owing to the decreased carboxylation catalytic efficiency, the quadruple-mutant is not a "better" enzyme. Nonetheless, because of its positive influence on specificity, the carboxy terminus, relatively far from the active site, may serve as a target for enzyme improvement via combinatorial approaches.

Background

Ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) (EC 4.1.1.39) is limited by a low carboxylation rate and competing oxygenase activity that initiates a wasteful photorespiratory pathway leading to the loss of fixed carbon [1]. The CO₂/O₂ specificity (Ω) of Rubisco is equal to the ratio of catalytic efficiencies (k_{cat}/K_m) for carboxylation (V_o/K_c) to oxygenation V_o/K_o [2]. Net photosynthetic CO₂ fixation would be increased if Rubisco could work faster with greater specificity for CO₂ [1,2]. Because there is natural variation in Rubisco kinetic constants [3], it may be possible to develop genetic engineer-

ing strategies aimed at improving the enzyme by focusing on regions responsible for this variation.

The green alga *Chlamydomonas reinhardtii* is an excellent model for studying plant-like Rubisco enzymes comprised of eight large subunits (~55 kDa, coded by the chloroplast *rbcl* gene) and eight small subunits (~15 kDa, coded by a family of nuclear *rbcs* genes) [1]. Mutants that lack Rubisco function can be maintained with acetate, and both the nuclear *rbcs* and chloroplast *rbcl* genes are amenable to transformation [4,5]. Because *Chlamydomonas* Rubisco has a faster carboxylation rate and lower Ω value

than Rubisco enzymes of flowering plants [3], but shares ~90% sequence identity with flowering-plant enzymes, a phylogenetic approach has been initiated to define the structural basis for differences in catalysis [6,7]. There are only 34 residues in the *Chlamydomonas* large subunit that differ from those of 500 flowering-plant sequences [6]. Changing five of these "phylogenetic" residues to those common to flowering plants, and introducing a spinach small-subunit loop, produced a *Chlamydomonas* Rubisco enzyme with catalytic properties more like those of flowering plants than *Chlamydomonas*, including a 12–17% increase in Ω [7]. However, because the Ω value was not identical to that of flowering-plant Rubisco, there must be other residues that also contribute to the differences in Rubisco catalytic properties.

The loop between β -strand 6 and α -helix 6 of the large-subunit α/β barrel folds over the transition-state analog carboxyarabinitol 1,5-bisphosphate (CABP) [8] (Fig. 1). In this closed conformation, the large-subunit carboxy terminus (residues Trp-462 to Leu-475) packs over loop 6 (Fig. 1) [9,10]. In *Chlamydomonas* Rubisco, substitution of conserved Asp-473, which was proposed to be an essential latch residue [11], with either Ala or Glu did not eliminate catalysis but caused a 14–17% decrease in Ω [12]. Deletion of 10 residues from the carboxy terminus of the *Synechococcus* (cyanobacterium) large subunit caused a 38% decrease in Ω , and a group of four plant-like substitutions E470P/T471A/K474T/L475V was reported to cause a 9% increase in Ω [13]. However, this small increase was close to the experimental error of the assays employed [13]. Lengthening the carboxy terminus beyond residue 475 had no effect on Ω [14].

The *Chlamydomonas* carboxy-terminal residues Asp-470, Thr-471, Ile-472, and Lys-474 comprise a set of previously identified "phylogenetic" residues (Fig. 1A) [6] that differ from the set common to flowering plants (Glu-470, Ala-471, Met-472, and Thr-474). The carboxy terminal residues of *Synechococcus*, studied previously [13], comprise a somewhat different set of residues (Thr-471, Lys-474, and Leu-475) relative to those of flowering plants (Ala-471, Thr-474, and Val-475). Therefore, the carboxy terminus of *Chlamydomonas* was changed via directed mutagenesis and chloroplast transformation to that of spinach Rubisco (Fig. 1B) to see whether the catalytic properties common to Rubisco of flowering plants might be obtained.

Results

Recovery and phenotype of the mutant strain

The residues equivalent to *Chlamydomonas* Asp-470, Thr-471, Ile-472, and Lys-474 are Glu, Ala, Met, and Thr, respectively, in most flowering-plant sequences [6]. The carboxy terminus of the spinach large subunit is identical in length to that of *Chlamydomonas*, and the availability of




Figure 1
Comparison of structural interactions at the carboxy-terminal/loop-6 interface in the large subunit of Rubisco from (A) *Chlamydomonas* (1GK8) [9] and (B) spinach (8RUC) [10]. Residues within 4 Å of the divergent carboxy-terminal residues 470, 471, 472, and 474 are shown as sticks. The carboxy terminus (yellow), loop 6 (red), and part of a loop in the amino-terminal domain of a neighboring large subunit (blue) are drawn as ribbons. Residues not in these three structural regions are colored green. The location of the active site is denoted by Lys-334 and the transition-state analog CABP. Potential hydrogen and ionic bonds are indicated by red and black dotted lines, respectively, connecting the participating atoms.

the spinach crystal structure [10] allows structural comparison (Fig. 1). However, whereas most plant Rubisco enzymes contain a Glu at position 470, this residue is a Pro in spinach Rubisco. Thus, a *Chlamydomonas* D470P/T471A/I472M/K474T mutant enzyme was created to mimic the large-subunit carboxy terminus of spinach Rubisco.

When the *rbcL*-deletion mutant MX3312 was transformed with the *rbcL*-D470P/T471A/I472M/K474T gene, photosynthesis-competent colonies were recovered. Thus, the substituted residues are not essential for Rubisco expression, assembly, or function *in vivo*. Furthermore, the mutant strain was indistinguishable from wild type with respect to growth at either 25 or 35°C, eliminating the possibility that the mutant is temperature conditional (data not shown) [15].

Mutant Rubisco holoenzyme stability

To assess the effect of the mutant substitutions on holoenzyme levels *in vivo*, protein extract was subjected to SDS-PAGE and western-blot analysis. As shown in Fig. 2, the D470P/T471A/I472M/K474T quadruple-mutant strain contains wild-type levels of Rubisco subunits. A wild-type level of holoenzyme could also be purified from the quadruple mutant when cell extract was fractionated on a




Figure 2
Protein expression and holoenzyme levels are unaltered in the D470P/T471A/I472M/K474T quadruple mutant. (A) SDS-polyacrylamide gel electrophoresis and (B) western-blot analysis of total soluble proteins (60 µg/lane) extracted from *Chlamydomonas* wild type (lane 1) and the D470P/T471A/I472M/K474T quadruple mutant (lane 2). The Rubisco large subunit (LS) and small subunit (SS) are indicated.

sucrose-density gradient (data not shown). No difference was detected between the mutant and wild-type enzymes when thermal stability experiments were performed *in vitro* (Fig. 3). Thus, analysis of the catalytic properties of the quadruple-mutant enzyme would not be expected to be influenced by structural instability during the assays.

Kinetic properties of the quadruple-mutant enzyme

To determine whether the spinach-like substitutions can confer catalytic properties mimicking those of the spinach enzyme, kinetic constants were determined for the purified mutant enzyme (Table 1). The Ω value of the D470P/T471A/I472M/K474T quadruple-mutant Rubisco enzyme is 10% higher than that of the wild-type *Chlamydomonas* enzyme. Nonetheless, this value is still 17% lower than that of spinach Rubisco (Table 1). Although the Ω value appears to be altered in the correct direction, the other kinetic constants remain quite different from those of spinach Rubisco (Table 1). The increase in Ω is likely the consequence of an increase in K_o/K_c (Table 1). Due to an apparent decrease in V_o/K_c , the D470P/T471A/I472M/K474T quadruple-mutant enzyme would not be a "better" Rubisco [1].

Discussion

Substitution of the *Chlamydomonas* carboxy-terminal residues with those of spinach, to produce the D470P/T471A/I472M/K474T quadruple mutant, did not influence the structural stability of the enzyme in assays performed *in vitro*.




Figure 3
Carboxy-terminal substitutions do not influence the structural stability of the enzyme in assays performed *in vitro*. Thermal inactivation of Rubisco purified from wild type (○) and the D470P/T471A/I472M/K474T quadruple mutant (●). Rubisco enzymes were incubated at each temperature for 10 min, and then assayed for RuBP carboxylase activity at 25°C. Activities were normalized to the specific activities measured after the 35°C incubation (wild type, 1.7 µmol/min/mg; D470P/T471A/I472M/K474T, 1.2 µmol/min/mg).

Table 1: Kinetic properties of Rubisco purified from *Chlamydomonas* wild type, mutant D470P/T471A/I472M/K474T, and spinach

Kinetic constant	Wild type	Mutant	Spinach
Ω^a ($V_c K_o / V_o K_c$)	61 ± 1	67 ± 2	81 ± 3
V_c^a ($\mu\text{mol/h/mg}$)	119 ± 11	100 ± 7	63 ± 6
k_{cat}^b (s^{-1})	2.7 ± 0.2	2.3 ± 0.2	1.4 ± 0.1
K_c^a ($\mu\text{M CO}_2$)	34 ± 2	40 ± 5	21 ± 1
K_o^a ($\mu\text{M O}_2$)	417 ± 62	611 ± 119	496 ± 40
V_c / K_c^b	3.5	2.5	3.0
K_o / K_c^b	12	15	24
V_c / V_o^b	5.1	4.5	3.4

^a Values are the mean ± (n-1) standard deviation of three or more separate enzyme preparations. ^b Calculated values.

I472M/K474T quadruple-mutant enzyme, causes a 10% increase in Ω , but the other kinetic constants are not like those of spinach Rubisco (Table 1). Thus, these engineered "phylogenetic" residues [6] do not by themselves account for the differences in catalysis between *Chlamydomonas* and spinach Rubisco. However, there should be other residues in the spinach enzyme that complement for the lower values of K_c and K_o (Table 1). Residue 341 (Val in *Chlamydomonas* and Ile in flowering-plant enzymes) is a "phylogenetic" residue in the loop 6, which is in van der Waals contact with the conserved Asp-473 of the carboxy terminus (Fig. 1). Another divergent residue, Arg-305, is also involved in van der Waals contact with the carboxy terminus. The analogous Lys-305 of the spinach enzyme interacts with the carboxy terminus via a single hydrogen-bond (Fig. 1) and a P305K substitution in *Chromatium vinosum* Rubisco causes an 80% increase in carboxylation catalytic efficiency [16]. It is thus likely that the addition of V341I and R305K substitutions to the D470P/T471A/I472M/K474T quadruple mutant would confer kinetic constants that are more like the flowering-plant enzymes.

Engineering of *Synechococcus* Rubisco to produce an E470P/T471A/K474T/L475V mutant enzyme with a spinach-like large-subunit carboxy terminus may also have resulted in a similar, relative increase in Ω [13], but the Ω value of that mutant *Synechococcus* enzyme is still ~10% lower than the Ω value of the *Chlamydomonas* quadruple-mutant enzyme analyzed in the present study. Variation in amino-acid identities at residues 472 and 475 may indicate that these residues are less important than the others. The loss of the E470-R131 and E470-K474 salt bridges in *Synechococcus* Rubisco [17] and D470-R131 and D470-K474 salt bridges in *Chlamydomonas* Rubisco [9] (Fig. 1) may be primarily responsible for the observed increases in Ω [18].

In a recent study, a set of three "phylogenetic" substitutions in the *Chlamydomonas* large subunit (C256F/K258R/I265V), which causes a 10% decrease in Ω [6], was complemented by the addition of two large-subunit phylogenetic substitutions (V221C/V235I) and the loop that resides between β -strands A and B of the spinach small subunit [7,19]. The resultant penta/ABS0 enzyme has a ~15% higher Ω value and all other catalytic properties similar to the spinach enzyme [7]. Although the D470P/T471A/I472M/K474T quadruple-mutant enzyme, analyzed in the present study, does not have kinetic constants like those of spinach Rubisco, it does have a 10% increase in Ω (Table 1). If these carboxy-terminal substitutions could increase the Ω value of the penta/ABS0 enzyme by 10% without substantially altering other kinetic constants, a *Chlamydomonas* enzyme would be generated with kinetic properties indistinguishable from those of spinach Rubisco [7].

Conclusion

Such combinatorial approaches may identify only a small number of regions responsible for the differences in kinetic properties between *Chlamydomonas* and spinach Rubisco. These regions may then serve as suitable targets for DNA shuffling and genetic selection aimed at improving Rubisco [20].

Methods

Strains and culture conditions

Chlamydomonas reinhardtii 2137 *mt+* is the wild-type strain [15]. Mutant MX3312, which has the *rbcl* coding region replaced with the *aadA* gene conferring spectinomycin resistance, was used as the host for chloroplast transformation [12,20]. All strains are maintained in darkness at 25°C on medium containing 10 mM acetate and 1.5% Bacto-agar [15]. For biochemical analysis, cells were grown in 500 ml of liquid acetate medium at 25°C on a rotary shaker at 220 rpm in darkness.

Directed mutagenesis and transformation

Using the plasmid pUC*rbcl*LP as a template [21], the *Chlamydomonas rbcl* gene was PCR-amplified, with the first base of the coding region numbered 1, using forward (bases -169 to -144) and reverse (bases 1404 to 1438) primers. The reverse primer was designed to change codons GAT to CCT, ACT to GCT, ATT to ATG, and AAA to ACA, thereby introducing amino-acid substitutions D470P, T471A, I472M, and K474T, respectively. The reverse primer also contained a 2-base change (TT to AA) following the *rbcl* stop codon that would introduce a *PacI* recognition site. After PCR amplification, the product was digested with *BspEI* and *PacI*, and cloned into the corresponding sites of a modified plasmid containing the same *PacI* site (pUC*rbcl*LP-*PacI*). Chloroplast-gene transformation was performed [22,23], and photosynthesis-compe-

tent colonies were selected on minimal medium with 80 $\mu\text{mol photons/m}^2/\text{s}$. Single-colony isolation, followed by PCR and restriction-enzyme analysis, was performed to ensure homoplasmy of the mutant gene, which was sequenced completely to confirm the mutations.

Biochemical analysis

Cell extracts were prepared from dark-grown cells and subjected to SDS-PAGE with a 7.5–15% polyacrylamide gradient [24,25]. The proteins in the gels were either stained with Coomassie blue [24] or transferred to nitrocellulose membrane and probed with rabbit anti-*Chlamydomonas* Rubisco IgG (0.5 $\mu\text{g/ml}$) [19]. Rubisco-subunit IgGs were detected with goat anti-rabbit IgG/alkaline phosphatase conjugate via chemiluminescence (Amersham Pharmacia Biotech) [25,19].

Rubisco holoenzyme was purified from cell extracts of *Chlamydomonas* and spinach (*Spinacea oleracea*, described previously [7]) by sucrose-gradient centrifugation in assay buffer (50 mM Bicine, pH 8.0, 10 mM NaHCO_3 , 10 mM MgCl_2 , 1 mM dithiothreitol) [26]. The kinetic constants of purified and activated enzymes were determined by measuring the incorporation of acid-stable ^{14}C from $\text{NaH}^{14}\text{CO}_3$ [27]. Ω values of the enzymes (10 $\mu\text{g}/0.5\text{-ml}$ reaction) were determined by assaying carboxylase and oxygenase activities simultaneously with 22 μM [^3H]RuBP (15.8 Ci/mol) and 2 mM $\text{NaH}^{14}\text{CO}_3$ (5 Ci/mol) in 30-min reactions at 25°C [28,29].

Rubisco thermal stability was assayed by incubating purified and activated enzyme (5 μg) at various temperatures for 10 min [30]. The samples were then cooled on ice, incubated at 25°C for 15 min, and used to initiate reactions in 0.5-ml standard assay mixtures containing $\text{NaH}^{14}\text{CO}_3$ (2 Ci/mol) and 0.4 mM RuBP [30].

Authors' contributions

SS contributed to the conceptualization and design of the study, carried out the experiments and data analysis, and drafted the manuscript. RJS conceived of the study, participated in its design and coordination, and made critical revisions to the draft for intellectual content. Both the authors have read and approve this final manuscript.

Acknowledgements

We thank Dr. S. R. Peddi for constructing pUCrbcLP-Pacl. This work was supported by USDA National Research Initiative Grant 2006-35318-17376 and the Nebraska Agricultural Research Division.

References

- Spreitzer RJ, Salvucci ME: **Rubisco: Structure, regulatory interactions, and possibilities for a better enzyme.** *Annu Rev Plant Biol* 2002, **53**:449-475.
- Laing WA, Ogren WL, Hageman RL: **Regulation of soybean net photosynthetic CO_2 fixation by the interaction of CO_2 , O_2 , and ribulose 1,5-diphosphate carboxylase.** *Plant Physiol* 1974, **54**:678-685.
- Jordan DB, Ogren WL: **Species variation in the specificity of ribulose biphosphate carboxylase/oxygenase.** *Nature* 1981, **291**:513-515.
- Spreitzer RJ: **Genetic engineering of Rubisco.** In *The Molecular Biology of Chloroplasts and Mitochondria in Chlamydomonas* Edited by: Rochaix JD, Goldschmidt-Clermont M, Merchant S. Dordrecht, Kluwer Academic Publishers; 1998:515-527.
- Spreitzer RJ: **Role of the small subunit in ribulose-1,5-bisphosphate carboxylase/oxygenase.** *Arch Biochem Biophys* 2003, **414**:141-149.
- Du YC, Peddi SR, Spreitzer RJ: **Assessment of structural and functional divergence far from the large subunit active site of ribulose-1,5-bisphosphate carboxylase/oxygenase.** *J Biol Chem* 2003, **278**:49401-49405.
- Spreitzer RJ, Peddi SR, Satagopan S: **Phylogenetic engineering at an interface between large and small subunits imparts land-plant kinetic properties to algal Rubisco.** *Proc Natl Acad Sci USA* 2005, **102**:17225-17230.
- Andersson I, Taylor TC: **Structural framework for catalysis and regulation in ribulose-1,5-bisphosphate carboxylase/oxygenase.** *Arch Biochem Biophys* 2003, **414**:130-140.
- Taylor TC, Backlund A, Bjorhall K, Spreitzer RJ, Andersson I: **First crystal structure of Rubisco from a green alga, *Chlamydomonas reinhardtii*.** *J Biol Chem* 2001, **276**:48159-48164.
- Andersson I: **Large structures at high resolution: The 1.6 Å crystal structure of spinach ribulose-1,5-bisphosphate carboxylase/oxygenase complexed with 2-carboxyarabinitol biphosphate.** *J Mol Biol* 1996, **259**:160-174.
- Duff AP, Andrews TJ, Curmi PM: **The transition between the open and closed states of Rubisco is triggered by the inter-phosphate distance of the bound biphosphate.** *J Mol Biol* 2000, **298**:903-916.
- Satagopan S, Spreitzer RJ: **Substitutions at the Asp-473 latch residue of *Chlamydomonas* ribulosebiphosphate carboxylase/oxygenase cause decreases in carboxylation efficiency and CO_2/O_2 specificity.** *J Biol Chem* 2004, **279**:14240-14244.
- Gutteridge S, Rhoades DF, Herrmann C: **Site-specific mutations in a loop region of the C-terminal domain of the large subunit of ribulose biphosphate carboxylase/oxygenase that influence substrate partitioning.** *J Biol Chem* 1993, **268**:7818-7824.
- Zhu G, Jensen RG, Bohnert HJ, Wildner GF, Schlitter J: **Dependence of catalysis and CO_2/O_2 specificity of Rubisco on the carboxy terminus of the large subunit at different temperatures.** *Photosynth Res* 1998, **57**:71-79.
- Spreitzer RJ, Mets L: **Photosynthesis-deficient mutants of *Chlamydomonas reinhardtii* with associated light-sensitive phenotypes.** *Plant Physiol* 1981, **67**:565-569.
- Uemura K, Shibata N, Anwaruzzaman M, Fujiwara M, Higuchi T, Kobayashi H, Kai Y, Yokota A: **The role of structural intersubunit microheterogeneity in the regulation of the activity in hysteresis of ribulose 1,5-bisphosphate carboxylase/oxygenase.** *J Biochem* 2000, **128**:591-599.
- Newman J, Gutteridge S: **The X-ray structure of *Synechococcus* ribulose-bisphosphate carboxylase/oxygenase-activated quaternary complex at 2.2-Å resolution.** *J Biol Chem* 1993, **268**:25876-25886.
- Burisch C, Wildner GF, Schlitter J: **Bioinformatic tools uncover the C-terminal strand of Rubisco's large subunit as hot-spot for specificity-enhancing mutations.** *FEBS Lett* 2007, **581**:741-748.
- Karkehabadi S, Peddi SR, Anwaruzzaman M, Taylor TC, Cederlund A, Genkov T, Andersson I, Spreitzer RJ: **Chimeric small subunits influence catalysis without causing global conformational changes in the crystal structure of ribulose-1,5-bisphosphate carboxylase/oxygenase.** *Biochemistry* 2005, **44**:9851-9861.
- Zhu G, Kurek I, True T, Zhang X, Majundar M, Liu L, Lassner M: **Enhancing photosynthesis by improving Rubisco carboxylase activity and specificity, and Rubisco activase thermostability through DNA shuffling.** In *Photosynthesis: Fundamental Aspects to Global Perspectives* Edited by: Van der Est A, Bruce D. Lawrence, KS, Allen Press; 2005:841-843.
- Du YC, Spreitzer RJ: **Suppressor mutations in the chloroplast-encoded large subunit improve the thermal stability of wild-type ribulose-1,5-bisphosphate carboxylase/oxygenase.** *J Biol Chem* 2000, **275**:19844-19847.

22. Finer JJ, Vain P, Jones MW, McMullen MD: **Development of the particle inflow gun for DNA delivery to plant cells.** *Plant Cell Rep* 1992, **11**:323-328.
23. Zhu G, Spreitzer RJ: **Directed mutagenesis of chloroplast ribulosebiphosphate carboxylase/oxygenase: Substitutions at large subunit asparagine 123 and serine 379 decrease CO₂/O₂ specificity.** *J Biol Chem* 1994, **269**:3952-3956.
24. Chua NH: **Electrophoretic analysis of chloroplast proteins.** *Methods Enzymol* 1980, **69**:434-446.
25. Thow G, Zhu G, Spreitzer RJ: **Complementing substitutions within loop regions 2 and 3 of the $\alpha\beta$ -barrel active site influence the CO₂/O₂ specificity of chloroplast ribulose-1,5-bisphosphate carboxylase/oxygenase.** *Biochemistry* 1994, **33**:5109-5114.
26. Spreitzer RJ, Chastain CJ: **Heteroplasmic suppression of an amber mutation in the *Chlamydomonas* chloroplast gene that encodes the large subunit of ribulosebiphosphate carboxylase/oxygenase.** *Curr Genet* 1987, **11**:611-616.
27. Chen Z, Chastain CJ, Al-Abed SR, Chollet R, Spreitzer RJ: **Reduced CO₂/O₂ specificity of ribulose-bisphosphate carboxylase/oxygenase in a temperature-sensitive chloroplast mutant of *Chlamydomonas*.** *Proc Natl Acad Sci USA* 1988, **85**:4696-4699.
28. Jordan DB, Ogren WL: **A sensitive assay procedure for simultaneous determination of ribulose-1,5-bisphosphate carboxylase and oxygenase activities.** *Plant Physiol* 1981, **67**:237-245.
29. Spreitzer RJ, Jordan DB, Ogren WL: **Biochemical and genetic analysis of an RuBP carboxylase/oxygenase-deficient mutant and revertants of *Chlamydomonas reinhardtii*.** *FEBS Lett* 1982, **148**:117-121.
30. Chen Z, Hong S, Spreitzer RJ: **Thermal instability of ribulose-1,5-bisphosphate carboxylase/oxygenase from a temperature-conditional chloroplast mutant of *Chlamydomonas reinhardtii*.** *Plant Physiol* 1993, **101**:1189-1194.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

