

University of Nebraska - Lincoln

## DigitalCommons@University of Nebraska - Lincoln

---

Historical Materials from University of  
Nebraska-Lincoln Extension

Extension

---

1974

### G74-113 A Quick Test for Atrazine Carryover (Revised March 1989)

Alex Martin

*University of Nebraska - Lincoln*, [amartin2@unl.edu](mailto:amartin2@unl.edu)


R.N. Stougaard

*University of Nebraska - Lincoln*

Patrick J. Shea

*University of Nebraska - Lincoln*, [pshea1@unl.edu](mailto:pshea1@unl.edu)

Follow this and additional works at: <https://digitalcommons.unl.edu/extensionhist>

 Part of the [Agriculture Commons](#), and the [Curriculum and Instruction Commons](#)

---

Martin, Alex; Stougaard, R.N.; and Shea, Patrick J., "G74-113 A Quick Test for Atrazine Carryover (Revised March 1989)" (1974). *Historical Materials from University of Nebraska-Lincoln Extension*. 1222.

<https://digitalcommons.unl.edu/extensionhist/1222>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



## A Quick Test for Atrazine Carryover

How can you tell if you have atrazine carryover in your fields? Plants grown in soil samples can tell.

---

*A. R. Martin, Extension Weed Specialist  
R. N. Stougaard, Extension Weed Specialist  
P.J. Shea, Associate Professor, Agronomy*

---

- [Plants Can Tell](#)
- [The Soil Sample](#)
- [A "No Atrazine" Sample Needed](#)
- [Planting and Growing](#)
- [The Symptoms and Crop Choices](#)
- [Why Limit Plant Population When Testing?](#)
- [Bioassay Not Limited to Atrazine](#)

Residues of atrazine may remain in the soil and affect some susceptible crops the next year. Crops most often affected include soybeans, field beans, sugarbeets, alfalfa, oats, wheat and many broadleaf horticultural crops.

Attempts to predict the extent of carryover and damage to sensitive crops the year following atrazine use have been only partially successful. The rate of atrazine disappearance and, therefore, the amount remaining the next year, is affected by soil texture, pH and organic matter content, as well as atrazine application rate and timing, rainfall, and plant growth the previous year.

Chemical analyses for atrazine are complicated, expensive and can be made only in specialized laboratories. In addition, results obtained from chemical analyses do not necessarily reveal whether your crop will be injured.

### Plants Can Tell



**Atrazine injury on wheat. Browning starts at the leaf tip and progresses toward the stem.**

Biological assays, using test crops similar to the field crop to be grown, are practical and can be done with simple equipment found in most homes or offices. The biological assay outlined as follows does not provide an exact measure of the amount of atrazine present in the soil, but it can show you whether sufficient atrazine remains to harm sensitive crops.

## **The Soil Sample**

Get a representative sample of soil from the field suspected of having atrazine residue. Take samples from the top two to three inches in several locations.

Atrazine residue often occurs in patches in a field. Sample enough places to avoid missing areas of high residue, such as turn-around areas, eroded knolls and terrace channels. Plants growing in such areas frequently show the most injury. You should take and analyze separately samples from areas where excessive residue is anticipated.

Collect your soil sample from at least six places in each area sampled. Remember that the assay is only as reliable or representative as the sample you begin with. A total of 10 pounds of soil is required for each sample assayed.

Assays should be run on the moist samples within two days after they are obtained from the field. Samples that cannot be assayed immediately should be stored in a freezer, if possible. Atrazine residue may decrease due to chemical and biological breakdown if samples are stored indoors under warm conditions.

Wet soil should be spread out and allowed to partially dry to a workable condition. If the soil is cloddy, crush so the clods are no larger than wheat seed, but do not pulverize.

### **A "No Atrazine" Sample Needed**

Divide the 10 pounds of soil into two equal portions. Beans or oats will be planted directly into one portion; the other portion will be treated with activated charcoal prior to planting oats or beans.

Add one level teaspoon of activated charcoal to half (five pounds) of the soil. Thoroughly mix the charcoal with the soil. Any atrazine in this soil will be bound to the carbon in the charcoal and will be inactivated. Soil treated in this way provides the equivalent of a soil not treated with atrazine for comparison purposes.

Activated carbon or charcoal can be obtained as a non-prescription item at most drug stores for a small charge. Many garden and hobby shops stock activated charcoal for use in terrariums. For best results use powdered charcoal. Granular charcoal can be powdered manually prior to use, but it is best to purchase powdered charcoal.

Potting soil or soil from a garden or fence row may be substituted for the charcoal-treated soil sample. Be certain the garden or fence row was not treated with a residual herbicide.

### **Planting and Growing**

Flower pots or half-gallon or larger milk cartons are satisfactory for planting and growing. If cartons are used, cut and lay them down for planting purposes. Use the same kind of container for both treatments. Punch holes in the bottoms for drainage. Fill one container using charcoal-treated or uncontaminated soil and another with soil as it came from the field. Properly identify each container.

Atrazine injury symptoms on seedlings should become apparent between eight and 20 days after the plants emerge. Less time is required if high levels of atrazine are present. With temperatures lower than 70°F it will take longer.

Plant six beans (soybeans, field beans or garden beans) or 10 oat seeds in each container. Press or punch the seeds no deeper than 1/4 inch into the soil. Thoroughly wet the soil. Germinate beans at 72°F. Lower temperatures are satisfactory for oats.

Cover the containers with plastic food wrap until plants begin to emerge to help insure favorable moisture conditions during germination. At emergence remove the plastic. Water as needed. Do not let the soil dry out.

Soon after emergence, thin the beans to three plants per container by clipping. Thinning should be done as soon as possible after emergence to reduce the amount of herbicide removed by discarded plants. It is not necessary to thin the oats.

Plant and thin so there is uniform distance between the seed and plants.

Keep the containers in a warm place (70 to 75°F) where they will get as much sunlight as possible. **Sunlight is essential for the development of atrazine injury symptoms.** Artificial light has much less intensity than sunlight and, therefore, is not entirely satisfactory.

## The Symptoms and Crop Choices

When low atrazine levels are present, atrazine symptoms normally appear in two weeks, providing moisture, temperature and light have been favorable. Leaf tips or edges of the older, more mature leaves become chlorotic (yellow-brown). At about the same time a brown speckling may appear near the soybean leaf edges. Eventually chlorosis may spread through the whole leaf. Plants growing in charcoal treated or uncontaminated soil should be normal. **NOTE:** If test plants die in both containers, disease could be the cause.

**Atrazine injury on soybeans. Yellowing around the leaf edge is the first sign. Leaves next turn brown at the outer edges; browning later extends toward the center of the leaf. New leaves appear normal at first but eventually brown with continued atrazine uptake and exposure to sunlight.**



If atrazine symptoms occur, plant the field from which the samples were obtained to an atrazine tolerant crop such as corn or sorghum.

## Why Limit Plant Population When Testing?

Research shows that with too many plants per container, low atrazine residues are not detected. **It is most important that plant numbers be kept low.** Three bean plants or 10 or less oat plants are all that should be used in five to 10 pounds of soil. These numbers more nearly approach field conditions, and provide a more accurate indication of low atrazine levels.

Simply put, when large numbers of plants are present, each plant extracts a very small amount of atrazine from the soil. The amount is so small that it may not be toxic to the bean or oat plants.

When fewer plants are grown in the same amount of soil, the roots have more room to spread out. With more soil per plant, each plant is able to extract more atrazine from the soil and will show greater injury symptoms, or even death, if the atrazine is present in harmful amounts.

## Bioassay Not Limited to Atrazine

The bioassay outlined has been directed to atrazine (AAtrex), the most widely used herbicide in Nebraska. Since atrazine is used so extensively and is relatively long lasting, more carry-over problems have occurred with atrazine.

The same bioassay procedure could be used to determine soil residues of metribuzin (Sencor/Lexone), simazine (Princep), linuron (Lorox), cyanazine (Bladex) and other triazine or substituted urea herbicides.

Similar procedures can be used to test for dinitroaniline herbicides (Prowl, Sonalan, Surflan, Treflan), although injury symptoms and susceptible crops will differ from the triazine and substituted ureas.

The information in this publication is supplied with the understanding that no discrimination and no endorsement by the Nebraska Cooperative Extension is intended.

---

***File G113 under: PESTICIDES, GENERAL***

***B-1, Herbicides***

*Revised March 1989; 12,000 printed.*

*Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Leo E. Lucas, Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources.*

*University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.*