

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Insecta Mundi

Center for Systematic Entomology, Gainesville,  
Florida

---

3-2010

## New species of Cerambycinae (Coleoptera, Cerambycidae) from South America

Maria Helena M. Galileo

*Museu de Ciências Naturais, Fundação Zoobotânica do Rio Grande do Sul, galileo@fzb.rs.gov.br*

Ubirajara R. Martins

*Museu de Zoologia, Universidade de São Paulo, urmsouza@usp.br*

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Entomology Commons](#)

---

Galileo, Maria Helena M. and Martins, Ubirajara R., "New species of Cerambycinae (Coleoptera, Cerambycidae) from South America" (2010). *Insecta Mundi*. 639.

<https://digitalcommons.unl.edu/insectamundi/639>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# INSECTA MUNDI

A Journal of World Insect Systematics

---

0115

New species of Cerambycinae (Coleoptera, Cerambycidae)  
from South America

Maria Helena M. Galileo

Museu de Ciências Naturais, Fundação Zoobotânica do Rio Grande do Sul  
CP 1188, 90001-970, Porto Alegre, RS, Brazil

Ubirajara R. Martins

Museu de Zoologia, Universidade de São Paulo  
CP 42494, 04218-970, São Paulo, SP, Brazil

Date of Issue: March 12, 2010

Maria Helena M. Galileo and Ubirajara R. Martins  
New species of Cerambycinae (Coleoptera, Cerambycidae) from South America  
*Insecta Mundi* 0115: 1-9

**Published in 2010 by**

Center for Systematic Entomology, Inc.  
P. O. Box 141874  
Gainesville, FL 32614-1874 U. S. A.  
<http://www.centerforsystematicentomology.org/>

**Insecta Mundi** is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod taxon. Manuscripts considered for publication include, but are not limited to, systematic or taxonomic studies, revisions, nomenclatural changes, faunal studies, book reviews, phylogenetic analyses, biological or behavioral studies, etc. **Insecta Mundi** is widely distributed, and referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc.

As of 2007, **Insecta Mundi** is published irregularly throughout the year, not as quarterly issues. As manuscripts are completed they are published and given an individual number. Manuscripts must be peer reviewed prior to submission, after which they are again reviewed by the editorial board to insure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

**Managing editor:** Paul E. Skelley, e-mail: [insectamundi@gmail.com](mailto:insectamundi@gmail.com)  
**Production editor:** Michael C. Thomas, e-mail: [insectamundi@gmail.com](mailto:insectamundi@gmail.com)  
**Editorial board:** J. H. Frank, M. J. Paulsen

**Printed copies deposited in libraries of:**

CSIRO, Canberra, ACT, Australia  
Museu de Zoologia, São Paulo, Brazil  
Agriculture and Agrifood Canada, Ottawa, Ontario, Canada  
The Natural History Museum, London, England  
Muzeum I Instytut Zoologii Pan, Warsaw, Poland  
National Taiwan University, Taipei, Taiwan  
California Academy of Sciences, San Francisco, CA, USA  
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA  
Field Museum of Natural History, Chicago, IL, USA  
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

**Electronic copies in PDF format:**

Printed CD mailed to all members at end of year.  
Florida Center for Library Automation: [purl.fcla.edu/fcla/insectamundi](http://purl.fcla.edu/fcla/insectamundi)  
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>  
Goethe Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

**Author instructions** available on the *Insecta Mundi* page at:  
<http://www.centerforsystematicentomology.org/insectamundi/>

Printed Copy	ISSN 0749-6737
On-Line	ISSN 1942-1354
CD-ROM	ISSN 1942-1362

New species of Cerambycinae (Coleoptera, Cerambycidae)  
from South America

Maria Helena M. Galileo

Museu de Ciências Naturais, Fundação Zoobotânica do Rio Grande do Sul  
CP 1188, 90001-970, Porto Alegre, RS, Brazil  
galileo@fzb.rs.gov.br

Ubirajara R. Martins

Museu de Zoologia, Universidade de São Paulo  
CP 42494, 04218-970, São Paulo, SP, Brazil  
urmsouza@usp.br

**Abstract.** New species are described in the tribe Piezocerini: *Gorybia rondonia* **sp. nov.** from Brazil (Rondônia) and *G. bahiensis* **sp. nov.** from Brazil (Bahia); in the tribe Hexoplonini: *Calycibidion rubricolle* **sp. nov.** from Brazil (Bahia); in the tribe Ibidionini, Tropicidina: *Tropidion argentina* **sp. nov.** from Argentina (La Rioja) and *T. boliviensis* **sp. nov.** from Bolivia (Santa Cruz); in the tribe Rhinotragini: *Ommata (Ommata) albitarsis* **sp. nov.** from Brazil (Rondônia); and in the tribe Rhopalophorini: *Cosmisoma viridescens* **sp. nov.** from Brazil (Bahia). To validate the tribal names, *Hexoplon* Thomson, 1864 and *Tropidion* Thomson, 1867, are here designated type genera of Hexoplonini and Tropicidina, respectively.

**Keywords.** Hexoplonini, Ibidionini, Piezocerini, Rhinotragini, Rhopalophorini.

**Resumo.** Novas espécies de Cerambycinae (Coleoptera, Cerambycidae) da América do Sul. Espécies novas descritas em Piezocerini: *Gorybia rondonia* sp. nov. do Brasil (Rondônia) e *G. bahiensis* sp. nov. do Brasil (Bahia). Em Hexoplonini: *Calycibidion rubricolle* sp. nov. do Brasil (Bahia). Em Ibidionini, Tropicidina: *Tropidion argentina* sp. nov. da Argentina (La Rioja) e *T. boliviensis* sp. nov. da Bolívia (Santa Cruz). Em Rhinotragini: *Ommata (Ommata) albitarsis* sp. nov. do Brasil (Rondônia). Em Rhopalophorini: *Cosmisoma viridescens* sp. nov. do Brasil (Rondônia). Para validar os nomes de tribo e subtribo, *Hexoplon* Thomson, 1864 e *Tropidion* Thomson, 1867, são aqui designados gênero-tipos de Hexoplonini e Tropicidina, respectivamente.

**Palavras-chave.** Hexoplonini, Ibidionini, Piezocerini, Rhinotragini, Rhopalophorini.

## Introduction

Based on material from various collections, new species within the tribes Piezocerini, Hexoplonini, Ibidionini (Tropicidina), Rhinotragini and Rhopalophorini are described.

Among Cerambycinae with finely faceted eyes, two species of *Gorybia* Pascoe, 1866 (Piezocerini) are described. This genus was reviewed by Martins (1976), with 32 species at the time. It was once again reviewed by Martins (2006), adding colored photographs for many species. Martins and Galileo (2007a) added *G. montana*, and in 2008, Galileo and Martins described *G. quadrispinosa*.

A new species is described in *Calycibidion* Martins, 1971 (Hexoplonini), which had only included two species so far. Hexoplonini was reviewed by Martins (2006).

Two species are described in *Tropidion* Thomson, 1867 (Ibidionini, Tropicidina) a speciose genus which was reviewed by Martins (1968), when it included 64 species; it was once again reviewed by Martins and Galileo (2007b), comprising 69 species in South America.

Hexoplonini Martins, 2006 and Tropicidina Martins and Galileo, 2007 are *nomina nuda* by article 16.2 of the ICZN (Bousquet et al., 2009). To validate these tribes, *Hexoplon* Thomson, 1864 and *Tropidion* Thomson, 1867, are designated as the type genera of Hexoplonini and Tropicidina, respectively.

In Rhinotragini, one species of *Ommata (Ommata)* White, 1855 is described. This subgenus had so far included seven species (Monné and Hovore, 2005). Concerning Rhopalophorini, one species was added to the genus *Cosmisoma* Audinet-Serville, 1834, which was reviewed by Monné and Magno (1988).

## Material

The acronyms throughout the text correspond to **ACMS**, American Coleoptera Museum, San Antonio, Texas, United States of America; **FSCA**, Florida State Collection of Arthropods, United States of America; **MNKM**, Museo Noel Kempff Mercado, Santa Cruz Department, Bolivia; **MZSP**, Museu de Zoologia da Universidade de São Paulo, São Paulo State, Brazil; **UEFS**, Universidade Estadual de Feira de Santana, Bahia State, Brazil.

## Piezocerini

### *Gorybia rondonia* Galileo and Martins, new species

(Figure 1)

**Description.** General integument dark reddish-brown (Figure 1). Frons and vertex with microsculptured alveoli. Upper ocular lobes with three rows of ommatidia. Antennal tubercles distant from each other and rounded at the top. Antennae almost reaching the elytral apex. Scape densely punctate. Antennomere III not projecting on the external apex; IV to X subserrate.

Prothorax rounded at sides. Pronotum (female), entirely alveolated except for a small longitudinal area at the basal groove; alveolus microsculptured. Prosternal process recurved. Mesosternum punctate and microsculptured. Metasternum shining at central area and microsculptured at the lateral-anterior regions.

Elytra with shining integument, densely punctate; at the basal one-half, gradually sparser towards the apex; erect hairs organized in five longitudinal rows. Elytral apices emarginated, having a short spine on the external side and wide projection on the internal one.

Anterior coxal cavities narrowly open behind. Meso- and metatibiae not expanded at the apex. Urosternites shining with microsculptured sides.

Measurements in mm. Total length, 11.8; prothorax length, 2.5; largest prothoracic width, 2.4; elytra length, 8.3; humeral width, 3.2.

**Type material.** Female holotype, BRAZIL, *Rondônia*: Mutum-Paraná (left margin of the Paraná River), 19.11.2003, without name of collector, light trap (MZSP).

**Etymology.** The specific name is a noun in apposition and refers to the State of Rondônia.

**Diagnosis.** *Gorybia rondonia* sp. nov. is similar to *G. quadrispinosa* Galileo and Martins, 2008 by emarginate elytral apices, dark-reddish brown coloring, antennomeres IV to X subserrated, and upper ocular lobes with three rows of ommatidia. It is distinguished by having linear middle and posterior tibiae, which do not expand towards the extremity (without an external projection). In *G. quadrispinosa*, the meso- and metatibiae projecting at the apex.


### *Gorybia bahiensis* Galileo and Martins, new species

(Figure 2)

**Description.** General coloring brown; antennae, dorsal area of elytra and urosternites, reddish-brown. Frons and vertex with microsculptured alveoli. Antennal tubercles slightly projecting and acute. Upper ocular lobes with three rows of ommatidia. Antennae reach the apical one-third of the elytra. Scape punctate and microsculptured. Antennomere III not expanded towards the apex; antennomeres IV to X with an external projection at the apex, which becomes increasingly pronounced towards the apical antennomeres; antennomere XI appendiculate.

Prothorax rounded at sides; basal constriction pronounced. Pronotum (Figure 2) completely alveolated, with long hairs and a shallow groove at the centre of the base. Prosternum smooth in the center and alveolated on the sides at the anterior regions; prosternal process not projecting. Mesosternum and mesepimera with shallow alveoli. Metasternum smooth in center; anterior region microsculptured at sides.





**Figure 1-5.** Habitus. 1) *Gorybia rondonia*, female holotype, length 11.8 mm. 2) *Gorybia bahiensis*, female holotype, length 6.4 mm. 3) *Calycibidion rubricolle*, male holotype, length 7.9 mm. 4) *Tropidion argentina*, male holotype, length 10.6 mm. 5) *Tropidion boliviensis*, female holotype, length 11.4 mm.

Elytra densely punctate at base, becoming gradually sparser towards the apex. Long and abundant hairs throughout the entire surface. Elytral apices emarginated with an external small spine and slightly projecting at sutural angle.

Meso- and metatibiae not expanded at apex. Urosternites with punctures and long, sparse hairs.

Measurements in mm. Total length, 6.4; prothorax length, 1.4; largest prothoracic width, 1.6; elytra length, 4.3; humeral width, 1.6.

**Type material.** Female holotype, BRAZIL, *Bahia*: São Félix do Coribe-Coribe, (km 24, 13°33'S 4415'W, 493 m), 07.XII.2007, F. Bravo col. (UEFS).

**Etymology.** The specific name is an adjective allusive to the State of Bahia.

**Diagnosis.** *Gorybia bahiensis* sp. nov. is similar to *G. adiaphora* Martins, 1976 in general coloring, the pronotum having a shallow center-basal groove, elytral apices with a small spine at sutural angle and metatibiae not expanded towards the apex. It is distinguished by its denser pilosity, with long elytra hairs, which cover the entire surface (over five rows); antennomere III not projecting at the external apex; and urosternites punctate with long hairs. On *G. adiaphora*, the elytral hairs are organized in five rows; antennomere III is projecting at the external extremity and urosternites are not punctate.

#### Hexoplonini Galileo and Martins, new tribe

Hexoplonini Martins, 2006: 22; Bousquet et al. 2009: 49 (*nomem nudum*).

**Type genus.** *Hexoplon* Thomson, 1864, **here designated.**

**Description.** Antennal scape curved inwards. Flagellomeres multicarinate or sulcate. Prothorax curved. Anterior coxal cavities with articular surface. Metafemurs spined at apex. The tribe is more completely described in Martins (2006).

#### *Calycibidion rubricolle* Galileo and Martins, new species

(Figure 3)

**Description.** Head black. Frons smooth with depressed areas in front of the antennal tubercles; longitudinal corrugations between the tubercles. Vertex sparsely punctate. Upper ocular lobes with three rows of ommatidia. Antennae black surpassing the elytral apices in the middle of antennomere IX. Scape microsculptured with sparse punctures.

Prothorax reddish. Pronotum with a brownish band along the anterior margin; punctures sparse. Prothorax and prosternum sides punctate. Mesosternum and metasternum reddish. Mesepisterna, mesepimera, metepisterna and a narrow band on the sides of the metasternum microsculptured. Metasternum central region punctate with long, sparse hairs.

Elytra (Figure 3) reddish-orange until slightly after the middle and black in the remaining portion. Coarsely, deeply and densely punctate throughout the entire surface, with erect and long hairs, organized in rows. Elytral apices emarginated; marginal spine longer and sharper than the sutural one.

Legs black, except coxae and base of femur, which are reddish; apical tarsomeres reddish-brown. Femurs with sparse punctures. Tibiae hairy and carinated.

Urosternites black, except at the basal half of I, which is reddish-brown; sparsely punctate with long hairs.

Measurements in mm. Total length, 7.9-8.6; prothorax length, 1.8-1.9; largest prothoracic width, 1.2-1.3; elytra length, 5.0-5.7; humeral width, 1.7-1.8.

**Type material.** Male holotype, BRAZIL, *Bahia*: São Félix do Coribe-Coribe, (km 24, 13°33'S 4415'W, 493 m), 07.XII.2007, A. M. Silva-Neto col. (UEFS); male paratype, same data of the holotype, F. Bravo col. (MZSP).

**Etymology.** Latin, *rubrus* = red; *collum* = neck; allusion to the red prothorax.

**Diagnosis.** *Calycibidion rubricolle* sp. nov. is similar to *C. turbidum* Napp and Martins, 1985 concerning the elytral apices with two spines. It is distinguished by its black head and reddish pronotum; black antennae; black legs with coxae and femur bases reddish. The head and prothorax of *C. turbidum* are black; the pedicels, the flagellomeres and the legs are brownish.

## Ibidionini

### **Tropidiina Galileo and Martins, new subtribe**

Tropidina Martins, 2007b: 7; Bousquet et al. 2009: 49 (*nomem nudum*).

**Type genus.** *Tropidion* Thomson, 1867, **here designated**.

**Description.** Antennal scape pyriform, sulcate basally. Flagellomeres not enlarged. Articles III and IV equal in length. Pronotum usually with tubercles. Anterior coxal cavities open behind. Anterior coxae without articular surface. The subtribe is more completely described in Martins (2007b).

### ***Tropidion argentina* Galileo and Martins, new species**

(Figure 4)

**Description.** Head reddish-brown, covered by a yellowish pilosity. Antennal tubercles projecting. Antennae reddish-brown, reaching elytral apices (males) at the tip of antennomere VII. Scape subpyriform, without a basal groove. Antennomere III carinate, slightly longer than IV.

Prothorax reddish-brown, longer than wide. Pronotum covered with sericeous pubescence, except in a center-longitudinal area and in a longitudinal band on each side; oblique branches begin at the anterior extremity of lateral bands, reaching the anterior one-third of the central-longitudinal band. Middle of pronotum with a central tubercle and a discrete gibosity on each side. Lateral portions of the prothorax smooth and shining. Prosternum with pubescence bands, prolonged up to the middle. Mesosternum and metasternum reddish. Scutellum covered by a whitish pubescence.

Elytra (Figure 4) reddish-brown; each one with a lateral yellowish spot on the anterior one-third. Numerous hairs throughout the entire elytra, poorly organized in rows. Elytral apices rounded.

Femora reddish-brown, clearer at the peduncles and sparsely pubescent. Urosternites reddish-brown.

Measurements in mm. Total length, 10.6; prothorax length, 2.4; largest prothoracic width, 1.6; elytra length, 7.0; humeral width, 2.0.

**Type material.** Male holotype, ARGENTINA, *La Rioja*: Castro Barros (Santa Vera Cruz, 28°67' S 66° 96'W, 1600 m), 15.II.2005, L. Stange col. (FSCA).

**Etymology.** Latin, *rubrus* = red; *collum* = neck; allusion to the red prothorax.


**Diagnosis.** *Tropidion argentina* sp. nov. is similar to *T. intermedium* (Martins, 1962) concerning unarmed elytral apices, but it is distinguished by its general reddish-brown coloring, having a yellow spot on the elytra, and by the pronotum, which presents three tubercles. On *T. intermedium*, the elytra are yellowish and without spots, the appendices are reddish and the pronotum presents five tubercles.

### ***Tropidion boliviensis* Galileo and Martins, new species**

(Figure 5)

**Description.** Head black, densely pubescent. Upper ocular lobes with three rows of ommatidia. Vertex punctate between the antennal tubercles and upper ocular lobes. Antennae (female) reach elytral apices





**Figure 6-7.** Habitus. 6) *Ommata (O.) albitarsis*, female holotype, length 8.5 mm. 7) *Cosmisoma viridescens*, male holotype, length 7.3 mm.

approximately at the tip of antennomere VII. Scape reddish-black, finely punctuate, pyriform, grooved at base.

Prothorax black and reddish at the base. Pronotum densely pubescent, except at the anterior one-fourth portion (where the pubescence is very sparse) and on the central tubercle. Five pronotal tubercles slightly apparent on a dorsal view, with the exception of the central tubercle, more visible through a side view. Prothorax sides pubescent, anterior margin shining. Prosternum with “V-shaped” pubescence at the posterior half. Mesosternum and metasternum reddish, covered by a whitish pubescence.

Elytra (Figure 5) with a reddish anterior half and black apical half; each having a whitish, oblique, dorsal spot on the anterior half and an oblique band between the dominant colours; this band turns forward near the margin, through a short prolongation. Rough punctures near the scutellum and along the suture. The reddish portion of the elytra is also punctate. Each elytron with five longitudinal rows of hairs. Elytral apices with an external long spine.

Femora reddish. Tips of metafemora slightly projecting. Tibiae and tarsi black. Urosternite I reddish; II to V black.

Measurements in mm. Total length, 11.4; prothorax length, 2.6; largest prothoracic width, 1.9; elytra length, 7.8; humeral width, 2.5.

**Type material.** Female holotype, BOLIVIA, *Santa Cruz*: Buena Vista (Hotel Flora and Fauna, 4,6 km SSE) 22-25.X.2007, Galileo and Martins col. (MNKM).

**Etymology.** The epithet is a Latin adjective referring to the native country (Bolivia).

**Diagnosis.** *Tropidion boliviensis* sp. nov. is similar to *Tropidion fairmairei* (Gounelle, 1909) by the coloring pattern and dense pronotum pubescence. It differs through its upper ocular lobes, with three rows of ommatidia, pronotum glabrous near the anterior margin, with a pronounced and glabrous centro-posterior tubercle, elytra with rough punctures and its anterior spot on the elytra, which prolongs towards the margin without reaching it.

In *T. fairmairei*, the superior ocular lobes present four rows of ommatidia, the pronotum is pubescent on the sides of the anterior one-third, the central tubercle is barely projecting, the elytra do not present rough punctures and the anterior elytra spot is rounded.

## Rhinotragini

### *Ommata (Ommata) albitarsis* Galileo and Martins, new species

(Figure 6)

**Description.** Head reddish; antennal tubercles black. Frons moderately punctate (32 x), covered by a shining, golden-yellow pubescence. Anterior region of the vertex, between the antennal tubercles, with same kind of pubescence. Occiput punctate. Malar area shorter than the lower ocular lobe (female).

Gula black on anterior portion and reddish on posterior region, with punctures more concentrated on the sides. Antennae reach elytral apices, almost at the apex of antennomere VIII. Scape black and punctate. Antennomeres III-VIII black and gradually thicker towards the apical segments; antennomeres IX-X white; antennomere XI with white basal half and a black apical one.

Prothorax (Figure 6) reddish. Gradually wider sides towards the base; the widest point is next to the basal constriction. Pronotum and sides of the prothorax punctate. Prosternum with a black band next to each procoxa; the posterior half is covered by a whitish pubescence. Mesosternum and metasternum black covered by a whitish pubescence, except metepisternum, with a slight metallic green shining. Sides of the metasternum, mesepisternum and metepisternum punctate. Scutellum black with a whitish pubescence.

Elytra metallic-green, indistinctly blue at the basal region; small reddish portions at sides of scutellum; elytral margins slightly narrower towards the apex; densely punctate throughout the entire surface. Elytral apices rounded and reaching the base of urosternite V.

Protrochanters reddish. Profemora black. Mesotrochanters reddish, mesofemora black with a narrow, reddish basal area. Metatrochanters reddish. Metafemora with white pedunculus and a black clava.

Tibiae, pro- and mesotarsi black. Metatarsomeres I-IV white; metatarsomere V white at the base and reddish at the apex. Urosternites black covered by a whitish pubescence.

Measurements in mm. Total length, 8.5; prothorax length, 2.1; largest prothoracic width, 1.6; anterior prothoracic width, 1.1; elytra length, 6.0; humeral width, 1.9.

**Type material.** Female holotype, BRAZIL, *Rondônia*: Porto Velho (Parque Ecológico), 3.VIII.2008, without name of collector Malaise trap, “terra firme” (MZSP).

**Etymology.** Latin, *albi* = white; Greek, *tarsos* = tarsi; allusion to white metatarsi.

**Diagnosis.** Concerning antennomeres IX-X white, prothorax reddish, elytra metallic-green and metatarsomeres I-IV white, *Ommata* (*O.*) *albitarsis* sp. nov. may be compared with *Ommata* (*O.*) *paradisiaca* Tippmann, 1953. It differs by antennomere XI white with apical half black; prosternum, with a black band next to each procoxa; mesofemora black with a narrow basal area reddish and elytra unicolor (excepting the reddish basal line beside the scutellum).

Zajciw (1960:94) described the female of *O. (O.) paradisiaca*, previously recognized only through the male, with the differential characters: antennomeres IV to VI black and antennae surpassing the elytral apices by their last two articles (the remaining are secondary sexual characters). In *O. (O.) paradisiaca*, antennomere XI is entirely white, the prosternum is entirely black, the basal halves of mesofemora are white and the elytra present a yellowish longitudinal band, which gradually narrows from the base to the middle.

## Rhopalophorini

### *Cosmisoma viridescens* Galileo and Martins, new species

(Figure 7)

**Description.** Head with a metallic-green integument. Frons and vertex punctate; punctures more sparse on the vertex. Antennae reach the elytra apices at the apical one third of antennomere V. Scape metallic-blue and irregularly punctate. Irregular fringe of long hairs on the internal edge of the scape, pedicel and antennomeres III-V. Apical region of antennomere V with a small tuft of long hairs on the internal edge.

Prothorax reddish with anterior constriction and narrow basal region metallic-green; sides discretely rounded. Pronotum glabrous, punctate except for a center-longitudinal band; the greenish areas transversely striate along the posterior edge and microcorrugations along the anterior edge. Scutellum punctate.

Elytra (Figure 7) metallic-green, punctate with a wide, dorsal and golden pubescence band, from the base to the apex. Lateral margins slightly narrowed towards the apex. Elytral apices acuminate.

Ventral surface metallic-green with dense sericeous pubescence. Metepisterna reddish. Legs with sparse sericeous pubescence and a few long hairs on the tibiae. Mesocoxae, metacoxae and bases of metafemora reddish. Metatibiae curved, without tuft of hairs.

Measurements in mm. Total length, 7.3; prothorax length, 1.7; largest prothoracic width, 1.4; elytra length, 4.7; humeral width, 1.8.

**Type material.** Male holotype, BRAZIL, *Bahia*: São Félix do Coribe-Coribe, (km 24, 13°33'S 44°15'W, 493 m), 07.XII.2007, F. Bravo col. (UEFS).

**Etymology.** Latin, *viridescens* = which becomes green; allusion to the integument color.

**Diagnosis.** *Cosmisoma viridescens* sp. nov. could be compared with *C. gratum* Monné and Magno, 1988 by antennomere V, with small hair tuft on the apex; prothorax reddish; antennae and legs black, with discrete metallic-blue reflexes; elytra unicolored with metallic-green gloss and the metatibiae with hair tufts. It is distinguished by its black head and reddish-orange pronotum with a metallic-green band along the margins. The head and prothorax of *C. gratum* are completely reddish.

## Acknowledgments

We thank Alexandre de Almeida e Silva (Universidade Federal de Rondônia) for donating the holotypes to the MZSP; Freddy Bravo (UEFS), James Wappes (ACMS) for the loan of specimens; Eleandro Moysés (fellowship IC/CNPq/FZB) for the photographs and Rejane Rosa for the color illustration, both from Museu de Ciências Naturais, Fundação Zoobotânica do Rio Grande do Sul; and Gino Nearn and Steve Lingafelter for revision of the manuscript and their helpful suggestions. To Conselho Nacional de Desenvolvimento Científico e Tecnológico for the fellowships.

## Literature Cited

- Bousquet, Y. D. J. Heffern, P. Bouchard, and E. H. Nearn. 2009.** Catalogue of family-group names in Cerambycidae (Coleoptera). *Zootaxa* 2321: 1-80.
- Galileo, M. H. M., and U. R. Martins. 2008.** Novos Cerambycinae (Cerambycidae) da Região Neotropical. *Papéis Avulsos Zoologia* 48(7): 49-54.
- Martins, U. R. 1968.** Monografia da tribo Ibdionini (Coleoptera, Cerambycinae). Parte II. *Arquivos de Zoologia* 16(2): 321-630.
- Martins, U. R. 1976.** Sistemática e evolução da tribo Piezocerini (Coleoptera, Cerambycidae). *Arquivos de Zoologia* 27(3-4): 165-370.
- Martins, U. R. 2006.** Tribo Hexoplioni. p. 21-211. *In*: U. R. Martins (org.). *Cerambycidae sul-americanos* (Coleoptera). *Taxonomia Volume 8*. Sociedade Brasileira de Entomologia; São Paulo. II+234 p.
- Martins, U. R., and M. H. M. Galileo. 2007a.** Novos Cerambycidae (Coleoptera) da Coleção Odette Morvan, Kaw, Guiana Francesa. *Papéis Avulsos Zoologia* 47(14): 175-179.
- Martins, U. R., and M. H. M. Galileo. 2007b.** Tribo Ibdionini, subtribo Tropicidina. p. 1-176. *In*: U. R. Martins (org.). *Cerambycidae sul-americanos* (Coleoptera). *Taxonomia Volume 9*. Sociedade Brasileira de Entomologia; São Paulo. II+350 p.
- Monné, M. A., and F. T. Hovore. 2006.** Checklist of the Cerambycidae, or longhorned wood-boring beetles, of the Western Hemisphere. *BioQuip; Rancho Dominguez*. 394 p.
- Monné, M. A., and P. R. Magno. 1988.** Gênero *Cosmisoma* Audinet-Serville, 1834: notas, novas taxa e chave para espécies sul-americanas (Coleoptera, Cerambycidae, Rhopalophorini). *Boletim do Museu Nacional* 319: 1-27.
- Zajciw, D. 1960.** Estudos sobre longicórneos neotrópicos II (Coleoptera, Cerambycidae). *Revista Brasileira de Biologia* 20(1): 93-98.

Received January 8, 2010; Accepted January 28, 2010.

