

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

7 - Seventh Eastern Wildlife Damage
Management Conference (1995)

Eastern Wildlife Damage Control Conferences

November 1995

PREY AVAILABILITY AND SELECTION BY MOUNTAIN LIONS IN ARAVAIPA-KLONDYKE AREA OF ARIZONA

Stan C. Cuning

Research Branch, Arizona Game and Fish Department

Follow this and additional works at: <https://digitalcommons.unl.edu/ewdcc7>



Part of the [Environmental Health and Protection Commons](#)

Cuning, Stan C., "PREY AVAILABILITY AND SELECTION BY MOUNTAIN LIONS IN ARAVAIPA-KLONDYKE AREA OF ARIZONA" (1995). *7 - Seventh Eastern Wildlife Damage Management Conference (1995)*. 5. <https://digitalcommons.unl.edu/ewdcc7/5>

This Article is brought to you for free and open access by the Eastern Wildlife Damage Control Conferences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 7 - Seventh Eastern Wildlife Damage Management Conference (1995) by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

PREY AVAILABILITY AND SELECTION BY MOUNTAIN LIONS IN ARAVAIPA-KLONDYKE AREA OF ARIZONA

STAN C. CUNNING, Research Branch, Arizona Game and Fish Department, 2221 W. Greenway Road, Phoenix, AZ
85023

ABSTRACT: Prey selection by mountain lions (*Felis concolor*) in the Aravaipa-Klondyke area (2,000 km²) in southeastern Arizona was studied from February 1991 through September 1993. Overall diet from frequency of occurrence as determined from 370 scats was: 48% deer (white-tailed and mule deer combined), 34% cattle, 17% javelins, 6% rabbit (cottontail and jackrabbit), 4% rodent, and 2% desert bighorn. Using a correction factor developed by Ackerman et al. (1984), we also estimated percent biomass and proportion of individuals killed. With respect to biomass consumed, cattle was 44%, deer 40%, javelins 10.9%, rabbits 2.9%, and rodents 0.02%. Based on weights of prey consumed, proportion of individuals eaten changed to rabbits 52.7%, deer 16.3%, rodents 12%, javelinas 10% cattle 8%, and desert bighorn 0.5%. Comparisons with availability as determined from four separate four-day double-count helicopter surveys found that lions selected calves (ate more than expected based on availability alone), killed and ate less deer than expected, and javelins as would be expected.

Proc. East. Wildl. Damage Meant, Conf: 7:130 1997.