

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications, Department of
Mathematics

Mathematics, Department of

2005

Matrix Model Superpotentials and Calabi–Yau Spaces: An A-D-E Classification

Carina Curto

University of Nebraska - Lincoln, ccurto2@math.unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/mathfacpub>



Part of the [Mathematics Commons](#)

Curto, Carina, "Matrix Model Superpotentials and Calabi–Yau Spaces: An A-D-E Classification" (2005).
Faculty Publications, Department of Mathematics. 41.
<https://digitalcommons.unl.edu/mathfacpub/41>

This Article is brought to you for free and open access by the Mathematics, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Mathematics by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Copyright © 2008 by Carina Curto
All rights reserved

MATRIX MODEL SUPERPOTENTIALS AND CALABI–YAU SPACES: AN A-D-E CLASSIFICATION

by

Carina Curto

Department of Mathematics
Duke University

Date: _____

Approved:

David R. Morrison, Supervisor

M. Ronen Plesser

Paul S. Aspinwall

William L. Pardon

Dissertation submitted in partial fulfillment of the
requirements for the degree of Doctor of Philosophy
in the Department of Mathematics
in the Graduate School of
Duke University

2005

ABSTRACT

(Mathematics)

MATRIX MODEL SUPERPOTENTIALS AND
CALABI-YAU SPACES: AN A-D-E CLASSIFICATION

by

Carina Curto

Department of Mathematics
Duke University

Date: _____

Approved:

David R. Morrison, Supervisor

M. Ronen Plesser

Paul S. Aspinwall

William L. Pardon

An abstract of a dissertation submitted in partial fulfillment of the
requirements for the degree of Doctor of Philosophy
in the Department of Mathematics
in the Graduate School of
Duke University

2005

Abstract

We use F. Ferrari’s methods relating matrix models to Calabi-Yau spaces in order to explain Intriligator and Wecht’s ADE classification of $\mathcal{N} = 1$ superconformal theories which arise as RG fixed points of $\mathcal{N} = 1$ SQCD theories with adjoints. The connection between matrix models and $\mathcal{N} = 1$ gauge theories can be seen as evidence for the Dijkgraaf–Vafa conjecture. We find that ADE superpotentials in the Intriligator–Wecht classification exactly match matrix model superpotentials obtained from Calabi-Yau’s with corresponding ADE singularities. Moreover, in the additional \widehat{O} , \widehat{A} , \widehat{D} and \widehat{E} cases we find new singular geometries. These ‘hat’ geometries are closely related to their ADE counterparts, but feature non-isolated singularities. As a byproduct, we give simple descriptions for small resolutions of Gorenstein threefold singularities in terms of transition functions between just two coordinate charts. To obtain these results we develop techniques for performing small resolutions and small blow-downs, including an algorithm for blowing down exceptional \mathbb{P}^1 ’s. In particular, we conjecture that small resolutions for isolated Gorenstein threefold singularities can be obtained by deforming matrix factorizations for simple surface singularities – and prove this in the length 1 and length 2 cases.

A mi tío, Pompilio Zigrino, por despertar mi interés en la física.
To my professor, Vincent Rodgers, for ten years of pedagogy and support.
A mis padres, Inés y Raúl, por darme el mundo y mucho más.

Acknowledgements

First and most importantly, I would like to thank my advisor, Dave Morrison – it has been amazing to work with you.

Thank you Ronen for introducing me to strings. Thanks to Ilarion and Sven for learning with me and for teaching me so many beautiful things. Thank you Anda and Janice – it would have been ten times harder without you! Thanks to Carly and Pedro for turning the whole world into playdough. And a grateful thanks to Ryan and Kris, who had to live with me. To Vladimir it goes without say.

Gracias a Julio Cortázar, Che Guevara, Frida Kahlo, Subcomandante Marcos, y Evita Perón por ser figuras facinantes y controversiales. Me han distraído cuando más lo he necesitado, y me mantienen conciente del continente.

Finally, I would like to acknowledge Shakira, Belle & Sebastian, Elliott Smith, Tori Amos, Gal Costa, Manu Chao, and all of the Russian music I listened to while writing.

This work is supported by NSF and VIGRE graduate fellowships.

Contents

Abstract	iv
Acknowledgements	vi
List of Tables	xi
List of Figures	xii
1 Introduction	1
2 Ferrari's construction	15
2.1 A bit about matrix models.	15
2.2 The geometric framework	16
2.2.1 Step 1: Construction of resolved Calabi-Yau	18
2.2.2 Step 2: Locating the \mathbb{P}^1 's	20
2.2.3 Step 3: Finding the blow-down map in Ferrari's examples	20
2.3 Example: A_k	23
3 Preliminary computations	25
3.1 Perturbations which contribute to superpotential	26
3.2 Ferrari's example 2	28
3.3 Superpotentials which change bundle structure	31
3.4 Classification of perturbation terms	34
3.5 Resolution of the ideal sheaf	35
4 Blowups for surface singularities	37
4.1 How to blow up a single node	37
4.2 The A_{n-1} story	39

4.2.1	Invariant theory	39
4.2.2	Matrix of relations	40
4.2.3	Summary of results	40
4.2.4	The blowup	41
4.3	The D_{n+2} story	43
4.3.1	Invariant theory	43
4.3.2	Matrix of relations	44
4.3.3	Summary of results	47
4.3.4	How to use this to blow up	48
4.3.5	The blowup	49
5	Small resolutions	52
5.1	“Bottom-up” approach	52
5.2	The A_{n-1} story	53
5.2.1	Invariant theory	53
5.2.2	Matrix factorization with deformation	54
5.2.3	The blowup	54
5.2.4	Identification of superpotential in A_1 case	55
5.3	The D_{n+2} story	58
5.3.1	Invariant theory	58
5.3.2	The Tyurina blowup	59
5.3.3	Comparison with non-deformed version	60
5.3.4	Matrix of relations for the full deformation	62
5.3.5	The blowup	66
5.4	Difficulties with the approach	69

6	Small blow-downs from global holomorphic functions	70
6.1	“Top-down” approach	70
6.2	The Intriligator–Wecht classification	71
6.3	The problem	72
6.4	Reformulation of the problem	73
6.5	The algorithm	74
6.5.1	A shortcut	76
6.5.2	Implementation	76
7	Length 1	77
7.1	The Case A_k	78
7.2	A puzzle	80
8	Length 2	82
8.1	The Case D_{k+2}	83
8.2	Transition functions for more general length 2 blowup	86
9	Length 3	89
9.1	The Case E_7	90
9.2	Matrix factorization for a length 3 singularity	93
10	Other ADE cases	94
10.1	The Case E_6	94
10.2	The Case E_8	96
11	The ‘Hat’ cases	98
11.1	The Case \widehat{O}	99
11.2	The Case \widehat{A}	100
11.3	The Case \widehat{D}	102

11.4 The Case \widehat{E}	104
11.5 Comparison with ADE cases	107
12 Future directions	109
A Maple code for blow-down algorithm	111
A.1 Description of the code	111
A.2 Maple code	113
Bibliography	118
Biography	120

List of Tables

1.1	Gorenstein threefold singularities in preferred versal form	6
1.2	Superpotentials corresponding to length 1, length 2, and length 3 singularities.	11
1.3	Geometries for the ‘Hat’ cases.	12
1.4	Geometrically equivalent superpotentials.	13
3.1	Bundle-changing perturbation terms in the 2-matrix model.	34
3.2	Summary of superpotential-contributing perturbation terms.	35
6.1	Identification of corresponding resolved geometries	72

List of Figures

1.1	The big picture	2
1.2	The 6 types of Gorenstein threefold singularities	7
1.3	The blowup of a node leaves residual surface singularities	8
1.4	Intriligator–Wecht Classification of RG Fixed Points	9
2.1	Ferrari’s diagram [12, page 631].	16
2.2	The triple of Calabi-Yau spaces corresponding to a single matrix model. . .	17
4.1	Residual singularities from the blowup of an A_6 node	42
4.2	Residual singularities from the blowup of a D_8 node	51
5.1	Small resolution via blowup of a length 1 node	55
5.2	The Tyurina blowup	61
5.3	Small resolution via blowup of a length 2 node	68
6.1	Intriligator–Wecht Classification of RG Fixed Points	71
A.1	An algorithm for constructing the blow down	112

Chapter 1

Introduction

Duality has long played an important role in string theory. Dualities between the five major classes of string theories led Witten to propose in 1995 the idea of an underlying 11-dimensional M-theory [25]. Earlier, the discovery of mirror symmetry as a duality of CFT's [15] motivated the proposed duality between type IIA and type IIB string theories, and gave birth to purely geometric versions of mirror symmetry with rich implications for the geometry of Calabi-Yau manifolds [16, 6]. By relating physical quantities (correlators, partition functions, spectra) between different theories with geometric input, dualities have uncovered many unexpected patterns in geometry. This has led to surprising conjectures (such as mirror symmetry and T-duality) which not only have important implications for physics, but are interesting and meaningful in a purely geometric light.

Recently, there has been a tremendous amount of work surrounding dualities which relate string theories to other classes of theories. Maldacena's 1997 AdS/CFT correspondence is perhaps the most famous example of such a duality [21]. The connection between Chern-Simons gauge theory and string theory was first introduced by Witten in 1992 [24]. In 1999, Gopakumar and Vafa initiated a program to study the relationship between large N limits of Chern-Simons theory (gauge theory) and type IIA topological string theory (geometry) by using ideas originally proposed by 't Hooft in the 1970's [14]. The resulting gauge theory/geometry correspondence led to a conjecture about extremal transitions, often referred to as the "geometric transition conjecture." In the case of conifold singularities, this is more or less understood. The conifold singularity can be resolved in two very different ways: (1) with a traditional blow up in algebraic geometry, in which the singular point gets replaced by an exceptional \mathbb{P}^1 , or (2) by a deformation of the algebraic equation which replaces the singular point with an S^3 whose size is controlled by the deformation parameter (see Figure 1.1). The physical degrees of freedom associated to D5 branes wrapping the \mathbb{P}^1 correspond to a 3-form flux through the S^3 . The geometric statement is that one can interpolate between the two kinds of resolutions.

In 2002, Dijkgraaf and Vafa expanded this program and proposed new dualities between type IIB topological strings on Calabi-Yau threefolds and matrix models [8, 9]. Due to the symmetry between type IIA and type IIB string theories, this may be viewed as "mirror"

to the Gopakumar-Vafa conjecture. By studying the conifold case, they found strong evidence for the matching of the string theory partition function with that of a matrix model whose potential is closely related to the geometry in question. In particular, a dual version of special geometry in Calabi-Yau threefolds is seen in the eigenvalue dynamics of the associated matrix model [8]. The proposed string theory/matrix model duality has led to an explosion of research on matrix models, a topic which had been dormant since the early 1990's, when it was studied in the context of 2D gravity [7]. The connection between string theory and matrix models is of very tangible practical importance, since many quantities which are difficult to compute in string theory are much easier to handle on the matrix model side.




Figure 1.1: The big picture

Inspired by these developments (summarized in Figure 1.1), in 2003 F. Ferrari was led to propose a direct connection between matrix models and the Calabi-Yau spaces of their dual string theories [12]. It is well known that the solution to a 1-matrix model can be characterized geometrically, in terms of a hyperelliptic curve. The potential for the matrix model serves as direct input into the algebraic equation for the curve, and the vacuum solutions (distributions of eigenvalues) can be obtained from the geometry of the curve and correspond to branch cuts on the Riemann surface. The work of Vafa and collaborators on the strongly coupled dynamics of four-dimensional $\mathcal{N} = 1$ supersymmetric gauge theories [14, 4, 5, 10] suggests that for multi-matrix models, higher-dimensional Calabi-Yau spaces might be useful. Ferrari pursues this idea in [12], finding evidence that certain multi-matrix

models can, indeed, be directly characterized in terms of higher-dimensional (non-compact) Calabi-Yau spaces.

By thinking of the matrix model potential $W(x_1, \dots, x_M)$ as providing constraints on the deformation space of an exceptional \mathbb{P}^1 within a smooth (resolved) Calabi-Yau $\widehat{\mathcal{M}}$, Ferrari outlines a precise prescription for constructing such smooth geometries directly from the potential. Specifically, the resolved geometry $\widehat{\mathcal{M}}$ is given by transition functions

$$\beta = 1/\gamma, \quad v_1 = \gamma^{-n}w_1, \quad v_2 = \gamma^{-m}w_2 + \partial_{w_1}E(\gamma, w_1), \quad (1.1)$$

between two coordinate patches (γ, w_1, w_2) and (β, v_1, v_2) , where β and γ are stereographic coordinates over an exceptional \mathbb{P}^1 . The perturbation comes from the “geometric potential” $E(\gamma, w)$, which is related to the matrix model potential W via

$$W(x_1, \dots, x_M) = \frac{1}{2\pi i} \oint_{C_0} \gamma^{-M-1} E(\gamma, \sum_{i=1}^M x_i \gamma^{i-1}) d\gamma, \quad (1.2)$$

where $M = n + 1 = -m - 1$. We explain this construction in detail in Chapter 2.

In the absence of the perturbation term $\partial_{w_1}E(\gamma, w_1)$, the transition functions (1.1) simply describe an $\mathcal{O}(M-1) \oplus \mathcal{O}(-M-1)$ bundle over a \mathbb{P}^1 . The matrix model superpotential W encodes the constraints on the sections x_1, \dots, x_M of the bundle due to the presence of $\partial_{w_1}E(\gamma, w_1)$. Note that this procedure is also invertible. In other words, given a matrix model superpotential $W(x_1, \dots, x_M)$, one can find a corresponding geometric potential $E(\gamma, w_1)$. As we will discuss in Chapter 3, not all perturbation terms $\gamma^j w_1^k$ contribute to the superpotential (1.2), so there may be many choices of geometric potential for a given W . Nevertheless, the associated geometry $\widehat{\mathcal{M}}$ is unique [12, page 634].

In 2000, S. Katz had already shown how to codify constraints on versal deformation spaces of curves in terms of a potential function,¹ in cases such as (1.1) where the constraints are integrable [18]. In 2001, F. Cachazo, S. Katz and C. Vafa constructed $\mathcal{N} = 1$ supersymmetric gauge theories corresponding to D5 branes wrapping 2-cycles of ADE fibered threefolds [5]. Ferrari studies the same kinds of geometries in a different context, by interpreting the associated potential as belonging to a matrix model, and proposing that the Calabi-Yau geometry encodes all relevant information about the matrix theory.² He is able to verify this in a few examples, and computes known resolvents of matrix models in terms of periods in the associated geometries. The matching results, as well as Ferrari’s solution of a previously unsolved matrix model, suggest that not only can matrix models simplify computations in string theory, but associated geometries from string theory can simplify computations in matrix models.

Many questions immediately arise from Ferrari’s construction. In particular, the matrix model resolvents are not directly encoded in the resolved geometry $\widehat{\mathcal{M}}$, but require knowing the corresponding singular geometry \mathcal{M}_0 obtained by blowing down the exceptional \mathbb{P}^1 . It

¹For a rigorous derivation of the D-brane superpotential, see [3].

²Specifically, it is the triple of spaces $\widehat{\mathcal{M}}, \mathcal{M}_0$, and \mathcal{M} that are conjectured to encode the matrix model quantities; the blow-down map $\pi : \widehat{\mathcal{M}} \rightarrow \mathcal{M}_0$ is the most difficult step towards performing the matrix model computations [12].

is not clear how to do this blow-down in general. It is also not obvious that a geometry constructed from a matrix model potential in this manner will indeed contain a \mathbb{P}^1 which can be blown down to become an isolated singularity.³ Just which matrix models can be “geometrically engineered” in this fashion? What are the corresponding geometries? Can different matrix model potentials correspond to the same geometry? If so, what common features of those models does the geometry encode? Ferrari asks many such questions at the end of his paper [12], and also wonders whether or not it might be possible to devise an algorithm which will automatically construct the blow-down given the initial resolved space.

Previously established results in algebraic geometry such as Laufer’s Theorem [20] and the classification of Gorenstein threefold singularities by S. Katz and D. Morrison [19] provide a partial answer to these questions.

Theorem (Laufer 1979). *Let \mathcal{M}_0 be an analytic space of dimension $D \geq 3$ with an isolated singularity at p . Suppose there exists a non-zero holomorphic D -form Ω on $\mathcal{M}_0 - \{p\}$.⁴ Let $\pi : \widehat{\mathcal{M}} \rightarrow \mathcal{M}_0$ be a resolution of \mathcal{M}_0 . Suppose that the exceptional set $A = \pi^{-1}(p)$ is one-dimensional and irreducible. Then $A \cong \mathbb{P}^1$ and $D = 3$. Moreover, the normal bundle of \mathbb{P}^1 in $\widehat{\mathcal{M}}$ must be either $\mathcal{N} = \mathcal{O}(-1) \oplus \mathcal{O}(-1)$, $\mathcal{O} \oplus \mathcal{O}(-2)$, or $\mathcal{O}(1) \oplus \mathcal{O}(-3)$.*

Laufer’s theorem immediately tells us that we can restrict our search of possible geometries to dimension 3, and that there are only three candidates for the normal bundle to our exceptional \mathbb{P}^1 . In Ferrari’s construction, the bundles $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$, $\mathcal{O} \oplus \mathcal{O}(-2)$, and $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ correspond to zero-, one-, and two-matrix models, respectively.⁵ Following the Dijkgraaf-Vafa correspondence, this puts a limit of 2 adjoint fields on the associated gauge theory, which is precisely the requirement for asymptotic freedom in $\mathcal{N} = 1$ supersymmetric gauge theories. This happy coincidence is perhaps our first indication that the Calabi-Yau geometry encodes information about the RG flow of its corresponding matrix model or gauge theory.

The condition $M \leq 2$ for the normal bundle $\mathcal{O}(M-1) \oplus \mathcal{O}(-M-1)$ in Laufer’s theorem is equivalent to asymptotic freedom, and reflects the fact that only for asymptotically free theories can we expect the \mathbb{P}^1 to be exceptional. In considering matrix model potentials with $M \geq 3$ fields, Ferrari points out that the normal bundle to the \mathbb{P}^1 changes with the addition of the perturbation $\partial_{w_1} E(\gamma, w_1)$, and makes the following conjecture [12, page 636]:

Conjecture (RG Flow, Ferrari 2003). *Consider the perturbed geometry for $m = -n - 2$ and associated superpotential W . Let \mathcal{N} be the normal bundle of a \mathbb{P}^1 that sits at a given critical point of W . Let r be the corank of the Hessian of W at the critical point. Then $\mathcal{N} = \mathcal{O}(r-1) \oplus \mathcal{O}(-r-1)$.*

³We will see later that the ‘hat’ potentials from the Intriligator-Wecht classification lead to geometries where an entire family of \mathbb{P}^1 ’s is blown down to reveal a space \mathcal{M}_0 with non-isolated singularities. It is interesting to wonder what the corresponding “geometric transition conjecture” should be for these cases.

⁴This is the Calabi-Yau condition.

⁵This is because these bundles have zero, one, and two independent global sections, respectively.

Ferrari proves the conjecture for $n = 1$, and limits himself to two-matrix models ($M = n + 1 = 2$) in the rest of his paper. Our first result gives evidence in support of the RG flow conjecture in a more general setting.

Proposition 1. *For $-M \leq r \leq M$, the addition of the perturbation term $\partial_{w_1} E(\gamma, w_1) = \gamma^{r+1} w_1$ in the transition functions*

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-M+1} w_1, \quad v_2 = \gamma^{M+1} w_2 + \gamma^{r+1} w_1,$$

changes the bundle from $\mathcal{O}(M-1) \oplus \mathcal{O}(-M-1)$ to $\mathcal{O}(r-1) \oplus \mathcal{O}(-r-1)$. In particular, the M -matrix model potential

$$W(x_1, \dots, x_M) = \frac{1}{2} \sum_{i=1}^{M-r} x_i x_{M-r+1-i}, \quad (r \geq 0)$$

is geometrically equivalent⁶ to the r -matrix model potential

$$W(x_1, \dots, x_r) = 0.$$

The proof is given in Section 3.3. The fact that the associated superpotential is purely quadratic is satisfying since for quadratic potentials we can often “integrate out” fields, giving a field-theoretic intuition for why the geometry associated to an M -matrix model can be equivalent to that of an r -matrix model, with $r < M$.

Laufer’s theorem constrains the dimension, the exceptional set and its normal bundle—but what are the possible singularity types? In the surface case (complex dimension two), the classification of simple singularities is a classic result.⁷ As hypersurfaces in \mathbb{C}^3 , the distinct geometries are given by:

$$\begin{aligned} A_k & : x^2 + y^2 + z^{k+1} = 0 \\ D_{k+2} & : x^2 + y^2 z + z^{k+1} = 0 \\ E_6 & : x^2 + y^3 + z^4 = 0 \\ E_7 & : x^2 + y^3 + yz^3 = 0 \\ E_8 & : x^2 + y^3 + z^5 = 0 \end{aligned} \tag{1.3}$$

In 1992, Katz and Morrison answered this question in dimension 3 when they characterized the full set of Gorenstein threefold singularities with irreducible small resolutions using invariant theory [19]. In order to do the classification, Katz and Morrison find it useful to think of threefolds as deformations of surfaces, where the deformation parameter t takes on the role of the extra dimension. The equations for the singularities can thus be written in so-called *preferred versal form*, as given in Table 1.1. The coefficients $\alpha_i, \delta_i, \gamma_i$, and ε_i are given by invariant polynomials, and are implicitly functions of the deformation parameter t .

⁶We will call two potentials *geometrically equivalent* if they yield the same geometry under Ferrari’s construction.

⁷An excellent reference for this and other results in singularity theory is [2]. For a more applications-oriented treatment (with many cute pictures!) see Arnol’d’s 1991 book [1]. For 15 characterizations of rational double points, see [11].

S	Preferred Versal Form
A_{n-1} ($n \geq 2$)	$-XY + Z^n + \sum_{i=2}^n \alpha_i Z^{n-i}$
D_n ($n \geq 3$)	$X^2 + Y^2Z - Z^{n-1} - \sum_{i=1}^{n-1} \delta_{2i} Z^{n-i-1} + 2\gamma_n Y$
E_4	$-XY + Z^5 + \varepsilon_2 Z^3 + \varepsilon_3 Z^2 + \varepsilon_4 Z + \varepsilon_5$
E_5	$X^2 + Y^2Z - Z^4 - \varepsilon_2 Z^3 - \varepsilon_4 Z^2 + 2\varepsilon_5 Y - \varepsilon_6 Z - \varepsilon_8$
E_6	$-X^2 - XZ^2 + Y^3 + \varepsilon_2 YZ^2 + \varepsilon_5 YZ + \varepsilon_6 Z^2 + \varepsilon_8 Y$ $+ \varepsilon_9 Z + \varepsilon_{12}$
E_7	$-X^2 - Y^3 + 16YZ^3 + \varepsilon_2 Y^2 Z + \varepsilon_6 Y^2 + \varepsilon_8 YZ + \varepsilon_{10} Z^2$ $+ \varepsilon_{12} Y + \varepsilon_{14} Z + \varepsilon_{18}$
E_8	$-X^2 + Y^3 - Z^5 + \varepsilon_2 YZ^3 + \varepsilon_8 YZ^2 + \varepsilon_{12} Z^3 + \varepsilon_{14} YZ$ $+ \varepsilon_{18} Z^2 + \varepsilon_{20} Y + \varepsilon_{24} Z + \varepsilon_{30}$

Table 1.1: Gorenstein threefold singularities in preferred versal form [19, page 465].

We will also find this representation of the singular threefolds useful, both in performing small resolutions explicitly (Chapter 5), and in identifying what kinds of singular geometries we get upon blowing down resolved geometries.

In contrast to what one might expect,⁸ there are only a finite number of families of Gorenstein threefold singularities with irreducible small resolutions. They are distinguished by the Kollár “length” invariant,⁹ and are resolved via small resolution of the appropriate length node in the corresponding Dynkin diagram. The precise statement of Katz and Morrison’s results are given by the following theorem and corollary [19, page 456]:

Theorem (Katz & Morrison 1992). *The generic hyperplane section of an isolated Gorenstein threefold singularity which has an irreducible small resolution defines one of*

⁸By taking hyperplane sections, one may get surface singularities corresponding to any of the ADE Dynkin diagrams. A priori, this could indicate that there is an infinite number of families of the threefold singularities with irreducible small resolutions. What Katz and Morrison discovered is that only a finite number of Dynkin diagrams can arise from “generic” hyperplane sections.

⁹**Defn:** Let $\pi : Y \rightarrow X$ irreducible small resolution of an isolated threefold singularity $p \in X$. Let $C = \pi^{-1}(p)$ be the exceptional set. The *length* of p is the length at the generic point of the scheme supported on C , with structure sheaf $\mathcal{O}_Y/\pi^{-1}(m_p, x)$.




Figure 1.2: The 6 types of Gorenstein threefold singularities

the primitive partial resolution graphs in Figure 1.2. Conversely, given any such primitive partial resolution graph, there exists an irreducible small resolution $Y \rightarrow X$ whose general hyperplane section is described by that partial resolution graph.

Corollary. *The general hyperplane section of X is uniquely determined by the length of the singular point P .*

We thus know that there are only a finite number of families of distinct geometries with the desired properties for Ferrari's construction, and they correspond to isolated threefold singularities with small resolutions. While much is known about the resolution of these singularities (they are obtained by blowing up divisors associated to nodes of the appropriate length in the corresponding Dynkin diagram), it is not easy to perform the small blowup explicitly.

In the case of simple surface singularities, the blowup of a single node in the associated Dynkin diagram can be obtained via an appropriate matrix factorization of the singular equation [13, 26]. Given a polynomial f defining a hypersurface, a pair of square matrices (ϕ, ψ) such that

$$\phi \cdot \psi = f \cdot \mathbf{1} \quad \text{and} \quad \psi \cdot \phi = f \cdot \mathbf{1}$$

is called a *matrix factorization* of f . A particular ADE singularity will have a different (indecomposable) matrix factorization for each node in the associated Dynkin diagram. The matrix factorization picks out the ideal corresponding to the node; with this the blowup is straightforward. The resulting surface usually contains residual singularities,




Figure 1.3: The blow up of the open length 2 node in the top Dynkin diagram causes the D_8 surface singularity to split into two lower-order surface singularities. The residual singularities are of types A_2 and D_5 , as suggested by the remaining parts of the diagram.

which are characterized by the remaining parts of the original Dynkin diagram once the blown up node has been removed (see Figure 1.3).

Because small resolutions of Gorenstein threefold singularities are obtained by blowing up nodes in the Dynkin diagrams of corresponding hyperplane section surface singularities, we are led to make the following conjecture:

Conjecture 1. *Small resolutions for Gorenstein threefold singularities can be obtained by deforming matrix factorizations for corresponding surface singularities.*

We prove this in the length 1 and length 2 cases, yielding the following theorem:

Theorem 1. *For Gorenstein threefold singularities of length 1 and length 2, with hyperplane section an A_n or D_{n+2} surface singularity, the small resolution is obtained by deforming the matrix factorization for a node of the same length in the corresponding Dynkin diagram.*

The proof is given in Chapter 5. We are thus able to explicitly perform the small resolutions in the A_n and D_{n+2} cases. However, even when the blow up has been found, it can be highly non-trivial to find the right coordinates for expressing the answer. The way in which we identify the superpotential from the resolved geometry requires that this geometry be represented in terms of transition functions over just two coordinate charts covering the exceptional \mathbb{P}^1 .¹⁰ In the length 2 case (D_{n+2}), this difficulty is already apparent, and performing the small resolution stops short of uncovering the corresponding matrix model potential.

The major obstacle in identifying which matrix model corresponds to each of the candidate singular geometries from [19] is the absence of a simple description in the form

¹⁰The techniques in [3] for computing the superpotential are not constrained in this way; perhaps those methods can be used to overcome this difficulty.

of (1.1) for their small resolutions. This frustration is also expressed in [5], where the same geometries are used to construct $\mathcal{N} = 1$ ADE quiver theories.¹¹ In the case where the normal bundle to the exceptional \mathbb{P}^1 is $\mathcal{O}(1) \oplus \mathcal{O}(-3)$, only Laufer’s example [20] and its extension by Pinkham and Morrison [22, page 368] was known. For us, the resolution to this problem came from a timely, albeit surprising, source.

In September, 2003 (just four days after Ferrari’s [12] came out on the preprint server!), Intriligator and Wecht posted their results on RG fixed points of $\mathcal{N} = 1$ SQCD with adjoints [17]. Using “a-maximization” and doing a purely field theoretic analysis, they classify all relevant adjoint superpotential deformations for 4d $\mathcal{N} = 1$ SQCD with \mathcal{N}_f fundamentals and $\mathcal{N}_a = 2$ adjoint matter chiral superfields, X and Y . The possible RG fixed points, together with the map of possible flows between fixed points, are summarized in Figure 1.4.




Figure 1.4: Intriligator–Wecht Classification of RG Fixed Points. The diagram on the right shows the map of possible flows between fixed points. Dotted lines indicate flow to a particular value of k . Note that $k' < k$. [17, pages 3-4]

Due to the form of the polynomials, Intriligator and Wecht name the relevant superpotential deformations according to the famous ADE classification of singularities in dimensions 1 and 2 (1.3). There is no geometry in their analysis, however, and they seem surprised to uncover a connection to these singularity types.¹²

¹¹“...the gauge theory description suggests a rather simple global geometric description of the blown up \mathbb{P}^1 for all cases. However such a mathematical construction is not currently known in the full generality suggested by the gauge theory. Instead only some explicit blown up geometries are known in detail...”[5, page 35]

¹²“On the face of it, this has no obvious connection to any of the other known ways in which Arnold’s singularities have appeared in mathematics or physics.”[17, page 3]

Naively, one may speculate that this is the answer!¹³ We make the following conjecture:

Conjecture 2. *The superpotentials in Intriligator and Wecht’s ADE classification for $\mathcal{N} = 1$ gauge theories (equivalently, the polynomials defining simple curve singularities), are precisely the matrix model potentials which yield small resolutions of Gorenstein threefold singularities under Ferrari’s construction (1.1).*

Armed with this new conjecture, we may now run the classification program backwards. Starting from the resolved space $\widehat{\mathcal{M}}$ given by transition functions over the exceptional \mathbb{P}^1 , we can verify the correspondence by simply performing the blow down and confirming that the resulting geometry has the right singularity type. In particular, the matrix model superpotentials (if correct) give us simple descriptions for the small resolutions of Gorenstein threefold singularities in terms of transition functions as in (1.1). Like other geometric insights stemming from dualities in string theory, such a result is of independent mathematical interest.

This still leaves us with some major challenges. As Ferrari points out, there is no known systematic way of performing the blow downs, and our first task is to devise an algorithm which will do so.

Proposition 2. *There is an algorithm for blowing down the exceptional \mathbb{P}^1 ’s when the resolved geometry is given by simple transition functions (1.1) as in Ferrari’s construction.*

We prove Proposition 2 by giving an explicit algorithm in Chapter 6. The algorithm can be implemented by computer,¹⁴ and searches for global holomorphic functions which can be used to construct the blow down. Any global holomorphic function on the resolved geometry $\widehat{\mathcal{M}}$ is necessarily constant on the exceptional \mathbb{P}^1 , so these functions are natural candidates for coordinates on the blown down geometry \mathcal{M}_0 , since the \mathbb{P}^1 must collapse to a point. We are unable to guarantee, however, that all (independent) global holomorphic functions are found by our algorithm.¹⁵ The resulting singular space \mathcal{M}_0 , whose defining equations are obtained by finding relations among the global holomorphic functions, must be checked. We can verify that we do, indeed, recover the original smooth space by inverting the blow down and performing the small resolution of the singular point.

We find that this program works perfectly in the A_k (length 1) and D_{k+2} (length 2) cases, lending credence to the idea that the Intriligator–Wecht classification is, indeed, the right answer! In the exceptional cases, however, a few mysteries arise. We are only able to find the blow down for the Intriligator–Wecht superpotential E_7 , and the resulting singular space has a length 3 singularity. We summarize these results in the following theorem:

¹³In particular, if the Dijkgraaf–Vafa conjecture holds, we should expect any classification of $\mathcal{N} = 1$ gauge theories to have a matrix model counterpart. Verifying such a correspondence thus provides a non-trivial consistency check on the proposed string theory/matrix model duality.

¹⁴See the appendix for Maple code.

¹⁵The issue is really about how long to run the code. Given a particular list of input monomials, the algorithm will find all independent global holomorphic functions that can be built using elements from this list. However, the list of possible monomials is, of course, infinite.

Theorem 2. Consider the 2-matrix model potentials $W(x, y)$ in the table below, with corresponding resolved geometries $\widehat{\mathcal{M}}$

$$\beta = 1/\gamma, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \partial_{w_1}E(\gamma, w_1),$$

given by perturbation terms $\partial_{w_1}E(\gamma, w_1)$. Blowing down the exceptional \mathbb{P}^1 in each $\widehat{\mathcal{M}}$ yields the following singular geometries \mathcal{M}_0 :

type	$W(x, y)$	$\partial_{w_1}E(\gamma, w_1)$	singular geometry \mathcal{M}_0
A_k	$\frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2$	$\gamma^2w_1^k + w_1$	$XY - T(Z^k - T) = 0$
D_{k+2}	$\frac{1}{k+1}x^{k+1} + xy^2$	$\gamma^2w_1^k + w_1^2$	$X^2 - Y^2Z + T(Z^{k/2} - T)^2 = 0, \quad k \text{ even}$ $X^2 - Y^2Z - T(Z^k - T^2) = 0, \quad k \text{ odd}$
E_7	$\frac{1}{3}y^3 + yx^3$	$\gamma^{-1}w_1^2 + \gamma w_1^3$	$X^2 - Y^3 + Z^5 + 3TYZ^2 + T^3Z = 0.$

Table 1.2: Superpotentials corresponding to length 1, length 2, and length 3 singularities.

By comparing the above singular geometries with the equations in preferred versal form (Table 1.1), we immediately identify the A_k and D_{k+2} superpotentials as corresponding to length 1 and length 2 threefold singularities, respectively. For the E_7 potential, we first note that the polynomial $X^2 - Y^3 + Z^5 + 3TYZ^2 + T^3Z$ is in preferred versal form for E_8 (with $\varepsilon_8 = -3T$ and $\varepsilon_{24} = -T^3$).¹⁶ On the other hand, using Proposition 4 in the proof of the Katz–Morrison classification [19, pages 499-500], we see that the presence of the monomial T^3Z constrains the threefold singularity type to length 3. We thus have the following corollary:

Corollary. The resolved geometries $\widehat{\mathcal{M}}$ given by the 2-matrix model potentials A_k , D_{k+2} , and E_7 in Table 1.2 are small resolutions for length 1, length 2, and length 3 singularities, respectively.

Although simple descriptions of the form (1.1) were previously known for small resolutions of length 1 and length 2 Gorenstein threefold singularities (Laufer’s example [20] in the length 2 case), it is striking that in no other case such a concrete representation for the blowup was known. Theorem 2, together with its Corollary, show a length 3 example where the small resolution also has an extremely simple form:

$$\beta = 1/\gamma, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^{-1}w_1^2 + \gamma w_1^3.$$

¹⁶ $T = 0$ yields a hyperplane section with E_8 surface singularity.

Chapters 7, 8, and 9 comprise the proof of Theorem 2. Missing are examples of length 4, length 5, and length 6 singularity types. For the moment, we are skeptical about whether or not these are describable using geometries that are simple enough to fit into Ferrari’s framework.

In some sense the Intriligator–Wecht classification does not contain enough superpotentials; only length 1, 2, and 3 singularities appear to be included. On the other hand, there are too many: the additional superpotentials \widehat{O} , \widehat{A} , \widehat{D} and \widehat{E} have no candidate geometries corresponding to the Katz–Morrison classification of Gorenstein threefold singularities! What kind of geometries do these new cases correspond to? And what (if any) is their relation to the original ADE classification? Using Ferrari’s framework and our new algorithmic blow down methods we are able to identify the geometries corresponding to these extra ‘hat’ cases. We summarize the results in the following theorem:

Theorem 3. *The singular geometries corresponding to the \widehat{O} , \widehat{A} , \widehat{D} and \widehat{E} cases in the Intriligator–Wecht classification of superpotentials are:*

type	$W(x, y)$	$\partial_{w_1} E(\gamma, w_1)$	singular geometry \mathcal{M}_0
\widehat{O}	0	0	$\mathbb{C}^3/\mathbb{Z}_3$
\widehat{A}	$\frac{1}{2}y^2$	$\gamma^2 w_1$	$\mathbb{C} \times \mathbb{C}^2/\mathbb{Z}_2$
\widehat{D}	xy^2	γw_1^2	$X^2 + Y^2 Z - T^3 = 0$
\widehat{E}	$\frac{1}{3}y^3$	$\gamma^2 w_1^2$	$\text{Spec}(\mathbb{C}[a, b, u, v]/\mathbb{Z}_2)/(b^4 - u^2 - av)$.

Table 1.3: Geometries for the ‘Hat’ cases.

The proof of Theorem 3 is the content of Chapter 11. We find that the resolved geometries have full families of \mathbb{P}^1 ’s which are blown down, and the resulting singular spaces have interesting relations to the ADE cases. The \widehat{A} geometry is a curve of A_1 singularities, while the equation for \widehat{D} looks like the equations for D_{k+2} where the k -dependent terms have been dropped. The identification of new, related geometries obtained by combining Ferrari’s framework with the Intriligator–Wecht classification turns out to be one of the most interesting parts of our story. The presence of these extra geometries may have implications for the relevant string dualities; perhaps the geometric transition conjecture can be expanded beyond isolated singularities.

It is surprising that even in the \widehat{O} case, with $W(x, y) = 0$ superpotential, the geometry is highly non-trivial. In fact, we find that it is the A_1, A_k , and \widehat{A} cases which are, in some sense, the “simplest.” Although the descriptions for the resolved geometries $\widehat{\mathcal{M}}$ in these cases make it appear as though the normal bundles to the exceptional \mathbb{P}^1 ’s are all $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ (as required by a 2-matrix model potential $W(x, y)$), these geometries can all

be described with fewer fields. A straightforward application of Proposition 1 shows that the \widehat{A} case is equivalent to a one-matrix model with $W(x) = 0$ superpotential. Similarly, we will see in Chapter 7 that

$$W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2 \quad \text{and} \quad W(x) = \frac{1}{k+1}x^{k+1}$$

are geometrically equivalent, so the A_k (length 1) cases are also seen to correspond to 1-matrix models, where the y field has been “integrated out.” This is a relief because we know that the exceptional \mathbb{P}^1 after blowing up an A_k singularity should have normal bundle $\mathcal{O} \oplus \mathcal{O}(-2)$. When $k = 1$, Proposition 1 further reduces the geometry to that of a 0-matrix model (no superpotential possible!), showing that A_1 is the most trivial case, with normal bundle $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$.¹⁷ These results are summarized in Table 1.4, and can be understood as evidence for Ferrari’s RG Conjecture. (Compare with Intriligator and Wecht’s map of possible RG flows in Figure 1.4.)

type	2 – matrix model	1 – matrix model	0 – matrix model
\widehat{O}	$W(x, y) = 0$		
\widehat{A}	$W(x, y) = \frac{1}{2}y^2$	$W(x) = 0$	
A_k	$W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2$	$W(x) = \frac{1}{k+1}x^{k+1}$	
A_1	$W(x, y) = \frac{1}{2}x^2 + \frac{1}{2}y^2$	$W(x) = \frac{1}{2}x^2$	$W = 0$

Table 1.4: Geometrically equivalent superpotentials.

Our analysis also indicates that the names are well-chosen: the \widehat{A} and \widehat{D} geometries are closely related to their A_k and D_{k+2} counterparts, and the same might be true for the \widehat{E} case. The relationship between the \widehat{A} , \widehat{D} and \widehat{E} geometries and the ADE singularities is worth exploring for purely geometric reasons. To summarize, string dualities have told us to enlarge the class of geometries considered in [19], and have pointed us to closely related ‘limiting cases’ of these geometries with non-isolated singularities.

The organization of this thesis is as follows. Chapter 2 reviews Ferrari’s construction of Calabi-Yau’s from matrix model superpotentials. Chapter 3 contains preliminary computations which are used throughout this work, as well as the proof of Proposition 1. Chapter 4 reviews how to blow up a node in the Dynkin diagram corresponding to a simple surface singularity. In the A_n and D_{n+2} cases, we rederive matrix factorizations and

¹⁷In contrast, the normal bundle in the \widehat{O} case is truly $\mathcal{O}(1) \oplus \mathcal{O}(-3)$, showing that this geometry requires a 2-matrix model description.

explicitly show how these are used to obtain the blowups. In Chapter 5 we prove Theorem 1, and explore small resolutions of Gorenstein threefold singularities as a “bottom-up” approach for identifying corresponding matrix model superpotentials. In Chapter 6 we develop a “top-down” approach to the problem: the Intriligator–Wecht classification provides us with “guesses” for the resolved geometries, and an algorithm is described which performs the blow-downs systematically (thereby proving Proposition 2). Chapters 7, 8, and 9 present our results in the length 1, 2, and 3 cases, and collectively form the proof of Theorem 2. Chapter 10 summarizes our partial knowledge in the case of the remaining E_6 and E_8 Intriligator–Wecht superpotentials. Chapter 11 shows our results in finding the new geometries associated to the \widehat{O} , \widehat{A} , \widehat{D} and \widehat{E} cases, proving Theorem 3. Finally, in Chapter 12 we discuss open questions and future directions for research. Appendix A contains our Maple code implementation of the blow-down algorithm.

Chapter 2

Ferrari's construction

In this chapter we review some basics of matrix models, and then present Ferrari's construction of Calabi-Yau spaces from matrix model superpotentials in detail. We illustrate the construction in several examples.

2.1 A bit about matrix models.

Here we briefly sketch the rudiments of the hermitian one-matrix model, and show the emergence of the hyperelliptic curve which geometrically encodes the solution. The general idea that one may be able to “solve” the matrix model by means of an algebraic variety motivates Ferrari to propose Calabi-Yau spaces as higher-dimensional analogues which can encode the solutions of multi-matrix models. An excellent reference for matrix models in physics is [7].

The partition function for a hermitian one-matrix model has the form

$$e^Z = \int dM e^{-\text{tr}W(M)} = \int \prod_{i=1}^N d\lambda_i \Delta^2(\lambda) e^{-\sum_i W(\lambda_i)},$$

where M is an $N \times N$ hermitian matrix, W a polynomial potential, and the λ_i 's are the N eigenvalues of M . In going from the first expression to the second, we diagonalize the matrices and pick up the Vandermonde determinant

$$\Delta(\lambda) = \prod_{i < j} (\lambda_i - \lambda_j).$$

To “solve” the matrix model means to find a distribution of eigenvalues which minimizes the effective action

$$S(\lambda) = \frac{1}{g_s} \sum_i W(\lambda_i) - 2 \sum_{i < j} \log(\lambda_i - \lambda_j).$$

The log term in the effective action comes from exponentiating the contribution from the Vandermonde determinant

$$\Delta(\lambda) = e^{\sum_{i < j} \log(\lambda_i - \lambda_j)},$$

which acts as a Coulomb repulsion between the eigenvalues.

In the large N limit the solution to the matrix model is given by a continuum of eigenvalues, with density ρ . The eigenvalues will fill a domain on the real axis; the different components of the domain are known as cuts. Because of this, the solution to the 1-matrix model can be expressed in terms of an algebraic equation called the spectral curve, a hyperelliptic curve in the (x, y) plane,

$$y^2 - W'(x)^2 + f(x) = 0,$$

where $f(x)$ depends on the potential.

2.2 The geometric framework

Here we review Ferrari's construction of non-compact Calabi-Yau's from matrix model superpotentials, following section 3 of [12].

The main idea behind the geometric setup is that deformations of the exceptional \mathbb{P}^1 in a resolved geometry $\widehat{\mathcal{M}}$ correspond to adjoint fields in the gauge theory [5]. Alternatively, the deformation space for a \mathbb{P}^1 wrapped by D-branes can be thought of in terms of matrix models. The number N of D-branes wrapping the \mathbb{P}^1 gives the size of the matrices ($N \times N$), while the number M of independent sections of the \mathbb{P}^1 normal bundle gives the number of matrices (an M -matrix model).




Figure 2.1: Ferrari's diagram [12, page 631].

Inspired by the string theory dualities (summarized in Figure 2.1), Ferrari develops a recipe to go straight from the matrix model to a Calabi-Yau space. If the dualities

hold, all of the matrix model quantities should be computable from the corresponding geometry. In this way, Ferrari's prescription provides a non-trivial consistency check on the Gopakumar-Vafa and Dijkgraaf-Vafa conjectures. Moreover, such a Calabi-Yau space provides a natural higher-dimensional analogue for the spectral (hyperelliptic) curve which encodes the solution to the hermitian one-matrix model.




Figure 2.2: The triple of Calabi-Yau spaces corresponding to a single matrix model.

For each theory, there are three relevant Calabi-Yau spaces: the resolved Calabi-Yau $\widehat{\mathcal{M}}$, the singular Calabi-Yau \mathcal{M}_0 , and the smooth deformed space \mathcal{M} (see Figure 2.2). In short, Ferrari's game consists of the following steps:

1. Start with an M -matrix model matrix model superpotential $W(x_1, \dots, x_M)$ and construct a smooth Calabi-Yau $\widehat{\mathcal{M}}$. The details of this construction are presented below.
2. Identify the exceptional \mathbb{P}^1 in the resolved space $\widehat{\mathcal{M}}$.
3. Blow down the exceptional \mathbb{P}^1 to get the singular \mathcal{M}_0 . The blow down map is $\pi : \widehat{\mathcal{M}} \rightarrow \mathcal{M}_0$.
4. Perturb the algebraic equation for \mathcal{M}_0 to get the smooth deformed space \mathcal{M} .
5. From the triple of geometries, compute matrix model quantities (resolvents).
6. Use standard matrix model techniques (loop equations) to check answers in cases where the matrix model solution is known.

In this framework, the matrix model superpotential is encoded in the transition functions defining the resolved geometry $\widehat{\mathcal{M}}$. Ferrari shows that a wide variety of matrix model superpotentials arise in this fashion, and that matrix model resolvents can be computed directly from the geometry. In other words, the solution to the matrix model is encoded in the corresponding triple of Calabi-Yau's.

The bottleneck to this program is Step 3, the construction of the blow-down map. While Ferrari's ad-hoc methods for constructing the blow-down are successful in his particular examples, he does not know how to construct the blow down in general. Moreover, it seems the calculation of the blow down map π is essentially equivalent to solving the associated matrix model, and hence it would be very useful to have an algorithm which computes π [12, page 655].

In this thesis, we are mainly concerned with Steps 1-3. Our goal is to show how to compute the blow-down map in a large class of examples, and therefore to understand better which singular geometries \mathcal{M}_0 arise from matrix models in Ferrari's framework. In the future, it would be nice to implement the deformation to \mathcal{M} and also to compute matrix model quantities for our examples (Steps 4-6). For the present, this is beyond our scope.

2.2.1 Step 1: Construction of resolved Calabi-Yau

We now turn to Step 1, the construction of the “upstairs” resolved space $\widehat{\mathcal{M}}$ given the matrix model superpotential $W(x_1, \dots, x_k)$. $\widehat{\mathcal{M}}$ is given by transition functions between just two coordinate charts over an exceptional \mathbb{P}^1 : (β, v_1, v_2) in the first chart, and (γ, w_1, w_2) in the second chart. β and γ should be thought of as stereographic coordinates for the \mathbb{P}^1 , with $\beta = \gamma^{-1}$. The other coordinates v_1, v_2 and w_1, w_2 span the normal directions to the \mathbb{P}^1 , and have non-trivial transition functions.

We first discuss the case where $W = 0$, in which the Calabi-Yau is the total space of a vector bundle over the exceptional \mathbb{P}^1 . We then show how a simple deformation of the transition functions leads to constraints on the sections of the bundle. The independent sections x_1, \dots, x_k correspond to matrix degrees of freedom (k independent sections for a k -matrix model). The constraints can be encoded in a potential $W(x_1, \dots, x_k)$. When W is non-zero, our geometry $\widehat{\mathcal{M}}$ is no longer a vector bundle – if the total space were a vector bundle, the sections x_1, \dots, x_k would be allowed to move freely and therefore satisfy no constraints. We shall refer to geometries with $W \neq 0$ as “deformed” or “constrained” bundles.

Pure $\mathcal{O}(n) \oplus \mathcal{O}(m)$ bundle

Consider the following $\widehat{\mathcal{M}}$ geometry for $n \geq 0$ and $m < 0$:

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-n} w_1, \quad v_2 = \gamma^{-m} w_2.$$

There is an $(n + 1)$ -dimensional family of \mathbb{P}^1 's that sit at

$$w_1(\gamma) = \sum_{i=1}^{n+1} x_i \gamma^{i-1}, \quad w_2(\gamma) = 0.$$

We have no freedom in the w_2 coordinate because $m < 0$ precludes $v_2(\beta)$ from being holomorphic whenever $w_2(\gamma)$ is. $w_1(\gamma)$ and $w_2(\gamma)$ define globally holomorphic sections, and in the β coordinate patch become

$$v_1(\beta) = \sum_{i=1}^{n+1} x_i \beta^{n-i+1}, \quad v_2(\beta) = 0.$$

The parameters x_i are precisely the fields in the associated superpotential, and they span the versal deformation space of the \mathbb{P}^1 's. In this case there are no constraints on the x_i 's, which corresponds to the fact that the superpotential is

$$W(x_1, \dots, x_{n+1}) = 0.$$

The geometry $\widehat{\mathcal{M}}$ is the total space of a vector bundle, which we might refer to as a “free” bundle because it is not constrained.

Deformed bundle; enter superpotential

Now consider the deformed geometry (with $n \geq 0$ and $m < 0$):

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-n}w_1, \quad v_2 = \gamma^{-m}w_2 + \partial_{w_1}E(\gamma, w_1),$$

where $E(\gamma, w_1)$ is a function of two complex variables which can be Laurent expanded in terms of entire functions E_i ,

$$E(\gamma, w_1) = \sum_{i=-\infty}^{\infty} E_i(w_1)\gamma^i.$$

We call E the “geometric potential.” The most general holomorphic section $(w_1(\gamma), w_2(\gamma))$ of the normal bundle \mathcal{N} to the \mathbb{P}^1 ’s still has

$$w_1(\gamma) = \sum_{i=1}^{n+1} x_i \gamma^{i-1}, \quad v_1(\beta) = \sum_{i=1}^{n+1} x_i \beta^{n-i+1},$$

but in order to ensure that $v_2(\beta)$ is holomorphic, the x_i ’s will have to satisfy some constraints. Since a holomorphic $w_2(\gamma)$ can only cancel poles in β^{-j} for $j \geq |m|$, the x_i ’s must satisfy $|m| - 1$ constraints in order to cancel remaining lower-order poles introduced by the perturbation. Hence the versal deformation space of the \mathbb{P}^1 is spanned by $n + 1$ parameters x_i satisfying $|m| - 1 = -m - 1$ constraints.

For the \mathbb{P}^1 to be isolated we need $n + 1 = -m - 1$, and we denote this quantity (the number of fields) by M . The constraints are integrable, and equivalent to the extremization $dW = 0$ of the corresponding superpotential $W(x_1, \dots, x_M)$. The \mathbb{P}^1 ’s then sit at the critical points of the superpotential, in the sense that for critical values of the x_i ’s, the pair $(w_1(\gamma), w_2(\gamma))$ will be a global holomorphic section defining a \mathbb{P}^1 .

Summary: General transition functions for $\widehat{\mathcal{M}}$

The resolved geometry $\widehat{\mathcal{M}}$ is described by two coordinate patches (γ, w_1, w_2) and (β, v_1, v_2) , with transition functions

$$\beta = 1/\gamma, \quad v_1 = \gamma^{-n}w_1, \quad v_2 = \gamma^{-m}w_2 + \partial_{w_1}E(\gamma, w_1).$$

In the absence of the $\partial_{w_1}E(\gamma, w_1)$ term, this would simply be an $\mathcal{O}(n) \oplus \mathcal{O}(m)$ bundle over the \mathbb{P}^1 parametrized by the stereographic coordinates γ and β . The perturbation comes from the “geometric potential” $E(\gamma, w)$, which can be expanded as

$$E(\gamma, w) = \sum_{i=-\infty}^{+\infty} E_i(w)\gamma^i.$$

The superpotential

The matrix model superpotential encodes the constraints on the sections x_1, \dots, x_M due to the presence of the perturbation term $\partial_{w_1} E(\gamma, w_1)$ in the defining transition functions for $\widehat{\mathcal{M}}$. It can be obtained directly from the geometric potential via

$$W(x_1, \dots, x_M) = \frac{1}{2\pi i} \oint_{C_0} \gamma^{-M-1} E(\gamma, \sum_{i=1}^M x_i \gamma^{i-1}) d\gamma,$$

where

$$M = n + 1 = -m - 1.$$

The contour integral is meant as a bookkeeping device; C_0 should be taken as a small loop encircling the origin. The integral is a compact notation used by Ferrari to encode all of the constraints at once. The general method for going from transition function perturbation (geometric potential) to superpotential was first presented in [18].

This procedure is also invertible. In other words, given a matrix model superpotential $W(x_1, \dots, x_M)$ one can find a corresponding geometric potential $E(\gamma, w_1)$, and therefore construct the associated geometry. E is not in general unique; from the expression for W one can see that terms can always be added to the geometric potential which will not contribute to the residue of the integrand, and hence will not affect the superpotential. Such terms have no effect on the geometry, however.

Going from W to E is essentially the implementation of Step 1 in Ferrari's game. We now turn to Step 2, which is to locate the exceptional \mathbb{P}^1 's.

2.2.2 Step 2: Locating the \mathbb{P}^1 's

The first task in constructing the blow-down maps is figuring out where the \mathbb{P}^1 's that we want to blow down are located. We will mostly be interested in the $M = 2$ case,

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \partial_{w_1} E(\gamma, w_1),$$

where we always have

$$w_1(\gamma) = x + \gamma y, \quad v_1(\beta) = \beta x + y,$$

with x and y critical points of $W(x, y)$ at the \mathbb{P}^1 's. Depending on the form of the perturbation $\partial_{w_1} E(\gamma, w_1)$, $w_2(\gamma)$ will be chosen to cancel poles of order ≥ 3 . The requirement that $v_2(\beta)$ be holomorphic will fix x and y values to be the same as for the critical points of $W(x, y)$.

2.2.3 Step 3: Finding the blow-down map in Ferrari's examples

As previously mentioned, Ferrari has no systematic way of constructing the blow-down map

$$\pi : \widehat{\mathcal{M}} \longrightarrow \mathcal{M}_0.$$

He successfully finds the blow-down in several examples, however, through clever but ad-hoc methods. Here we show how Ferrari finds the blow-down by discussing the main examples from his paper [12].

Example 1 [12, pages 636-7]

The resolved geometry $\widehat{\mathcal{M}}$ given by transition functions

$$\beta = \gamma^{-1}, \quad v_1 = w_1, \quad v_2 = \gamma^2 w_2 + \gamma P(w_1),$$

engineers the one-matrix model with potential $W(x)$ such that $W'(x) = P(x)$. The \mathbb{P}^1 's sit at

$$w_1(\gamma) = x = v_1(\beta), \quad w_2(\gamma) = 0 = v_2(\beta),$$

with $P(x) = 0$. In this case, global holomorphic functions are not difficult to find:

$$\begin{aligned} \pi_1 &= w_1 = v_1, \\ \pi_2 &= 2\gamma w_2 + P(w_1) = 2\beta v_2 - P(v_1), \\ \pi_3 &= w_2 - \gamma P(w_1) - \gamma^2 w_2 = -v_2 - \beta P(v_1) + \beta^2 v_2, \\ \pi_4 &= w_2 + \gamma P(w_1) + \gamma^2 w_2 = v_2 - \beta P(v_1) + \beta^2 v_2. \end{aligned}$$

These functions collapse the isolated \mathbb{P}^1 's to points, and provide a birational isomorphism outside the set of \mathbb{P}^1 's. The singular blown down geometry is uncovered by observing that the π_i 's satisfy the algebraic equation

$$\mathcal{M}_0 : \pi_4^2 = \pi_3^2 + \pi_2^2 - P^2(\pi_1).$$

Taking hyperplane sections $\pi_1 = x_c$, for critical points of the potential $P(x_c) = 0$, we see that we have A_1 surface singularities which are simultaneously resolved upon inverting the blow-down.

Example 2 [12, pages 637-8]

Consider the superpotential

$$W(x, y) = V(x) + U(y) - xy,$$

with resolved geometry $\widehat{\mathcal{M}}$ of the form

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \gamma Q(w_1/\gamma) + \gamma^2 P(w_1) - \gamma w_1,$$

where $U'(y) = Q(y)$ and $V'(x) = P(x)$. The blow-down is constructed by trial and error. Starting with the ansatz

$$\pi_1 = -\beta v_2 + \dots = -\gamma^2 w_2 - Q(w_1/\gamma) + w_1 - \gamma P(w_1) + \dots,$$

we can move the non-holomorphic term $Q(w_1/\gamma)$ to the left-hand-side, yielding

$$\pi_1 = -\beta v_2 + Q(v_1) = -\gamma^2 w_2 + w_1 - \gamma P(w_1).$$

Similarly we get

$$\pi_2 = \gamma w_2 + P(w_1) = v_1 - \beta Q(v_1) + \beta^2 v_2.$$

π_3 and π_4 require a bit more creativity, and suitable functions are

$$\begin{aligned}\pi_3 &= -\beta^3 v_2 + \beta^2 Q(v_1) + \beta P(\pi_1) - \beta v_1 = \frac{P(\pi_1) - P(w_1)}{\gamma} - w_2, \\ \pi_4 &= \gamma^3 w_2 + \gamma^2 P(w_1) + \gamma Q(\pi_2) - \gamma w_1 = \frac{Q(\pi_2) - Q(v_1)}{\beta} + v_2.\end{aligned}$$

The singular Calabi–Yau is

$$\mathcal{M}_0 : \pi_3 \pi_4 = (P(\pi_1) - \pi_2)(Q(\pi_2) - \pi_1).$$

As in Example 1, this is a length 1 threefold singularity. Why does the normal bundle to the \mathbb{P}^1 in the resolved geometry look like $\mathcal{O}(1) \oplus \mathcal{O}(-3)$? We know that for A_1 singularities it should be $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$, and for A_k ($k > 1$) the normal bundle is $\mathcal{O} \oplus \mathcal{O}(-2)$. We will discuss this puzzle in detail in Section 3.2.

Example 3 [12, pages 638-9]

This is a generalization of Laufer’s example [20], which can also be found in [5]. The superpotential is

$$W(x, y) = xy^2 + V(x) + yU(y^2),$$

and the resolved geometry $\widehat{\mathcal{M}}$ has the form¹

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \gamma Q(w_1/\gamma) + \gamma^2 P(w_1) + w_1^2,$$

where

$$\begin{aligned}V'(x) &= P(x) = -F_1(x^2) - xF_2(x^2), \\ Q(y) &= U(y^2) + 2y^2 U'(y^2) = -G(y^2).\end{aligned}$$

Again, the blow-down map is constructed using the tricks in Examples 1 and 2, although in this case π_1, \dots, π_4 have a rather more complicated form (see [12, page 639] for details). The singular geometry is as usual determined by finding a relation among the π_i ’s,

$$\mathcal{M}_0 : \pi_2 [(\pi_2 - F_1(\pi_1))^2 - \pi_1 F_2^2(\pi_1)] = \pi_4^2 - \pi_1 \pi_3^2 - G(\pi_2) [(\pi_2 - F_1(\pi_1))\pi_3 + \pi_4 F_2(\pi_1)].$$

Despite the complicated form, special cases reveal that this threefold has length 2 (type D_{n+2}) singularities, as in Laufer’s original example.

¹Note that this is almost identical to the previous example, except for the last perturbation term w_1^2 . The geometry in this case, however, is quite different and corresponds to a length 2 threefold singularity.

2.3 Example: A_k

We now illustrate Steps 1-3 in a simple example. Consider the matrix model potential

$$W(x) = \frac{1}{k+1}x^{k+1}.$$

Since there is only 1 field, the resolved geometry $\widehat{\mathcal{M}}$ is given by transition functions

$$\beta = \gamma^{-1}, \quad v_1 = w_1, \quad v_2 = \gamma^2 w_2 + \gamma w_1^k,$$

for an $\mathcal{O} \oplus \mathcal{O}(-2)$ bundle over the exceptional \mathbb{P}^1 . To locate the \mathbb{P}^1 , we first note that

$$w_1(\gamma) = x = v_1(\beta)$$

are the only holomorphic sections for the \mathcal{O} line bundle. Substituting this into the transition function for v_2 yields

$$v_2(\beta) = \beta^{-2} w_2(\gamma) + \beta^{-1} x^k,$$

which is only holomorphic if $x^k = w_2(\gamma) = 0$. Therefore we have a single \mathbb{P}^1 located at

$$w_1(\gamma) = w_2(\gamma) = 0, \quad v_1(\beta) = v_2(\beta) = 0.$$

Note that the position of the \mathbb{P}^1 corresponds exactly to the critical point of the superpotential:

$$dW = x^k dx = 0 \quad \implies \quad x^k = 0.$$

The blow-down

To find the blow down map π , we must look for global holomorphic functions (which will necessarily be constant on the \mathbb{P}^1). We can immediately write down

$$\begin{aligned} \pi_1 &= v_1 = w_1, \\ \pi_2 &= v_2 = \gamma^2 w_2 + \gamma w_1^k, \end{aligned}$$

which are independent. Moreover, notice the combination $\beta v_2 - v_1^k = \gamma w_2$. This gives us

$$\begin{aligned} \pi_3 &= \beta v_2 - v_1^k = \gamma w_2, \\ \pi_4 &= \beta^2 v_2 - \beta v_1^k = w_2. \end{aligned}$$

Since $\beta = \pi_4/\pi_3$, we have immediately from the definition of π_3 the relation

$$\mathcal{M}_0: \pi_3^2 = \pi_4 \pi_2 - \pi_3 \pi_1^k.$$

This corresponds to an A_k (length 1) singularity!

The blowup

We check our computation by inverting the blow-down. If we define

$$v_3 = \beta v_2 - v_1^k, \quad \text{and} \quad w_3 = \gamma w_2 + w_1^k,$$

we can write

$$\begin{aligned}\pi_1 &= v_1 = w_1, \\ \pi_2 &= v_2 = \gamma w_3, \\ \pi_3 &= v_3 = \gamma w_2, \\ \pi_4 &= \beta v_3 = w_2.\end{aligned}$$

In particular

$$\beta = \pi_4/\pi_3 = \gamma^{-1}.$$

This suggests that to recover the small resolution $\widehat{\mathcal{M}}$ we should blow up

$$\pi_3 = \pi_4 = 0 \quad \text{in} \quad \mathcal{M}_0 : \pi_3^2 = \pi_4\pi_2 - \pi_3\pi_1^k.$$

Denoting the \mathbb{P}^1 coordinates by $[\beta : \gamma]$ and imposing the relation

$$\beta\pi_3 = \gamma\pi_4$$

in the blowup, we find in each chart

$(\gamma = 1)$	$(\beta = 1)$
$\pi_4 = \beta\pi_3$	$\pi_3 = \gamma\pi_4$
$\pi_3 = \beta\pi_2 - \pi_1^k$	$\gamma^2\pi_4 = \pi_2 - \gamma\pi_1^k$
(β, π_1, π_2)	(γ, π_1, π_4)

The transition functions between the two charts are easily found to be

$$\beta = \gamma^{-1}, \quad \pi_1 = \pi_1, \quad \pi_2 = \gamma^2\pi_4 + \gamma\pi_1^k.$$

Identifying with the original coordinates, we find

$$\beta = \gamma^{-1}, \quad v_1 = w_1, \quad v_2 = \gamma^2 w_2 + \gamma w_1^k.$$

This is exactly what we started with!

Note that for $k = 1$, we have a bundle-changing superpotential (see Section 3.3), and the normal bundle to the exceptional \mathbb{P}^1 is $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ instead of $\mathcal{O} \oplus \mathcal{O}(-2)$.

Chapter 3

Preliminary computations

In this chapter we do some preliminary calculations in order to investigate more carefully the interplay between perturbation terms, superpotentials, and bundle structure. Due to Laufer's theorem (see Introduction), we focus on the one- and two-matrix model cases. After computing which perturbation terms come from superpotentials, we turn our attention to a puzzle that arises from Ferrari's example 2: a \mathbb{P}^1 with what seems to be an $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ normal bundle is blown down to reveal a geometry which should have come from blowing down a \mathbb{P}^1 with normal bundle $\mathcal{O} \oplus \mathcal{O}(-2)$. We resolve this apparent contradiction in Section 3.3 by investigating terms in the superpotential which not only constrain the sections of the normal bundle, but actually change the local structure. This leads us to prove Proposition 1:

Proposition 1. *For $-M \leq r \leq M$, the addition of the perturbation term $\partial_{w_1} E(\gamma, w_1) = \gamma^{r+1} w_1$ in the transition functions*

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-M+1} w_1, \quad v_2 = \gamma^{M+1} w_2 + \gamma^{r+1} w_1,$$

changes the bundle from $\mathcal{O}(M-1) \oplus \mathcal{O}(-M-1)$ to $\mathcal{O}(r-1) \oplus \mathcal{O}(-r-1)$. In particular, the M -matrix model potential

$$W(x_1, \dots, x_M) = \frac{1}{2} \sum_{i=1}^{M-r} x_i x_{M-r+1-i}, \quad (r \geq 0)$$

is geometrically equivalent¹ to the r -matrix model potential

$$W(x_1, \dots, x_r) = 0.$$

We finish by presenting a resolution of the ideal sheaf, which introduces new coordinates which will be useful for understanding some later examples.

¹We will call two potentials *geometrically equivalent* if they yield the same geometry under Ferrari's construction.

3.1 Perturbations which contribute to superpotential

Consider the transition functions for a (putative) $\mathcal{O}(M-1) \oplus \mathcal{O}(-M-1)$ bundle with perturbation:

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{1-M} w_1, \quad v_2 = \gamma^{M+1} w_2 + \partial_{w_1} E(\gamma, w_1).$$

Not all possible perturbation terms

$$\partial_{w_1} E(\gamma, w_1) = \gamma^n w_1^m, \quad n \in \mathbb{Z}, \quad m \in \mathbb{Z}_{\geq 0}$$

will contribute to the superpotential. Here we compute the superpotential associated to such perturbations

$$\begin{aligned} W(x_1, \dots, x_M) &= \frac{1}{2\pi i} \oint_{C_0} \gamma^{-M-1} E(\gamma, \sum_{i=1}^M x_i \gamma^{i-1}) d\gamma \\ &= \frac{1}{2\pi i} \oint_{C_0} \frac{d\gamma}{m+1} \gamma^{n-M-1} (\sum_{i=1}^M x_i \gamma^{i-1})^{m+1} \end{aligned}$$

to find exactly which terms contribute in the 1-matrix and 2-matrix models. This will provide a useful reference for future computations.

Linear perturbations

In the case of linear perturbations ($m=1$), the terms

$$\partial_{w_1} E(\gamma, w_1) = \gamma^n w_1$$

yield purely quadratic superpotentials:

$$\begin{aligned} W(x_1, \dots, x_M) &= \frac{1}{2\pi i} \oint_{C_0} \frac{1}{2} \gamma^{n-M-1} (\sum_{i=1}^M x_i \gamma^{i-1})^2 d\gamma \\ &= \frac{1}{2} \sum_{i=1}^M x_i x_{M-n+2-i} \end{aligned}$$

We will see later that these are the terms which can change bundle structure.

1-matrix model

The expression for the potential is

$$\begin{aligned} W(x) &= \frac{1}{2\pi i} \oint_{C_0} \frac{d\gamma}{m+1} \gamma^{n-2} x^{m+1} \\ &= \begin{cases} 0 & n \neq 1 \\ \frac{1}{m+1} x^{m+1} & n = 1 \end{cases} \end{aligned}$$

The only perturbation terms

$$\partial_{w_1} E(\gamma, w_1) = \gamma^n w_1^m$$

which contribute satisfy

$$n = 1.$$

In particular, a superpotential of the form

$$W(x) = \frac{1}{m+1} x^{m+1}$$

corresponds to a perturbation term of the form

$$\partial_{w_1} E(\gamma, w_1) = \gamma w_1^m.$$

2–matrix model

The expression for the potential is

$$\begin{aligned} W(x, y) &= \frac{1}{2\pi i} \oint_{C_0} \frac{d\gamma}{m+1} \gamma^{n-3} (x + \gamma y)^{m+1} \\ &= \frac{1}{m+1} \binom{m+1}{2-n} x^{m+n-1} y^{2-n} \end{aligned}$$

The only perturbation terms

$$\partial_{w_1} E(\gamma, w_1) = \gamma^n w_1^m$$

which contribute satisfy $0 \leq 2 - n \leq m + 1$, equivalently

$$1 - m \leq n \leq 2.$$

In particular, a superpotential of the form

$$W(x, y) = \frac{1}{j+k} \binom{j+k}{k} x^j y^k$$

corresponds to a perturbation term of the form

$$P(j, k) \equiv \partial_{w_1} E(\gamma, w_1) = \gamma^{2-k} w_1^{j+k-1}.$$

Remark

The change of variables

$$x \leftrightarrow y,$$

in the superpotential is equivalent to interchanging

$$j \leftrightarrow k.$$

A quick computation shows that

$$P(k, j) = \gamma^{2-j} w_1^{j+k-1} = \gamma^{k-j} P(j, k).$$

In other words,

$$W(x, y) \mapsto W(y, x) \implies P(j, k) \mapsto \gamma^{k-j} P(j, k).$$

3.2 Ferrari's example 2

Recall Ferrari's second example from Section 2.2.3, where from $\widehat{\mathcal{M}}$ transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma Q(w_1/\gamma) + \gamma^2P(w_1) - \gamma w_1,$$

he performs the blow-down to get singular geometry \mathcal{M}_0 :

$$XY = (P(t) - Z)(Q(Z) - t).$$

We will check the result by blowing up the Weil divisor

$$Y = P(t) - Z = 0,$$

to verify that the resolved space is, indeed, what Ferrari started with. We introduce coordinates $[\gamma : \beta]$ on the exceptional \mathbb{P}^1 , and the relation

$$\gamma Y = \beta(P(t) - Z).$$

The blowup is summarized in two charts

(γ, Y, t)	(β, Z, X)
$\gamma Y = P(t) - Z$	$Y = \beta(P(t) - Z)$
$X = \gamma(Q(Z) - t)$	$\beta X = Q(Z) - t$

with transition functions

$\beta = 1/\gamma$	$\gamma = 1/\beta$
$Z = -\gamma Y + P(t)$	$Y = -\beta Z + \beta P(t)$
$X = -\gamma t + \gamma Q(Z)$	$t = -\beta X + Q(Z).$

The puzzle

Our blowup clearly yields a geometry with normal bundle $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ over the exceptional \mathbb{P}^1 ! How is it that Ferrari represents such a space by transition functions for an $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ bundle? In order to investigate this question, we first need to identify what the change of coordinates is between the two representations for the resolved geometry.

Identification with Ferrari's coordinates

We begin the identification by noting that

$$\pi_1 = t, \quad \pi_2 = Z, \quad \pi_3 = Y, \quad \pi_4 = X.$$

From Ferrari (3.40) [12, page 638], we can explicitly compute

$$\begin{aligned}\gamma &= \frac{\pi_4}{Q(\pi_2) - \pi_1} = \gamma, \\ w_1 &= \pi_1 + \frac{\pi_2\pi_4}{Q(\pi_2) - \pi_1} = t - \gamma^2 Y + \gamma P(t), \\ w_2 &= \frac{(\pi_2 - P(w_1))\pi_3}{P(\pi_1) - \pi_2} = -Y + \gamma^{-1}(P(t) - P(w_1)).\end{aligned}$$

Using our transition functions along with Ferrari's (3.30) [12, page 637]

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3 w_2 + \gamma Q(w_1/\gamma) + \gamma^2 P(w_1) - \gamma w_1,$$

we complete the identifications

$(\gamma, Y, t) \longrightarrow (\gamma, w_1, w_2)$	$(\beta, Z, X) \longrightarrow (\beta, v_1, v_2)$
$\begin{aligned}\gamma &= \gamma \\ w_1 &= t - \gamma^2 Y + \gamma P(t) \\ w_2 &= -Y + \gamma^{-1}(P(t) - P(w_1))\end{aligned}$	$\begin{aligned}\beta &= \beta \\ v_1 &= Z - \beta^2 X + \beta Q(Z) \\ v_2 &= X - \beta^{-1}(Q(Z) - Q(v_1)).\end{aligned}$

Jacobian determinants

To check that these are valid coordinate changes, we compute the Jacobians in each chart. In each case, the straightforward identification of the \mathbb{P}^1 coordinate means that we only need the remaining 2×2 minor to check that the determinant does not vanish.

In the γ chart we have

$$\begin{aligned}\frac{\partial w_1}{\partial Y} &= -\gamma^2, & \frac{\partial w_2}{\partial Y} &= -1 + \gamma P'(w_1), \\ \frac{\partial w_1}{\partial t} &= 1 + \gamma P'(t), & \frac{\partial w_2}{\partial t} &= \gamma^{-1}(P'(t) - P'(w_1)) - P'(w_1)P'(t).\end{aligned}$$

The determinant is

$$\begin{aligned}\det &= \frac{\partial w_1}{\partial Y} \frac{\partial w_2}{\partial t} - \frac{\partial w_2}{\partial Y} \frac{\partial w_1}{\partial t} \\ &= -\gamma(P'(t) - P'(w_1)) + \gamma^2 P'(w_1)P'(t) + (1 - \gamma P'(w_1))(1 + \gamma P'(t)) \\ &= 1.\end{aligned}$$

In the β chart we have

$$\begin{aligned}\frac{\partial v_1}{\partial Z} &= 1 + \beta Q'(Z), & \frac{\partial v_2}{\partial Z} &= \beta^{-1}(Q'(v_1) - Q'(Z)) + Q'(v_1)Q'(Z), \\ \frac{\partial v_1}{\partial X} &= -\beta^2, & \frac{\partial v_2}{\partial X} &= 1 - \beta Q'(v_1).\end{aligned}$$

The determinant is

$$\begin{aligned}
\det &= \frac{\partial v_1}{\partial Z} \frac{\partial v_2}{\partial X} - \frac{\partial v_2}{\partial Z} \frac{\partial v_1}{\partial X} \\
&= (1 + \beta Q'(Z))(1 - \beta Q'(v_1)) + \beta(Q'(v_1) - Q'(Z)) + \beta^2 Q'(v_1)Q'(Z) \\
&= 1.
\end{aligned}$$

This shows that we have a valid change of coordinates.

Resolution of the puzzle

How can it be that via a valid change of coordinates, one can go from a normal bundle which is $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ to one which is $\mathcal{O}(1) \oplus \mathcal{O}(-3)$? The resolution of this apparent paradox lies in the fact that Ferrari never had an $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ normal bundle to begin with. One of his superpotential terms actually *changes the bundle structure*, and despite appearances, his original $\widehat{\mathcal{M}}$ transition functions also correspond to an $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ bundle.

Consider the linearized version of the transition functions,

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 - \gamma w_1.$$

The γw_1 term disrupts the $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ structure, and suggests the following change of coordinates

$$\widetilde{w}_1 = w_1 - \gamma^2w_2, \quad \widetilde{v}_1 = v_1 + \beta^2v_2.$$

With these new coordinates we find the transition functions to be

$$\beta = 1/\gamma, \quad \widetilde{v}_1 = \gamma w_2, \quad v_2 = -\gamma \widetilde{w}_1,$$

which clearly exhibits an $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ structure.

With an $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ bundle structure, $M = 0$. We therefore can't have any nontrivial superpotential, even if there's a geometric potential. However, by changing coordinates Ferrari achieved a setup where $M = 2$, and the superpotential was

$$W(x, y) = V(x) + U(y) - xy,$$

with $V'(x) = P(x)$ and $U'(y) = Q(y)$. In particular, it is the xy term in the superpotential which leads to the change in the normal bundle structure. (Recall that we're assuming from the beginning that $x^2|P(x)$ and $x^2|Q(x)$. Otherwise our $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ identification is invalid, as the polynomials contribute to the linearized transition functions.)

In this example, Ferrari is implicitly using the superpotential term to artificially introduce fields x and y into a resolved geometry that should correspond to $M = 0$ fields. The 2-matrix model theory he gets, therefore, should be related to the "0-matrix model" corresponding to the geometry with an $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ bundle.

3.3 Superpotentials which change bundle structure

What we see in Ferrari's example 2 is actually part of a more general phenomenon, which we were led to uncover in trying to understand precisely this puzzle. In this section we describe a family of superpotentials which change the underlying bundle structure, thus proving Proposition 1.

Consider an $\mathcal{O}(n) \oplus \mathcal{O}(m)$ normal bundle over a \mathbb{P}^1 with geometric potential

$$E(\gamma, w_1) = \frac{1}{2}\gamma^k w_1^2, \quad k \in \mathbb{Z}.$$

The perturbed transition functions are

$$\beta = 1/\gamma, \quad v_1 = \gamma^{-n} w_1, \quad v_2 = \gamma^{-m} w_2 + \gamma^k w_1.$$

If (n, m, k) satisfies

$$-n \leq k \leq -m,$$

we can perform the following change of coordinates:

$$\begin{aligned} \widetilde{w}_1 &= w_1 + \gamma^{-m-k} w_2, & \widetilde{v}_1 &= v_2, \\ \widetilde{w}_2 &= w_2, & \widetilde{v}_2 &= -v_1 + \beta^{n+k} v_2. \end{aligned}$$

Notice that

$$\begin{aligned} \widetilde{v}_1 &= \gamma^{-m} w_2 + \gamma^k w_1 = \gamma^k \widetilde{w}_1, \\ \widetilde{v}_2 &= -\gamma^{-n} w_1 + \gamma^{-n-k} (\gamma^{-m} w_2 + \gamma^k w_1) = \gamma^{-n-m-k} \widetilde{w}_2, \end{aligned}$$

and so the new transition functions are

$$\beta = 1/\gamma, \quad \widetilde{v}_1 = \gamma^k \widetilde{w}_1, \quad \widetilde{v}_2 = \gamma^{-n-m-k} \widetilde{w}_2.$$

The geometric potential has changed our $\mathcal{O}(n) \oplus \mathcal{O}(m)$ bundle into an $\mathcal{O}(-k) \oplus \mathcal{O}(n+m+k)$ bundle, with no superpotential. The corresponding superpotential can be computed for $n+m = -2$, and depends on the number of fields $M = n+1 = -m-1$:

$$\begin{aligned} W(x_1, \dots, x_M) &= \frac{1}{2\pi i} \oint_{C_0} \gamma^{-M-1} E(\gamma, \sum_{i=1}^M x_i \gamma^{i-1}) d\gamma \\ &= \frac{1}{2} \sum_{i=1}^M x_i x_{M-k-i+2} \end{aligned}$$

Note that all of these bundle-changing superpotentials are purely quadratic! In the cases of interest, where $n+m = -2$, the condition for the change of coordinates to be valid becomes

$$-n \leq k \leq n+2.$$

For allowed pairs (n, k) we can thus get

$$\mathcal{O}(n) \oplus \mathcal{O}(-n-2) \longrightarrow \mathcal{O}(-k) \oplus \mathcal{O}(k-2)$$

by means of the perturbation. Alternatively, we can think of these examples as “true” $\mathcal{O}(-k) \oplus \mathcal{O}(k-2)$ bundles which can be rewritten to “look like” $\mathcal{O}(n) \oplus \mathcal{O}(-n-2)$ plus a superpotential term.

RG conjecture

In order to make contact with Ferrari’s RG conjecture (see Introduction), we change notation a bit from the previous discussion:

$$M = n + 1, \quad r = k - 1.$$

The perturbation term in the following transition functions

$$\beta = \gamma^{-1}, \quad w'_1 = \gamma^{-M+1}w_1, \quad w'_2 = \gamma^{M+1}w_2 + \gamma^{r+1}w_1,$$

changes the bundle

$$\mathcal{O}(M-1) \oplus \mathcal{O}(-M-1) \longrightarrow \mathcal{O}(r-1) \oplus \mathcal{O}(-r-1), \quad \text{for } -M \leq r \leq M.$$

The change of coordinates:

$$\begin{aligned} v_1 &= w_1 + \gamma^{M-r}w_2, & v'_1 &= w'_2, \\ v_2 &= w_2, & v'_2 &= -w'_1 + \beta^{M+r}w'_2, \end{aligned}$$

yields new transition functions

$$\beta = \gamma^{-1}, \quad v'_1 = \gamma^{r+1}v_1, \quad v'_2 = \gamma^{1-r}v_2.$$

The superpotential

The superpotential corresponding to the perturbation

$$\partial_{w_1} E(\gamma, w_1) = \gamma^{r+1}w_1$$

is given by

$$W_r(x_1, \dots, x_M) = \begin{cases} \frac{1}{2} \sum_{i=1}^{M-r} x_i x_{M-r+1-i}, & \text{for } r \geq 0, \\ \frac{1}{2} \sum_{i=1-r}^M x_i x_{M-r+1-i}, & \text{for } r \leq 0. \end{cases}$$

This completes the proof of Proposition 1.

Notice that the case $r = M$ is not interesting, as the bundle remains unchanged and the superpotential vanishes. Moreover, the symmetry $r \mapsto -r$ in the bundle expression interchanges two different superpotentials, but this amounts to a simple change of coordinates. To see this, first note that:

$$\begin{aligned} r \geq 0: \quad W_r(x_1, \dots, x_M) &= \frac{1}{2} \sum_{i=1}^{M-|r|} x_i x_{M-|r|+1-i} \\ r \leq 0: \quad W_r(x_1, \dots, x_M) &= \frac{1}{2} \sum_{i=1+|r|}^M x_i x_{M+|r|+1-i} = \frac{1}{2} \sum_{i=1}^{M-|r|} x_{i+|r|} x_{M+1-i} \end{aligned}$$

The direction of the coordinate shift depends on the sign of r :

$$\begin{aligned} r \geq 0: \quad r \mapsto -r &\text{ is equivalent to } x_i \mapsto x_{i+|r|} \\ r \leq 0: \quad r \mapsto -r &\text{ is equivalent to } x_i \mapsto x_{i-|r|} \end{aligned}$$

i.e. $r \mapsto -r$ on the bundle side is equivalent to a simple coordinate change for the corresponding superpotential.

We summarize the first few examples in the following table:

r	-3	-2	-1	0	1	2	3
$M = 1$				$\frac{1}{2}x_1^2$			
$M = 2$			$\frac{1}{2}x_2^2$	x_1x_2	$\frac{1}{2}x_1^2$		
$M = 3$		$\frac{1}{2}x_3^2$	x_2x_3	$x_1x_3 + \frac{1}{2}x_2^2$	x_1x_2	$\frac{1}{2}x_1^2$	
$M = 4$	$\frac{1}{2}x_4^2$	x_3x_4	$x_2x_4 + \frac{1}{2}x_3^2$	$x_1x_4 + x_2x_3$	$x_1x_3 + \frac{1}{2}x_2^2$	x_1x_2	$\frac{1}{2}x_1^2$

The Hessian

We compute the partial derivatives of our bundle-changing superpotentials:

$$\begin{aligned} r \geq 0: \quad \frac{\partial W_r}{\partial x_j} &= x_{M-r+1-j}, \quad \frac{\partial^2 W_r}{\partial x_k \partial x_j} = \delta_{k, M-r+1-j} \quad \text{for } 1 \leq j \leq M-r. \\ r \leq 0: \quad \frac{\partial W_r}{\partial x_j} &= x_{M-r+1-j}, \quad \frac{\partial^2 W_r}{\partial x_k \partial x_j} = \delta_{k, M-r+1-j} \quad \text{for } 1-r \leq j \leq M. \end{aligned}$$

In each case, there is only one k for every j which yields a non-zero second-partial. This means the Hessian matrix has at most one non-zero entry in each row and in each column. The corank of the Hessian is thus easy to compute, and is equal to the number of rows (or columns) comprised entirely of zeroes. In both the $r \geq 0$ and $r \leq 0$ cases, the corank of the Hessian is r (see the ranges for j values). This is consistent with what we expect from Ferrari's RG conjecture.

3.4 Classification of perturbation terms

1–matrix model

We consider the case with $M = 1$ fields, which gives an $\mathcal{O} \oplus \mathcal{O}(-2)$ bundle with perturbation:

$$\beta = \gamma^{-1}, \quad v_1 = w_1, \quad v_2 = \gamma^2 w_2 + \partial_{w_1} E(\gamma, w_1).$$

The only perturbation terms which contribute to the superpotential are those of the form

$$\partial_{w_1} E(\gamma, w_1) = \gamma w_1^m.$$

All other terms can be added freely to the RHS of the v_2 transition function without affecting the superpotential. The linear term γw_1 changes the bundle structure to $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$, which we will see corresponds to the A_1 case, with potential $W(x) = \frac{1}{2}x^2$.

2–matrix model

We consider the case with $M = 2$ fields, which gives an $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ bundle with perturbation:

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \partial_{w_1} E(\gamma, w_1).$$

The only perturbation terms which contribute to the superpotential are those of the form

$$\partial_{w_1} E(\gamma, w_1) = \gamma^n w_1^m, \quad \text{s.t. } -m < n < 3.$$

All other terms can be added freely to the RHS of the v_2 transition function without affecting the superpotential. Interchanging x and y in the superpotential leads to

$$\partial_{w_1} E(\gamma, w_1) = \gamma^n w_1^m \mapsto \gamma^{3-m-n} w_1^m.$$

Again, the linear ($m = 1$) perturbation terms are the ones which change the bundle structure. We have $w_1, \gamma w_1$, and $\gamma^2 w_1$. The first and third terms are related by interchanging x and y , and yield an $\mathcal{O} \oplus \mathcal{O}(-2)$ bundle with zero potential. The γw_1 term reduces the bundle all the way to $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$. The effect of these terms is summarized in Table 3.1.

$\partial_{w_1} E(\gamma, w_1)$	$W(x, y)$	new bundle	type
w_1	$\frac{1}{2}y^2$	$\mathcal{O} \oplus \mathcal{O}(-2)$	\widehat{A}
γw_1	xy	$\mathcal{O}(-1) \oplus \mathcal{O}(-1)$	A_1 (Ferrari's Example 2)
$\gamma^2 w_1$	$\frac{1}{2}x^2$	$\mathcal{O} \oplus \mathcal{O}(-2)$	\widehat{A}

Table 3.1: Bundle-changing perturbation terms in the 2-matrix model.

Summary

Table 3.2 summarizes everything we need to know about perturbation terms which contribute to the superpotential in the one- and two-matrix model cases.

	potential	$\partial_{w_1} E(\gamma, w_1)$	bundle – changing
1 – matrix	$W(x) = \sum \frac{a_m}{m+1} x^{m+1}$	$\sum a_m \gamma w_1^m$	γw_1
2 – matrix	$W(x, y) = \sum \frac{a_{jk}}{j+k} \binom{j+k}{k} x^j y^k$	$\sum a_{jk} \gamma^{2-k} w_1^{j+k-1}$	$w_1, \gamma w_1, \gamma^2 w_1$

Table 3.2: Summary of superpotential-contributing perturbation terms.

3.5 Resolution of the ideal sheaf

Here we present a resolution of the ideal sheaf over the exceptional \mathbb{P}^1 by introducing additional coordinates w_3 and v_3 which simplify the transition functions. In some cases, this strategy allows us to immediately write down global holomorphic functions which help to perform the blow-down of the \mathbb{P}^1 . However, what we find most useful in this framework is the identification of natural, additional w_3, v_3 coordinates. In the case of the E_7 Intriligator–Wecht superpotential (see Chapter 9), these extra coordinates will help us to do the blowup and verify that the singular geometry we find is the correct one.

Starting from the transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \partial_{w_1} E(\gamma, w_1),$$

with perturbation term

$$\partial E = \sum c_i \gamma^{2-a_i} w_1^{a_i+b_i-1} = \sum c_i \beta^{-b_i-1} v_1^{a_i+b_i-1},$$

let

$$a = \max a_i, \quad b = \max b_i.$$

Consider the simpler transition functions

$$\begin{aligned} \beta &= \gamma^{-1}, \\ v_1 &= \gamma^{-1} w_1, \\ v_2 &= \gamma^{2-a} w_3, \\ v_3 &= \gamma^{2-b} w_2, \end{aligned}$$

where the new functions w_3 and v_3 are defined by

$$\begin{aligned} w_3 &= \gamma^{a+1} w_2 + \gamma^{a-2} \partial E = \gamma^{a+1} w_2 + p(\gamma, w_1) w_1, \\ v_3 &= \beta^{b+1} v_2 - \beta^{b+1} \partial E = \beta^{b+1} v_2 - q(\beta, v_1) v_1, \end{aligned}$$

and the functions p and q are given by

$$\begin{aligned} p(\gamma, w_1) &= \sum c_i \gamma^{a-a_i} w_1^{a_i+b_i-2}, \\ q(\beta, v_1) &= \sum c_i \beta^{b-b_i} v_1^{a_i+b_i-2}. \end{aligned}$$

In the framework of the

$$\mathcal{O}(1) \oplus \mathcal{O}(a-2) \oplus \mathcal{O}(b-2)$$

bundle, we can easily write down global holomorphic functions as long as either $a < 2$ or $b < 2$. Unfortunately this only covers the \widehat{A} , \widehat{D} , and \widehat{E} cases. However, this strategy also yields some insight in the other cases.

We have an exact sequence

$$0 \longrightarrow \mathcal{O}(-a-b) \longrightarrow \mathcal{O}(1-a) \oplus \mathcal{O}(1-b) \oplus \mathcal{O}(1-a-b) \longrightarrow \mathcal{O}(2-a) \oplus \mathcal{O}(-1) \oplus \mathcal{O}(2-b) \longrightarrow \mathcal{O},$$

with maps in the first chart,

$$\begin{bmatrix} 1 & \gamma^{a+1} & w_1 \end{bmatrix} \begin{bmatrix} w_1 & -w_3 & 0 \\ 0 & w_2 & -w_1 \\ -1 & p(\gamma, w_1) & \gamma^{a+1} \end{bmatrix} \begin{bmatrix} w_3 \\ w_1 \\ w_2 \end{bmatrix},$$

and in the second chart,

$$\begin{bmatrix} \beta^{b+1} & 1 & v_1 \end{bmatrix} \begin{bmatrix} v_1 & -v_2 & 0 \\ 0 & v_3 & -v_1 \\ -\beta^{b+1} & q(\beta, v_1) & 1 \end{bmatrix} \begin{bmatrix} v_2 \\ v_1 \\ v_3 \end{bmatrix}.$$

For reference, we also record the transformation properties of the maps:

$$\begin{bmatrix} \mathcal{O}(b+1) & \mathcal{O}(a+1) & \mathcal{O}(1) \end{bmatrix} \begin{bmatrix} \mathcal{O}(1) & \mathcal{O}(a-2) & \mathcal{O}(1+a-b) \\ \mathcal{O}(1+b-a) & \mathcal{O}(b-2) & \mathcal{O}(1) \\ \mathcal{O}(b+1) & \mathcal{O}(a+b-2) & \mathcal{O}(a+1) \end{bmatrix}.$$

Chapter 4

Blowups for surface singularities

In this chapter we will begin by describing in general terms how to blow up a single node of a Dynkin diagram. This will require some sophisticated techniques, including use of the McKay correspondence in order to identify the Weil divisor associated to each particular node. Once the appropriate matrix factorization has been found, the blowup is fairly straightforward. Here we focus on blowing up nodes in A_{n-1} and D_{n+2} diagrams. We will use these results in Chapter 5, where corresponding small resolutions will be obtained by deforming the matrix factorizations in the surface case.

Recall that given a polynomial f defining a hypersurface, a pair of square matrices (ϕ, ψ) such that

$$\phi \cdot \psi = f \cdot \mathbf{1} \quad \text{and} \quad \psi \cdot \phi = f \cdot \mathbf{1}$$

is called a *matrix factorization*¹ of f .

4.1 How to blow up a single node

The main ideas in this section come from [13] and [26].

We would like to blow up an arbitrary node in the Dynkin diagram for a simple surface singularity. The McKay correspondence will allow us to identify which Weil divisor corresponds to which node. The main idea is to find the module corresponding to the representation for a particular node. The presence of the singularity is reflected in that this module is not locally free. Relations among the generators can be arranged in a matrix which, together with its matrix of syzygies, gives a matrix factorization for the singular equation. Each node yields a different (indecomposable) matrix factorization for the original equation. Once the matrix factorization is found, the blowup is obtained by trying to make the module locally free.

Let $\Gamma \subset SU(2)$ be a finite subgroup, and \mathbb{C}^2/Γ the corresponding singularity. The McKay correspondence says that there is a 1 – 1 correspondence between irreducible representations of Γ and nodes on the extended Dynkin diagram, where the trivial representation

¹For everything you ever wanted to know about matrix factorizations, see [26].

corresponds to the added node. The degree (dimension) of an irreducible representation ρ is given by the coefficient of that node in the largest root. Moreover, ρ and $\tilde{\rho}$ are adjacent in the diagram iff $\tilde{\rho}$ appears in the decomposition of $\rho \otimes \rho_0$, where ρ_0 is the defining 2-dimensional representation (from the induced action on \mathbb{C}^2).

In order to find the Weil divisor for a particular node, we will look at the corresponding representation ρ of Γ . Identifying the action of Γ on the ring $\mathbb{C}[x, y]$, we will find that the elements transforming in the ρ representation form a $\mathbb{C}[x, y]^\Gamma$ -module which is not locally free. Relations among the generators give a matrix factorization for the original equation. This is then used to do the blowup.

Using matrix factorizations to blow up a node in an ADE Dynkin diagram is a natural continuation of the work of Gonzalez-Sprinberg & Verdier [13]. We rederive matrix factorizations for A_{n-1} and D_{n+2} surface singularities in a similar, but slightly different way. In the A_{n-1} and D_{n+2} cases, the finite groups Γ are:

$$\begin{aligned} \mathbb{Z}_n &= \left\langle g = \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix} \right\rangle, & \xi &= e^{2\pi i/n}, & \xi^n &= 1, \\ \mathbb{B}_{4n} &= \left\langle g = \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix}, h = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\rangle, & \zeta &= e^{\pi i/n}, & \zeta^{2n} &= 1. \end{aligned}$$

$\Gamma \subset SU(2)$ acts on $\mathbb{C}[x, y]$, and its irreducible representations (in theory) can all be found there. Note that x and y should be regarded as coordinate functions on \mathbb{C}^2 :

$$x = \begin{pmatrix} 1 & 0 \end{pmatrix}, \quad y = \begin{pmatrix} 0 & 1 \end{pmatrix}.$$

General recipe

The general procedure for blowing up a single node in a Dynkin diagram thus consists of the following steps:

1. Identify the group action for the diagram type.
2. Study the invariant theory under this action to identify the $\mathbb{C}[x, y]^\Gamma$ -module containing the irreducible representations corresponding to the chosen node.
3. Find relations among the generators of this module, arrange in a matrix. This, together with the matrix of syzygies, gives a matrix factorization.
4. From the matrix of relations identify the Weil divisor corresponding to the node.
5. Blow up this divisor!

In the fully deformed cases (see Chapter 5), we will take the matrix of relations for generators of the module from the non-deformed case, and cleverly “extend” the entries to the full deformation.

4.2 The A_{n-1} story

4.2.1 Invariant theory

\mathbb{Z}_n Group action

The action of \mathbb{Z}_n on $\mathbb{C}[x, y]$ is characterized by

$$\begin{aligned} g \cdot x &= \begin{pmatrix} 1 & 0 \\ 0 & \xi \end{pmatrix} \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix} = \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix} = \xi x, \\ g \cdot y &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \xi & 0 \\ 0 & \xi^{-1} \end{pmatrix} = \begin{pmatrix} 0 & \xi^{-1} \\ \xi & 0 \end{pmatrix} = \xi^{-1} y, \end{aligned}$$

from which we induce the action on monomials of higher degree:

$$\begin{aligned} g \cdot x^m &= (g \cdot x)^m = \xi^m x^m, \\ g \cdot y^m &= (g \cdot y)^m = \xi^{-m} y^m, \\ g \cdot x^k y^l &= (g \cdot x)^k (g \cdot y)^l = \xi^{k-l} x^k y^l. \end{aligned}$$

\mathbb{Z}_n Invariant theory

Since $\xi^n = 1$, the invariant polynomials under \mathbb{Z}_n are generated by:

$$x^n, xy, y^n.$$

If we define

$$X = x^n, Y = y^n, Z = xy,$$

then X, Y and Z are invariant under Γ . In particular, X, Y and Z satisfy the equation for an A_{n-1} surface singularity:

$$XY - Z^n = 0.$$

The irreducible representations $\{\rho_j\}$ of \mathbb{Z}_n are labeled by the characters $\{\chi_j\}$, where $\chi_j(g) = \xi^j$, $j = 1 \dots n$. From now on we will simply refer to ξ^j as the character. ρ_n is the trivial representation; all the irreducible representations are one-dimensional. For instance, the defining representation ρ_0 splits as $\rho_0 = \rho_1 \oplus \rho_{n-1}$. Within the ring $\mathbb{C}[x, y]$, the irreducible representations with character ξ^m are given by the one-dimensional vector spaces spanned by the monomials

$$\xi^m : \{x^j y^k\} \text{ s.t. } j - k \equiv m \pmod{n}.$$

Let $\mathbb{C}[x, y]^\Gamma$ denote the subring of invariant functions. It is generated by X, Y and Z , and can also be written as

$$\mathbb{C}[x, y]^\Gamma = \mathbb{C}[X, Y, Z]/(XY - Z^n).$$

By definition Γ acts trivially on $\mathbb{C}[x, y]^\Gamma$, so multiplying a monomial by an element of the invariant ring will yield a new representation with the same character. Consider the representation $x^j y^k$ where $j - k = m + qn$. If $q \geq 0$ we can write

$$x^j y^k = x^{j-k} Z^q = x^m X^q Z^k \in x^m \mathbb{C}[x, y]^\Gamma.$$

Similarly, for $q \leq -1$ we can write

$$x^j y^k = y^{k-j} Z^j = y^{-m-qn} Z^j = y^{n-m} Y^{-q-1} Z^j \in y^{n-m} \mathbb{C}[x, y]^\Gamma.$$

In other words, all of the irreducible representations with character ξ^m can be found in the $\mathbb{C}[x, y]^\Gamma$ -module

$$\langle x^m, y^{n-m} \rangle \mathbb{C}[x, y]^\Gamma.$$

Importantly, this is not a free module. Let $s_1 = x^m$, and $s_2 = y^{n-m}$. Then any element in the module can be written as $as_1 + bs_2$, where $a, b \in \mathbb{C}[x, y]^\Gamma$. However, there are relations, and the above representation is not unique (if elements in the module could be uniquely specified by ordered pairs (a, b) , the module would be free). For example,

$$\begin{aligned} Ys_1 &= Z^m s_2, \\ Xs_2 &= Z^{n-m} s_1. \end{aligned}$$

These relations are related to a factorization of the defining equation:

$$X \cdot Y - Z^m \cdot Z^{n-m} = 0.$$

Blowing up the singularity renders the module (locally) free.

4.2.2 Matrix of relations

The relations among the generators s_1 and s_2 of the $\mathbb{C}[x, y]^\Gamma$ -module corresponding to the m th node (the \mathbb{Z}_n representation of character ξ^m) can be put into a matrix

$$\mathcal{R} = \begin{pmatrix} Y & -Z^m \\ Z^{n-m} & -X \end{pmatrix}$$

which has a corresponding matrix of syzygies

$$\mathcal{S} = \begin{pmatrix} X & -Z^m \\ Z^{n-m} & -Y \end{pmatrix}.$$

Together, the pair $(\mathcal{R}, \mathcal{S})$ gives a matrix factorization for the A_{n-1} equation:

$$\mathcal{S}\mathcal{R} = \mathcal{R}\mathcal{S} = (XY - Z^n) \mathbf{1}_{2 \times 2}.$$

4.2.3 Summary of results

$\Gamma = \mathbb{Z}_n$, $X = \text{Spec } \mathbb{C}[x, y]^\Gamma$. If we let

$$X = x^n, \quad Y = y^n, \quad Z = xy,$$

then the ring of invariant polynomials can be written

$$\mathbb{C}[x, y]^\Gamma = \mathbb{C}[X, Y, Z]/(XY - Z^n),$$

which we recognize as a space with A_{n-1} singularity at the origin. The sheaf of \mathcal{O}_X -modules we are interested in is generated by s_1 and s_2 :

$$\mathcal{F} = \{s_1 = x^m, s_2 = y^{n-m}\} \cdot \mathcal{O}_X.$$

We can view these as providing a map

$$\begin{aligned} (s_1, s_2) : \mathcal{O}_X^{\oplus 2} &\longrightarrow \mathcal{F}. \\ (f, g) &\longmapsto f \cdot s_1 + g \cdot s_2 \end{aligned}$$

\mathcal{F} corresponds to elements of $\mathbb{C}[x, y]$ which transform as a ξ^m representation of \mathbb{Z}_n . We find relations:

$$\begin{aligned} Y s_1 - Z^m s_2 &= 0, \\ Z^{n-m} s_1 - X s_2 &= 0. \end{aligned}$$

These provide a matrix factorization

$$\mathcal{R} = \begin{pmatrix} Y & -Z^m \\ Z^{n-m} & -X \end{pmatrix}, \quad \mathcal{S} = \begin{pmatrix} X & -Z^m \\ Z^{n-m} & -Y \end{pmatrix},$$

and correspond to a factorization of the defining equation:

$$X \cdot Y - Z^m \cdot Z^{n-m} = 0.$$

Away from the origin the two relations are the same, and allow us to solve for one generator in terms of the other:

$$s_2 = \frac{Y}{Z^m} s_1 = \frac{Z^{n-m}}{X} s_1.$$

This suggests that for purposes of figuring out the blow-up, it might be okay to consider just one of the two relations.

4.2.4 The blowup

The original equation in \mathbb{C}^3 is

$$XY = Z^n.$$

We have the exact sequence:

$$\begin{array}{ccccccc} \mathcal{O}_X & \xrightarrow{(Y, -Z^m)} & \mathcal{O}_X^{\oplus 2} & \xrightarrow{(s_1, s_2)} & \mathcal{F} & \longrightarrow & 0. \\ f & \longmapsto & (fY, -fZ^m) & \longmapsto & f(Y s_1 - Z^m s_2) = 0 & & \\ & & (f, g) & \longmapsto & f s_1 + g s_2 & & \end{array}$$

In particular, we see that the map to \mathcal{F} has a one-dimensional kernel—implying that \mathcal{F} has rank 1—except at

$$Y = Z^m = 0,$$

where the rank of the $\mathcal{O}_X \rightarrow \mathcal{O}_X^{\oplus 2}$ map drops, the relation becomes vacuous, and the rank of \mathcal{F} jumps to two. We blow up the Weil divisor $Y = Z^m = 0$, which corresponds to the ideal

$$I = \langle Y, Z^m \rangle.$$

We introduce coordinates $[\mu, \nu]$ for \mathbb{P}^1 , and the relation

$$\mu Y = \nu Z^m.$$

We find the new relation and proper transform of the original curve in two coordinate charts:

$\mu = 1$	$\nu = 1$
$Y = \nu Z^m$	$\mu Y = Z^m \rightarrow (A_{m-1})$
$(A_{n-m-1}) \leftarrow X\nu = Z^{n-m}$	$X = \mu Z^{n-m}$
(ν, X, Z)	(μ, Y, Z)

The proper transform in the first chart is the equation for an A_{n-m-1} singularity, whereas the new relation in the second chart describes an A_{m-1} singularity! We can thus see how the blowup breaks $A_{n-1} \rightarrow A_{n-m-1} \oplus A_{m-1}$.




Figure 4.1: An A_6 surface singularity splits into a pair of singularities of type A_2 and A_3 after the open node in the A_6 diagram is blown up.

4.3 The D_{n+2} story

4.3.1 Invariant theory

\mathbb{B}_{4n} Group action

The action of \mathbb{B}_{4n} on $\mathbb{C}[x, y]$ is characterized by

$$\begin{aligned} g \cdot x &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix} = \begin{pmatrix} \zeta & 0 \\ 0 & 1 \end{pmatrix} = \zeta x, \\ g \cdot y &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix} = \begin{pmatrix} 0 & \zeta^{-1} \\ \zeta & 0 \end{pmatrix} = \zeta^{-1} y, \\ h \cdot x &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = y, \\ h \cdot y &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = -x. \end{aligned}$$

Extending the action to arbitrary monomials we have:

$$\begin{aligned} g \cdot x^j y^k &= \zeta^{j-k} x^j y^k, \\ h \cdot x^j y^k &= (-1)^k x^k y^j. \end{aligned}$$

\mathbb{B}_{4n} Invariant theory

Since $\zeta^{2n} = 1$, the invariant polynomials under $\mathbb{Z}_{2n} \subset \mathbb{B}_{4n}$ are generated by:

$$x^{2n}, \quad xy, \quad y^{2n}.$$

However, $h \cdot x^{2n} = y^{2n}$, $h \cdot y^{2n} = x^{2n}$, and $h \cdot xy = -xy$. If we define

$$X = \frac{i}{2}xy(x^{2n} - y^{2n}), \quad Y = \frac{1}{2}(x^{2n} + y^{2n}), \quad Z = x^2 y^2,$$

then X, Y and Z are invariant under $\Gamma = \mathbb{B}_{4n}$. In particular, X, Y and Z satisfy the equation:

$$X^2 + Y^2 Z - Z^{n+1} = 0,$$

which is preferred versal form for a D_{n+2} singularity.

To construct irreducible representations of \mathbb{B}_{4n} in $\mathbb{C}[x, y]$, we start with one-dimensional representations under \mathbb{Z}_{2n} and complete the orbit under \mathbb{B}_{4n} . Since $h^2 = -1$, the pairs $(x^j y^k, h \cdot x^j y^k)$ span complete and irreducible representations of \mathbb{B}_{4n} . Observe that when $j \neq k$ the representations are two-dimensional, while $j = k$ yields one-dimensional representations. Again, we label the representations by the character evaluated on g . For character $\zeta^m + \zeta^{-m}$, $m = 1 \dots 2n$, we have:

$$\zeta^m + \zeta^{-m} : \{(x^j y^k, (-1)^k x^k y^j)\} \quad \text{s.t. } j - k \equiv m \pmod{2n}.$$

Let $\mathbb{C}[x, y]^\Gamma$ denote the subring of invariant functions. It is generated by X, Y and Z , and can also be written as

$$\mathbb{C}[x, y]^\Gamma = \mathbb{C}[X, Y, Z]/(X^2 + Y^2 Z - Z^{n+1}).$$

By definition Γ acts trivially on $\mathbb{C}[x, y]^\Gamma$, so multiplying a monomial by an element of the invariant ring will yield a new representation with the same character. Consider the $\zeta^m + \zeta^{-m}$ representations,

$$(x^j y^k, (-1)^k x^k y^j) \text{ where } j - k = m + 2nq.$$

We split these into four cases:

$$\begin{aligned} q \geq 0, k \text{ even} & : & (x^j y^k, x^k y^j) &= (x^{j-k}, y^{j-k}) Z^{k/2} \\ & & &\in (x^{m+2nq}, y^{m+2nq}) \mathbb{C}[x, y]^\Gamma, \\ q \geq 0, k \text{ odd} & : & (x^j y^k, -x^k y^j) &= (x^{j-k+1} y, -y^{j-k+1} x) Z^{(k-1)/2} \\ & & &\in (x^{m+1+2nq} y, -y^{m+1+2nq} x) \mathbb{C}[x, y]^\Gamma, \\ q \leq -1, j \text{ even} & : & (x^j y^k, (-1)^k x^k y^j) &= (y^{k-j}, (-1)^k x^{k-j}) Z^{j/2} \\ & & &\in (y^{-m-2nq}, (-1)^m x^{-m-2nq}) \mathbb{C}[x, y]^\Gamma, \\ q \leq -1, j \text{ odd} & : & (x^j y^k, (-1)^k x^k y^j) &= (y^{k-j+1} x, (-1)^k x^{k-j+1} y) Z^{(j-1)/2} \\ & & &\in (y^{-m-2nq+1} x, (-1)^{m+1} x^{-m-2nq+1} y) \mathbb{C}[x, y]^\Gamma. \end{aligned}$$

It seems at first glance that we have four infinite families of generators for the $\mathbb{C}[x, y]^\Gamma$ -module corresponding to the $\zeta^m + \zeta^{-m}$ representations:

$$\begin{aligned} s_1^{(q)} &= \begin{pmatrix} x^{m+2nq} \\ y^{m+2nq} \end{pmatrix}, & s_2^{(q)} &= \begin{pmatrix} y^{-2nq-m} \\ (-1)^m x^{-2nq-m} \end{pmatrix}, \\ s_3^{(q)} &= \begin{pmatrix} x^{m+1+2nq} y \\ -y^{m+1+2nq} x \end{pmatrix}, & s_4^{(q)} &= \begin{pmatrix} y^{-2nq-m+1} x \\ (-1)^{m+1} x^{-2nq-m+1} y \end{pmatrix}. \end{aligned}$$

Since neither x^{2n} nor y^{2n} are invariant functions, it is not obvious that we don't need the generators for all values of q : $q \geq 0$ for $s_1^{(q)}, s_3^{(q)}$ and $q \leq -1$ for $s_2^{(q)}, s_4^{(q)}$.

The module

It seems that in the end there will only be four generators for this module, with relations. We conjecture is that the following four representations (one from each of the four families) generate the module:

$$s_1 = \begin{pmatrix} x^m \\ y^m \end{pmatrix}, \quad s_2 = \begin{pmatrix} y^{2n-m} \\ (-1)^m x^{2n-m} \end{pmatrix}, \quad s_3 = \begin{pmatrix} x^{m+1} y \\ -y^{m+1} x \end{pmatrix}, \quad s_4 = \begin{pmatrix} y^{2n-m+1} x \\ (-1)^{m+1} x^{2n-m+1} y \end{pmatrix}.$$

Note that $s_1 = s_1^{(0)}, s_2 = s_2^{(-1)}, s_3 = s_3^{(0)}$, and $s_4 = s_4^{(-1)}$.

4.3.2 Matrix of relations

Cramer's rule

The module is not free, however; there are relations among the generators. We can compute these using Cramer's rule, which states that the solution to a system of equations

$$Av = b,$$

is given by

$$v_j = \frac{\det M_j}{\det A},$$

where M_j is the matrix obtained from A by replacing the j th column with the vector b . If we want to solve for s_3 and s_4 in terms of s_1 and s_2 , our systems of equations will look like

$$\begin{pmatrix} s_1 & s_2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = s_3,$$

$$\begin{pmatrix} s_1 & s_2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = s_4.$$

In order to implement Cramer's rule, we first compute all of the pairwise determinants $\det(s_i s_j)$. By restricting to m even or m odd we can also write these in terms of the invariant functions X, Y and Z .

$\det(s_i s_j)$	any m	m even	m odd
$\det(s_1 s_2)$	$(-1)^m x^{2n} - y^{2n}$	$-2iX/\sqrt{Z}$	$-2Y$
$\det(s_1 s_3)$	$-2x^{m+1}y^{m+1}$	$-2\sqrt{Z}Z^{m/2}$	$-2Z^{(m+1)/2}$
$\det(s_1 s_4)$	$(-1)^{m+1}x^{2n+1}y - y^{2n+1}x$	$-2\sqrt{Z}Y$	$-2iX$
$\det(s_2 s_3)$	$(-1)^{m+1}x^{2n+1}y - y^{2n+1}x$	$-2\sqrt{Z}Y$	$-2iX$
$\det(s_2 s_4)$	$2(-1)^{m+1}x^{2n-m+1}y^{2n-m+1}$	$-2\sqrt{Z}Z^{n-m/2}$	$2Z^{n+1-(m+1)/2}$
$\det(s_3 s_4)$	$(-1)^{m+1}x^{2n+2}y^2 + y^{2n+2}x^2$	$2i\sqrt{Z}X$	$2ZY$

For m even,

$$b = s_3 : \quad v_1 = \frac{\det(s_3 s_2)}{\det(s_1 s_2)} = \frac{iYZ}{X}, \quad v_2 = \frac{\det(s_1 s_3)}{\det(s_1 s_2)} = \frac{-iZ^{m/2+1}}{X}.$$

$$b = s_4 : \quad v_1 = \frac{\det(s_4 s_2)}{\det(s_1 s_2)} = \frac{iZ^{n+1-m/2}}{X}, \quad v_2 = \frac{\det(s_1 s_4)}{\det(s_1 s_2)} = \frac{-iYZ}{X}.$$

For m odd,

$$b = s_3 : \quad v_1 = \frac{\det(s_3 s_2)}{\det(s_1 s_2)} = \frac{-iX}{Y}, \quad v_2 = \frac{\det(s_1 s_3)}{\det(s_1 s_2)} = \frac{Z^{(m+1)/2}}{Y}.$$

$$b = s_4 : \quad v_1 = \frac{\det(s_4 s_2)}{\det(s_1 s_2)} = \frac{Z^{n+1-(m+1)/2}}{Y}, \quad v_2 = \frac{\det(s_1 s_4)}{\det(s_1 s_2)} = \frac{iX}{Y}.$$

Matrices of relations

In the case of m odd, we find the following matrix of relations:

$$m \text{ odd} \quad \begin{pmatrix} s_1 & s_2 & s_3 & s_4 \\ -iX & Z^{(m+1)/2} & -Y & 0 \\ Z^{n+1-(m+1)/2} & iX & 0 & -Y \\ -YZ & 0 & -iX & Z^{(m+1)/2} \\ 0 & -YZ & Z^{n+1-(m+1)/2} & iX \end{pmatrix}$$

For m even, we initially find the matrix:

$$m \text{ even} \quad \begin{pmatrix} s_1 & s_2 & s_3 & s_4 \\ iYZ & -iZ^{m/2+1} & -X & 0 \\ iZ^{n+1-m/2} & -iYZ & 0 & -X \\ -X & 0 & iY & -iZ^{m/2} \\ 0 & -iX & -Z^{n-m/2} & Y \end{pmatrix}$$

Switching the top two rows with the bottom two

$$m \text{ even} \quad \begin{pmatrix} s_1 & s_2 & s_3 & s_4 \\ -X & 0 & iY & -iZ^{m/2} \\ 0 & -iX & -Z^{n-m/2} & Y \\ iYZ & -iZ^{m/2+1} & -X & 0 \\ iZ^{n+1-m/2} & -iYZ & 0 & -X \end{pmatrix}$$

and multiplying the first and third rows by i , the second row by -1 , and the fourth row by $-i$ (we can do this because the rows just reflect relations among the generators s_1, \dots, s_4 and can thus be shuffled and scaled independently):

$$m \text{ even} \quad \begin{pmatrix} s_1 & s_2 & s_3 & s_4 \\ -iX & 0 & -Y & Z^{m/2} \\ 0 & iX & Z^{n-m/2} & -Y \\ -YZ & Z^{m/2+1} & -iX & 0 \\ Z^{n+1-m/2} & -YZ & 0 & iX \end{pmatrix}$$

Now the m even matrix matches the m odd one!

Matrix factorizations

Now in both the even and odd cases, we can manipulate the relations to find syzygies, modulo the defining equation $-X^2 - Y^2Z + Z^{n+1}$. There are four syzygies which are easy to determine; these can also be represented as a 4×4 matrix which multiplies by the relations matrix to give zero (modulo the equation).

In the case of m odd, the syzygies \mathcal{S} and relations \mathcal{R} matrices are

$$\mathcal{S}_{\text{odd}} = \begin{pmatrix} -iX & Z^{(m+1)/2} & Y & 0 \\ Z^{n+1-(m+1)/2} & iX & 0 & Y \\ YZ & 0 & -iX & Z^{(m+1)/2} \\ 0 & YZ & Z^{n+1-(m+1)/2} & iX \end{pmatrix}$$

$$\mathcal{R}_{\text{odd}} = \begin{pmatrix} -iX & Z^{(m+1)/2} & -Y & 0 \\ Z^{n+1-(m+1)/2} & iX & 0 & -Y \\ -YZ & 0 & -iX & Z^{(m+1)/2} \\ 0 & -YZ & Z^{n+1-(m+1)/2} & iX \end{pmatrix}$$

The product of these matrices is $(-X^2 - Y^2Z + Z^{n+1})$ times the 4×4 identity matrix.

In the case of m even, the syzygies and relations matrices are

$$\mathcal{S}_{\text{even}} = \begin{pmatrix} -iX & 0 & Y & Z^{m/2} \\ 0 & iX & Z^{n-m/2} & Y \\ YZ & Z^{m/2+1} & -iX & 0 \\ Z^{n+1-m/2} & YZ & 0 & iX \end{pmatrix}$$

$$\mathcal{R}_{\text{even}} = \begin{pmatrix} -iX & 0 & -Y & Z^{m/2} \\ 0 & iX & Z^{n-m/2} & -Y \\ -YZ & Z^{m/2+1} & -iX & 0 \\ Z^{n+1-m/2} & -YZ & 0 & iX \end{pmatrix}$$

The product of these matrices is again $(-X^2 - Y^2Z + Z^{n+1})$ times the 4×4 identity matrix.

4.3.3 Summary of results

$\Gamma = \mathbb{B}_{4n}$, $X = \text{Spec } \mathbb{C}[x, y]^\Gamma$. If we let

$$X = \frac{i}{2}xy(x^{2n} - y^{2n}), \quad Y = \frac{1}{2}(x^{2n} + y^{2n}), \quad Z = x^2y^2,$$

then the ring of invariant polynomials can be written

$$\mathbb{C}[x, y]^\Gamma = \mathbb{C}[X, Y, Z]/(X^2 + Y^2Z - Z^{n+1}),$$

which we recognize as a space with D_{n+2} singularity at the origin. The sheaf of \mathcal{O}_X -modules of interest is:

$$\mathcal{F} = \{s_1, s_2, s_3, s_4\} \cdot \mathcal{O}_X,$$

where

$$s_1 = \begin{pmatrix} x^m \\ y^m \end{pmatrix}, \quad s_2 = \begin{pmatrix} y^{2n-m} \\ (-1)^m x^{2n-m} \end{pmatrix}, \quad s_3 = \begin{pmatrix} x^{m+1}y \\ -y^{m+1}x \end{pmatrix}, \quad s_4 = \begin{pmatrix} y^{2n-m+1}x \\ (-1)^{m+1}x^{2n-m+1}y \end{pmatrix}.$$

The relations among these generators depend on whether m is even or odd. In each case we have found 4×4 matrices of relations. As in the A_{n-1} case, we only need half of the relations to do the blowup. In both the m even and the m odd cases we find that it is simplest to choose the top two relations, which express s_3, s_4 in terms of s_1, s_2 .

In the case of m odd, the relations matrix is

$$\mathcal{R}_{\text{odd}} = \begin{pmatrix} s_1 & s_2 & s_3 & s_4 \\ -iX & Z^{(m+1)/2} & -Y & 0 \\ Z^{n+1-(m+1)/2} & iX & 0 & -Y \\ -YZ & 0 & -iX & Z^{(m+1)/2} \\ 0 & -YZ & Z^{n+1-(m+1)/2} & iX \end{pmatrix}$$

In the case of m even, the relations matrix is

$$\mathcal{R}_{\text{even}} = \begin{pmatrix} s_1 & s_2 & s_3 & s_4 \\ -iX & 0 & -Y & Z^{m/2} \\ 0 & iX & Z^{n-m/2} & -Y \\ -YZ & Z^{m/2+1} & -iX & 0 \\ Z^{n+1-m/2} & -YZ & 0 & iX \end{pmatrix}$$

4.3.4 How to use this to blow up

Our “relations” matrix \mathcal{R} is a 4×4 matrix which describes a map $A^4 \rightarrow A^4$, where A is the coordinate ring of our space M (a D_{n+2} singularity). We have $A = \mathbb{C}[X, Y, Z]/E$, where E is the equation given above.

We calculated that for the syzygies matrix \mathcal{S} and the relations matrix \mathcal{R} , we have

$$\mathcal{S}\mathcal{R} = EI_4.$$

This implies that in the quotient field,

$$\mathcal{R}^{-1} = \frac{1}{E}\mathcal{S}$$

but also, since $\det \mathcal{R} = E^2$, the matrix

$$E^2\mathcal{R}^{-1} = E\mathcal{S}$$

is the matrix of cofactors of \mathcal{R} , i.e., the matrix whose entries are all the 3×3 sub-determinants of \mathcal{R} .

It follows that when restricted to our space $E = 0$, the matrix of cofactors vanishes identically. Thus, \mathcal{R} generically has rank at most 2.

We see, therefore, that \mathcal{R} defines a module over A which can be generated by 4 elements yet has rank 2 (unless the entries in \mathcal{R} are not generic, in which case the rank could be 1 or 0). To find the smallest blowup of our space on which this module becomes locally free, we use a variant of the blowup construction based on a Grassmannian rather than projective space. That is, we consider the closure of the set

$$\{(m, g) \in M \times G(2, 4) \mid g = \text{fiber of our module over } m\}.$$

(At points at which the rank of the cokernel of \mathcal{R} is not 2, this definition does not apply: that is why we need the closure.)

We can explicitly describe points in $G(2, 4)$ by means of Plücker coordinates, and as in the usual blowup, we can take these to be proportional to certain quantities formed out of the matrix \mathcal{R} . The Plücker coordinates correspond to 2×2 minors of the original matrix, and we’ll end up with relations like:

$$[\alpha_1, \dots, \alpha_6] = [X^2 - Z^{n+1}, YZ^{n+1-(m+1)/2}, iXY, \dots]$$

(where the equality is evaluated in \mathbb{P}^5). In addition, we need to impose the equations satisfied by the Grassmannian $G(2, 4) \subset \mathbb{P}^5$ which in this case is a single quadratic equation.

To see the equation of $G(2, 4) \subseteq \mathbb{P}^5$, let v_1, v_2, v_3, v_4 span a 2-dimensional subspace of \mathbb{C}^2 . Cramer’s rule tells us that

$$(v_1 \wedge v_2)v_3 + v_1(v_2 \wedge v_3) + (v_3 \wedge v_1)v_2 = 0.$$

Wedging the above equation with v_4 yields

$$(v_1 \wedge v_2)(v_3 \wedge v_4) + (v_1 \wedge v_4)(v_2 \wedge v_3) - (v_1 \wedge v_3)(v_2 \wedge v_4) = 0.$$

In terms of the Plücker coordinates

$$\gamma = v_1 \wedge v_2, \quad \beta = v_1 \wedge v_3, \quad \delta = v_1 \wedge v_4, \quad \alpha = v_2 \wedge v_3, \quad \varepsilon = v_2 \wedge v_4, \quad \varphi = v_3 \wedge v_4,$$

the equation satisfied by $G(2, 4)$ is

$$\gamma\varphi + \delta\alpha - \beta\varepsilon = 0. \tag{4.1}$$

4.3.5 The blowup

We can find additional relations among the Plücker coordinates by restricting attention to the 2×2 minors in the top two rows of the relations matrices \mathcal{R}_{odd} and $\mathcal{R}_{\text{even}}$. Together with the relation (4.1)

$$\gamma\varphi + \delta\alpha - \beta\varepsilon = 0,$$

this gives us the blowup.

m odd

The top two rows of \mathcal{R}_{odd} gives

$$\begin{pmatrix} -iX & Z^{(m+1)/2} & -Y & 0 \\ Z^{n+1-(m+1)/2} & iX & 0 & -Y \end{pmatrix}$$

and has 2×2 minors $[ij]$ (where i and j denote columns):

$$\begin{aligned} \gamma &= [12] = X^2 - Z^{n+1}, & \varphi &= [34] = Y^2, \\ \beta &= [13] = YZ^{n+1-(m+1)/2}, & \varepsilon &= [24] = -YZ^{(m+1)/2}, \\ \delta &= [14] = iXY, & \alpha &= [23] = iXY. \end{aligned}$$

We find additional Plücker coordinate relations

$$\begin{aligned} \delta &= \alpha, \\ \beta &= -Z^{n-m}\varepsilon, \\ \gamma &\equiv -Z\varphi \pmod{\text{equation}}, \end{aligned}$$

where the last relation holds modulo the equation for the singular surface

$$X^2 + Y^2Z - Z^{n+1} = 0.$$

m even

The top two rows of $\mathcal{R}_{\text{even}}$ gives

$$\begin{pmatrix} -iX & 0 & -Y & Z^{m/2} \\ 0 & iX & Z^{n-m/2} & -Y \end{pmatrix}$$

and has 2×2 minors $[ij]$ (where i and j denote columns):

$$\begin{aligned} \gamma &= [12] = X^2, & \varphi &= [34] = Y^2 - Z^n, \\ \beta &= [13] = -iXZ^{n-m/2}, & \varepsilon &= [24] = -iXZ^{m/2}, \\ \delta &= [14] = iXY, & \alpha &= [23] = iXY. \end{aligned}$$

Again we find Plücker coordinate relations

$$\begin{aligned} \delta &= \alpha, \\ \beta &= Z^{n-m}\varepsilon, \\ \gamma &\equiv -Z\varphi \pmod{\text{equation}}. \end{aligned}$$

Identification of residual singularities

We can combine the two cases by writing the Plücker coordinate relations as

$$\begin{aligned}\delta &= \alpha, \\ \beta &= (-1)^m Z^{n-m} \varepsilon, \\ \gamma &\equiv -Z\varphi \pmod{\text{equation}},\end{aligned}$$

The relation for the grassmanian (4.1) becomes

$$\alpha^2 - \varphi^2 Z + (-1)^{m+1} \varepsilon^2 Z^{n-m} = 0.$$

The interesting charts for the blowup are $\varphi = 1$ and $\varepsilon = 1$.

m odd

($\varphi = 1$) In this chart we have

$$\varepsilon = \varphi \varepsilon_1, \quad \alpha = \varphi \alpha_1, \quad \alpha_1^2 - Z + \varepsilon_1^2 Z^{n-m} = 0.$$

To see what kind of singularity this gives, define $G(Z, \alpha_1)$ by

$$Z^{(m+1)/2} - \alpha_1^{m+1} = (Z - \alpha_1^2)G(Z, \alpha_1),$$

and notice that

$$-Z^{(m+1)/2} = \varepsilon_1 Y, \quad \text{and} \quad Z - \alpha_1^2 = \varepsilon_1^2 Z^{n-m}.$$

We are thus left with

$$-\alpha_1^{m+1} = \varepsilon_1 (Y + \varepsilon_1 Z^{n-m} G(Z, \alpha_1)) \stackrel{\text{def}}{=} \varepsilon_1 \tilde{Y},$$

which is the equation for an A_m singularity.

($\varepsilon = 1$) In this chart

$$\alpha = \varepsilon \alpha_2, \quad \varphi = \varepsilon \varphi_2, \quad \alpha_2^2 - \varphi_2^2 Z + Z^{n-m} = 0,$$

which we immediately recognize as a D_{n-m+1} singularity.

m even

($\varphi = 1$) In this chart we have

$$\varepsilon = \varphi \varepsilon_1, \quad \alpha = \varphi \alpha_1, \quad \alpha_1^2 - Z - \varepsilon_1^2 Z^{n-m} = 0.$$

To see what kind of singularity this gives, define $G(Z, \alpha_1)$ by

$$Z^{m/2} - \alpha_1^m = (Z - \alpha_1^2)G(Z, \alpha_1),$$

and notice that

$$-Z^{m/2}\alpha_1 = \varepsilon_1 Y, \quad \text{and} \quad Z - \alpha_1^2 = -\varepsilon_1^2 Z^{n-m}.$$

We are thus left with

$$-\alpha_1^{m+1} = \varepsilon_1 (Y - \alpha_1 \varepsilon_1 Z^{n-m} G(Z, \alpha_1)) \stackrel{\text{def}}{=} \varepsilon_1 \tilde{Y},$$

which is the equation for an A_m singularity.

($\varepsilon = 1$) In this chart

$$\alpha = \varepsilon \alpha_2, \quad \varphi = \varepsilon \varphi_2, \quad \alpha_2^2 - \varphi_2^2 Z - Z^{n-m} = 0,$$

which we immediately recognize as a D_{n-m+1} singularity.

We have thus seen that in both the odd and even cases, the blowup breaks the D_{n+2} surface singularity into two “lower order” singularities

$$D_{n+2} \longrightarrow D_{n-m+1} \oplus A_m.$$

This is exactly what remains in the diagram once the blown up node has been removed!




Figure 4.2: The blow up of the open length 2 node in the top Dynkin diagram splits the D_8 surface singularity into two lower-order surface singularities: A_2 and D_5 .

Chapter 5

Small resolutions

5.1 “Bottom-up” approach

In light of the algebraic geometry results of Laufer, Katz, and Morrison [20, 19] (see Introduction for more details), we know that there are only 6 types of geometries with isolated singularities which fit into Ferrari’s framework. These singular threefolds may have hyperplane sections pertaining to any of the ADE Dynkin diagrams, but the *generic hyperplane section* is determined by the *length* of the singular point. The small resolution can be obtained by blowing up any node of the same length in any of the diagrams corresponding to a hyperplane section, even if it’s not the generic one.

In the following “bottom up” approach, we will tackle the problem of starting from one of the known singular threefolds, blowing up an arbitrary node in the associated A_n or D_{n+2} Dynkin diagram (this corresponds to a hyperplane section which is not generic for $n > 1$), and then identifying the associated matrix model superpotential. In this way we hope to classify all of the matrix models which can be engineered by Calabi-Yau geometries following Ferrari’s construction. The basic outline of the strategy is as follows:

- Start with a singular Calabi-Yau \mathcal{M}_0 with isolated Gorenstein singularity p which admits an irreducible small resolution.
- Blow up a node of a corresponding Dynkin diagram to get the small resolution $\widehat{\mathcal{M}}$.
- Write the smooth space $\widehat{\mathcal{M}}$ in terms of transition functions in two charts over exceptional \mathbb{P}^1 .
- Identify superpotential $W(x_1, \dots, x_M)$!

We find that explicitly performing the small resolution is already a challenge. Already in the length 2 case (D_{n+2}) this is highly non-trivial. We are able to find the small resolutions by deforming the matrix factorizations from Chapter 4. Thus, in this section we prove Theorem 1:

Theorem 1. *For Gorenstein threefold singularities of length 1 and length 2, with hyperplane section an A_n or D_{n+2} surface singularity, the small resolution is obtained by deforming the matrix factorization for a node of the same length in the corresponding Dynkin diagram.*

Perhaps surprisingly, once we are able to find the small resolution it is not easy to identify the corresponding superpotential! Already in the length 2 case, we find the task of identifying the right coordinates for a description of the blowup commensurate with Ferrari's framework to be too difficult without further help. In the length 1 case, we can identify transition functions but only when we restrict ourselves to the A_1 case.

5.2 The A_{n-1} story

Here we discuss the length 1 case. The invariant theory can be found in [19, pages 467–469]. The small resolution via deformation of the A_{n-1} matrix factorization proves the first part of Theorem 1.

5.2.1 Invariant theory

We start with the equation for an A_{n-1} singularity in preferred versal form,

$$\Phi_{A_{n-1}} : -XY + Z^n + \sum_{i=2}^n \alpha_i Z^{n-i}.$$

Introducing the distinguished polynomial

$$f_{n-1}(U, t) = \prod_{t=1}^n (U + t_i), \quad \sum_i t_i = 0,$$

we can rewrite the equation as

$$\Phi_{A_{n-1}} : -XY + f_{n-1}(Z, t).$$

If we are to break

$$A_{n-1} \longrightarrow A_{n-m-1} \oplus A_{m-1},$$

we expect the original deformed equation to factor in a way which preserves the residual symmetries. Rewriting f_{n-1} as¹

$$f_{n-1}(U, t) = \prod_{t=1}^{n-m} (U + t_i) \prod_{t=1}^m (U + t_i) = f'_{n-m-1}(U, t) f''_{m-1}(U, t),$$

we find the factorization

$$\Phi_{A_{n-1}} : -XY + f'_{n-m-1}(Z, t) f''_{m-1}(Z, t). \tag{5.1}$$

¹Note that primes do not indicate derivatives!

5.2.2 Matrix factorization with deformation

In order to “guess” a deformation for the matrix factorization from Chapter 4

$$\mathcal{R} = \begin{pmatrix} Y & -Z^m \\ Z^{n-m} & -X \end{pmatrix}, \quad \mathcal{S} = \begin{pmatrix} X & -Z^m \\ Z^{n-m} & -Y \end{pmatrix},$$

first note that in the absence of deformation ($t = 0$) the allowed invariant polynomials become

$$f'_{n-m-1}(Z) = Z^{n-m}, \quad f''_{m-1}(Z) = Z^m.$$

This leads us naturally to try the deformed matrix factorization

$$\mathcal{R}_{\text{def}} = \begin{pmatrix} Y & -f''_{m-1}(Z, t) \\ f'_{n-m-1}(Z, t) & -X \end{pmatrix}, \quad \mathcal{S}_{\text{def}} = \begin{pmatrix} X & -f''_{m-1}(Z, t) \\ f'_{n-m-1}(Z, t) & -Y \end{pmatrix}.$$

We find that this is a valid matrix factorization for the deformed equation (5.1), as

$$\mathcal{R}_{\text{def}}\mathcal{S}_{\text{def}} = \mathcal{S}_{\text{def}}\mathcal{R}_{\text{def}} = (XY - f'_{n-m-1}(Z, t)f''_{m-1}(Z, t))\mathbf{1}_{2 \times 2}.$$

5.2.3 The blowup

We blow up the Weil divisor $Y = f''_{m-1}(Z, t) = 0$, which corresponds to the ideal

$$I = \langle Y, f''_{m-1}(Z, t) \rangle.$$

We introduce coordinates $[\mu, \nu]$ for \mathbb{P}^1 , and the relation

$$\mu Y = \nu f''_{m-1}(Z, t).$$

We find the new relation and proper transform of the original curve in two coordinate charts:

$\mu = 1$	$\nu = 1$
$Y = \nu f''_{m-1}(Z, t)$	$\mu Y = f''_{m-1}(Z, t) \rightarrow (A_{m-1})$
$(A_{n-m-1}) \leftarrow X\nu = f'_{n-m-1}(Z, t)$	$X = \mu f'_{n-m-1}(Z, t)$
(ν, X, Z, t)	(μ, Y, Z, t)

The proper transform in the first chart is the equation for an A_{n-m-1} singularity, whereas the new relation in the second chart describes an A_{m-1} singularity! We can thus see how (restricted to a hyperplane) the blowup breaks $A_{n-1} \rightarrow A_{n-m-1} \oplus A_{m-1}$.




Figure 5.1: An A_6 surface singularity splits into a pair of singularities of type A_2 and A_3 after the open node in the A_6 diagram is blown up. The length 1 singularity in the threefold can be thought of as a deformation of this surface singularity, which is in turn given by a (non-generic) hyperplane section of the total space. As far as the hypersurface is concerned, there are still remaining residual singularities after the blowup. In contrast, the total space of the threefold is smooth upon blowing up a single length 1 node in the diagram.

5.2.4 Identification of superpotential in A_1 case

We now focus on threefold singularities which are deformations of the case A_1 ($n = 2, m = 1$). By considering higher order deformations t^k for $k > 1$, we recover superpotentials for all A_k .

Higher order deformations

We start with the equation for an A_1 singularity with higher order deformations:

$$XY = Z^2 - t^{2k} = (Z + t^k)(Z - t^k),$$

for $k \geq 1$. We blow up the ideal $I = \langle Y, Z + t^k \rangle$, corresponding to the Weil divisor

$$Y = Z + t^k = 0.$$

We introduce coordinates $[\gamma, \beta]$ for the \mathbb{P}^1 , and the relation

$$\beta Y = \gamma(Z + t^k).$$

The blowup is described in two charts:

$$\left. \begin{array}{l} \gamma \text{ chart } (\gamma, X, Z, t) \\ Y = \gamma(Z + t^k) \\ H = \gamma X - Z + t^k = 0 \end{array} \right| \begin{array}{l} \beta \text{ chart } (\beta, X, Y, t) \\ Z + t^k = \beta Y \\ F = X - \beta^2 Y + 2\beta t^k = 0 \end{array}$$

The transition functions

$$\beta = \gamma^{-1}, \quad X = X, \quad Y = \gamma(Z + t^k), \quad F = \gamma^{-1}H,$$

suggest that the exact sequence of normal bundles

$$0 \longrightarrow N_{C/T} \longrightarrow N_{C/F} \longrightarrow N_{T/F} \longrightarrow 0$$

for the curve C within the threefold T , inside the fourfold F , is given by

$$\begin{array}{l} \gamma \text{ chart} \quad (t, X, Z) \quad \begin{bmatrix} 0 \\ \gamma \\ -1 \end{bmatrix} \quad H_0 = \gamma X - Z \\ \\ 0 \quad \longrightarrow \quad N_{C/T} \quad \xrightarrow{?} \quad \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(-1) \quad \longrightarrow \quad \mathcal{O}(1) \quad \longrightarrow \quad 0 \\ \\ \beta \text{ chart} \quad (t, X, Y) \quad \begin{bmatrix} 0 \\ 1 \\ -\beta^2 \end{bmatrix} \quad F_0 = X - \beta^2 Y \end{array}$$

where H_0 and F_0 are linearized versions of H and F . In homogeneous coordinates, the $N_{C/F} \longrightarrow N_{T/F}$ map becomes

$$\mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(-1) \quad \begin{bmatrix} 0 \\ \gamma \\ -\beta^2 \end{bmatrix} \quad \longrightarrow \quad \mathcal{O}(1).$$

We easily identify the following elements in the kernel:

$$[1, 0, 0], \quad [0, \beta^2, \gamma].$$

Since the corresponding matrix for the map $N_{C/T} \longrightarrow N_{C/F}$ has type

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & \beta^2 & \gamma \end{bmatrix} = \begin{bmatrix} \mathcal{O} & * & * \\ * & \mathcal{O}(2) & \mathcal{O}(1) \end{bmatrix},$$

we can conclude that

$$N_{C/T} = \mathcal{O} \oplus \mathcal{O}(-2).$$

The exact sequence with all the maps in each chart is thus,

$$\begin{array}{l} \gamma \text{ chart} \quad (t, r) \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & \gamma \end{bmatrix} \quad (t, X, Z) \quad \begin{bmatrix} 0 \\ \gamma \\ -1 \end{bmatrix} \quad H_0 = \gamma X - Z \\ \\ 0 \quad \longrightarrow \quad \mathcal{O} \oplus \mathcal{O}(-2) \quad \longrightarrow \quad \mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(-1) \quad \longrightarrow \quad \mathcal{O}(1) \quad \longrightarrow \quad 0. \\ \\ \beta \text{ chart} \quad (t, r') \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & \beta^2 & 1 \end{bmatrix} \quad (t, X, Y) \quad \begin{bmatrix} 0 \\ 1 \\ -\beta^2 \end{bmatrix} \quad F_0 = X - \beta^2 Y \end{array}$$

In particular, the maps

$$\begin{aligned}(t, r) &\longmapsto (t, r, \gamma r), \\ (t, r') &\longmapsto (t, \beta^2 r', r'),\end{aligned}$$

imply that

$$r = X, \quad r' = Y.$$

Using the full nonlinear transition functions, we see that

$$Y = \gamma(Z + t^k) = \gamma(\gamma X + 2t^k) = \gamma^2 X + 2\gamma t^k.$$

In other words, r, r' have transition function

$$r' = \gamma^2 r + 2\gamma t^k,$$

which has a superpotential correction to the $\mathcal{O}(-2)$ structure.

The superpotential

All together, the $\mathcal{O} \oplus \mathcal{O}(-2)$ transition functions are

$$\beta = \gamma^{-1}, \quad t = t, \quad r' = \gamma^2 r + 2\gamma t^k.$$

Comparing with Ferrari's setup,

$$\beta = \gamma^{-1}, \quad s' = \gamma^{-n} s, \quad r' = \gamma^{-m} r + \partial_s E(\gamma, s),$$

we find that $n = 0$, $m = -2$, and $M = 1$. The geometric potential is given by

$$E(\gamma, t) = \gamma \frac{2}{k+1} t^{k+1}.$$

The superpotential for our deformed A_1 case is thus,

$$\begin{aligned}W(x) &= \frac{1}{2\pi i} \oint_{C_0} \gamma^{-2} E(\gamma, x) d\gamma \\ &= \frac{2}{k+1} x^{k+1}.\end{aligned}$$

Note that the normal bundle to the \mathbb{P}^1 is $\mathcal{O} \oplus \mathcal{O}(-2)$ for $k > 1$, but for $k = 1$ we have a bundle-changing superpotential, and the normal bundle is $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$. In particular, the different types of A_1 deformations yield all of the A_k geometries and superpotentials.

Ferrari's example 1

We recover Ferrari's example if we let

$$t^k \longmapsto \frac{1}{2} P(t),$$

for some polynomial $P(t)$. The original deformed equation becomes

$$XY = Z^2 - P(t)^2/4,$$

and the r, r' transition function is

$$r' = \gamma^2 r + \gamma P(t).$$

If \widetilde{W} is a polynomial such that $\widetilde{W}'(t) = P(t)$, then the geometric potential is

$$E(\gamma, t) = \gamma \widetilde{W}(t),$$

and the superpotential is given by

$$\begin{aligned} W(x) &= \frac{1}{2\pi i} \oint_{C_0} \gamma^{-2} E(\gamma, x) d\gamma \\ &= \widetilde{W}(x). \end{aligned}$$

5.3 The D_{n+2} story

Now we discuss the length 2 case. The invariant theory and the Tyurina blowup (a length 1 resolution!) can be found in [19, pages 469–471]. The small resolution is obtained by deforming the matrix factorization for a length 2 node in the D_{n+2} diagram, and thus completes the proof of Theorem 1.

5.3.1 Invariant theory

We start with the equation for a D_{n+2} singularity in preferred versal form,

$$\Phi_{D_{n+2}} : X^2 + Y^2 Z - Z^{n+1} - \sum_{i=1}^{n+1} \delta_{2i} Z^{n+1-i} + 2\gamma_{n+2} Y.$$

Let t_1, \dots, t_{n+2} be coordinates on \mathbb{C}^{n+2} , and $\mathcal{W}(D_{n+2}) = (\mathbb{Z}/2)^{n+1} \ltimes S_{n+2}$ the Weyl group of the associated lie algebra. Then the above coefficients are invariant polynomials in $\mathbb{C}^{n+2}/\mathcal{W}(D_{n+2})$:

$$\begin{aligned} \gamma_{n+2} &:= t_1 \cdots t_{n+2} = s_{n+2}, \\ \delta_{2i} &:= \text{the } i\text{th elementary symmetric function of } t_1^2, \dots, t_{n+2}^2. \end{aligned}$$

Define δ by

$$\delta^2 \equiv \delta_{2n+4} = \gamma_{n+2}^2.$$

The functions $\delta, \delta_2, \dots, \delta_{2n+2}$ generate $\mathbb{C}[V]^{\mathcal{W}}$.

Factorization

In order to factor the above D_{n+2} equation, we introduce distinguished polynomials:

$$f(U, t) = \prod_{i=1}^{n+2} (U + t_i) = U^{n+2} + U \sum_{i=1}^{n+1} s_i U^{n+1-i} + \delta$$

$$g(Z, t) = \prod_{i=1}^{n+2} (Z + t_i^2) = Z^{n+2} + Z \sum_{i=1}^{n+1} \delta_{2i} Z^{n+1-i} + \delta^2.$$

Note that $f(0, t) = \delta$. We can now rewrite the D_{n+2} equation as

$$\Phi_{D_{n+2}} : X^2 + Y^2 Z - 1/Z(g(Z, t) - \delta^2) + 2\delta Y. \quad (5.2)$$

Decomposing $f(U, t)$ into odd and even parts,

$$\begin{aligned} f(U, t) &= UP(-U^2, t) + Q(-U^2, t), \\ g(-U^2, t) &= f(U, t)f(-U, t) = (Q + UP)(Q - UP) = Q^2 - U^2 P^2, \\ g(Z, t) &= ZP(Z, t)^2 + Q(Z, t)^2. \end{aligned}$$

Finally, define $h(Z, t)$, $S(Z, t)$ and $G(Z, U, t)$ by

$$h(Z, t) = \frac{1}{Z}(g(Z, t) - \delta^2), \quad (5.3)$$

$$S(Z, t) = \frac{1}{Z}(Q(Z, t) - \delta), \quad (5.4)$$

$$UP(Z, t) + Q(Z, t) = (Z + U^2)G(Z, U, t) + f(U, t). \quad (5.5)$$

Again we rewrite the D_{n+2} equation (5.2) as

$$\begin{aligned} \Phi_{D_{n+2}} : & X^2 + Y^2 Z - 1/Z(ZP(Z, t)^2 + Q(Z, t)^2 - \delta^2) + 2\delta Y \\ &= X^2 + Y^2 Z - 1/Z(ZP(Z, t)^2 + (ZS(Z, t) + \delta)^2 - \delta^2) + 2\delta Y \\ &= X^2 + Y^2 Z - 1/Z(ZP(Z, t)^2 + ZS(Z, t)(ZS(Z, t) + 2\delta)) + 2\delta Y \\ &= X^2 + Y^2 Z - P^2 - S(ZS + 2\delta) + 2\delta Y. \end{aligned}$$

5.3.2 The Tyurina blowup

Tyurina's blow up for a D_{n+2} surface singularity [19, 23] results from a very clever factorization of the deformed equation.

The miracle

Our D_{n+2} equation now factors!

$$\Phi_{D_{n+2}} : (X + P)(X - P) + (Y - S)(YZ + SZ + 2\delta).$$

This is the factorization needed for the Tyurina blowup.

The blowup

Our refactored equation is

$$\Phi_{D_{n+2}} : (X + P)(X - P) + (Y - S)(YZ + SZ + 2\delta).$$

We blow up the top node in the Dynkin Diagram for D_{n+2} , corresponding to the Weil divisor $X + P = Y - S = 0$. This leaves us with an A_{n+1} singularity, because the remaining diagram has n connected nodes.

The blow up yields

$$(X - P) = (Y - S)U, \quad \tilde{\Phi}_{D_{n+2}} : (X + P)U + (YZ + SZ + 2\delta).$$

Since $X + P = (Y - S)U + 2P$, and $Q = SZ + \delta$, we can rewrite the $\tilde{\Phi}_{D_{n+2}}$ equation as

$$\tilde{\Phi}_{D_{n+2}} : (Y - S)U^2 + 2UP + (Y - S)Z + 2(SZ + \delta) = (Y - S)(Z + U^2) + 2(UP + Q).$$

Using the definition of G we find

$$\tilde{\Phi}_{D_{n+2}} : (Y - S)(Z + U^2) + 2(Z + U^2)G + 2f(U, t) = (Z + U^2)(Y - S + 2G) + 2f(U, t).$$

Defining

$$\begin{aligned} \tilde{Z} &= Z + U^2, \\ \tilde{Y} &= Y - S + 2G, \end{aligned}$$

we see that the proper transform $\tilde{\Phi}_{D_{n+2}}$ is of the form

$$\tilde{Y}\tilde{Z} = -2f_{n+2}(U, t),$$

which corresponds to an A_{n+1} singularity.

Unfortunately, we see that this corresponds to blowing up a length 1 node in the associated Dynkin diagram, which will **not** yield a small resolution for the length 2 threefold singularity type. In the next section, we show how to blow up length 2 node in the non-deformed case (i.e. for a surface singularity). The results will help us to later do the blow up in the fully deformed case.

5.3.3 Comparison with non-deformed version

To ensure that our deformed equations match the non-deformed ones in the appropriate ($t = 0$) limit, we explicitly compute the newly defined polynomials in the non-deformed case:

$$\begin{aligned} f(U, t) &= UP(-U^2, t) + Q(-U^2, t), \\ g(-U^2, t) &= f(U, t)f(-U, t) = (Q + UP)(Q - UP) = Q^2 - U^2P^2, \\ g(Z, t) &= ZP(Z, t)^2 + Q(Z, t)^2. \end{aligned}$$




Figure 5.2: The Tyurina blowup. The special factorization of the (deformed) equation for a D_{n+2} singularity allows us to blow up the open length 1 node in the top Dynkin diagram. The resulting space has a residual A_{n+1} surface singularity. Unfortunately, this is not the right node to blow up when resolving a length 2 threefold singularity. The small resolution for this singularity type can only come from blowing up a length 2 node in the diagram for a corresponding hyperplane section.

Setting the deformation parameter $t = 0$,

$$f(U, t) = U^{n+2}.$$

The computation of $P(Z), Q(Z)$ breaks up into four cases:

$$\begin{aligned} n = 0 \quad \text{mod } 4 : & \quad P(Z) = 0, & \quad Q(Z) = -Z^{(n+2)/2}, \\ n = 1 \quad \text{mod } 4 : & \quad P(Z) = -Z^{(n+1)/2}, & \quad Q(Z) = 0, \\ n = 2 \quad \text{mod } 4 : & \quad P(Z) = 0, & \quad Q(Z) = Z^{(n+2)/2}, \\ n = 3 \quad \text{mod } 4 : & \quad P(Z) = Z^{(n+1)/2}, & \quad Q(Z) = 0. \end{aligned}$$

Moreover, in each of these cases we find

$$g(Z) = Z^{n+2}.$$

In particular, after doing the $D_{n+2} \longrightarrow D_{n-m+1} \oplus A_m$ blowup, the invariant functions $g'_{n-m+1}(Z)$ and $f''_{m+1}(U)$ in the non-deformed case are given by:

$$\begin{aligned} m = 1 \quad \text{mod } 4 : & \quad P''(Z) = 0, & \quad Q''(Z) = -Z^{(m+1)/2}, \\ m = 2 \quad \text{mod } 4 : & \quad P''(Z) = -Z^{m/2}, & \quad Q''(Z) = 0, \\ m = 3 \quad \text{mod } 4 : & \quad P''(Z) = 0, & \quad Q''(Z) = Z^{(m+1)/2}, \\ m = 0 \quad \text{mod } 4 : & \quad P''(Z) = Z^{m/2}, & \quad Q''(Z) = 0, \end{aligned}$$

and

$$f''(U) = U^{m+1}, \quad g'(Z) = Z^{n-m+1}.$$

5.3.4 Matrix of relations for the full deformation

Invariant theory

If we are to break

$$D_{n+2} \longrightarrow D_{n-m+1} \oplus A_m,$$

we expect the original equation with deformations to factor in a certain way. Recalling the definition

$$h(Z, t) = \frac{1}{Z}(g(Z, t) - \delta^2),$$

we see that we can write the deformed D_{n+2} equation (5.2) as

$$\Phi_{D_{n+2}} : X^2 + Y^2 Z - h(Z, t) + 2\delta Y, \quad (5.6)$$

where

$$\begin{aligned} f_{n+2}(U, t) &= \prod_{i=1}^{n+2} (U + t_i) = f'_{n-m+1}(U) f''_{m+1}(U), \\ g_{n+2}(Z, t) &= \prod_{i=1}^{n+2} (Z + t_i^2) = g'_{n-m+1}(Z) g''_{m+1}(Z), \\ h_{n+1}(Z, t) &= \frac{1}{Z}(g - \delta^2). \end{aligned}$$

(Note that primes do not indicate derivatives!) Recall that

$$\delta = f(0, t) = \prod_{i=1}^{n+2} t_i, \quad \delta' = \sqrt{g'(0)} = f'(0).$$

The allowed invariant functions after breaking to $D_{n-m+1} \oplus A_m$ will be:

$$g'_{n-m+1}(Z), \quad f''_{m+1}(U), \quad \text{and} \quad \delta'.$$

We can also replace $g'_{n-m+1}(Z)$ with

$$h'_{n-m}(Z) = \frac{1}{Z}(g'_{n-m+1}(Z) - \delta'^2).$$

Recalling the P, Q decomposition for $f(U)$ into odd and even parts, we have $f''_{m+1}(U) = UP''(-U^2) + Q''(-U^2)$. This yields

$$g''_{m+1} = ZP''(Z)^2 + Q''(Z)^2.$$

The importance of this is that we cannot directly use g''_{m+1} , but since f''_{m+1} is legal after the symmetry breaking, P'' and Q'' are. The legal decomposition of g_{n+2} after the blowup is then,

$$g_{n+2}(Z) = g'_{n-m+1}(Z)(ZP''(Z)^2 + Q''(Z)^2).$$

Plugging this into the deformed D_{n+2} equation yields:

$$\Phi_{D_{n+2}} : X^2 + Y^2 Z - \frac{1}{Z}(g'(Z)P''^2 + Q''^2) - \delta'^2 \delta''^2 + 2\delta' \delta'' Y.$$

Note that without deformations ($t = 0$),

$$\begin{aligned} m \text{ even} &\implies g'(Z) = Z^{n-m+1}, \quad P''(Z) = (-1)^{m/2} Z^{m/2}, \quad Q''(Z) = 0. \\ m \text{ odd} &\implies g'(Z) = Z^{n-m+1}, \quad P''(Z) = 0, \quad Q''(Z) = (-1)^{(m+1)/2} Z^{(m+1)/2}. \end{aligned}$$

Quasi-homogeneity of the matrices

Recalling the matrices of relations for these two cases: in the case of m odd, the relations matrix is

$$\begin{pmatrix} -iX & Z^{(m+1)/2} & -Y & 0 \\ Z^{n+1-(m+1)/2} & iX & 0 & -Y \\ -YZ & 0 & -iX & Z^{(m+1)/2} \\ 0 & -YZ & Z^{n+1-(m+1)/2} & iX \end{pmatrix},$$

and in the case of m even, the relations matrix is

$$\begin{pmatrix} -iX & 0 & -Y & Z^{m/2} \\ 0 & iX & Z^{n-m/2} & -Y \\ -YZ & Z^{m/2+1} & -iX & 0 \\ Z^{n+1-m/2} & -YZ & 0 & iX \end{pmatrix},$$

we are tempted to make some natural guesses for the deformed analogues. We need some clues from the quasi-homogeneity of the matrix. We assign weights in the equation

$$X^2 + Y^2 Z - Z^{n+1}$$

as follows:

$$\deg Z = 2, \quad \deg X = n + 1, \quad \deg Y = n, \quad \deg(\text{eqn}) = 2n + 2.$$

A combined matrix which reduces to each of these in the m odd and m even cases will have in each entry the degrees

$$\begin{pmatrix} n+1 & m+1 & n & m \\ 2n-m+1 & n+1 & 2n-m & n \\ n+2 & m+2 & n+1 & m+1 \\ 2n-m+2 & n+2 & 2n-m+1 & n+1 \end{pmatrix}$$

We know that $g'(Z) = Z^{n-m+1} + \dots$, and so $\deg g'(Z) = 2n - 2m + 2$. Hence, from the deformed equation:

$$X^2 + Y^2 Z - g'(Z)P''(Z)^2 - \frac{g'(Z)Q''(Z)^2}{Z},$$

we see that

$$\deg P''(Z) = m, \quad \deg Q''(Z) = m + 1.$$

We record for reference the degree of each element:

$$\begin{array}{c} \text{eqn} \\ 2n+2 \end{array} \left| \begin{array}{c} X \\ n+1 \end{array} \right| \left| \begin{array}{c} Y \\ n \end{array} \right| \left| \begin{array}{c} Z \\ 2 \end{array} \right| \left| \begin{array}{c} g' \\ 2n-2m+2 \end{array} \right| \left| \begin{array}{c} h' \\ 2n-2m \end{array} \right| \left| \begin{array}{c} h'' \\ 2m \end{array} \right| \left| \begin{array}{c} \delta' \\ n-m+1 \end{array} \right| \left| \begin{array}{c} \delta'' \\ m+1 \end{array} \right| \left| \begin{array}{c} P'' \\ m \end{array} \right| \left| \begin{array}{c} S'' \\ m-1 \end{array} \right| \left| \begin{array}{c} Q'' \\ m+1 \end{array} \right|$$

The fully deformed equation for the D_{n+2} singularity is given by

$$\text{eqn} : X^2 + Y^2Z - h + 2\delta Y = 0,$$

where

$$\begin{aligned} Q &= SZ + \delta, \\ h &= (g - \delta^2)/Z = P^2 + (Q^2 - \delta^2)/Z = P^2 + ZS^2 + 2\delta S \\ &= Zh'h'' + h'\delta''^2 + h''\delta'^2, \\ h'' &= P''^2 + ZS''^2 + 2\delta''S'', \\ Q'' &= S''Z + \delta'', \\ \delta &= \delta'\delta''. \end{aligned}$$

The degrees of the various elements that can go in the matrix are

$$\begin{array}{c} \text{eqn} \\ 2n+2 \end{array} \left| \begin{array}{c} X \\ n+1 \end{array} \right| \left| \begin{array}{c} Y \\ n \end{array} \right| \left| \begin{array}{c} Z \\ 2 \end{array} \right| \left| \begin{array}{c} h' \\ 2n-2m \end{array} \right| \left| \begin{array}{c} \delta' \\ n-m+1 \end{array} \right| \left| \begin{array}{c} P'' \\ m \end{array} \right| \left| \begin{array}{c} S'' \\ m-1 \end{array} \right| \left| \begin{array}{c} Q'' \\ m+1 \end{array} \right|$$

Note that when $t = 0$ we have $\delta' = 0$ and

$$\begin{array}{l} m \text{ even} \\ m \text{ odd} \end{array} \left| \begin{array}{l} h'(Z) = Z^{n-m}, \quad P''(Z) = i^m Z^{m/2}, \quad Q''(Z) = 0. \\ h'(Z) = Z^{n-m}, \quad P''(Z) = 0, \quad Q''(Z) = i^{m+1} Z^{(m+1)/2}. \end{array} \right.$$

Finding fully deformed matrix of relations

The fully deformed matrix of relations is found by requiring that (1) it reduce properly to the non-deformed case, (2) the entries respect quasi-homogeneity, (3) the relations in the ideal of 2×2 minors match those found in the non-deformed case.

The most general matrix with these degrees and which matches at $t = 0$ is thus

$$\left(\begin{array}{cccc} -iX + a_1\delta'P'' & i^{\pm(m+1)}Q'' & -Y + a_2\delta'S'' & i^{\pm m}P'' \\ i^{\pm(m+1)}h'Q'' + b_1\delta'Y & iX + b_2\delta'P'' & i^{\pm m}h'P'' & -Y + b_3\delta'S'' \\ -YZ + c_1\delta'Q'' + c_2\delta'S''Z & i^{\pm m}ZP'' & -iX + c_3\delta'P'' & i^{\pm(m+1)}Q'' \\ i^{\pm m}h'P''Z + d_1\delta'X + d_2\delta'^2P'' & -YZ + d_3\delta'Q'' + d_4\delta'S''Z & i^{\pm(m+1)}h'Q'' + d_5\delta'Y & iX + d_6\delta'P'' \end{array} \right).$$

To find the right coefficients, we will want the minors of this matrix to satisfy the same relations that the minors in the $t = 0$ case satisfy. For the top two rows and the bottom two rows, these are

1. $M_{12}^{14} = M_{12}^{23}$, or $[3] = [4]$. This requires

- $b_3 = a_2$

- $b_2 = -a_1$
 - $b_1 = 0$
2. $M_{12}^{13} = (-1)^m h' M_{12}^{24}$, or $[2] = (-1)^m h'[5]$.
 3. $M_{12}^{12} + Z M_{12}^{34} = \text{eqn}$, or $[1] + Z[6] = \text{eqn}$.
 4. $M_{34}^{14} = M_{34}^{23}$, or $[3b] = [4b]$.
 5. $M_{34}^{13} = (-1)^m h' M_{34}^{24}$, or $[2b] = (-1)^m h'[5b]$.
 6. $M_{34}^{12} + Z M_{34}^{34} = Z \text{eqn}$, or $[1b] + Z[6b] = Z \text{eqn}$.

Imposing the first condition $[3] = [4]$, the matrix becomes

$$\begin{pmatrix} -iX + a_1 \delta' P'' & i^{\pm(m+1)} Q'' & -Y + a_2 \delta' S'' & i^{\pm m} P'' \\ i^{\pm(m+1)} h' Q'' & iX - a_1 \delta' P'' & i^{\pm m} h' P'' & -Y + a_2 \delta' S'' \\ -YZ + c_1 \delta' Q'' + c_2 \delta' S'' Z & i^{\pm m} Z P'' & -iX + c_3 \delta' P'' & i^{\pm(m+1)} Q'' \\ i^{\pm m} h' P'' Z + d_1 \delta' X + d_2 \delta'^2 P'' & -YZ + d_3 \delta' Q'' + d_4 \delta' S'' Z & i^{\pm(m+1)} h' Q'' + d_5 \delta' Y & iX + d_6 \delta' P'' \end{pmatrix}.$$

The top two rows suggest that we can make the change of coordinates

$$X \mapsto \tilde{X} = X + ia_1 \delta' P'', \quad Y \mapsto \tilde{Y} = Y - a_2 \delta' S'',$$

which would additionally require

$$c_3 = a_1, \quad d_6 = -a_1, \quad c_2 = a_2, \quad \text{and} \quad d_4 = a_2.$$

Computing

$$\begin{aligned} \tilde{X}^2 &= X^2 + 2ia_1 \delta' X P'' - a_1^2 \delta'^2 P''^2 = X^2 + 2ia_1 \delta' (\tilde{X} - ia_1 \delta' P'') P'' - a_1^2 \delta'^2 P''^2 \\ &= X^2 + 2ia_1 \delta' \tilde{X} P'' + a_1^2 \delta'^2 P''^2, \\ \tilde{Y}^2 Z &= Y^2 Z - 2a_2 \delta' Y S'' Z + a_2^2 \delta'^2 S''^2 Z, \end{aligned}$$

and expanding the original deformed equation

$$\begin{aligned} 0 &= X^2 + Y^2 Z - h + 2\delta Y \\ &= X^2 + Y^2 Z - h'(Z P''^2 + Q''^2) - \delta'(2Y Z S'' - 2Y Q'') - \delta'^2(P''^2 - Z S''^2 + 2S'' Q'') \end{aligned}$$

we see that the coordinate change only makes sense if

$$a_1 = \pm i, \quad \text{and} \quad a_2 = 1.$$

In this case the equation becomes

$$0 = \tilde{X}^2 + \tilde{Y}^2 Z - h'(Z P''^2 + Q''^2) + 2\delta'(\tilde{Y} Q'' \pm \tilde{X} P'')$$

and the matrix is

$$\begin{pmatrix} -i\tilde{X} & i^{\pm(m+1)} Q'' & -\tilde{Y} & i^{\pm m} P'' \\ i^{\pm(m+1)} h' Q'' & i\tilde{X} & i^{\pm m} h' P'' & -\tilde{Y} \\ -\tilde{Y} Z + c_1 \delta' Q'' & i^{\pm m} Z P'' & -i\tilde{X} & i^{\pm(m+1)} Q'' \\ i^{\pm m} h' P'' Z + d_1 \delta' \tilde{X} & -\tilde{Y} Z + d_3 \delta' Q'' & i^{\pm(m+1)} h' Q'' + d_5 \delta' \tilde{Y} & i\tilde{X} \end{pmatrix}.$$

Requiring conditions 2 and 3 forces us to choose $a_1 = (-1)^{m+1}i$, giving change of coords

$$X \mapsto \tilde{X} = X + (-1)^m \delta' P'', \quad Y \mapsto \tilde{Y} = Y - \delta' S'',$$

and deformed equation

$$0 = \tilde{X}^2 + \tilde{Y}^2 Z - h'(ZP''^2 + Q''^2) + 2\delta'(\tilde{Y}Q'' + (-1)^{m+1}\tilde{X}P''). \quad (5.7)$$

Fixing the signs (and the remaining coefficients to zero) our matrix is now

$$\mathcal{R} = \begin{pmatrix} -i\tilde{X} & i^{-(m+1)}Q'' & -\tilde{Y} & i^m P'' \\ i^{(m+1)}h'Q'' & i\tilde{X} & i^{-m}h'P'' & -\tilde{Y} \\ -\tilde{Y}Z & i^m P''Z & -i\tilde{X} & i^{-(m+1)}Q'' \\ i^{-m}h'P''Z & -\tilde{Y}Z & i^{(m+1)}h'Q'' & i\tilde{X} \end{pmatrix}. \quad (5.8)$$

Note that the odd and even cases are now treated simultaneously. Together with the syzygies matrix \mathcal{S} , this yields a matrix factorization for the deformed equation (5.6).

5.3.5 The blowup

We can find additional relations among the Plücker coordinates by restricting attention to the 2×2 minors in the top two rows of the relations matrix \mathcal{R} (5.8). Together with the relation (4.1)

$$\gamma\varphi + \delta\alpha - \beta\varepsilon = 0,$$

this gives us the blowup.

The 2×2 minors for the top two rows of this matrix are

$$\begin{aligned} [1] &= \tilde{X}^2 - h'Q''^2, & [6] &= \tilde{Y}^2 - h'P''^2 \\ [2] &= i^{-m+3}h'(\tilde{X}P'' + i^{2m-2}\tilde{Y}Q''), & [5] &= i^{m+3}(\tilde{X}P'' + i^{2m-2}\tilde{Y}Q'') \\ [3] &= i\tilde{X}\tilde{Y} - i^{2m+1}h'P''Q'', & [4] &= i\tilde{X}\tilde{Y} - i^{2m+1}h'P''Q'' \end{aligned}$$

and they satisfy the relations

- [3] = [4]
- [2] = $(-1)^m h'[5]$, and
- [1] + $Z[6]$ + $2i^{(m-1)}\delta'[5]$ = [eqn].

The 2 by 2 minors for the bottom two rows are

$$\begin{aligned} [6b] &= \tilde{X}^2 - h'Q''^2, & [1b] &= Z^2(\tilde{Y}^2 - h'P''^2) \\ [2b] &= i^{m+3}h'Z(\tilde{Y}Q'' + i^{2m-2}\tilde{X}P''), & [5b] &= i^{-m+3}Z(\tilde{Y}Q'' + i^{2m-2}\tilde{X}P'') \\ [3b] &= -i\tilde{X}\tilde{Y} + i^{2m+1}h'P''Q'', & [4b] &= -i\tilde{X}\tilde{Y} + i^{2m+1}h'P''Q'' \end{aligned}$$

and they satisfy almost the same relations

- $[3b] = [4b]$
- $[2b] = (-1)^m h' [5b]$, and
- $[1b] + Z[6b] + 2i^{(m+1)} \delta' [5b] = Z[\text{eqn}]$.

This motivates us to restrict attention to the Plücker coordinates

$$\alpha = [4] = i\tilde{X}\tilde{Y} - i^{2m+1}h'P''Q'' \quad (5.9)$$

$$\varepsilon = [5] = i^{m+3}\tilde{X}P'' + i^{-m+1}\tilde{Y}Q'' \quad (5.10)$$

$$\varphi = [6] = \tilde{Y}^2 - h'P''^2 \quad (5.11)$$

and the equation for the Grassmanian $G(2, 4) \subseteq \mathbb{P}^5$ ($[1]\varphi - [2]\varepsilon + [3]\alpha = 0$) becomes

$$(-Z\varphi - 2i^{m-1}\delta'\varepsilon)\varphi + (-1)^{m+1}h'\varepsilon^2 + \alpha^2 = 0.$$

It will be useful to rewrite this as

$$\alpha^2 - \varphi^2 Z + (-1)^{m+1}h'\varepsilon^2 + 2i^{m+1}\delta'\varepsilon\varphi = 0. \quad (5.12)$$

Identification of residual singularities in the hyperplane

The interesting charts for the blowup are $\varphi = 1$ and $\varepsilon = 1$.

$\varphi = 1$ In this case we have $\alpha = \varphi\alpha_1$, $\varepsilon = \varphi\varepsilon_1$, and the equation (5.12) becomes

$$\alpha_1^2 - Z + (-1)^{m+1}h'\varepsilon_1^2 + 2i^{m+1}\delta'\varepsilon_1 = 0. \quad (5.13)$$

To see what kind of singularity this gives, define $G''(Z, i\alpha_1, t)$ as in (5.5) by

$$i\alpha_1 P''(Z, t) + Q''(Z, t) = (Z - \alpha_1^2)G''(Z, i\alpha_1, t) + f''_{m+1}(i\alpha_1, t),$$

and notice that from (5.13) we have

$$Z - \alpha_1^2 = (-1)^{m+1}h'\varepsilon_1^2 + 2i^{m+1}\delta'\varepsilon_1.$$

Moreover, notice from (5.11) (dropping tildes) that

$$i\alpha_1 P'' + Q'' = \frac{i\alpha P'' + \varphi Q''}{\varphi} = \frac{-XY P'' + i^{2m}h'P''^2 Q'' + Y^2 Q'' - h'P''^2 Q''}{Y^2 - h'P''^2}.$$

When m is even, this simplifies to

$$i\alpha_1 P'' + Q'' = i^{m+3}Y\varepsilon_1,$$

and we have

$$i^{m+3}Y\varepsilon_1 = \varepsilon_1((-1)^{m+1}h'\varepsilon_1 + 2i^{m+1}\delta')G'' + f''_{m+1}.$$

Defining

$$\hat{Y} = i^{m+3}Y - ((-1)^{m+1}h'\varepsilon_1 + 2i^{m+1}\delta')G'',$$

we find that our equation in this chart is

$$\widehat{Y} \varepsilon_1 = f''_{m+1}(i\alpha_1, t),$$

which is precisely the equation for a deformed A_m singularity!

$\varepsilon = 1$ In this case we have $\alpha = \varepsilon\alpha_2$, $\varphi = \varepsilon\varphi_2$, and the equation (5.12) becomes

$$\alpha_2^2 - \varphi_2^2 Z + (-1)^{m+1} h' + 2i^{m+1} \delta' \varphi_2 = 0,$$

which we immediately recognize as the preferred versal form for a deformed D_{n-m+1} singularity!

We thus see the expected residual singularities, since in the appropriate hyperplane the blowup breaks $D_{n+2} \longrightarrow D_{n-m+1} \oplus A_m$. Notice, however, that the total space for the resolved threefold

$$\alpha^2 - \varphi^2 Z + (-1)^{m+1} h' \varepsilon^2 + 2i^{m+1} \delta' \varepsilon \varphi = 0,$$

is smooth.




Figure 5.3: The blow up of the open length 2 node in the top Dynkin diagram provides a small resolution for the length 2 threefold singularity. When restricted to the hyperplane section, the D_8 surface singularity splits into two lower-order surface singularities: A_2 and D_5 . Nevertheless, the total space for the blow up (the threefold) is smooth.

Transition functions

The coordinates in our two charts are $(\alpha_1 = \alpha/\varphi, \varepsilon_1 = \varepsilon/\varphi, Z, t)$ for the $\varphi = 1$ chart, and $(\alpha_2 = \alpha/\varepsilon, \varphi_2 = \varphi/\varepsilon, Z, t)$ for the $\varepsilon = 1$ chart. The transition functions between them can be written

$$\varphi_2 = \varepsilon_1^{-1}, \tag{5.14}$$

$$\alpha_2 = \varepsilon_1^{-1} \alpha_1, \tag{5.15}$$

$$Z = (-1)^{m+1} \varepsilon_1^2 h'(Z, t) + \alpha_1^2 + 2i^{m+1} \delta' \varepsilon_1, \tag{5.16}$$

$$t = t. \tag{5.17}$$

5.4 Difficulties with the approach

Although we were able to successfully perform explicit small resolutions for deformed A_k and D_{k+2} equations, it is not at all clear how to use these blowups for finding the associated matrix model superpotentials. The results do not come to us packaged in such a way that comparison with Ferrari's framework can be made. Only in the A_1 case were we able to make the connection, and here the answer was already known from Ferrari's example 1.

The next section will develop an alternative "top-down" approach, beginning with the Intriligator–Wecht superpotentials, which we have conjectured provide the answer. This will prove to be a more fruitful strategy, although we will still need to use our small resolution techniques in order to show that we get the right singular space in the length 2 case. The algorithm we devise for performing the blow-downs cannot guarantee that we have all global holomorphic functions, and hence each case must be checked by blowing back up and recovering the original space.

Chapter 6

Small blow-downs from global holomorphic functions

6.1 “Top-down” approach

We have seen in Chapter 5 the difficulties that arise in the “bottom-up” approach. It is hard to perform small resolutions explicitly (thus far we have only done so in the A_{n-1} and D_{n+2} cases), but even this is not enough. Only in the A_1 case were we able to identify the superpotential from our small resolution, and in this case we already knew the answer!

While we know from the Katz–Morrison result [19] what the singular geometries should be, it is hard to fit their small resolutions into Ferrari’s framework, and hence we are unable to find a corresponding superpotential. On the other hand, if we could guess the form of the superpotentials, then showing which singular geometries they correspond to might not be so difficult. As in Ferrari’s original examples, we could start with the resolved geometry, find global holomorphic functions, and then use these to blow down the exceptional \mathbb{P}^1 ’s. We will call this the “top-down” approach:

- Guess W , then compute transition functions for resolved geometry $\widehat{\mathcal{M}}$.
- Find global holomorphic functions (hard, need algorithm!).
- Use these to construct the blow-down; relations among the global holomorphic functions describe the singular space \mathcal{M}_0 .
- Blow back up to check that we recover the original resolved geometry $\widehat{\mathcal{M}}$.

The last step is necessary because we do not yet know a way of determining whether or not we have found all (independent) global holomorphic functions. This is because the algorithm searches for global holomorphic functions which can be constructed from a given list of monomials. While we can be certain we haven’t “missed” any functions which can be made out of monomials in the list, it is, of course, impossible to check all possible

monomials. It is thus only in blowing back up that can we confirm that the singular space is correct. This will require using the small resolution methods developed in Chapter 5.

Our guesses for $W(x, y)$ come from the Intriligator–Wecht classification of RG fixed points of $\mathcal{N} = 1$ SQCD with adjoints. We make the most naive conjecture possible: that the ADE superpotentials in the classification correspond (under Ferrari’s construction) to geometries with ADE singularities. Moreover, in order to perform the blow-downs we devise an algorithm which searches for global holomorphic functions, thereby proving Proposition 2:

Proposition 2. *There is an algorithm for blowing down the exceptional \mathbb{P}^1 ’s when the resolved geometry is given by simple transition functions (1.1) as in Ferrari’s construction.*

6.2 The Intriligator–Wecht classification

Here we recall (see Introduction) the Intriligator–Wecht classification of RG fixed points of $\mathcal{N} = 1$ SQCD with adjoints [17]. Using “a-maximization” and doing a purely field theoretic analysis, they classify all relevant adjoint superpotential deformations for 4d $\mathcal{N} = 1$ SQCD with \mathcal{N}_f fundamentals and $\mathcal{N}_a = 2$ adjoint matter chiral superfields, X and Y . The possible RG fixed points, together with the map of possible flows between fixed points, are summarized below:




Figure 6.1: Intriligator–Wecht Classification of RG Fixed Points. The diagram on the right shows the map of possible flows between fixed points. Dotted lines indicate flow to a particular value of k . Note that $k' < k$. [17, pages 3-4]

Using Table 3.2 from Chapter 3 we identify the corresponding resolved geometries, and these are shown in Table 6.1. There are two possible perturbation terms $\partial_{w_1} E(\gamma, w_1)$ for

type	$W(x, y)$	$\partial_{w_1} E(\gamma, w_1)$	$\mathbb{P}^1 : (w_1, w_2), (v_1, v_2)$
\widehat{O}	0	0	$(x + \gamma y, 0), (\beta x + y, 0)$
\widehat{A}	$\frac{1}{2}y^2$	$w_1 \leftrightarrow \gamma^2 w_1$	$(x, 0), (\beta x, x)$
\widehat{D}	xy^2	$w_1^2 \leftrightarrow \gamma w_1^2$	$(x, 0), (\beta x, x^2)$
\widehat{E}	$\frac{1}{3}y^3$	$\gamma^{-1} w_1^2 \leftrightarrow \gamma^2 w_1^2$	$(x, 0), (\beta x, \beta x^2)$
A_k	$\frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2$	$\gamma^2 w_1^k + w_1 \leftrightarrow \gamma^{1-k} w_1^k + \gamma^2 w_1$	$(0, 0), (0, 0)$
D_{k+2}	$\frac{1}{k+1}x^{k+1} + xy^2$	$\gamma^2 w_1^k + w_1^2 \leftrightarrow \gamma^{1-k} w_1^k + \gamma w_1^2$	$(0, 0), (0, 0)$
E_6	$\frac{1}{3}y^3 + \frac{1}{4}x^4$	$\gamma^{-1} w_1^2 + \gamma^2 w_1^3 \leftrightarrow \gamma^2 w_1^2 + \gamma^{-2} w_1^3$	$(0, 0), (0, 0)$
E_7	$\frac{1}{3}y^3 + yx^3$	$\gamma^{-1} w_1^2 + \gamma w_1^3 \leftrightarrow \gamma^2 w_1^2 + \gamma^{-1} w_1^3$	$(0, 0), (0, 0)$
E_8	$\frac{1}{3}y^3 + \frac{1}{5}x^5$	$\gamma^{-1} w_1^2 + \gamma^2 w_1^4 \leftrightarrow \gamma^2 w_1^2 + \gamma^{-3} w_1^4$	$(0, 0), (0, 0)$

Table 6.1: Identification of corresponding resolved geometries

each case due to the $x \leftrightarrow y$ symmetry, which exchanges

$$\gamma^n w_1^m \leftrightarrow \gamma^{3-n-m} w_1^m.$$

All of these examples are in the case $M = 2$, with transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \partial_{w_1} E(\gamma, w_1),$$

which gives a deformed $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ bundle. In the \widehat{A} and A_k cases, the linear perturbation term w_1 indicates that the normal bundle to the \mathbb{P}^1 is not really $\mathcal{O}(1) \oplus \mathcal{O}(-3)$, but rather a funny description for $\mathcal{O} \oplus \mathcal{O}(-2)$, or $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ in the A_1 case.

6.3 The problem

We have seen from Ferrari's examples in Chapter 2 that we can blow down exceptional \mathbb{P}^1 's by finding global holomorphic functions (ghf's) on the resolved space $\widehat{\mathcal{M}}$. Global

holomorphic functions must necessarily be constant on the \mathbb{P}^1 , so they provide natural coordinates for the blow-down. Beginning with transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-n}w_1, \quad v_2 = \gamma^{-m}w_2 + \partial_{w_1}E(\gamma, w_1),$$

the idea is to find polynomials in $\mathbb{C}[\beta, v_1, v_2]$ which are also polynomials in $\mathbb{C}[\gamma, w_1, w_2]$.

For example, in the case of E_6 :

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^3 + \gamma^{-1}w_1^2.$$

we immediately see from the transition functions that neither β, v_1 or v_2 are ghf's. However, if we consider βv_2 we have

$$\beta v_2 = \gamma^2w_2 + \gamma w_1^3 + \gamma^{-2}w_1^2.$$

The non-holomorphic term on the right can be moved to the left, using $v_1 = \gamma^{-1}w_1$, and so we construct the ghf

$$\beta v_2 - v_1^2 = \gamma^2w_2 + \gamma w_1^3.$$

Multiplying by β, v_1 or v_2 yields new ghf's

$$\begin{aligned} \beta(\beta v_2 - v_1^2) &= \gamma w_2 + w_1^3, \\ v_1(\beta v_2 - v_1^2) &= \gamma w_1 w_2 + w_1^4, \\ v_2(\beta v_2 - v_1^2) &= (\gamma^4 w_2 + \gamma^3 w_1^3 + w_1^2)(\gamma w_2 + w_1^3). \end{aligned}$$

Unfortunately, these functions all vanish on $\beta v_2 = v_1^2$, which is more than the \mathbb{P}^1 we are trying to blow down (the defining equations for the \mathbb{P}^1 are $v_1 = v_2 = 0$). We must find a collection of ghf's which collapse the exceptional \mathbb{P}^1 to a point, but provide a birational isomorphism everywhere else.

This rather ad-hoc method for finding global holomorphic functions turns out to be quite limited, as cleverness quickly runs dry with polynomials of higher degree. We would like to make the search for ghf's more systematic.

6.4 Reformulation of the problem

We reformulate our problem of finding global holomorphic functions as an ideal membership problem. We begin by illustrating the reformulation in the case of E_6 :

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^3 + \gamma^{-1}w_1^2.$$

We can think of this as describing a variety in \mathbb{C}^6 , defined by the following ideal $I \subset \mathbb{C}[\gamma, w_1, w_2, \beta, v_1, v_2]$:

$$I = \langle \beta\gamma - 1, v_1 - \beta w_1, v_2 - \gamma^3 w_2 - \gamma^2 w_1^3 - \beta w_1^2 \rangle.$$

In order to blow down the exceptional \mathbb{P}^1 , we must find functions which are holomorphic in each coordinate chart, and will therefore be constant on the \mathbb{P}^1 . Such global holomorphic functions correspond to elements of the ideal I that can be written in the form¹

$$f - g \in I, \quad \text{where } f \in \mathbb{C}[\beta, v_1, v_2], \quad g \in \mathbb{C}[\gamma, w_1, w_2].$$

For each such element, the global holomorphic function is $f = g$.

In general, we begin with transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-n}w_1, \quad v_2 = \gamma^{-m}w_2 + \partial_{w_1}E(\gamma, w_1),$$

and form the ideal

$$I = \langle \beta\gamma - 1, v_1 - \beta^n w_1, v_2 - \beta^m w_2 - \partial_{w_1} \tilde{E}(\gamma, w_1) \rangle,$$

where $\tilde{E}(\gamma, w_1)$ is obtained from $E(\gamma, w_1)$ by replacing all instances of γ^{-1} with β .

6.5 The algorithm

Consider a monomial $\beta^i v_1^j v_2^k \in \mathbb{C}[\beta, v_1, v_2]$. Using Groebner basis techniques, we can easily reduce this modulo the ideal

$$I = \langle \beta\gamma - 1, v_1 - \beta^n w_1, v_2 - \beta^m w_2 - \partial_{w_1} \tilde{E}(\gamma, w_1) \rangle,$$

which is determined by our particular geometry. In general, we will find

$$\beta^i v_1^j v_2^k \stackrel{\text{mod } I}{\equiv} \text{pure}(\beta, v_1, v_2) + \text{pure}(\gamma, w_1, w_2) + \text{mixed},$$

where “pure” and “mixed” stand for pure and mixed terms² in the appropriate variables. We can then bring the $\text{pure}(\beta, v_1, v_2)$ terms to the left hand side, “updating” our initial monomial to the polynomial

$$\beta^i v_1^j v_2^k - \text{pure}(\beta, v_1, v_2) \stackrel{\text{mod } I}{\equiv} \text{pure}(\gamma, w_1, w_2) + \text{mixed}.$$

Now the challenge is to find a linear combination f of such polynomials in $\mathbb{C}[\beta, v_1, v_2]$ such that the mixed terms cancel, and we are left with

$$f \stackrel{\text{mod } I}{\equiv} g, \quad \text{where } f \in \mathbb{C}[\beta, v_1, v_2], \quad g \in \mathbb{C}[\gamma, w_1, w_2].$$

The central idea (as in the Euclidean division algorithm) is to put a term order on the mixed terms we are trying to cancel. In this way, we can make sure we are cancelling mixed terms in an efficient manner, and the cancellation procedure terminates. Because mixed terms (such as βw_1) correspond to “poles” in the γ coordinate chart (such as $\gamma^{-1}w_1$), we use the weighted degree term order

$$> \text{TP} := \text{wdeg}([1, 1, 1, -1, 0, 0], [\text{b}, \text{v}[1], \text{v}[2], \text{g}, \text{w}[1], \text{w}[2]]):$$

which keeps track of the degree of the poles in γ .

Beginning with the superpotential, our algorithm thus consists of the following steps:

¹Note that in our E_6 example none of the defining generators of I are of this form!

²We will refer to any monomial in $\mathbb{C}[\gamma, w_1, w_2, \beta, v_1, v_2]$ which does not belong to either $\mathbb{C}[\beta, v_1, v_2]$ or $\mathbb{C}[\gamma, w_1, w_2]$ as a mixed term.

1. Compute transition functions following Ferrari's framework. This gives an ideal

$$I = \langle \beta\gamma - 1, v_1 - \beta^n w_1, v_2 - \beta^m w_2 - \partial_{w_1} \tilde{E}(\gamma, w_1) \rangle \subset \mathbb{C}[\gamma, w_1, w_2, \beta, v_1, v_2].$$

2. Find a Groebner basis G for the ideal I , with respect to a term order T .
3. Generate a list L of monomials in β, v_1 , and v_2 (up to some degree).
4. Reduce monomial $L[j]$ mod I , using G . What you have is

$$L[j] = \beta^i v_1^j v_2^k \stackrel{\text{mod } I}{\equiv} \text{pure}(\beta, v_1, v_2) + \text{pure}(\gamma, w_1, w_2) + \text{mixed}$$

where “pure” and “mixed” stand for pure and mixed terms in the appropriate variables. Bring the $\text{pure}(\beta, v_1, v_2)$ terms over to the LHS to make a polynomial

$$\beta^i v_1^j v_2^k - \text{pure}(\beta, v_1, v_2) \in \mathbb{C}[\beta, v_1, v_2].$$

- Record this polynomial in the array F as $F[j, 1]$.
- Record the leading term (with respect to the term order TP) of the “mixed” part as $F[j, 2]$, and store the leading coefficient as $F[j, 3]$.

5. Reduction routine

- Cycle through the list of previous polynomials $F[1..j - 1, *]$ and cancel leading mixed terms as much as possible.
 - The result is a new “updated” polynomial $F[j, 1]$ which is *reduced* in the sense that its leading mixed term is as low as possible (with respect to the term order TP) due to cancellation with leading mixed terms from previous polynomials.
 - Reduce the “updated” $F[j, 1]$ modulo the ideal I to update $F[j, 2]$ and $F[j, 3]$.
 - If the new leading mixed term $F[j, 2]$ is 0, we have a **global holomorphic function!**
6. Determine which global holomorphic functions are “new,” so that the final list isn't redundant.
 - Check that the new global holomorphic function X_l is not in the ring $\mathbb{C}[X_1, \dots, X_{l-1}]$ generated by the previous functions!
 - To do this we find a Groebner basis for the ideal $\langle X_1, \dots, X_l \rangle$ and compute partials to make sure we can't solve for the new function in terms of the previous ones.
 7. Find relations among the global holomorphic functions. These will determine the (singular) geometry of the blow down.

This completes the proof of Proposition 2.

6.5.1 A shortcut

In all of the Intriligator–Wecht cases, we can find weights for the variables β, v_1, v_2 and γ, w_1, w_2 such that the transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \partial_{w_1}E(\gamma, w_1),$$

are quasi-homogeneous. In the E_6 case, for example, the transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^3 + \gamma^{-1}w_1^2,$$

are quasi-homogeneous if we assign the weights

$$\beta \left| \begin{array}{c} v_1 \\ 1 \end{array} \right| v_2 \left| \begin{array}{c} v_2 \\ 7 \end{array} \right| \gamma \left| \begin{array}{c} w_1 \\ -1 \end{array} \right| w_2 \left| \begin{array}{c} w_2 \\ 3 \\ 10 \end{array} \right|.$$

In particular, this means all elements of the ideal

$$I = \langle \beta\gamma - 1, v_1 - \beta w_1, v_2 - \gamma^3w_2 - \gamma^2w_1^3 - \beta w_1^2 \rangle$$

are quasi-homogeneous in these weights, and all terms in the expression

$$\beta^i v_1^j v_2^k \stackrel{\text{mod } I}{\equiv} \text{pure}(\beta, v_1, v_2) + \text{pure}(\gamma, w_1, w_2) + \text{mixed},$$

will have the same weight.

This immediately tells us that only combinations of monomials *of the same weight* can be used to cancel mixed terms – i.e. the global holomorphic functions we build will themselves be quasi-homogeneous. This observation cuts computational time immensely, since it means that in the **reduction routine** we need only cycle through lists of polynomials of the same weight in order to reduce the order of the mixed terms. In particular, we can run the algorithm in parallel for different weights, restricting ourselves to lists of monomials in $\mathbb{C}[\beta, v_1, v_2]$ which are all in the same weighted degree.

6.5.2 Implementation


At this point, we have only used the blow-down algorithm in order to find global holomorphic functions in the Intriligator–Wecht cases. Our Maple implementation, therefore, takes full advantage of the “shortcut” 6.5.1, and has been coded specifically for the 2-matrix model case – with deformed $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ bundle. The complete Maple code, along with a brief description of all functions and procedures, can be found in the appendix.

All of the global holomorphic functions listed in all of the cases we analyze (Chapters 7, 8, 9, 10, and 11) have been found using this implementation of the algorithm.

It is important to remember that the algorithm does not tell us when we are done looking for global holomorphic functions. The only way to check that we have the right blow-down \mathcal{M}_0 is by inverting the procedure (blowing back up) and verifying that we recover the original transition functions for the resolved geometry $\widehat{\mathcal{M}}$. It is because of this that we must still rely on the small resolution techniques developed in Chapter 5.

Chapter 7

Length 1



In this chapter we prove the first part of **Theorem 2**:

The resolved geometry corresponding to the Intriligator–Wecht superpotential

$$W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2,$$

corresponds to the singular geometry

$$XY - T(Z^k - T) = 0.$$

We will also discover that this potential is geometrically equivalent to

$$W(x) = \frac{1}{k+1}x^{k+1}.$$

7.1 The Case A_k

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator–Wecht superpotential

$$W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^k + w_1.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}w_2 + \beta^{-2}(x + \beta^{-1}y)^k + x + \beta^{-1}y.$$

If we choose

$$w_2(\gamma) = \frac{x^k - (x + \gamma y)^k}{\gamma},$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}x^k + \beta^{-1}y + x.$$

This is only holomorphic if

$$x^k = y = 0,$$

and so we have a single \mathbb{P}^1 located at

$$w_1(\gamma) = w_2(\gamma) = 0, \quad v_1(\beta) = v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = x^k dx + y dy = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^k + w_1,$$

are quasi-homogeneous if we assign the weights

$$\begin{array}{c|c|c|c|c|c} \beta & v_1 & v_2 & \gamma & w_1 & w_2 \\ k-1 & k+1 & 2 & 1-k & 2 & 3k-1 \end{array}.$$

The global holomorphic functions will thus necessarily be quasi-homogeneous in these weights. We find the following global holomorphic functions:

$$\begin{array}{c|c|c|c|c|c} 2 & y_1 & = & v_2 & = & \gamma^3w_2 + \gamma^2w_1^k + w_1 \\ k+1 & y_2 & = & \beta v_2 - v_1 & = & \gamma^2w_2 + \gamma w_1^k \\ 2k & y_3 & = & \beta^2v_2 - \beta v_1 & = & \gamma w_2 + w_1^k \\ 3k-1 & y_4 & = & v_2^{k-1}v_1 - \beta^3v_2 + \beta^2v_1 & = & \end{array}$$

These are the first 4 “distinct” functions produced by our algorithm, in the sense that none is contained in the ring generated by the other 3.

The singular geometry \mathcal{M}_0

We conjecture that the ring of global holomorphic functions is generated by y_1, y_2, y_3 and y_4 , subject to the degree $4k$ relation

$$\mathcal{M}_0 : y_2 y_4 + y_3^2 + y_2^2 y_1^{k-1} - y_3 y_1^k = 0.$$

The functions y_i give us a blow-down map whose image \mathcal{M}_0 has an isolated A_k singularity. To see this, consider the change of variables

$$\tilde{y}_4 = y_4 + y_2 y_1^{k-1} = \beta v_2^k - \beta^3 v_2 + \beta^2 v_1.$$

Note that like y_4 , \tilde{y}_4 is also quasi-homogeneous of degree $3k - 1$. The functions y_1, y_2, y_3 and \tilde{y}_4 now satisfy the simpler relation

$$\mathcal{M}_0 : y_2 \tilde{y}_4 + y_3(y_3 - y_1^k) = 0.$$

The blowup

We now verify that we have identified the right singular space \mathcal{M}_0 by inverting the blow-down. In the β and γ charts we find

$$\begin{array}{l} \beta = \\ v_1 = \\ v_2 = \end{array} \begin{array}{l} y_3/y_2 \\ \beta y_1 - y_2 = (y_3 y_1 - y_2^2)/y_2 \\ y_1 \end{array} \left| \begin{array}{l} \gamma = \\ w_1 = \\ w_2 = \\ = \end{array} \begin{array}{l} y_2/y_3 \\ y_1 - \gamma y_2 = (y_1 y_3 - y_2^2)/y_3 \\ \beta(y_3 - w_1^k) \\ -\tilde{y}_4 + \frac{y_1^k - (y_1 - \gamma y_2)^k}{\gamma} \end{array} \right.$$

This suggests that we should blow up

$$y_2 = y_3 = 0,$$

for the small resolution of \mathcal{M}_0 . We introduce \mathbb{P}^1 coordinates $[\beta : \gamma]$ such that $\beta y_2 = \gamma y_3$. The blow up in each chart is then

$$\begin{array}{l} (\gamma = 1) \\ \\ \tilde{y}_4 = \beta(y_1^k - \beta y_2) \\ \text{coords : } (y_1, y_2, \beta) \end{array} \left| \begin{array}{l} (\beta = 1) \\ \\ y_2 = \gamma y_3 \\ y_3 = y_1^k - \gamma \tilde{y}_4 \\ \text{coords : } (y_1, \tilde{y}_4, \gamma) \end{array} \right.$$

Transition functions

The transition functions between the β and γ charts are

$$\beta = \gamma^{-1}, \quad y_1 = y_1, \quad y_2 = \gamma(y_1^k - \gamma\tilde{y}_4) = -\gamma^2\tilde{y}_4 + \gamma y_1^k.$$

Note that for $k > 1$, this is an $\mathcal{O} \oplus \mathcal{O}(-2)$ bundle over the exceptional \mathbb{P}^1 , and corresponds to a superpotential with a single field ($M = 1$):

$$W(x) = \frac{1}{k+1}x^{k+1}.$$

(For $k = 1$ the bundle is actually $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ and $W = 0$.)

In terms of the original coordinates, the transition functions become

$$\beta = \gamma^{-1}, \quad v_2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1, \quad \beta v_2 - v_1 = \gamma^2 w_2 + \gamma w_1^k.$$

Substituting the second transition function into the third reveals $v_1 = \gamma^{-1} w_1$, and so we recover our original transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1,$$

which define an $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ bundle deformed by the two field ($M = 2$) superpotential

$$W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2.$$

7.2 A puzzle

The problem

We saw the A_k case in Section 2.3, with geometry $\widehat{\mathcal{M}}$ given by the superpotential

$$W(x) = \frac{1}{k+1}x^{k+1},$$

and hence corresponding to a deformed $\mathcal{O} \oplus \mathcal{O}(-2)$ bundle over the \mathbb{P}^1 , with one field. However, Intriligator and Wecht identify

$$W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2,$$

as corresponding to an A_k -type singularity, with an extra field y which requires that the transition functions look like

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1.$$

In particular, the geometry $\widehat{\mathcal{M}}$ looks like that of an $\mathcal{O}(1) \oplus \mathcal{O}(-3)$ bundle over the exceptional \mathbb{P}^1 ! What's going on here?

Resolution of the problem

For $n = 1$ and $k = 0$, Proposition 1 tells us that the superpotential

$$W(x, y) = \frac{1}{2}y^2$$

changes the bundle

$$\mathcal{O}(1) \oplus \mathcal{O}(-3) \longrightarrow \mathcal{O} \oplus \mathcal{O}(-2).$$

Hence the extra field y from the Intriligator–Wecht potential (with purely quadratic contribution to the superpotential) can be “integrated out.” Its effect is to change the bundle for A_k from $\mathcal{O}(1) \oplus \mathcal{O}(-3)$, which is necessary for a two-field description, to reveal the true underlying $\mathcal{O} \oplus \mathcal{O}(-2)$ structure. In other words,

$$W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2 \quad \text{and} \quad W(x) = \frac{1}{k+1}x^{k+1}$$

are geometrically equivalent.

Beyond Proposition 1

Note that this is not just a straightforward application of Proposition 1, which implies that $W(x, y) = \frac{1}{2}y^2$ and $W(x) = 0$ are geometrically equivalent. Beginning with transition

functions for the Intriligator–Wecht A_k superpotential $W(x, y) = \frac{1}{k+1}x^{k+1} + \frac{1}{2}y^2$,

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^k + w_1,$$

the change of coordinates suggested in the proof of Proposition 1

$$\begin{aligned} \widetilde{v}_1 &= v_2, & \widetilde{w}_1 &= w_1 + \gamma^3w_2, \\ \widetilde{v}_2 &= -v_1 + \beta v_2, & \widetilde{w}_2 &= w_2, \end{aligned}$$

does not yield the appropriate new transition functions. Instead, the more complicated change of coordinates

$$\begin{aligned} \widetilde{v}_1 &= v_2, & \widetilde{w}_1 &= w_1 + \gamma^3w_2 + \gamma^2w_1^k, \\ \widetilde{v}_2 &= -v_1 + \beta v_2, & \widetilde{w}_2 &= w_2 - \gamma^{-1}[(\gamma^3w_2 + \gamma^2w_1^k + w_1)^k - w_1^k], \end{aligned}$$

is needed to give new transition functions


$$\beta = \gamma^{-1}, \quad \widetilde{v}_1 = \widetilde{w}_1, \quad \widetilde{v}_2 = \gamma^2\widetilde{w}_2 + \gamma\widetilde{w}_1^k,$$

corresponding to the superpotential $W(x) = \frac{1}{k+1}x^{k+1}$.

It would be interesting to try to generalize Proposition 1 to include examples such as this, where there are additional terms in the superpotential besides the quadratic pieces which suggest a change in bundle structure. Trying to understand what makes the change of coordinates possible in this case may give hints as to how the geometric picture for RG flow might be extended. The ultimate goal would be to understand how “integrating out” the y coordinate in a potential of the form $W(x, y) = f(x, y) + y^2$ affects other terms involving y .

Chapter 8

Length 2



In this chapter we prove the second part of **Theorem 2**:

The resolved geometry corresponding to the Intriligator–Wecht superpotential

$$W(x, y) = \frac{1}{k+1}x^{k+1} + xy^2$$

corresponds to the singular geometry

$$X^2 - Y^2Z + T(Z^{k/2} - T)^2 = 0.$$

8.1 The Case D_{k+2}

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator-Wecht superpotential

$$W(x, y) = \frac{1}{k+1}x^{k+1} + xy^2,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^k + w_1^2.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}w_2 + \beta^{-2}(x + \beta^{-1}y)^k + (x + \beta^{-1}y)^2.$$

If we choose

$$w_2(\gamma) = \frac{x^k - (x + \gamma y)^k}{\gamma},$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}(x^k + y^2) + \beta^{-1}(2xy) + x^2.$$

This is only holomorphic if

$$x^k + y^2 = 2xy = 0,$$

and so we have a single \mathbb{P}^1 located at

$$w_1(\gamma) = w_2(\gamma) = 0, \quad v_1(\beta) = v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = (x^k + y^2)dx + 2xy dy = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^k + w_1^2,$$

are quasi-homogeneous if we assign the weights

$$\begin{array}{c} \beta \\ k-2 \end{array} \left| \begin{array}{c} v_1 \\ k \end{array} \right| \begin{array}{c} v_2 \\ 4 \end{array} \left| \begin{array}{c} \gamma \\ 2-k \end{array} \right| \begin{array}{c} w_1 \\ 2 \end{array} \left| \begin{array}{c} w_2 \\ 3k-2 \end{array} \right|.$$

The global holomorphic functions will thus necessarily be quasi-homogeneous in these weights. We find the following global holomorphic functions for k even:

$$\begin{array}{c} 4 \\ 2k \\ 3k-2 \\ 3k \end{array} \left| \begin{array}{c} X \\ Y \\ Z \\ U \end{array} \right| = \begin{array}{c} v_2 \\ \beta^2v_2 - v_1^2 \\ \beta(X^{k/2} - Y) \\ v_1(X^{k/2} - Y) \end{array} = \begin{array}{c} \gamma^3w_2 + \gamma^2w_1^k + w_1^2 \\ \gamma w_2 + w_1^k \\ \gamma^{-1}(X^{k/2} - Y) \\ \gamma^{-1}w_1(X^{k/2} - Y), \end{array}$$

and a similar set of global holomorphic functions for k odd:

$$\begin{array}{l|l} 4 & X = v_2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2 \\ 2k & Y = \beta^2 v_2 - v_1^2 = \gamma w_2 + w_1^k \\ 3k-2 & Z = v_1 X^{\lfloor k/2 \rfloor} - \beta Y = \gamma^{-1}(w_1 X^{\lfloor k/2 \rfloor} - Y) \\ 3k & U = \beta X^{\lceil k/2 \rceil} - v_1 Y = \gamma^{-1}(X^{\lceil k/2 \rceil} - w_1 Y). \end{array}$$

The singular geometry \mathcal{M}_0

We conjecture that the ring of global holomorphic functions is generated by X, Y, Z and U , subject to the degree $6k$ relation

$$\begin{array}{l} \mathcal{M}_0 : \quad U^2 - XZ^2 + Y(X^{k/2} - Y)^2 = 0, \quad k \text{ even,} \\ \mathcal{M}_0 : \quad U^2 - XZ^2 - Y(X^k - Y^2) = 0, \quad k \text{ odd.} \end{array}$$

The functions X, Y, Z and U give us a blow-down map whose image \mathcal{M}_0 has an isolated D_{k+2} singularity.

Review of length 2 blowup

Before doing the blowup to see that we have the right blow down, we review some results from Chapter 5. There we found small resolutions of length 2 singularities by using deformations of matrix factorizations for D_{n+2} surface singularities.

Recall (dropping tildes!) the deformed D_{n+2} equation (5.7)

$$0 = X^2 + Y^2 Z - h'(ZP''^2 + Q''^2) + 2\delta'(YQ'' + (-1)^{m+1}XP''),$$

where for $t = 0$ we have $\delta' = 0$ and

$$\begin{array}{l} m \text{ even} \left| \begin{array}{l} h'(Z) = Z^{n-m}, \quad P''(Z) = i^m Z^{m/2}, \quad Q''(Z) = 0. \\ h'(Z) = Z^{n-m}, \quad P''(Z) = 0, \quad Q''(Z) = i^{m+1} Z^{(m+1)/2}. \end{array} \right. \\ m \text{ odd} \end{array}$$

The blowup was given by the equation for the Grassmanian $G(2, 4) \subseteq \mathbb{P}^5$

$$\alpha^2 - \varphi^2 Z + (-1)^{m+1} h' \varepsilon^2 + 2i^{m+1} \delta' \varepsilon \varphi = 0,$$

in terms of Plücker coordinates

$$\begin{aligned} \alpha &= iXY - i^{2m+1} h' P'' Q'' \\ \varepsilon &= i^{m+3} X P'' + i^{-m+1} Y Q'' \\ \varphi &= Y^2 - h' P''^2. \end{aligned}$$

The interesting charts were $\varphi = 1$ and $\varepsilon = 1$, with transition functions

$$\varphi_2 = \varepsilon_1^{-1}, \tag{8.1}$$

$$\alpha_2 = \varepsilon_1^{-1} \alpha_1, \tag{8.2}$$

$$Z = (-1)^{m+1} \varepsilon_1^2 h'(Z, t) + \alpha_1^2 + 2i^{m+1} \delta' \varepsilon_1, \tag{8.3}$$

$$t = t. \tag{8.4}$$

The blowup

As usual, we verify that we have the right singular space by inverting the blow down. A simple change of coordinates shows our \mathcal{M}_0 equations to be

$$\begin{aligned} X^2 + Y^2 Z - T(Z^{k/2} - T)^2 &= 0 & (k \text{ even}) \\ X^2 + Y^2 Z + T(Z^k - T^2) &= 0 & (k \text{ odd}). \end{aligned}$$

k even

For k even, the equation for \mathcal{M}_0 corresponds to

$$\delta' = 0, \quad h' = T, \quad P'' = 0, \quad Q'' = i^k(Z^{k/2} - T), \quad n = m = k - 1,$$

and Plücker coordinates

$$\alpha = iXY, \quad \varepsilon = -Y(Z^{k/2} - T), \quad \varphi = Y^2.$$

The connection with the original transition function coordinates is

$$\begin{array}{l|l} 4 & Z = v_2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2 \\ 2k & T = \beta^2 v_2 - v_1^2 = \gamma w_2 + w_1^k \\ 3k - 2 & Y = \beta(Z^{k/2} - T) = \gamma^{-1}(Z^{k/2} - T) \\ 3k & iX = v_1(Z^{k/2} - T) = \gamma^{-1} w_1(Z^{k/2} - T). \end{array}$$

Note that

$$\begin{aligned} \beta &= \frac{Y}{Z^{k/2} - T} = -\frac{\varphi}{\varepsilon} = -\varphi_2 \\ v_1 &= \frac{iX}{Z^{k/2} - T} = -\frac{\alpha}{\varepsilon} = -\alpha_2 \\ v_2 &= \frac{Z}{Z^{k/2} - T} = -\frac{\varepsilon}{\varphi} = -\varepsilon_1 \\ \gamma &= \frac{Z^{k/2} - T}{Y} = -\frac{\varepsilon}{\varphi} = -\varepsilon_1 \\ w_1 &= \frac{iX}{Y} = \frac{\alpha}{\varphi} = \alpha_1 \\ w_2 &= \beta(T - w_1^k) = -\varepsilon_2^{-1}(T - \alpha_2^k) \end{aligned}$$

In particular, the transition functions (8.1) become

$$\begin{aligned} \beta &= \gamma^{-1}, \\ v_1 &= \gamma^{-1} w_1, \\ v_2 &= \gamma^2 T + w_1^2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2. \end{aligned}$$

k odd

Here the equation for \mathcal{M}_0 corresponds to

$$\delta' = 0, \quad h' = -T, \quad P'' = i^{k-1} Z^{(k-1)/2}, \quad Q'' = iT, \quad n = m = k - 1,$$

and Plücker coordinates

$$\alpha = iXY + i^{3k+3}T^2Z^{(k-1)/2}, \quad \varepsilon = i^{2k+1}XZ^{(k-1)/2} + i^{3k+3}TY, \quad \varphi = Y^2 + i^{2k+2}TZ^{k-1}.$$

$$\begin{array}{l|l} 4 & Z = v_2 = \gamma^3w_2 + \gamma^2w_1^k + w_1^2 \\ 2k & T = \beta^2v_2 - v_1^2 = \gamma w_2 + w_1^k \\ 3k-2 & Y = v_1Z^{(k-1)/2} - \beta T = \gamma^{-1}(w_1Z^{(k-1)/2} - T) \\ 3k & iX = \beta Z^{(k-1)/2} - v_1T = \gamma^{-1}(Z^{(k-1)/2} - w_1T). \end{array}$$

Note that

$$\begin{aligned} iX + Y &= (Z^{(k-1)/2} - T)(\beta + v_1) \\ \alpha + \phi &= (iX + Y)Y + (Z^{(k-1)/2} + i^{k+1}T)Z^{(k-1)/2}T, \end{aligned}$$

and recall that for m even

$$i\alpha_1P'' + Q'' = i^{m+3}Y\varepsilon_1.$$

8.2 Transition functions for more general length 2 blowup

Generalization to $P'' = 0$.

Note that $P'' = 0$ implies m odd. The deformed equation simplifies to

$$0 = X^2 + Y^2Z - h'Q''^2 + 2\delta'YQ'',$$

with Plücker coordinates

$$\alpha = iXY, \quad \varepsilon = i^{-m+1}YQ'', \quad \varphi = Y^2.$$

Following the example of the Intriligator–Wecht case for k even (and m odd), we conjecture that the connection with the transition function coordinates in the blowup is

$$\begin{array}{l|l} 4 & Z = v_2 = \gamma^3w_2 + \gamma^2w_1^k + w_1^2 \\ 2k & T = \beta^2v_2 - v_1^2 = \gamma w_2 + w_1^k \\ 3k-2 & Y = i^{-m-1}\beta Q'' = i^{-m-1}\gamma^{-1}Q'' \\ 3k & iX = i^{-m-1}v_1Q'' = i^{-m-1}\gamma^{-1}w_1Q'', \end{array}$$

and we can express the upstairs coordinates as

$$\begin{aligned} \beta &= i^{m+1}Y/Q'' = -\varphi/\varepsilon = -\varphi_2 \\ v_1 &= i^{m+2}X/Q'' = -\alpha/\varepsilon = -\alpha_2 \\ v_2 &= Z \\ \gamma &= i^{-m-1}Q''/Y = -\varepsilon/\varphi = -\varepsilon_1 \\ w_1 &= iX/Y = \alpha/\varphi = \alpha_1 \\ w_2 &= \beta((-1)^{m+1}h' - w_1^k - 2i^{m+1}\beta\delta') \\ &= -\varepsilon_2^{-1}((-1)^{m+1}h' - \alpha_2^k + 2i^{m+1}\varepsilon_2^{-1}\delta'). \end{aligned}$$

From the original deformed D_{n+2} equation

$$0 = X^2 + Y^2Z - h'(ZP''^2 + Q''^2) + 2\delta'(YQ'' + (-1)^{m+1}XP''),$$

we know that ($P'' = 0$)

$$\begin{aligned}
v_2 = Z &= \frac{h'Q''^2 - X^2 - 2\delta'YQ''}{Y^2} \\
&= \gamma^2 h' + w_1^2 - 2\delta' i^{m+1} \gamma \\
&= \gamma^2 (\gamma w_2 + w_1^k + 2i^{m+1} \gamma^{-1} \delta') + w_1^2 - 2\delta' i^{m+1} \gamma \\
&= \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2.
\end{aligned}$$

Equivalently, the equation for the Grassmannian $G(2, 4) \subseteq \mathbb{P}^5$

$$(-Z\varphi - 2i^{m-1}\delta'\varepsilon)\phi + (-1)^{m+1}h'\varepsilon^2 + \alpha^2 = 0,$$

tells us that

$$\begin{aligned}
v_2 = Z &= \frac{\alpha^2 + (-1)^{m+1}h'\varepsilon^2 + 2i^{m+1}\delta'\varepsilon\varphi}{\varphi^2} \\
&= \left(\frac{\alpha}{\varphi}\right)^2 + (-1)^{m+1}h'\left(\frac{\varepsilon}{\varphi}\right)^2 + 2i^{m+1}\delta'\left(\frac{\varepsilon}{\varphi}\right) \\
&= w_1^2 + (-1)^{m+1}h'\gamma^2 - 2i^{m+1}\gamma\delta' \\
&= \gamma^2 (\gamma w_2 + w_1^k + 2i^{m+1} \gamma^{-1} \delta') + w_1^2 - 2\delta' i^{m+1} \gamma \\
&= \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2.
\end{aligned}$$

Notice that this upstairs transition functions does not depend on h' or Q'' ! However, since

$$\delta' = \delta'(T), \quad \text{and} \quad h' = h'(Z, T),$$

it is not clear that we can in general write this as transition functions between two \mathbb{C}^3 charts.

Assuming that these coordinates suffice to describe the blowup, we get exactly the Intriligator–Wecht transition functions:

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2.$$

In other words, the actual values for m , Q'' , δ' and h' don't matter, as long as the corresponding upstairs coordinates are legitimate (and this is where they may matter).

Full generality

We have Plücker coordinates

$$\begin{aligned}
\alpha &= iXY - i^{2m+1}h'P''Q'' \\
\varepsilon &= i^{m+3}XP'' + i^{-m+1}YQ'' \\
\varphi &= Y^2 - h'P''^2,
\end{aligned}$$

and equation for the Grassmanian $G(2, 4) \subseteq \mathbb{P}^5$

$$(-Z\varphi - 2i^{m-1}\delta'\varepsilon)\varphi + (-1)^{m+1}h'\varepsilon^2 + \alpha^2 = 0.$$

It is easy to check that this is exactly equivalent to the original deformed equation

$$0 = X^2 + Y^2 Z - h'(ZP''^2 + Q''^2) + 2\delta'(YQ'' + (-1)^{m+1}XP'').$$

Defining upstairs coordinates

$$\begin{array}{lcl} \beta & = & -\varphi/\varepsilon \quad \left| \quad \gamma & = & -\varepsilon/\varphi \\ v_1 & = & -\alpha/\varepsilon \quad \left| \quad w_1 & = & \alpha/\varphi \\ v_2 & = & Z & \left| \quad w_2 & = & \beta[(-1)^{m+1}h' - w_1^k - 2i^{m+1}\beta\delta'] \end{array}$$

and using the equation for $G(2, 4)$ yields transition function

$$\begin{aligned} v_2 = Z &= \frac{\alpha^2 + (-1)^{m+1}h'\varepsilon^2 + 2i^{m+1}\delta'\varepsilon\varphi}{\varphi^2} \\ &= \left(\frac{\alpha}{\varphi}\right)^2 + (-1)^{m+1}h' \left(\frac{\varepsilon}{\varphi}\right)^2 + 2i^{m+1}\delta' \left(\frac{\varepsilon}{\varphi}\right) \\ &= w_1^2 + (-1)^{m+1}h'\gamma^2 - 2i^{m+1}\gamma\delta' \\ &= \gamma^2(\gamma w_2 + w_1^k + 2i^{m+1}\gamma^{-1}\delta') + w_1^2 - 2\delta'i^{m+1}\gamma \\ &= \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2. \end{aligned}$$


Note that the choice of k was arbitrary in the definition of the upstairs coordinates. Moreover, we don't know if these are enough coordinates to fully describe the resolved geometry. However, in cases where it is enough (and this may depend on $k!$), we see that the upstairs transition functions are all the same as in the Intriligator–Wecht case:

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3 w_2 + \gamma^2 w_1^k + w_1^2.$$

The question is whether we can solve for X, Y and T in the $\varepsilon = 1$ chart (β, v_1, v_2) and similarly can we restrict to γ, w_1 and w_2 in the $\varphi = 1$ chart. In the k even Intriligator–Wecht case, with $\delta' = P'' = 0$, and $h' = T$, this certainly seems to be the case.

Chapter 9

Length 3



In this chapter we prove the third part of **Theorem 2**:

The resolved geometry corresponding to the Intriligator–Wecht superpotential

$$W(x, y) = \frac{1}{3}y^3 + yx^3$$

corresponds to the singular geometry

$$X^2 - Y^3 + Z^5 + 3TYZ^2 + T^3Z = 0.$$

9.1 The Case E_7

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator-Wecht superpotential

$$W(x, y) = \frac{1}{3}y^3 + yx^3,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma w_1^3 + \gamma^{-1}w_1^2.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}w_2 + \beta^{-1}(x + \beta^{-1}y)^3 + \beta(x + \beta^{-1}y)^2.$$

If we choose

$$w_2(\gamma) = \frac{x^3 + 3\gamma x^2 y - (x + \gamma y)^3}{\gamma^2},$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}(3x^2 y) + \beta^{-1}(x^3 + y^2) + 2xy + \beta x^2.$$

This is only holomorphic if

$$3x^2 y = x^3 + y^2 = 0,$$

and so we have a single \mathbb{P}^1 located at

$$w_1(\gamma) = w_2(\gamma) = 0, \quad v_1(\beta) = v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = 3x^2 y dx + (y^2 + x^3)dy = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma w_1^3 + \gamma^{-1}w_1^2,$$

are quasi-homogeneous if we assign the weights

$$\begin{array}{c|c|c|c|c|c} \beta & v_1 & v_2 & \gamma & w_1 & w_2 \\ \hline 1 & 3 & 5 & -1 & 2 & 8 \end{array}.$$

The global holomorphic functions will thus necessarily be quasi-homogeneous in these weights. We find the following global holomorphic functions:

$$\begin{array}{l|l} 6 & X = \beta v_2 - v_1^2 \\ 8 & Y = v_1 v_2 - \beta^2 X \\ 10 & Z = v_2^2 - \beta v_1 X \\ 15 & F = v_2^3 - 2v_1^3 X + (\beta^3 - 3v_1)X^2. \end{array}$$

The singular geometry \mathcal{M}_0

We conjecture that the ring of global holomorphic functions is generated by X, Y, Z and F , subject to the degree 30 relation

$$\mathcal{M}_0: F^2 - Z^3 + X^5 + 3X^2YZ + XY^3 = 0.$$

Do the functions X, Y, Z and F give us a blow-down map whose image \mathcal{M}_0 has an isolated E_7 singularity?

The blowup

We now verify that we have identified the right singular space \mathcal{M}_0 by inverting the blow-down. In the β and γ charts we find

$$\begin{array}{l|l} \beta = \frac{Y^2 + XZ}{F} & \gamma = \frac{F}{Y^2 + XZ} \\ v_1 = \frac{X^3 + YZ}{F} & w_1 = \frac{X^3 + YZ}{Y^2 + XZ} \\ v_2 = \frac{Z^2 - X^2Y}{F} & w_2 = \frac{X^4 - Y^3}{Y^2 + XZ} = w_1X - Y. \end{array}$$

This suggests that we should rewrite the equation for \mathcal{M}_0 as

$$\mathcal{M}_0: F^2 + XY(Y^2 + XZ) + X^2(X^3 + YZ) - Z(Z^2 - X^2Y) = 0,$$

and that we can obtain $\widehat{\mathcal{M}}$ by blowing up

$$F = Y^2 + XZ = X^3 + YZ = Z^2 - X^2Y = 0.$$

The locus \mathcal{C}

Let \mathcal{S} denote the surface

$$\mathcal{S}: F = Y^2 + XZ = 0.$$

Our $\widehat{\mathcal{M}}$ coordinate patches (β, v_1, v_2) and (γ, w_1, w_2) cover everything except the locus

$$\mathcal{C} = \mathcal{S} \cap \mathcal{M}_0.$$

The intersection of \mathcal{S} with the 3-fold \mathcal{M}_0 yields the new equation

$$X^5 + 2X^2YZ - Z^3 = 0.$$

(This was obtained by finding the Groebner basis for the ideal generated by $F, Y^2 + XZ$, and the equation for \mathcal{M}_0 .)

For $X \neq 0$ we can write

$$Z = -\frac{Y^2}{X},$$

and so the equations for \mathcal{C} become

$$\mathcal{C} : F = Y^2 + XZ = (X^4 - Y^3)^2 = 0, \quad (X \neq 0).$$

We can parametrize this curve by

$$\begin{aligned} X &= t^3, \\ Y &= t^4, \\ Z &= -t^5, \\ F &= 0. \end{aligned}$$

From this we see that blowing up $\mathcal{C} \subset \mathcal{M}_0$ is equivalent to blowing up

$$F = Y^2 + XZ = X^3 + YZ = Z^2 - X^2Y = X^4 - Y^3 = 0.$$

In this case we would have additional coordinates v_3, w_3 for the blowup:

$$\begin{array}{l|l} \beta = \frac{Y^2 + XZ}{F} & \gamma = \frac{F}{Y^2 + XZ} \\ v_1 = \frac{X^3 + YZ}{F} & w_1 = \frac{X^3 + YZ}{Y^2 + XZ} \\ v_2 = \frac{Z^2 - X^2Y}{F} & w_2 = \frac{X^4 - Y^3}{Y^2 + XZ} \\ v_3 = \frac{X^4 - Y^3}{F} & w_3 = \frac{Z^2 - X^2Y}{Y^2 + XZ}. \end{array}$$

Note that both v_3 and w_3 add nothing new, as we can solve for them in terms of the other coordinates:

$$\begin{aligned} v_3 &= \gamma^{-1}w_2 = \beta^4v_2 - \beta^3v_1^2 - v_1^3, \\ w_3 &= \beta^{-1}v_2 = \gamma^4w_2 + \gamma^2w_1^3 + w_1^2. \end{aligned}$$

These are precisely the additional coordinates we introduced in our resolution of the ideal sheaf (see Section 3.5). Finally, with these identifications we find transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma w_1^3 + \gamma^{-1}w_1^2,$$


which are exactly the ones we started with.

9.2 Matrix factorization for a length 3 singularity

Our singular geometry is in preferred versal form of type E_8 , and the presence of the T^3Z term in the polynomial¹

$$X^2 - Y^3 + Z^5 + 3TYZ^2 + T^3Z = 0,$$

tells us that our geometry has a length 3 singularity which can be resolved by blowing up the open node in the following E_8 diagram:²



An alternative method for finding the small resolution for this length 3 singularity should thus be possible by deforming the E_8 matrix factorization corresponding to this node. After some trial and error (and help from Maple), we find the following deformed matrix factorization for our singular equation:

$$\begin{pmatrix} \psi & -xI \\ xI & \varphi \end{pmatrix}, \quad \begin{pmatrix} \varphi & xI \\ -xI & \psi \end{pmatrix},$$

where we have used Knörrer's periodicity [26, Chapter 12] and the fact that

$$\varphi = \begin{pmatrix} -y & z^2 & tz \\ t & -y & z^2 \\ z & t & -y \end{pmatrix}, \quad \psi = \begin{pmatrix} y^2 - tz^2 & yz^2 + t^2z & z^4 + tyz \\ z^3 + ty & y^2 - tz^2 & yz^2 + t^2z \\ yz + t^2 & z^3 + ty & y^2 - tz^2 \end{pmatrix},$$

gives a matrix factorization for

$$-y^3 + z^5 + 3tyz^2 + t^3z = 0.$$

Using the approach in Chapter 5, this should allow us to perform the small resolution, and hence to extend Theorem 1 to include a length 3 singularity. We leave this for future work.

¹Note the change of coordinates!

²This uses results in the proof of the Katz–Morrison classification [19], as discussed in the Introduction.

Chapter 10

Other ADE cases

There are two ADE cases left in the Intriligator–Wecht classification: E_6 and E_8 . Using the algorithm from Chapter 6, we were able to find several global holomorphic functions in each case, but not enough to perform the blow-down. It is not clear whether or not these cases will yield singular spaces of lengths 4, 5 or 6, and for the moment we are skeptical that these small resolutions admit such simple descriptions. Nevertheless, we record here our progress, in the hope that it might be useful for future work.

10.1 The Case E_6

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator–Wecht superpotential

$$W(x, y) = \frac{1}{3}y^3 + \frac{1}{4}x^4,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^3 + \gamma^{-1}w_1^2.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}w_2 + \beta^{-2}(x + \beta^{-1}y)^3 + \beta(x + \beta^{-1}y)^2.$$

If we choose

$$w_2(\gamma) = \frac{x^3 - (x + \gamma y)^3}{\gamma},$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}x^3 + \beta^{-1}y^2 + 2xy + \beta x^2.$$

This is only holomorphic if

$$x^3 = y^2 = 0,$$

and so we have a single \mathbb{P}^1 located at

$$w_1(\gamma) = w_2(\gamma) = 0, \quad v_1(\beta) = v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = x^3 dx + y^2 dy = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3 w_2 + \gamma^2 w_1^3 + \gamma^{-1} w_1^2,$$

are quasi-homogeneous if we assign the weights

$$\begin{array}{c|c|c|c|c|c} \beta & v_1 & v_2 & \gamma & w_1 & w_2 \\ \hline 1 & 4 & 7 & -1 & 3 & 10 \end{array}.$$

The global holomorphic functions will thus necessarily be quasi-homogeneous in these weights. We find (so far!) the following global holomorphic functions:

$$\begin{array}{c|c|c} 8 & y_1 & = & \beta v_2 - v_1^2 \\ 9 & y_2 & = & \beta y_1 \\ 12 & y_3 & = & v_1 y_1 \\ 15 & y_4 & = & v_2 y_1 \\ 19 & y_5 & = & (v_1 v_2 - \beta^3 y_1) y_1 \\ 22 & y_6 & = & (v_2^2 - \beta^2 v_1 y_1) y_1 \\ 29 & y_7 & = & (v_2^3 - 2\beta v_1^3 y_1 + \beta^5 y_1^2) y_1 \end{array}$$

But there may be many more...

The singular geometry \mathcal{M}_0

The functions y_1, \dots, y_7 satisfy 21 independent relations. The first few are

$$\begin{array}{c|c|c} 24 & r_1 & = & y_1^3 - y_2 y_4 + y_3^2 = 0 \\ 27 & r_2 & = & y_2^3 - y_3 y_4 + y_1 y_5 = 0 \\ 30 & r_3 & = & -y_4^2 + y_1 y_6 + y_3 y_2^2 = 0 \\ 31 & r_4 & = & -y_6 y_2 + y_1^2 y_4 + y_3 y_5 = 0 \\ 34 & r_5 & = & -y_3 y_6 + y_4 y_5 + y_1^2 y_2^2 = 0 \\ 37 & r_6 & = & y_6 y_4 - y_1 y_7 - y_5 y_2^2 + 2y_2 y_3 y_1^2 = 0 \\ 38 & r_7 & = & -y_2 y_7 + y_5^2 - 2y_1^2 y_6 + 3y_1 y_4^2 = 0 \\ 39 & r_8 & = & -y_4^2 y_2 + y_4 y_3^2 + y_1 y_6 y_2 - y_1 y_3 y_5 = 0 \end{array}$$

and the remaining occur in degrees

$$41, 44, 45, 46, 52, 58, 62, 64, 66, 70, 94, 100, 138.$$

From a simple generating function we can generate all the monomials in each degree. Here we list only up to degree 40, and only in cases where there is more than one monomial in that degree.

24	y_1^3, y_3^2, y_2y_4	34	$y_1^2y_2^2, y_4y_5, y_3y_6$
27	y_2^3, y_3y_4, y_1y_5	35	$y_2^3, y_1y_3y_4, y_1^2y_5$
28	$y_1^2y_3, y_2y_5$	36	$y_2^4, y_3y_1^3, y_3^3, y_2y_3y_4, y_1y_2y_5$
29	$y_1y_2y_3, y_7$	37	$y_2y_3y_1^2, y_5y_2^2, y_4y_6, y_1y_7$
30	$y_3y_2^2, y_4^2, y_1y_6$	38	$y_2^2y_1y_3, y_1y_4^2, y_5^2, y_1^2y_6, y_2y_7$
31	$y_1^2y_4, y_3y_5, y_2y_6$	39	$y_2^3y_3, y_4y_1^3, y_4y_3^2, y_2y_4^2, y_1y_3y_5, y_1y_2y_6$
32	$y_1^4, y_1y_3^2, y_1y_2y_4$	40	$y_1^5, y_1^2y_3^2, y_2y_1^2y_4, y_2y_3y_5, y_2^2y_6$
33	$y_1^3y_2, y_3^2y_2, y_2^2y_4$		

Note that in almost every case, the relations r_i involve every possible monomial in their respective degrees.

10.2 The Case E_8

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator-Wecht superpotential

$$W(x, y) = \frac{1}{3}y^3 + \frac{1}{5}x^5,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^4 + \gamma^{-1}w_1^2.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}w_2 + \beta^{-2}(x + \beta^{-1}y)^4 + \beta(x + \beta^{-1}y)^2.$$

If we choose

$$w_2(\gamma) = \frac{x^4 - (x + \gamma y)^4}{\gamma},$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}x^4 + \beta^{-1}y^2 + 2xy + \beta x^2.$$

This is only holomorphic if

$$x^4 = y^2 = 0,$$

and so we have a single \mathbb{P}^1 located at

$$w_1(\gamma) = w_2(\gamma) = 0, \quad v_1(\beta) = v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = x^4 dx + y^2 dy = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^4 + \gamma^{-1}w_1^2,$$

are quasi-homogeneous if we assign the weights

$$\beta \left| \begin{array}{c|c} v_1 & v_2 \\ \hline 2 & 5 \end{array} \right| \gamma \left| \begin{array}{c|c} w_1 & w_2 \\ \hline -2 & 3 \end{array} \right| \left| \begin{array}{c} 14 \\ 14 \end{array} \right| .$$

The global holomorphic functions will thus necessarily be quasi-homogeneous in these weights. We find (so far!) the following global holomorphic functions:

$$\begin{array}{l|l} 10 & y_1 = \beta v_2 - v_1^2 \\ 12 & y_2 = \beta y_1 \\ 15 & y_3 = v_1 y_1 \\ 18 & y_4 = v_2 y_1 \\ 26 & y_5 = (v_2^2 - \beta^3 y_1) y_1 \end{array}$$

But there may be many more...

The singular geometry \mathcal{M}_0

The functions y_1, \dots, y_5 satisfy 2 independent relations

$$\begin{array}{l|l} 30 & r_1 = y_1^3 - y_2 y_4 + y_3^2 = 0 \\ 36 & r_2 = y_2^3 - y_4^2 + y_1 y_5 = 0 \end{array}$$

Remarks

- Each relation uses every possible monomial in its degree.
- r_1 and r_2 are almost identical to the first two relations in the E_6 case. Moreover, the first two relations r_1, r_2 in the E_6 case are the only relations we would have if we only considered the first five functions y_1, \dots, y_5 .
- The functions y_1, \dots, y_5 are almost identical to the first five functions in the E_6 case!

Chapter 11

The ‘Hat’ cases

In this section we analyze the extra ‘hat’ cases in the Intriligator–Wecht classification of superpotentials. In particular, we prove **Theorem 3**:

The singular geometries corresponding to the \widehat{O} , \widehat{A} , \widehat{D} and \widehat{E} cases in the Intriligator–Wecht classification of superpotentials are:

type	$W(x, y)$	$\partial_{w_1} E(\gamma, w_1)$	singular geometry \mathcal{M}_0
\widehat{O}	0	0	$\mathbb{C}^3/\mathbb{Z}_3$
\widehat{A}	$\frac{1}{2}y^2$	$\gamma^2 w_1$	$\mathbb{C} \times \mathbb{C}^2/\mathbb{Z}_2$
\widehat{D}	xy^2	γw_1^2	$X^2 + Y^2 Z - T^3 = 0$
\widehat{E}	$\frac{1}{3}y^3$	$\gamma^2 w_1^2$	$\text{Spec}(\mathbb{C}[a, b, u, v]/\mathbb{Z}_2)/(b^4 - u^2 - av)$.

11.1 The Case \widehat{O}

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator-Wecht superpotential

$$W(x, y) = 0,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}w_2.$$

If we choose

$$w_2(\gamma) = 0,$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = 0.$$

This is holomorphic for all x and y , and so we have a 2-parameter family of \mathbb{P}^1 s located at

$$w_1(\gamma) = x + \gamma y, w_2(\gamma) = 0, \quad v_1(\beta) = \beta x + y, v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2$$

are quasi-homogeneous if we assign the weights

$$\beta \left| \begin{array}{c} v_1 \\ d+1 \end{array} \right| v_2 \left| \begin{array}{c} e-3 \end{array} \right| \gamma \left| \begin{array}{c} w_1 \\ d \end{array} \right| w_2 \left| \begin{array}{c} e \end{array} \right|.$$

Notice the freedom in choosing d and e : there is a two-dimensional lattice of possible weight assignments. The global holomorphic functions will necessarily be quasi-homogeneous in these weights. We find global holomorphic functions:

$$X_{ij} = \beta^i v_1^j v_2 = \gamma^{3-i-j} w_1^j w_2, \quad i, j \geq 0, \quad i + j \leq 3.$$

The singular geometry \mathcal{M}_0

If we rewrite our functions in a homogeneous manner as

$$\tilde{X}_{ij} = a^{3-i-j}b^i c^j, \quad i, j \geq 0, \quad i + j \leq 3,$$

we can now identify the ring of global holomorphic functions as homogeneous polynomials of degree 3 in 3 variables. In other words, the ring is isomorphic to

$$\mathbb{C}[a, b, c]^{\mathbb{Z}_3},$$

and our singular variety is simply

$$\mathcal{M}_0 : \quad \mathbb{C}^3/\mathbb{Z}_3.$$

The blowup

We now verify that we have identified the right singular space \mathcal{M}_0 by inverting the blow-down. In the β and γ charts we find

$$\begin{array}{l|l} \beta = X_{10}/X_{00} = \tilde{X}_{10}/\tilde{X}_{00} = b/a & \gamma = a/b \\ v_1 = X_{01}/X_{00} = \tilde{X}_{01}/\tilde{X}_{00} = c/a & w_1 = c/b \\ v_2 = X_{00} = \tilde{X}_{00} = a^3 & w_2 = b^3 \end{array}$$

which gives transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2.$$

These are precisely what we started with!

Remark

In the resolved $\widehat{\mathcal{M}}$ geometry, what we have is a \mathbb{P}^1 inside a \mathbb{P}^2 (or any other del Pezzo surface). If you have a \mathbb{P}^2 inside a Calabi-Yau and blow it down, you get $\mathbb{C}^3/\mathbb{Z}^3$ as the singular point.

11.2 The Case \widehat{A}

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator-Wecht superpotential

$$W(x, y) = \frac{1}{2}x^2,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}(w_2 + y) + \beta^{-2}x.$$

If we choose

$$w_2(\gamma) = -y,$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}x.$$

This is only holomorphic $x = 0$. Since y is free, we have a 1-parameter family of \mathbb{P}^1 s located at

$$w_1(\gamma) = \gamma y, w_2(\gamma) = -y, \quad v_1(\beta) = y, v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = xdx = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1$$

are quasi-homogeneous if we assign the weights

$$\beta \left| \begin{array}{c} v_1 \\ 1 \end{array} \right|_{d+1} \left| \begin{array}{c} v_2 \\ d-2 \end{array} \right|_{d-2} \left| \begin{array}{c} \gamma \\ -1 \end{array} \right|_{-1} \left| \begin{array}{c} w_1 \\ d \end{array} \right|_d \left| \begin{array}{c} w_2 \\ d+1 \end{array} \right|_{d+1}.$$

Notice the freedom in choosing d : there is a one-dimensional lattice of possible weight assignments. The global holomorphic functions will necessarily be quasi-homogeneous in these weights. We find global holomorphic functions:

$$\begin{array}{l} d-2 \left| \begin{array}{l} y_1 \\ y_2 \\ y_3 \\ y_4 \end{array} \right. = \begin{array}{l} v_2 = \gamma^3w_2 + \gamma^2w_1 \\ \beta v_2 = \gamma^2w_2 + \gamma w_1 \\ \beta^2 v_2 = \gamma w_2 + w_1 \\ \beta^3 v_2 - v_1 = w_2 \end{array} \end{array}$$

The singular geometry \mathcal{M}_0

The functions y_i satisfy the single degree $2d - 2$ relation

$$y_2^2 - y_1 y_3 = 0,$$

with y_4 free. In other words, our singular geometry \mathcal{M}_0 is a curve of A_1 singularities, parametrized by y_4 .

The blowup

We now verify that we have identified the right singular space \mathcal{M}_0 by inverting the blow-down. In the β and γ charts we find

$$\begin{array}{l|l} \beta = & y_2/y_1 \\ v_1 = & (y_2y_3 - y_1y_4)/y_1 \\ v_2 = & y_1 \end{array} \quad \left| \quad \begin{array}{l} \gamma = & y_1/y_2 \\ w_1 = & (y_2y_3 - y_1y_4)/y_2 \\ w_2 = & y_4. \end{array} \right.$$

This gives transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1,$$

as expected.

11.3 The Case \widehat{D}

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator-Wecht superpotential

$$W(x, y) = x^2y,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma w_1^2.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}(w_2 + y^2) + \beta^{-2}(2xy) + \beta^{-1}x^2.$$

If we choose

$$w_2(\gamma) = -y^2,$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}(2xy) + \beta^{-1}x^2.$$

This is only holomorphic if $x = 0$. Since y is free, we have a 1-parameter family of \mathbb{P}^1 s located at

$$w_1(\gamma) = \gamma y, w_2(\gamma) = -y^2, \quad v_1(\beta) = y, v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = 2xy dx + x^2dy = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma w_1^2$$

are quasi-homogeneous if we assign the weights

$$\beta \left| \begin{array}{c} v_1 \\ d+1 \end{array} \right| \quad v_2 \left| \begin{array}{c} 2d-1 \end{array} \right| \quad \gamma \left| \begin{array}{c} w_1 \\ d \end{array} \right| \quad w_2 \left| \begin{array}{c} 2d+2 \end{array} \right|.$$

Notice the freedom in choosing d : there is a one-dimensional lattice of possible weight assignments. The global holomorphic functions will necessarily be quasi-homogeneous in these weights. We find global holomorphic functions:

$$\begin{array}{l} 2d-1 \\ 2d \\ 3d \\ 2d+2 \end{array} \left| \begin{array}{l} y_1 \\ y_2 \\ y_3 \\ y_4 \end{array} \right| = \begin{array}{l} v_2 = \gamma^3w_2 + \gamma w_1^2 \\ \beta v_2 = \gamma^2w_2 + w_1^2 \\ v_1v_2 = \gamma^2w_1w_2 + w_1^3 \\ \beta^3v_2 - v_1^2 = w_2 \end{array}$$

The singular geometry \mathcal{M}_0

The functions y_i satisfy the single degree $6d$ relation

$$y_3^2 - y_2^3 + y_1^2y_4 = 0.$$

The blowup

We now verify that we have identified the right singular space \mathcal{M}_0 by inverting the blow-down. In the β and γ charts we find

$$\begin{array}{l} \beta \\ v_1 \\ v_2 \end{array} = \begin{array}{l} y_2/y_1 \\ y_3/y_1 \\ y_1 \end{array} \left| \begin{array}{l} \gamma \\ w_1 \\ w_2 \end{array} \right| = \begin{array}{l} y_1/y_2 \\ y_3/y_2 \\ y_4 \end{array},$$

with transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma w_1^2.$$

11.4 The Case \widehat{E}

The resolved geometry $\widehat{\mathcal{M}}$

From the Intriligator-Wecht superpotential

$$W(x, y) = \frac{1}{3}x^3,$$

we compute the resolved geometry $\widehat{\mathcal{M}}$ in terms of transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^2.$$

To find the \mathbb{P}^1 s, we substitute $w_1(\gamma) = x + \gamma y$ into the v_2 transition function

$$v_2(\beta) = \beta^{-3}(w_2 + 2xy + \gamma y^2) + \beta^{-2}x^2.$$

If we choose

$$w_2(\gamma) = -2xy - \gamma y^2,$$

in the β chart the section is

$$v_1(\beta) = \beta x + y, \quad v_2(\beta) = \beta^{-2}x^2.$$

This is only holomorphic if $x = 0$. Since y is free, we have a 1-parameter family of \mathbb{P}^1 s located at

$$w_1(\gamma) = \gamma y, w_2(\gamma) = -\gamma y^2, \quad v_1(\beta) = y, v_2(\beta) = 0.$$

This is exactly what we expect from computing critical points of the superpotential

$$dW = x^2 dx = 0.$$

Global holomorphic functions

The transition functions

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1}w_1, \quad v_2 = \gamma^3w_2 + \gamma^2w_1^2$$

are quasi-homogeneous if we assign the weights

$$\beta \left| \begin{array}{c} v_1 \\ d+1 \end{array} \right| v_2 \left| \begin{array}{c} 2d-2 \\ 2d-2 \end{array} \right| \gamma \left| \begin{array}{c} w_1 \\ -1 \end{array} \right| w_2 \left| \begin{array}{c} d \\ 2d+1 \end{array} \right|.$$

Notice the freedom in choosing d : there is a one-dimensional lattice of possible weight assignments. The global holomorphic functions will necessarily be quasi-homogeneous in these weights. We find global holomorphic functions:

$$\begin{array}{r|l} 2d-2 & y_1 = v_2 \\ 2d-1 & y_2 = \beta v_2 \\ 2d & y_3 = \beta^2 v_2 \\ 3d-1 & y_4 = v_1 v_2 \\ 3d & y_5 = \beta v_1 v_2 \\ 4d & y_6 = v_1^2 v_2 \\ 4d+1 & y_7 = \beta(\beta^4 v_2 - v_1^2)v_2 = \beta v_3 v_2 \\ 5d+1 & y_8 = v_1(\beta^4 v_2 - v_1^2)v_2 = v_1 v_3 v_2 \\ 6d+2 & y_9 = (\beta^4 v_2 - v_1^2)^2 v_2 = v_3^2 v_2 \end{array}$$

where we have defined

$$v_3 = \beta^4 v_2 - v_1^2 = \gamma^{-1} w_2.$$

Note that this is the same definition made in our resolution of the ideal sheaf (Section 3.5).

The singular geometry \mathcal{M}_0

The functions y_i satisfy a total of 20 distinct relations, most of which are obvious. To simplify things, consider the monomial mapping

$$\beta^i v_1^j v_3^k v_2 \longmapsto a^{2-i-j-k} b^i c^j f^k.$$

Our functions now become

$$\begin{array}{r|l} 2d-2 & y_1 = a^2 \\ 2d-1 & y_2 = ab \\ 2d & y_3 = b^2 \\ 3d-1 & y_4 = ac \\ 3d & y_5 = bc \\ 4d & y_6 = c^2 \\ 4d+1 & y_7 = bf \\ 5d+1 & y_8 = cf \\ 6d+2 & y_9 = f^2 \end{array}$$

Note that

$$\begin{array}{l} \beta = y_2/y_1 = b/a \\ v_1 = y_4/y_1 = c/a \\ v_2 = y_1 = a^2 \\ v_3 = y_7/y_2 = f/a, \end{array}$$

so the relation defining v_3 becomes

$$af = b^4 - c^2.$$

This means we can add the function af to our list, together with the relation:

$$y_{10} = af = b^4 - c^2.$$

Now the functions y_1, \dots, y_{10} are exactly the 10 monomials of degree 2 in 4 variables, together with the above relation. The ring of global holomorphic functions is thus

$$(\mathbb{C}[a, b, c, f]/\mathbb{Z}_2)/(af - b^4 + c^2),$$

where the \mathbb{Z}_2 acts diagonally as -1. In other words, we have a hypersurface in a \mathbb{Z}_2 quotient space:

$$(b^2 + c)(b^2 - c) = af \quad \text{in } \mathbb{C}^4/\mathbb{Z}_2.$$

We can immediately see from this equation that a small resolution, where we blow up an ideal of the form

$$b^2 + c = a = 0,$$

won't work, since the \mathbb{Z}_2 action interchanges $b^2 + c$ and $b^2 - c$. We will need to do a big blow up of the origin instead.

The blowup

We now verify that we have identified the right singular space \mathcal{M}_0 by inverting the blow-down. In the β and γ charts we find

$$\begin{array}{l|l} \beta = b/a & \gamma = a/b \\ v_1 = c/a & w_1 = c/b \\ v_2 = a^2 & w_2 = f/b \\ v_3 = f/a & w_3 = b^2 \end{array}$$

We will perform the big blowup of the origin, with corresponding \mathbb{P}^3 coordinates:

$$\begin{array}{cccc} a & = & b & = & c & = & f & = & 0. \\ \alpha & & \delta & & \rho & & \nu & & \end{array}$$

Note that all eight coordinates switch sign under the \mathbb{Z}_2 action.

The blowup has four coordinate charts

$$\begin{array}{c|c|c|c} \alpha = 1 & \delta = 1 & \rho = 1 & \nu = 1 \\ \hline a = a & a = \alpha_2 b & a = \alpha_3 c & a = \alpha_4 f \\ b = \delta_1 a & b = b & b = \delta_3 c & b = \delta_4 f \\ c = \rho_1 a & c = \rho_2 b & c = c & c = \rho_4 f \\ f = \nu_1 a & f = \nu_2 b & f = \nu_3 c & f = f \\ \hline \nu_1 = \delta_1^4 a^2 - \rho_1^2 & b^2 = \alpha_2 \nu_2 + \rho_2^2 & \alpha_3 \nu_3 = \delta_3^4 c^2 - 1 & \alpha_4 = \delta_4^4 f^2 - \rho_4^2 \\ \hline (\alpha^2, \delta_1, \rho_1) & (\alpha_2, \rho_2, \nu_2) & (\alpha_3, \delta_3, c^2, \nu_3) & (\delta_4, \rho_4, f^2) \end{array}$$

Remarks

- The functions $\alpha_i, \delta_i, \rho_i,$ and ν_i are all invariant under the \mathbb{Z}_2 action, since they are all ratios of functions which change sign:

$$\delta_1 = \delta/\alpha, \quad \rho_2 = \rho/\delta, \quad \dots \text{ etc.}$$

- Because $a, b, c,$ and f all change sign under the \mathbb{Z}_2 action, we must take their invariant counterparts $a^2, b^2, c^2,$ and f^2 when we list the final coordinates for each chart.
- In the $\delta = 1$ chart, we solve for b^2 instead of b , because b is not an invariant function.

- The blow up is nonsingular. In the $\alpha = 1, \delta = 1,$ and $\nu = 1$ charts we see this because we are left with three coordinates and no relations, so these charts are all isomorphic to \mathbb{C}^3 . In the $\rho = 1$ chart, we have a hypersurface in \mathbb{C}^4 defined by the non-singular equation

$$\alpha_3 \nu_3 = \delta_3^4 c^2 - 1.$$

Transition functions

Between the first two charts $\alpha = 1$ and $\delta = 1,$ we have transition functions

$$\begin{aligned} \delta_1 &= \delta/\alpha = \alpha_2^{-1} \\ \rho_1 &= \rho/\alpha = (\delta/\alpha)(\rho/\delta) = \alpha_2^{-1} \rho_2 \\ a^2 &= \alpha_2^2 b^2 = \alpha_2^2 (\alpha_2 \nu_2 + \rho_2^2) \\ &= \alpha_2^3 \nu_2 + \alpha_2^2 \rho_2^2 \end{aligned}$$

Notice that

$$\begin{array}{l|l} \delta_1 = b/a = \beta & \alpha_2 = a/b = \gamma \\ \rho_1 = c/a = v_1 & \rho_2 = c/b = w_1 \\ a^2 = a^2 = v_2 & \nu_2 = f/b = w_2, \end{array}$$

and so our transition functions are really

$$\beta = \gamma^{-1}, \quad v_1 = \gamma^{-1} w_1, \quad v_2 = \gamma^3 w_2 + \gamma^2 w_1^2.$$

These are exactly the \widehat{E} transition functions we started with!

11.5 Comparison with ADE cases

We have seen that the singular geometries corresponding to Intriligator and Wecht's 'hat' cases are given by

$$\begin{array}{l|l|l} \widehat{O} & W(x, y) = 0 & \mathbb{C}^3/\mathbb{Z}_3 \\ \widehat{A} & W(x, y) = \frac{1}{2}y^2 & \mathbb{C}[X, Y, Z, T]/(XY - Z^2) \cong \mathbb{C} \times \mathbb{C}^2/\mathbb{Z}_2 \\ & & \text{geometry has curve of } A_1 \text{ singularities} \\ \widehat{D} & W(x, y) = xy^2 & y_1^2 - y_2^3 + y_3^2 y_4 = 0 \\ & & \text{recall geometry for } D_{k+2} : \\ & & y_1^2 + y_2^3 + y_3^2 y_4 + y_4^k y_2 = 0 \\ \widehat{E} & W(x, y) = \frac{1}{3}y^3 & (\mathbb{C}[a, b, u, v]/\mathbb{Z}_2)/(b^4 - u^2 - av) \\ & & \text{This is a hypersurface in } \mathbb{C}^4/\mathbb{Z}_2. \end{array}$$

Note that in both the \widehat{A} and \widehat{D} cases, the resulting equations can be obtained from the A_k and D_{k+2} equations by dropping the k -dependent terms. In other words, we are

tempted to think of \widehat{A} and \widehat{D} as the $k \rightarrow \infty$ limit. Perhaps in trying to come up with an analogous statement for \widehat{E} we can learn something about the “missing” E_6 and E_8 cases. In particular, it will be interesting to understand the role of these spaces in a geometric model for RG flow.

Chapter 12

Future directions

We have learned many things about the interplay between string dualities and Gorenstein threefold singularities. The full story, however, is far from complete. We end by posing a series of questions for the future which are beyond the scope of this work.

From the Intriligator–Wecht classification, we still need to understand the E_6 and E_8 cases. Using our algorithm we have found many global holomorphic functions, but not enough to give us the blow-down. There are also questions which arise from the extra ‘hat’ cases. What is the interpretation of the \widehat{O} , \widehat{A} , \widehat{D} , and \widehat{E} geometries from the string theory perspective? The \mathbb{P}^1 s are no longer isolated; do they correspond to D-branes wrapping families of \mathbb{P}^1 s? Moreover, what is the role of higher order terms in the superpotential?

Do we have a geometric model for RG flow? Proposition 1 suggests that the geometry might encode something about the RG fixed points of the corresponding matrix models or gauge theories. Can Proposition 1 be extended? The discussion in Section 7.2 suggests that this should be possible, since we can find an appropriate change of coordinates to make it work in the A_k cases. Finding more general coordinate changes which can show how the rest of the terms in the superpotential are affected when a bundle-changing coordinate is “integrated out” is a necessary step in developing this kind of geometric picture.

Furthermore, Intriligator and Wecht have a chart of all possible flows between the RG fixed points. We can make a similar chart based on our geometric framework. Do they match? Finally, what is the role of fundamentals? Our entire analysis involves only adjoint fields, which correspond geometrically to parameters of the \mathbb{P}^1 deformation space. Intriligator and Wecht only find the ADE classification for superpotentials involving 2 adjoint fields, but their paper also analyzes many cases with fundamentals. Is it possible to have a geometric interpretation for these fields?

As far as Ferrari’s construction is concerned, there are many open ends to be explored. Can we generalize Ferrari’s framework to include perturbation terms for both v_1 and v_2 transition functions? Can we generalize for cases where the geometry is specified by more than two charts? This would enable more flexibility in identifying superpotentials in a “bottom-up” approach. On the other hand, the techniques developed in [3] in principle allow computation of the superpotential in general. In cases where the superpotential

cannot be easily identified in the transition functions, perhaps this approach should be used instead.

Moreover, in all of our new cases there is still work to be done to complete the remaining steps in Ferrari's program. For example, what is the solution to the matrix model corresponding to the length 3 singularity? And what can we learn about the matrix models corresponding to the 'hat' cases? Although the singularities are no longer isolated, is it still possible to compute resolvents from the geometry? If Ferrari's conjecture about the Calabi-Yau geometry encoding the solution to the matrix model is correct, we should now be able to solve the matrix models corresponding to the length 3 and 'hat' cases. If solutions are already known (or can be computed using traditional matrix model techniques), these examples will provide new tests to the conjecture.

Finally, there is the idea of Conjecture 1 – that small resolutions for Gorenstein threefold singularities might be obtainable by deforming matrix factorizations in the surface case. In Theorem 1 we were able to prove this in the length 1 and length 2 cases, and with the help of the Intriligator–Wecht E_7 case we are very close to showing this in length 3 as well. An ambitious, but tangible goal is to extend this result to lengths 4, 5 and 6. In particular, this may lead us to an alternative (and more concrete) proof of the original Katz–Morrison classification [19].

Appendix A

Maple code for blow-down algorithm

A.1 Description of the code

The actual maple routine has the following parts:

- Input information
 - perturbation terms for ADE cases
 - weights for β, v_1, v_2 in each case
 - initial (empty) seed array X .
- Ideal-related definitions
 - ideal in $\mathbb{C}[\beta, v_1, v_2, \gamma, w_1, w_2]$
 - term order T (lexdeg, eliminate v_2)
 - term order TP (wdeg, degrees reflect poles)
 - groebner basis G for ideal, wrt T
 - `modG`, `arraymodG`, `shift` and `mix`, all compute reductions modulo the ideal G
- Lists of monomials
 - ‘`genfun`’ creates generating function of q-homog monomials
 - ‘`List`’ extracts all monomials in weighted degree n (uses ‘`genfun`’)
- Procedures on arrays (used exclusively for ‘`newfun`’ and ‘`invert`’)
 - ‘`concat`’ takes an array and adds an element to the end
 - ‘`yarray`’ creates an array of indeterminates $[y_1, \dots, y_n]$
 - ‘`equal0`’ converts an array A into a list $\{A_i = 0\}$




Figure A.1: An algorithm for constructing the blow down

- ‘**subtract**’ takes the difference between two arrays
- ‘**partials**’ takes the derivative of all in functions in an array A wrt a given variable, then evaluates at a point ‘ P ’ (a list of equations) and returns a list
- ‘**zero**’ creates a list of m elements which are all 0
- ‘**extract**’ picks out the elements which are pure in γ, w_1, w_2 from an array G .

- Main routines

- ‘**xreduce**’ is *the* main routine: it inputs a function and a list F of other functions $F[i,1]$ together with their leading mixed terms $F[i,2]$ and leading mixed coefficients $F[i,3]$. The function is reduced as much as possible by F , and what is returned is a polynomial in which the original function is modified by terms $F[i,1]$ which are subtracted in order to reduce the leading mixed term. (Uses **shift**, **mix**, and **TP**.)
- ‘**polysearch**’ receives an array ‘ L ’ of polynomials and a list X of global holomorphic functions as input. It uses ‘**xreduce**’ to cycle through each polynomial and maximally reduce its leading mixed term. It then uses the results to build the array $F[i,j]$ of maximally reduced polynomials and their mixed terms, all in the same weighted degree. When the mixed term is 0, ‘**newfun**’ is used to check if the global homogeneous function is in the ring generated by the list Y

of prior ghf's. The final output is the updated list Y . (Uses `shift`, `mix`, `TP`, `xreduce`, and `newfun`.)

- `newfun` checks if the global holomorphic function 'f' is new, in the sense that it is not in the ring generated by previous ghf's. It returns the (potentially updated) list of truly independent ghf's. (Uses `yarray`, `concat`, `subtract`, `extract`, `parcials`, `equal0`, `zero`.)

- Execution commands

- `findpoly` searches for global holomorphic functions within certain bounds on the weighted degree. (Uses `List` and `polysearch`)
- `invert` solves for original variables (β, v_1, v_2 or γ, w_1, w_2) in terms of the new ones: global holomorphic functions fed in through the array A . (Uses `yarray` and `subtract`.)

A.2 Maple code

```
> with(Groebner):
> extraterm:=0:
```

THE FOLLOWING MUST BE CHANGED ACCORDING TO ADE CASE

These are the perturbation terms for each of the ADE cases.

```
> Ohat:= 0: Ahat:= g^2*w[1]: Dhat:= g*w[1]^2: Ehat:= g^2*w[1]^2:
> Ak:= k -> g^2*w[1]^k + w[1]: Dk:= k -> g^2*w[1]^k + w[1]^2:
> E6:= b*w[1]^2 + g^2*w[1]^3: E7:= b*w[1]^2 + g*w[1]^3: E8:= b*w[1]^2 + g^2*w[1]^4:
```

These are the weights $[\text{deg}(b), \text{deg}(v[1]), \text{deg}(v[2])]$ in each of the cases.

```
> deg_Ohat:= [1,1,1]: deg_Ahat:= [1,4,1]: deg_Dhat:= [1,2,1]: deg_Ehat:= [1,3,2]:
> deg_Ak:= k -> [k-1,k+1,2]: deg_Dk:= k -> [k-2,k,4]:
> deg_E6:= [1,4,7]: deg_E7:= [1,3,5]: deg_E8:= [2,5,8]:
```

This is the perturbation term in the transition functions, and what distinguishes our different cases.

```
> pterm:=Ak(4);
```

"d" captures the weights for $b, v[1], v[2]$ in order to make the transition functions quasi-homogeneous.

```
> d:=deg_Ak(4);
```

The first holomorphic function goes into the "seed" array X .

```
> X:=Array([]):
```

IDEAL AND REDUCTION MODULO IDEAL

We define the ideal coming from the transition functions, in which we will look for global holomorphic functions.

```
> ideal:= [b*g-1, v[1]-b*w[1], v[2]-g^3*w[2]-pterm];
```

Our chosen term order for computing groebner basis and reduction mod the ideal.

```
> T:=lexdeg([v[2]], [w[2], v[1], w[1], b, g]):
```

The term order for keeping track of the polar degree, which we use to order the cancellations.

```
> TP:=wdeg([1,1,1,-1,0,0], [b,v[1],v[2],g,w[1],w[2]]):
```

We compute the groebner basis of 'ideal' with respect to the term order T .

```
> G:=gbasis(ideal,T);
```

The function "modG" computes the reduction of a function f modulo the ideal G .

```
> modG:= f -> normalf(f,G,T):
```

The procedure "arraymodG" computes the reduction of every function in the array "A" modulo the ideal G.

```
> arraymodG:=proc(A::Array)
> local n,i,B;
> n:=ArrayNumElems(A);
> B:=Array(1..n);
> for i to n do
>   B[i]:=modG(A[i]);
> end do;
> return B;
> end:
```

The function "mix" extracts the mixed terms from a polynomial, by subtracting off the pure terms from each chart.

```
> mix:= f -> modG(f) - subs(g=0,w[1]=0,w[2]=0,modG(f)) - subs(b=0,v[1]=0,v[2]=0,modG(f)):
```

The function "shift" just brings terms in b,v[1],v[2] over to the left-hand-side.

```
> shift:= f -> f - subs(g=0,w[1]=0,w[2]=0,modG(f)):
```

GENERATING MONOMIALS OF SAME WEIGHTED HOMOGENEOUS DEGREE

The function "genfun" creates a generating function for the quasi-homogeneous monomials in each degree up to n.

```
> genfun:= n ->
  convert(map(expand,series(1/((1-b*u^d[1])*(1-v[1]*u^d[2])*(1-v[2]*u^d[3])),u,n+1)),polynom):
```

The function "List" uses the above generating function to extract the monomials in weighted degree n.

```
> List:= n -> Array([op(1+coeff(genfun(n),u,n))]):
```

PROCEDURES ON ARRAYS

The "concat" procedure takes an array "A" and tacks on an extra element "a" onto the end.

```
> concat:= proc(A::Array, a)
> local B,n;
> n:= ArrayNumElems(A);
> B:= Array(1..n+1);
> B[1..n]:= A[1..n];
> B[n+1]:= a;
> return B;
> end:
```

The "yarray" procedure creates an array with indeterminates y[1],...,y[n].

```
> yarray:= proc(n::integer)
> local Y,i;
> Y:= Array(1..n);
> for i from 1 to n do
>   Y[i]:= y[i];
> end do;
> return Y;
> end:
```

The procedure "equal0" converts an array into a list A[i]=0.

```
> equal0:= proc(A::Array)
> local B,i,n;
> n:=ArrayNumElems(A);
> B:=Array(1..n);
> for i from 1 to n do
>   B[i]:= A[i]=0;
> end do;
> return convert(B,'list');
> end:
```

The "subtract" procedure takes the difference between two arrays.

```
> subtract:= proc(A::Array, B::Array)
> local C,i,n;
> n:= ArrayNumElems(A);
> C:= Array(1..n);
> for i from 1 to n do
>   C[i]:= A[i]-B[i]
> end do;
> return C;
> end:
```

The procedure "partials" takes the derivative of all of the functions in A w.r.t. the variable "var", then evaluates at "P".

```
> partials:= proc(A::Array,P::list,var)
> local i,n,t;
> n:=ArrayNumElems(A); t:=Array(1..n);
> for i from 1 to n do
>   t[i]:=eval(subs(P,diff(A[i],var)));
> end do;
> return convert(t,'list');
> end:
```

The procedure "zero" creates an array of 0's.

```
> zero:=proc(m::integer)
> local A,i; A:=Array(1..m);
> for i to m do
>   A[i]:=0;
> end do;
> return convert(A,'list');
> end:
```

The procedure "extract" picks out elements which are pure in (g,w[1],w[2]) from the array G.

```
> extract:=proc(G::Array)
> local n,i,j,h,H;
> n:=ArrayNumElems(G); H:=Array(1..n); j:=1;
> for i to n do
>   h:= subs(v[2]=0,v[1]=0,b=0,G[i]);
>   if G[i]=h then
>     H[j]:=G[i];
>     j:=j+1;
>   end if;
> end do;
> return H[1..j-1];
> end:
```

THE MAIN REDUCTION ROUTINES

The procedure "xreduce" is the main routine. It reduces the function mod the array "F", returning a modified function "F" whose mixed terms have lower degree.

```
> xreduce:= proc (function,F::Array)
> local i,n,f,h,lt,lc,subtraction;
> n:= ArrayNumElems(F)/3;
> f:= shift(function);
> h:= mix(f);
> while h <> 0 do
>   lt:= leadterm(h,TP);
```



```

> lc:= leadcoeff(h,TP);
> subtraction:=0;
> for i from n to 1 by -1 do
>   if F[i,2] = lt then
>     f:= f-lc/F[i,3]*F[i,1];
>     h:= mix(f);
>     subtraction:= 1;
>   end if;
>   if subtraction = 1 then
>     break;
>   end if;
> end do;
> if subtraction = 0 then
>   h:= h - lc*lt;
> end if;
> end do;
> return f;
> end:

```

The procedure "polysearch" receives an array "L" of polynomials as input, uses "xreduce", and finds the global holomorphic functions.

```

> polysearch:= proc(L:Array, X:Array, dw::integer)
> local i,j,n,f,F,s1,nf,Y,t;
> n:= ArrayNumElems(L);
> F:= Array(1..n,1..3);
> F[1,1]:= r(L[1]);
> s1:= s(F[1,1]);
> F[1,2]:= leadterm(s1,TP);
> F[1,3]:= leadcoeff(s1,TP);
> Y:= X;
> print('n'=n);
> j:= 2;
> for i from 2 to n do
>   print(i,j); print('wdeg'=dw);
>   f:= xreduce(L[i],F[1..j-1,1..3]);
>   if f <> 0 then
>     s1:= mix(f);
>     F[j,1]:= f;
>     if s1 = 0 then
>       print('f'=F[j,1]);
>       print('holomorphic');
>       nf:= newfun(F[j,1],Y);
>       Y:= nf[1]; t:= nf[2];
>       print(t);
>     else
>       F[j,2]:= leadterm(s1,TP);
>       F[j,3]:= leadcoeff(s1,TP);
>       print('h'=F[j,3]*F[j,2]);
>     end if;
>     j:= j+1;
>   end if;
> end do;
> return Y;

```

```
> end:
```

The procedure "newfun" checks if the global holomorphic function "f" is new, in the sense that it is not a multiple of previous functions. It returns the (potentially updated) list of truly distinct global holomorphic functions Y, which was fed in through X.

```
> newfun:= proc (f,X::Array)
> local Y,yideal,TY,G,GY,A,n,m,t;
> n:=ArrayNumElems(X)+1;
> Y:=yarray(n); A:=concat(X,f);
> TY:=lexdeg([v[2],v[1],b],convert(Y,'list'));
> yideal:= convert(subtract(Y,A),'list');
> G:= gbasis(yideal,TY); GY:=extract(Array(G)); m:=ArrayNumElems(GY);
t:=parcials(GY,equal0(Y),y[n]);
> if t = zero(m) then return [A,'New! ',GY,t]; else return [X,'not_new',GY,t]; end if;
> end:
```

EXECUTION COMMANDS

The procedure "findpoly" searches for global holomorphic polynomials between certain bounds ("m" and "n") on the weighted degree. It takes as input the array "X", which contains previously found functions.

```
> findpoly:= proc(start,step,n,X)
> local i, j, Y, L;
> Y:= X;
> for i from 0 to n do
>   j:= start+step*i;
>   print('o');
>   print('o');
>   print('wdeg' = j);
>   print('o');
>   L:= List(j);
>   Y:= polysearch(L,Y,j);
>   print('Y'=Y);
> end do;
> return Y;
> end:

> invert:=proc(A::Array, type)
> local n,TA,yideal,G;
> n:=ArrayNumElems(A);
> if type = beta then TA:=lexdeg([v[2],v[1],b],convert(yarray(n),'list')); else
TA:=lexdeg([w[2],w[1],g],convert(yarray(n),'list')); end if;
> yideal:= convert(subtract(yarray(n),A),'list');
> G:= gbasis(yideal,TA);
> return G;
> end:
```

SAMPLE COMPUTATION

```
> A:= findpoly(1,1,11,X);
> nf:=newfun(O,A);
> invert(A,beta);
> B:=invert(arraymodG(A),gamma);
> ArrayNumElems(Array(B));
> solve(B[15]=0,w[2]);
```

Bibliography

- [1] V. I. Arnol'd. *The theory of singularities and its applications*. Lezioni Fermiane. [Fermi Lectures]. Accademia Nazionale dei Lincei, Rome, 1991.
- [2] V. I. Arnol'd, S. M. Guseĭn-Zade, and A. N. Varchenko. *Singularities of differentiable maps. Vol. I*, volume 82 of *Monographs in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1985. The classification of critical points, caustics and wave fronts, Translated from the Russian by Ian Porteous and Mark Reynolds.
- [3] P. Aspinwall and S. Katz. Computation of superpotentials for D-branes. *hep-th/0412209*, 2004.
- [4] F. Cachazo, K. Intriligator, and C. Vafa. A large N duality via a geometric transition. *Nuclear Phys. B*, 603(1-2):3–41, 2001.
- [5] F. Cachazo, S. Katz, and C. Vafa. Geometric transitions and $\mathcal{N} = 1$ quiver theories. *hep-th/0108120*, 2001.
- [6] D. A. Cox and S. Katz. *Mirror symmetry and algebraic geometry*, volume 68 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [7] P. Di Francesco, P. Ginsparg, and J. Zinn-Justin. 2D gravity and random matrices. *Phys. Rep.*, 254(1-2):133, 1995.
- [8] R. Dijkgraaf and C. Vafa. Matrix models, topological strings, and supersymmetric gauge theories. *Nuclear Phys. B*, 644(1-2):3–20, 2002.
- [9] R. Dijkgraaf and C. Vafa. On geometry and matrix models. *Nuclear Phys. B*, 644(1-2):21–39, 2002.
- [10] R. Dijkgraaf and C. Vafa. A perturbative window into non-perturbative physics. *hep-th/0208048*, 2002.
- [11] A. H. Durfee. Fifteen characterizations of rational double points and simple critical points. *Enseign. Math. (2)*, 25(1-2):131–163, 1979.
- [12] F. Ferrari. Planar diagrams and Calabi-Yau spaces. *Adv. Theor. Math. Phys.*, 7(4):619–665, 2003.
- [13] G. González-Sprinberg and J.-L. Verdier. Points doubles rationnels et représentations de groupes. *C. R. Acad. Sci. Paris Sér. I Math.*, 293(2):111–113, 1981.

- [14] R. Gopakumar and C. Vafa. On the gauge theory/geometry correspondence. *Adv. Theor. Math. Phys.*, 3(5):1415–1443, 1999.
- [15] B. R. Greene and M. R. Plesser. Duality in Calabi-Yau moduli space. *Nuclear Phys. B*, 338(1):15–37, 1990.
- [16] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil, and E. Zaslow. *Mirror symmetry*, volume 1 of *Clay Mathematics Monographs*. American Mathematical Society, Providence, RI, 2003. With a preface by Vafa.
- [17] K. Intriligator and B. Wecht. RG fixed points and flows in SQCD with adjoints. *Nuclear Phys. B*, 677(1-2):223–272, 2004.
- [18] S. Katz. Versal deformations and superpotentials for rational curves in smooth threefolds. *math.AG/0010289*, 2000.
- [19] S. Katz and D. R. Morrison. Gorenstein threefold singularities with small resolutions via invariant theory for Weyl groups. *J. Algebraic Geom.*, 1(3):449–530, 1992.
- [20] H. B. Laufer. On \mathbf{CP}^1 as an exceptional set. In *Recent developments in several complex variables (Proc. Conf., Princeton Univ., Princeton, N. J., 1979)*, volume 100 of *Ann. of Math. Stud.*, pages 261–275. Princeton Univ. Press, Princeton, N.J., 1981.
- [21] J. M. Maldacena. The large N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998.
- [22] H. C. Pinkham. Factorization of birational maps in dimension 3. In *Singularities, Part 2 (Arcata, Calif., 1981)*, volume 40 of *Proc. Sympos. Pure Math.*, pages 343–371. Amer. Math. Soc., Providence, RI, 1983.
- [23] G. N. Tyurina. Resolution of singularities of flat deformations of rational double points. *Functional Anal. Appl.*, 4:68–73, 1970.
- [24] E. Witten. Chern-Simons gauge theory as a string theory. In *The Floer memorial volume*, volume 133 of *Progr. Math.*, pages 637–678. Birkhäuser, Basel, 1995.
- [25] E. Witten. String theory dynamics in various dimensions. *Nuclear Phys. B*, 443(1-2):85–126, 1995.
- [26] Y. Yoshino. *Cohen-Macaulay modules over Cohen-Macaulay rings*, volume 146 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1990.

Biography

Carina Curto was born on April 15, 1978, of Argentine parents. She was raised in Iowa City, IA. In 2000 she graduated from Harvard, and received an NSF graduate fellowship to support her PhD studies at Duke.