

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications -- Chemistry Department

Published Research - Department of Chemistry

1992

The eukaryotic initiation factor 2-associated 67-kDa polypeptide (p⁶⁷) plays a critical role in regulation of protein synthesis initiation in animal cells

Manas K. Ray

University of Nebraska - Lincoln

Bansidhar Datta

University of Nebraska at Lincoln

Arup Chakraborty

University of Nebraska - Lincoln

Ansuman Chattopadhyay

University of Nebraska - Lincoln

Susanne Meza-Keuthen

University of Nebraska - Lincoln

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/chemfacpub>

 Part of the [Chemistry Commons](#)

Ray, Manas K.; Datta, Bansidhar; Chakraborty, Arup; Chattopadhyay, Ansuman; Meza-Keuthen, Susanne; and Gupta, Naba K., "The eukaryotic initiation factor 2-associated 67-kDa polypeptide (p⁶⁷) plays a critical role in regulation of protein synthesis initiation in animal cells" (1992). *Faculty Publications -- Chemistry Department*. 47.

<https://digitalcommons.unl.edu/chemfacpub/47>

This Article is brought to you for free and open access by the Published Research - Department of Chemistry at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications -- Chemistry Department by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Manas K. Ray, Bansidhar Datta, Arup Charkraborty, Ansuman Chattopadhyay, Susanne Meza-Keuthen, and Naba K. Gupta

The eukaryotic initiation factor 2-associated 67-kDa polypeptide (p⁶⁷) plays a critical role in regulation of protein synthesis initiation in animal cells

(translational regulation/peptide chain initiation/protein phosphorylation)

MANAS K. RAY, BANSIDHAR DATTA*, ARUP CHAKRABORTY, ANSUMAN CHATTOPADHYAY, SUSANNE MEZA-KEUTHEN, AND NABA K. GUPTA†

Department of Chemistry, University of Nebraska, Lincoln, NE 68588-0304

Communicated by Myron K. Brakke, October 18, 1991 (received for review August 1, 1991)

ABSTRACT The eukaryotic initiation factor 2 (eIF-2)-associated 67-kDa polypeptide (p⁶⁷) isolated from reticulocyte lysate protects the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation and promotes protein synthesis in the presence of active eIF-2 kinases. We have now studied the roles of p⁶⁷ and eIF-2 kinases in regulation of protein synthesis using several animal cell lysates and an animal cell line (KRC-7) in culture under various growth conditions. The results are as follows. (i) Both p⁶⁷ and eIF-2 kinase(s) are present in active forms in all animal cells under normal growth conditions and p⁶⁷ protects the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation, thus promoting protein synthesis in the presence of active eIF-2 kinases. (ii) In heme-deficient reticulocyte lysates and in serum-starved KRC-7 cells in culture, p⁶⁷ is deglycosylated and subsequently degraded. This leads to eIF-2 kinase-catalyzed eIF-2 α -subunit phosphorylation and thus to protein synthesis inhibition. (iii) Addition of a mitogen (namely, phorbol 12-myristate 13-acetate) to serum-starved KRC-7 cells in culture induces an increase of p⁶⁷ and thus increases protein synthesis. These results suggest the following conclusions. (i) Protein synthesis inhibition in a heme-deficient reticulocyte lysate is not due to the activation of an eIF-2 kinase (heme-regulated inhibitor), as is generally believed, but is due to degradation of p⁶⁷. The heme-regulated inhibitor is present in an active form and possibly in equal amounts in both heme-deficient and heme-supplemented reticulocyte lysates but cannot phosphorylate eIF-2 α subunit because of the presence of p⁶⁷. (ii) p⁶⁷ is essential for protein synthesis as it protects the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation and promotes protein synthesis in the presence of one or more active eIF-2 kinases present in all animal cells. (iii) p⁶⁷ is both degradable and inducible. Only the p⁶⁷ level correlates directly with the protein synthesis activity of the cell, indicating that p⁶⁷ is a critical factor in protein synthesis regulation in animal cells.

A widely used protein synthesis regulatory mechanism in animal cells involves phosphorylation of the α subunit of a key peptide chain initiation factor, eukaryotic initiation factor 2 (eIF-2), by one or more eIF-2 kinases, which inhibits protein synthesis. In several cases studied, these inhibitors appear to remain in an inactive form and are activated under certain physiological conditions. Numerous reports indicate that in reticulocyte lysates one inhibitor, heme-regulated inhibitor (HRI), is activated in the absence of hemin and another inhibitor, double-stranded-RNA-activated inhibitor (dsI), is activated in the presence of double-stranded RNA (for reviews, see refs. 1 and 2). In both cases the activated inhibitor specifically phosphorylates the eIF-2 α subunit and thus inactivates eIF-2 activity and inhibits protein synthesis.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

As noted earlier, this regulatory mechanism involving eIF-2 α -subunit phosphorylation is widely used in animal cells to regulate protein synthesis under various physiological conditions, which include nutritional deprivation (3, 4), heat shock (4–6), and viral infection (7–14).

We have observed that a 67-kDa polypeptide, p⁶⁷, which remains associated with reticulocyte eIF-2 through several steps of purification, protects the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation, thus promoting protein synthesis in the presence of activated eIF-2 kinases (15). p⁶⁷ is a glycoprotein and contains multiple O-linked GlcNAc residues (16). There are indications that the glycosyl residues on p⁶⁷ may be necessary to protect the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation. Based on our results, we postulated that protein synthesis in animal cells may be regulated by eIF-2 kinases and by the availability of p⁶⁷. eIF-2 kinases phosphorylate the eIF-2 α subunit and thus inhibit protein synthesis, and the increased availability of p⁶⁷ renders the eIF-2 α subunit resistant to eIF-2 kinase-catalyzed phosphorylation, which promotes protein synthesis in the presence of eIF-2 kinases. Two reports from our laboratory are in agreement with this hypothesis (17, 18).

In the present study, we have examined the roles of p⁶⁷ and the eIF-2 kinases in regulation of protein synthesis using several animal cell lysates and also an animal cell line (KRC-7) in culture under various growth conditions. We provide evidence that p⁶⁷ and one or more eIF-2 kinases are present in active forms in all animal cells under normal growth conditions. However, eIF-2 kinases cannot phosphorylate the eIF-2 α subunit because of the presence of p⁶⁷. Under certain physiological conditions, such as during heme deficiency in reticulocyte lysates and also in serum-starved animal cells in culture, p⁶⁷ is degraded and thus eIF-2 kinase(s) are able to phosphorylate the eIF-2 α subunit and inhibit protein synthesis. p⁶⁷ is both inducible and degradable. Also, p⁶⁷ levels in the cells correlate directly with the protein synthesis activity of the cells, suggesting that p⁶⁷ plays a critical role in regulation of protein synthesis.

MATERIALS AND METHODS

Most of the materials used in this study were as described (15–17). The cloned cell line KRC-7 derived from Reuber H35 rat hepatoma cells was kindly provided by John Koontz (University of Tennessee, Knoxville). A sample of dsI prepared according to Kudlicki *et al.* (19) was generously

Abbreviations: eIF-2, eukaryotic initiation factor 2; HRI, heme-regulated inhibitor; dsI, double-stranded-RNA-activated inhibitor; WGA, wheat germ agglutinin; PMA, phorbol 12-myristate 13-acetate.

*Present address: Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT 06510.

†To whom reprint requests should be addressed.

donated by Gisela Kramer (Clayton Foundation, Biochemistry Institute, Austin, TX). A 10,000 × *g* cell supernatant from rat liver and from rat brain was prepared as described by Pain *et al.* (20).

Preparation of Antibodies. Polyclonal antibodies against eIF-2 α or β subunits or against p⁶⁷ were prepared using a modification of the procedure described (15, 16). The isolated polypeptides were injected into mice. When the antibody titer in the serum was >1:1000, the mouse was injected with TG-180 sarcoma cells (obtained from American Type Culture Collection) for generation of ascites fluid. Antibodies from the ascites fluid were purified on DEAE-Affi-Gel Blue, according to the procedure supplied by the manufacturer (Bio-Rad), and were characterized by NaDodSO₄/PAGE followed by immunoblot analysis. dsI was purified from interferon- β -induced HeLa cells by the procedure of Ray *et al.* (21) and was injected into rabbit. Immune serum from the rabbit was purified on DEAE-Affi-Gel Blue and characterized as above. The monoclonal antibodies against p⁶⁷ were prepared as described (16). As reported these monoclonal antibodies specifically recognize the GlcNAc moieties on p⁶⁷ (16). All the antibodies were monospecific and reacted only with their corresponding antigens.

Cell Culture. KRC-7 cells were grown in Dulbecco's modified Eagle's medium (DMEM), containing 5% (vol/vol) fetal calf serum and 5% (vol/vol) calf serum. The experimental procedures were essentially similar to those described by Trevillyan *et al.* (22). Serum-starved cells were prepared by washing cell monolayers with Hanks' balanced salt solution followed by culturing in serum-free DMEM for an additional 3 days. The serum-starved cells were stimulated by addition of 1.5 μ M phorbol 12-myristate 13-acetate (PMA) (Sigma). For analysis, 6 × 10⁵ cells were seeded onto 100-mm tissue culture dishes containing 10 ml of medium.

Cell Lysate Preparation. Cells were harvested, washed twice with phosphate-buffered saline (PBS, GIBCO), and lysed with lysis buffer [20 mM Hepes, pH 7.5/10 mM KCl/1.5 mM Mg(OAc)₂/2 mM dithiothreitol]. Cell lysates were centrifuged at 10,000 rpm in a Beckman Microfuge B centrifuge for 10 min. Supernatants were stored at -70°C. The concentration of proteins in the lysate was measured by Bio-Rad protein assay kit using a bovine serum albumin standard.

RESULTS

Assays for eIF-2 Kinase and p⁶⁷ Activities. To assay both eIF-2 kinase(s) and p⁶⁷ activities in a mixture, we developed an experimental procedure that measures eIF-2 α -subunit phosphorylation in the mixture before and after inhibition of p⁶⁷ activity by preincubation with either p⁶⁷ antibodies or wheat germ agglutinin (WGA). As reported (16), WGA binds to the glycosyl residues on p⁶⁷ and interferes with the p⁶⁷ activity to protect the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation.

The results of this assay are shown in Fig. 1. As before, HRI efficiently phosphorylated the three-subunit eIF-2 (lane 1) and addition of p⁶⁷ protected the eIF-2 α subunit from phosphorylation (lane 2). However, the ability of p⁶⁷ to protect the eIF-2 α subunit was almost completely inhibited by preincubation of the mixture of eIF-2 and p⁶⁷ with either p⁶⁷ antibodies (lane 3) or WGA (lane 6). Under similar conditions, preincubation of the mixture of eIF-2 and p⁶⁷ with either eIF-2 α -subunit or β -subunit antibodies had no significant effect (lanes 4 and 5). None of the antibodies (p⁶⁷ or eIF-2 α or β subunits), when added alone in the absence of p⁶⁷, inhibited HRI-catalyzed phosphorylation of the eIF-2 α subunit (data not shown). We also analyzed various antibody-treated incubation mixtures for possible degradation of p⁶⁷ by a standard immunoblot




FIG. 1. Assay for eIF-2 kinases and p⁶⁷ activities. Reaction conditions for HRI-catalyzed phosphorylation of eIF-2 α subunit were as described (15). A 25- μ l reaction mixture contained 20 mM Tris-HCl (pH 7.8), 100 mM KCl, 10 μ g of bovine serum albumin, 2 mM dithiothreitol, 3 μ g of eIF-2 (fraction V), 1.6 μ g of p⁶⁷, and where indicated 10 μ g of polyclonal antibodies against p⁶⁷ or against eIF-2 α or β subunits or 10 μ g of WGA. Reaction mixtures were incubated on ice for 1 hr, then 0.02 μ g of HRI and 10 μ Ci [γ -³²P]ATP were added, and the mixtures were further incubated at 37°C for 10 min. The radioactively labeled eIF-2 α subunit was then analyzed by NaDodSO₄/PAGE followed by autoradiography. Lanes: 1, eIF-2 alone; 2, eIF-2 plus p⁶⁷; 3, eIF-2, p⁶⁷, plus p⁶⁷ antibodies; 4, eIF-2, p⁶⁷, plus eIF-2 α -subunit antibodies; 5, eIF-2, p⁶⁷, plus eIF-2 β -subunit antibodies; 6, eIF-2, p⁶⁷, plus WGA.

procedure. p⁶⁷ remained undegraded during incubation (data not shown).

We carried out similar experiments using purified dsI in place of HRI and the results were similar. p⁶⁷ inhibited dsI-catalyzed phosphorylation of eIF-2 α subunits. Also p⁶⁷ antibodies and WGA inhibited this p⁶⁷ activity, but eIF-2 α - and β -subunit antibodies had no effect.

Studies Using Animal Cell Lysates. We used the above assay to study eIF-2 kinase(s) and p⁶⁷ activities in four cell lysates; heme-deficient rabbit reticulocyte lysates, heme-supplemented rabbit reticulocyte lysates, and also cell lysates prepared from rat liver and rat brain (Fig. 2). In the absence of p⁶⁷ antibodies, significant eIF-2 α -subunit phosphorylation was observed only with heme-deficient reticulocyte lysate. In contrast, the heme-supplemented reticulocyte lysate and also the lysates from liver and brain showed very little or no phosphorylation. However, when the cell lysates were preincubated with p⁶⁷ antibodies significant phosphorylation of the eIF-2 α subunit was observed with all of the cell lysates. The extent of phosphorylation, with both heme-deficient and heme-supplemented reticulocyte lysates, was comparable. However, it should be noted that, although significant changes in the phosphorylation pattern in all of the cell lysates con-




FIG. 2. Analysis of eIF-2 kinases and p⁶⁷ activities in various cell lysates. Heme-deficient and heme-supplemented reticulocyte lysates and also cell lysates from rat brain and rat liver were used. Heme-deficient and heme-supplemented reticulocyte lysates prepared as described (15) were preincubated at 37°C for 10 min before use. Approximately 20 μ l of reticulocyte lysate or 120 μ g of protein in brain or liver extract was used. eIF-2 α -subunit phosphorylation by endogenous eIF-2 kinase(s) in the presence or absence of p⁶⁷ antibodies was carried out as described in Fig. 1 and phosphorylated proteins were analyzed by NaDodSO₄/PAGE and autoradiography. Lanes: 1, heme-deficient reticulocyte lysate; 2, heme-deficient reticulocyte lysate-preincubated with p⁶⁷ antibodies; 3, heme-supplemented reticulocyte lysate; 4, heme-supplemented reticulocyte lysate preincubated with p⁶⁷ antibodies; 5, lysate from rat brain; 6, lysate from rat brain preincubated with p⁶⁷ antibodies; 7, lysate from rat liver; 8, lysate from rat liver preincubated with p⁶⁷ antibodies.

taining p⁶⁷ antibodies was observed with the eIF-2 α subunit, one polypeptide of \approx 200 kDa or more polypeptides present in reticulocyte lysates and in rat brain and rat liver lysates showed a similar change in phosphorylation pattern. The significance of this observation will be discussed later.

These results, however, show that both eIF-2 kinase(s) and p⁶⁷ are present in active forms in all of the cell lysates and p⁶⁷ inhibits eIF-2 α -subunit phosphorylation by active eIF-2 kinases. Removal of p⁶⁷ by p⁶⁷ antibodies facilitated eIF-2 kinase-catalyzed phosphorylation of the eIF-2 α subunit.

The results presented in Fig. 3 show that p⁶⁷ is indeed present in heme-supplemented reticulocyte lysate; and this polypeptide is partially deglycosylated in heme-deficient reticulocyte lysate. For these experiments the lysates were preincubated with p⁶⁷ monoclonal (Fig. 3A) or polyclonal (Fig. 3B) antibodies and the antigen-antibody complexes were precipitated with protein A-agarose (23). The precipitates were then assayed on an immunoblot using p⁶⁷ polyclonal antibodies. p⁶⁷ is present in heme-supplemented reticulocyte lysate (Fig. 3 A and B, lanes 2) and also in heme-deficient reticulocyte lysate (Fig. 3 A and B, lanes 1). However, the level of this polypeptide in heme-deficient reticulocyte lysate was significantly lower. In experiments described in Fig. 3A, p⁶⁷ present in the lysates was immunoprecipitated using p⁶⁷ monoclonal antibodies, which reacted only with glycosylated p⁶⁷. The p⁶⁷ level in heme-deficient lysate (lane 1) was $<20\%$ of that present in heme-supplemented lysate (lane 2), indicating extensive deglycosylation of p⁶⁷ in heme-deficient lysate.

These results thus provide evidence that p⁶⁷ is present in both heme-supplemented and heme-deficient reticulocyte lysates. In heme-deficient lysate, p⁶⁷ is partially deglycosylated. This allows eIF-2 kinases to phosphorylate eIF-2 and thus inhibit protein synthesis. Using a similar immunoblot experiment, we also observed that p⁶⁷ is present in glycosy-

lated form in both rat liver and rat brain extracts (data not shown).

Studies Using a Tumor Hepatoma Cell Line (KRC-7). We used a tumor hepatoma cell line (KRC-7) and analyzed the roles of p⁶⁷ and eIF-2 kinase(s) in regulating protein synthesis under various growth conditions. We grew the cells to confluency, then added serum-depleted medium for 3–4 days, and stimulated the serum-starved cells with a mitogen, PMA. We measured the protein synthesis rate, the levels of various polypeptides, and also activation states of eIF-2 kinase(s) at various times during confluency, after serum starvation, and after PMA addition. Protein synthesis rate was measured by incorporation of [³⁵S]methionine into cellular proteins as described (17). The protein synthesis rate was maximum in confluent cells and was reduced to 30% of the maximum level after 100 hr of serum starvation. Upon PMA addition the serum-starved cells regained 70% of the original protein synthesis activity of the confluent cells within 4 hr.

Levels of Various Polypeptides. Immunoblot analysis was used to measure levels of various polypeptides (p⁶⁷, dsI, and eIF-2 β and α subunits) at various times, after serum starvation and after PMA addition (Fig. 4). p⁶⁷ levels were measured using both monoclonal and polyclonal antibodies. dsI and eIF-2 β - and α -subunit levels were measured using polyclonal antibodies.

As shown in Fig. 4, levels of eIF-2 β and α subunits and, more importantly, the level of dsI remained essentially unchanged in confluent (–72 hr), serum-starved, and PMA-stimulated cells. On the other hand, p⁶⁷ levels changed dramatically; p⁶⁷ was prominent in confluent cells, disappeared rapidly after serum starvation, and became prominent after PMA addition. A significant observation is that p⁶⁷ level decreased faster after serum depletion and appeared more slowly after PMA addition when measured using monoclonal antibodies than when measured using polyclonal antibodies. These results, in agreement with our observation with heme-deficient reticulocyte lysates (Fig. 3), suggest that the first step after serum depletion may be deglycosylation of p⁶⁷ and that deglycosylated p⁶⁷ is subsequently degraded.

eIF-2 Kinase Activity. The results presented in Fig. 4 show that the dsI polypeptide level remains essentially unchanged under various growth conditions. However, the eIF-2 kinases, such as HRI and dsI, may remain in either active or inactive forms. We have, therefore, analyzed eIF-2 kinase




FIG. 3. Analysis of p⁶⁷ in heme-deficient and heme-supplemented reticulocyte lysates. Reticulocyte lysates (40 μ l) without or with hemin (25 μ l) were incubated for 10 min at 37°C. (A) The lysates were mixed with 25 μ g of p⁶⁷ monoclonal antibodies. (B) The lysates were mixed with 25 μ g of p⁶⁷ polyclonal antibodies. The reaction mixtures were incubated at 4°C overnight and antigen-antibody complexes were precipitated by adding 30 μ l of protein A-agarose (GIBCO) followed by gentle tapping of the suspension for 1 hr at room temperature. The precipitates were then washed three times with 1 \times PBS containing 0.05% Nonidet P-40. The polypeptides in the precipitates were then separated by NaDodSO₄/PAGE on 15% gels and subsequently transferred electrophoretically to a nitrocellulose sheet. Lanes were excised from the blot and incubated with p⁶⁷ antibodies as described by Bio-Rad (Bio-Rad immunoblot assay kit). The antigen-antibody complexes were then detected using goat anti-mouse IgG conjugated to horseradish peroxidase (Bio-Rad). (A) Lanes: 1, heme-deficient lysate; 2, heme-supplemented lysate. (B) Lanes: 1, heme-deficient lysate; 2, heme-supplemented lysate. The lower molecular mass bands represent the IgG heavy chain.




FIG. 4. Immunoblot analysis of p⁶⁷, dsI, eIF-2 β subunit (p⁵⁴), and eIF-2 α subunit (p³⁸) in tumor hepatoma cells (KRC-7) that are confluent, quiescent, or PMA-induced. Cells grown to confluency (lane –72 hr) were transferred to serum-depleted medium and after 72 hr (lane 0 hr) were treated with PMA. Approximately 120 μ g of protein in cell lysates was analyzed on an immunoblot using p⁶⁷ monoclonal antibodies (row a), p⁶⁷ polyclonal antibodies (row b), dsI polyclonal antibodies (row c), eIF-2 β -subunit polyclonal antibodies (row d), and eIF-2 α -subunit polyclonal antibodies (row e). The immunoblot procedure for detection of p⁶⁷ and eIF-2 α and eIF-2 β subunits was the same as described in Fig. 3. The immunoblot procedure for detection of dsI was the same except that goat anti-rabbit IgG conjugated to horseradish peroxidase was used to detect dsI rabbit antibodies.

activities by analyzing eIF-2 α -subunit phosphorylation of endogenous eIF-2 using cell extracts at various times after the assay as described in Fig. 1. eIF-2 α -subunit phosphorylation was low with extracts from confluent and PMA-induced cells but was significantly increased when a similar extract from quiescent cell was used (Fig. 5A). However, as described in Fig. 1, when the same cell extracts were first preincubated with p^{67} polyclonal antibodies to remove endogenous p^{67} (Fig. 5B) or with WGA to inhibit the eIF-2- p^{67} interaction (Fig. 5C), the extent of eIF-2 α -subunit phosphorylation using all the cell extracts was significantly increased to approximately the same level.

DISCUSSION

Data presented in this report provide evidence that all animal cells contain one or more eIF-2 kinases in the active form and the activities of these eIF-2 kinases may or may not be evident, depending on the variable presence of p^{67} . p^{67} protects the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation and thus promotes protein synthesis in the presence of active eIF-2 kinase(s).

In separate experiments we observed that p^{67} specifically protects the eIF-2 α subunit from both HRI- and dsI-catalyzed phosphorylation but does not inhibit dsI-catalyzed phosphorylation of histones or casein kinase catalyzed phosphorylation of the eIF-2 β subunit (data not shown). However, the results presented in Fig. 2 indicate that p^{67} also inhibits phosphorylation of one or more additional proteins (30 and 110 kDa) present in reticulocyte lysates and also in rat brain and rat liver lysates. It may be that under *in vitro* conditions, the glycosyl residues of p^{67} may bind to these additional proteins and thus inhibit their phosphorylation. On the other hand, the possibility that p^{67} may inhibit phosphorylation of a select group of closely related proteins cannot be ruled out. However, abundant evidence reported previously and also presented in this paper suggests that a major function of p^{67} is to protect the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation and thus to promote protein synthesis in the presence of active eIF-2 kinases.




FIG. 5. eIF-2 α -subunit phosphorylation using a lysate from KRC-7 cells grown under various conditions. (A) Approximately 60 μ g of protein in lysates from confluent (lanes -72 hr), serum-starved (lanes -64 hr through 0 hr), and PMA-induced (lanes 1/2, 1, 2, and 4 hr) KRC-7 cells was used and phosphorylation of the endogenous eIF-2 α subunit by endogenous eIF-2 kinases was carried out as described in Fig. 1. (B) The cell lysates were preincubated with p^{67} polyclonal antibodies. (C) The lysates were preincubated with WGA and were then phosphorylated.

Two widely studied eIF-2 kinases are HRI and dsI. In heme-deficient reticulocyte lysates, only HRI is active, as dsI requires double-stranded RNA and ATP for activation. In this report, we have provided evidence that, in both heme-supplemented and heme-deficient lysates, HRI is present in active form and possibly in equal amounts. In heme-supplemented lysate, this HRI activity is not evident due to the presence of p^{67} , whereas in heme-deficient lysates p^{67} appears to be degraded, thus allowing HRI to actively phosphorylate eIF-2 α subunit. Removal of p^{67} from both heme-supplemented and heme-deficient reticulocyte lysates with p^{67} antibodies resulted in similar extents of eIF-2 α -subunit phosphorylation. This indicates that the eIF-2 kinase in active form may be present at the same level in both systems. These results suggest that eIF-2 α -subunit phosphorylation, observed in heme-deficient reticulocyte lysate, is not due to activation of HRI, as is widely believed, but rather to degradation of p^{67} . We also provide evidence that p^{67} degradation involves partial deglycosylation in heme-deficient lysates. It may, therefore, be assumed that hemin, present in heme-supplemented lysates, prevents p^{67} deglycosylation, possibly by inhibiting the deglycosylating enzyme(s).

Reticulocyte lysates also contain another eIF-2 kinase, dsI, in inactive form, and this eIF-2 kinase is activated in the presence of double-stranded RNA and ATP (1, 2). Activated dsI inhibits protein synthesis even in the presence of hemin and presumably in the presence of p^{67} . The mechanism of eIF-2 α -subunit phosphorylation by dsI in the presence of p^{67} is not clear. Further work will be necessary to examine p^{67} activity during dsI inhibition of protein synthesis in heme-supplemented reticulocyte lysate. It may be possible, however, that dsI is formed in relatively large excess in the presence of double-stranded RNA and ATP and at a very high concentration can phosphorylate the eIF-2 α subunit even in the presence of equimolar concentrations of p^{67} .

Like heme-supplemented reticulocyte lysates, the cell lysates prepared from rat brain, rat liver, and confluent and PMA-induced tumor hepatoma cells (KRC-7) contain both p^{67} and eIF-2 kinase(s) in active forms, and eIF-2 kinase activity becomes evident only after removal of p^{67} with p^{67} antibodies. However, these eIF-2 kinases have not been characterized. The amount of dsI polypeptide is essentially unchanged in tumor hepatoma cells under various growth conditions. This includes confluency, serum starvation, and mitogen stimulation (Fig. 4). Again, it is not clear whether this dsI is in an active form. Our results show that these cells contain at least one eIF-2 kinase in an active form; however, this eIF-2 kinase has not been identified as yet as dsI. As noted, dsI exists in an inactive form in the cells and requires double-stranded RNA and ATP for activation. The cell extracts used in this study were not treated with double-stranded RNA and ATP. Also, our present work suggests that the HRI may be a normal cell constituent, as this inhibitor is present in an active form and in approximately equal amounts in both heme-supplemented and heme-deficient reticulocyte lysates. Possibly a HRI-like eIF-2 kinase is present in an active form in all animal cells and dsI activation serves specialized functions under certain physiological conditions. As postulated earlier, this inhibitor (dsI) may be formed in large excess upon activation under certain physiological conditions, such as virus infections, and may inhibit protein synthesis in the presence of normal p^{67} concentrations.

The roles and requirements for p^{67} in KRC-7 cells under various growth conditions are clearly evident. This polypeptide is present in high concentrations in confluent cells and disappears rapidly from the cells after serum depletion with an accompanying decrease in the protein synthesis rate in the cells. Also, this polypeptide reappears soon after PMA addition to the serum-starved resting cells as the cells regain

protein synthesis activity. Thus, there is a clear correlation between the p⁶⁷ level and the protein synthesis activity of the cells, indicating a direct involvement of p⁶⁷ in protein synthesis. Furthermore, a comparison of the p⁶⁷ levels studied using monoclonal (Fig. 4, row a) and polyclonal (Fig. 4, row b) antibodies reveals (in agreement with our previous observation with the heme-deficient reticulocyte lysates, Fig. 3) that the initial event in p⁶⁷ degradation involves deglycosylation of p⁶⁷ and deglycosylated p⁶⁷ is unstable and is subsequently degraded. These results, therefore, establish the following facts. (i) p⁶⁷ is necessary to protect the eIF-2 α subunit from eIF-2 kinase-catalyzed phosphorylation and thus to promote protein synthesis in the presence of active eIF-2 kinase(s). (ii) At least one eIF-2 kinase is present in all of the cells in active form under various growth conditions. (iii) p⁶⁷ is degraded rapidly in serum-depleted cells thus allowing eIF-2 kinase(s) to phosphorylate eIF-2 and inhibit protein synthesis. (iv) The initial event of p⁶⁷ degradation involves deglycosylation of the p⁶⁷ polypeptide.

An important aspect of p⁶⁷ regulation is that this is the only polypeptide presently studied that is both easily degradable and inducible. The levels of this polypeptide correlate directly to the protein synthesis activities of the cells. As evident in this study, the initial event for degradation involves deglycosylation of p⁶⁷. The deglycosylated p⁶⁷ is unstable and subsequently degraded. Again, addition of a mitogen, such as PMA, induces increased appearance of p⁶⁷, presumably by transcription activation of the p⁶⁷ gene. Thus, p⁶⁷ level in the cell is regulated both transcriptionally and posttranscriptionally. This factor is essential for protein synthesis and thus plays a critical role in protein synthesis regulation.

This investigation was supported by a Nebraska State Grant for Cancer and Smoking Disease (91-21) and National Institutes of Health Grant GM 22079.

1. Gupta, N. K., Ahmad, M. F., Chakrabarti, D. & Nasrin, N.

- (1987) in *Translation Regulation in Gene Expression*, ed. Ilan, J. (Plenum, New York), pp. 287–334.
2. Hershey, J. W. B. (1991) *Annu. Rev. Biochem.* **60**, 717–755.
 3. Duncan, R. & Hershey, J. W. B. (1985) *J. Biol. Chem.* **260**, 5493–5497.
 4. Scorsone, K. A., Panniers, R., Rowlands, A. G. & Henshaw, E. (1987) *J. Biol. Chem.* **262**, 14538–14543.
 5. Duncan, R. & Hershey, J. W. B. (1984) *J. Biol. Chem.* **259**, 11882–11884.
 6. Benedetti, A. D. & Baglioni, C. (1987) *J. Biol. Chem.* **262**, 338–342.
 7. Reichel, P. A., Merrick, W. C., Siekierka, J. & Mathews, M. B. (1985) *Nature (London)* **313**, 196–200.
 8. Schneider, R. J., Safer, B., Munemitsu, S. M., Samuel, C. E. & Shenk, T. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 4321–4325.
 9. Siekierka, J., Mariano, T., Reichel, P. A. & Mathews, M. B. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 1259–1263.
 10. Whitaker-Dowling, P. A. & Younger, J. S. (1984) *Virology* **137**, 171–181.
 11. Rice, A. P. & Kerr, I. M. (1984) *J. Virol.* **50**, 229–236.
 12. Akkaraju, G. R., Whitaker-Dowling, P., Younger, J. S. & Jagus, R. (1989) *J. Biol. Chem.* **264**, 10321–10325.
 13. Ransoni, L. J. & Dasgupta, A. (1987) *J. Virol.* **61**, 1781–1787.
 14. Ransoni, L. J. & Dasgupta, A. (1988) *J. Virol.* **62**, 3551–3588.
 15. Datta, B., Chakrabarti, D., Roy, A. L. & Gupta, N. K. (1988) *Proc. Natl. Acad. Sci. USA* **85**, 3324–3328.
 16. Datta, B., Ray, M. K., Chakrabarti, D., Wylie, D. & Gupta, N. K. (1989) *J. Biol. Chem.* **264**, 20620–20624.
 17. Datta, B., Ray, M. K., Chakrabarti, D. & Gupta, N. K. (1988) *Indian J. Biochem.* **25**, 478–482.
 18. Gupta, N. K., Datta, B., Roy, A. L. & Ray, M. K. (1990) in *Post Transcriptional Control of Gene Expression*, eds. McCarthy, J. E. G. & Tuite, M. F. (Springer, Berlin), pp. 511–520.
 19. Kudlicki, W., Fullilove, S., Read, R., Kramer, G. & Hardesty, B. (1987) *J. Biol. Chem.* **262**, 9695–9701.
 20. Pain, V. M., Lewis, J. A., Hovos, P., Henshaw, C. E. & Clemens, J. M. (1980) *J. Biol. Chem.* **255**, 1486–1491.
 21. Ray, P., Chen, J. & London, I. M. (1988) *Proc. Natl. Acad. Sci. USA* **85**, 1427–1431.
 22. Trevillyan, J. M., Kulkarni, R. K. & Byers, C. V. (1984) *J. Biol. Chem.* **259**, 897–902.
 23. Reed, R., Griffith, J. & Maniatis, T. (1988) *Cell* **53**, 949–969.