

University of Nebraska - Lincoln

## DigitalCommons@University of Nebraska - Lincoln

---

Kenneth Bloom Publications

Research Papers in Physics and Astronomy

---

June 2008

### Search for scalar top quarks in the acoplanar charm jets and missing transverse energy final state in $pp$ collisions at $\sqrt{s} = 1.96$ TeV

V. M. Abazov

*Joint Institute for Nuclear Research, Dubna, Russia*

Kenneth A. Bloom

*University of Nebraska-Lincoln, kbloom2@unl.edu*

Gregory Snow

*University of Nebraska-Lincoln, gsnow1@unl.edu*

D0 Collaboration

Follow this and additional works at: <https://digitalcommons.unl.edu/physicsbloom>

 Part of the [Physics Commons](#)

---

Abazov, V. M.; Bloom, Kenneth A.; Snow, Gregory; and Collaboration, D0, "Search for scalar top quarks in the acoplanar charm jets and missing transverse energy final state in  $pp$  collisions at  $\sqrt{s} = 1.96$  TeV" (2008). *Kenneth Bloom Publications*. 244.

<https://digitalcommons.unl.edu/physicsbloom/244>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



## Search for scalar top quarks in the acoplanar charm jets and missing transverse energy final state in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

DØ Collaboration

V.M. Abazov<sup>aj</sup>, B. Abbott<sup>bw</sup>, M. Abolins<sup>bm</sup>, B.S. Acharya<sup>ac</sup>, M. Adams<sup>ay</sup>, T. Adams<sup>aw</sup>, E. Aguilo<sup>f</sup>, S.H. Ahn<sup>ae</sup>, M. Ahsan<sup>bg</sup>, G.D. Alexeev<sup>aj</sup>, G. Alkhazov<sup>an</sup>, A. Alton<sup>bl,1</sup>, G. Alverson<sup>bk</sup>, G.A. Alves<sup>b</sup>, M. Anastasoie<sup>ai</sup>, L.S. Ancu<sup>ai</sup>, T. Andeen<sup>ba</sup>, S. Anderson<sup>as</sup>, B. Andrieu<sup>q</sup>, M.S. Anzels<sup>ba</sup>, M. Aoki<sup>ax</sup>, Y. Arnoud<sup>n</sup>, M. Arov<sup>bh</sup>, M. Arthaud<sup>r</sup>, A. Askew<sup>aw</sup>, B. Åsman<sup>ao</sup>, A.C.S. Assis Jesus<sup>c</sup>, O. Atramentov<sup>aw</sup>, C. Avila<sup>h</sup>, C. Ay<sup>x</sup>, F. Badaud<sup>m</sup>, A. Baden<sup>bi</sup>, L. Bagby<sup>ax</sup>, B. Baldin<sup>ax</sup>, D.V. Bandurin<sup>bg</sup>, P. Banerjee<sup>ac</sup>, S. Banerjee<sup>ac</sup>, E. Barberis<sup>bk</sup>, A.-F. Barfuss<sup>o</sup>, P. Bargassa<sup>cb</sup>, P. Baringer<sup>bf</sup>, J. Barreto<sup>b</sup>, J.F. Bartlett<sup>ax</sup>, U. Bassler<sup>r</sup>, D. Bauer<sup>aq</sup>, S. Beale<sup>f</sup>, A. Bean<sup>bf</sup>, M. Begalli<sup>c</sup>, M. Begel<sup>bu</sup>, C. Belanger-Champagne<sup>ao</sup>, L. Bellantoni<sup>ax</sup>, A. Bellavance<sup>ax</sup>, J.A. Benitez<sup>bm</sup>, S.B. Beri<sup>aa</sup>, G. Bernardi<sup>q</sup>, R. Bernhard<sup>w</sup>, I. Bertram<sup>ap</sup>, M. Besançon<sup>r</sup>, R. Beuselinck<sup>aq</sup>, V.A. Bezzubov<sup>am</sup>, P.C. Bhat<sup>ax</sup>, V. Bhatnagar<sup>aa</sup>, C. Biscarat<sup>t</sup>, G. Blazey<sup>az</sup>, F. Blekman<sup>aq</sup>, S. Blessing<sup>aw</sup>, D. Bloch<sup>s</sup>, K. Bloom<sup>bo</sup>, A. Boehnlein<sup>ax</sup>, D. Boline<sup>bj</sup>, T.A. Bolton<sup>bg</sup>, G. Borissov<sup>ap</sup>, T. Bose<sup>by</sup>, A. Brandt<sup>bz</sup>, R. Brock<sup>bm</sup>, G. Brooijmans<sup>br</sup>, A. Bross<sup>ax</sup>, D. Brown<sup>cc</sup>, N.J. Buchanan<sup>aw</sup>, D. Buchholz<sup>ba</sup>, M. Buehler<sup>cc</sup>, V. Buescher<sup>v</sup>, V. Bunichev<sup>al</sup>, S. Burdin<sup>ap,2</sup>, S. Burke<sup>as</sup>, T.H. Burnett<sup>cd</sup>, C.P. Buszello<sup>aq</sup>, J.M. Butler<sup>bj</sup>, P. Calfayan<sup>y</sup>, S. Calvet<sup>p</sup>, J. Cammin<sup>bs</sup>, W. Carvalho<sup>c</sup>, B.C.K. Casey<sup>ax</sup>, H. Castilla-Valdez<sup>ag</sup>, S. Chakrabarti<sup>r</sup>, D. Chakraborty<sup>az</sup>, K. Chan<sup>f</sup>, K.M. Chan<sup>bc</sup>, A. Chandra<sup>av</sup>, F. Charles<sup>s,x</sup>, E. Cheu<sup>as</sup>, F. Chevallier<sup>n</sup>, D.K. Cho<sup>bj</sup>, S. Choi<sup>af</sup>, B. Choudhary<sup>ab</sup>, L. Christofek<sup>by</sup>, T. Christoudias<sup>aq</sup>, S. Cihangir<sup>ax</sup>, D. Claes<sup>bo</sup>, Y. Coadou<sup>f</sup>, M. Cooke<sup>cb</sup>, W.E. Cooper<sup>ax</sup>, M. Corcoran<sup>cb</sup>, F. Couderc<sup>r</sup>, M.-C. Cousinou<sup>o</sup>, S. Crépe-Renaudin<sup>n</sup>, D. Cutts<sup>by</sup>, M. Ćwiok<sup>ad</sup>, H. da Motta<sup>b</sup>, A. Das<sup>as</sup>, G. Davies<sup>aq</sup>, K. De<sup>bz</sup>, S.J. de Jong<sup>ai</sup>, E. De La Cruz-Burelo<sup>bl</sup>, C. De Oliveira Martins<sup>c</sup>, J.D. Degenhardt<sup>bl</sup>, F. Déliot<sup>r</sup>, M. Demarteau<sup>ax</sup>, R. Demina<sup>bs</sup>, D. Denisov<sup>ax</sup>, S.P. Denisov<sup>am</sup>, S. Desai<sup>ax</sup>, H.T. Diehl<sup>ax</sup>, M. Diesburg<sup>ax</sup>, A. Dominguez<sup>bo</sup>, H. Dong<sup>bt</sup>, L.V. Dudko<sup>al</sup>, L. Duflost<sup>p</sup>, S.R. Dugad<sup>ac</sup>, D. Duggan<sup>aw</sup>, A. Duperrin<sup>o</sup>, J. Dyer<sup>bm</sup>, A. Dyshkant<sup>az</sup>, M. Eads<sup>bo</sup>, D. Edmunds<sup>bm</sup>, J. Ellison<sup>av</sup>, V.D. Elvira<sup>ax</sup>, Y. Enari<sup>by</sup>, S. Eno<sup>bi</sup>, P. Ermolov<sup>al</sup>, H. Evans<sup>bb</sup>, A. Evdokimov<sup>bu</sup>, V.N. Evdokimov<sup>am</sup>, A.V. Ferapontov<sup>bg</sup>, T. Ferbel<sup>bs</sup>, F. Fiedler<sup>x</sup>, F. Filthaut<sup>ai</sup>, W. Fisher<sup>ax</sup>, H.E. Fisk<sup>ax</sup>, M. Fortner<sup>az</sup>, H. Fox<sup>ap</sup>, S. Fu<sup>ax</sup>, S. Fuess<sup>ax</sup>, T. Gadfort<sup>br</sup>, C.F. Galea<sup>ai</sup>, E. Gallas<sup>ax</sup>, C. Garcia<sup>bs</sup>, A. Garcia-Bellido<sup>cd</sup>, V. Gavrillov<sup>ak</sup>, P. Gay<sup>m</sup>, W. Geist<sup>s</sup>, D. Gelé<sup>s</sup>, C.E. Gerber<sup>ay</sup>, Y. Gershtein<sup>aw</sup>, D. Gillberg<sup>f</sup>, G. Ginter<sup>bs</sup>, N. Gollub<sup>ao</sup>, B. Gómez<sup>h</sup>, A. Goussiou<sup>cd</sup>, P.D. Grannis<sup>bt</sup>, H. Greenlee<sup>ax</sup>, Z.D. Greenwood<sup>bh</sup>, E.M. Gregores<sup>d</sup>, G. Grenier<sup>t</sup>, Ph. Gris<sup>m</sup>, J.-F. Grivaz<sup>p</sup>, A. Grohsjean<sup>y</sup>, S. Grünendahl<sup>ax</sup>, M.W. Grünewald<sup>ad</sup>, F. Guo<sup>bt</sup>, J. Guo<sup>bt</sup>, G. Gutierrez<sup>ax</sup>, P. Gutierrez<sup>bw</sup>, A. Haas<sup>br</sup>, N.J. Hadley<sup>bi</sup>, P. Haefner<sup>y</sup>, S. Hagopian<sup>aw</sup>, J. Haley<sup>bp</sup>, I. Hall<sup>bm</sup>, R.E. Hall<sup>au</sup>, L. Han<sup>g</sup>, K. Harder<sup>ar</sup>, A. Harel<sup>bs</sup>, R. Harrington<sup>bk</sup>, J.M. Hauptman<sup>be</sup>, R. Hauser<sup>bm</sup>, J. Hays<sup>aq</sup>, T. Hebbeker<sup>u</sup>, D. Hedin<sup>az</sup>, J.G. Hegeman<sup>ah</sup>, J.M. Heinmiller<sup>ay</sup>, A.P. Heinson<sup>av</sup>, U. Heintz<sup>bj</sup>, C. Hensel<sup>bf</sup>, K. Herner<sup>bt</sup>, G. Hesketh<sup>bk</sup>, M.D. Hildreth<sup>bc</sup>, R. Hirosky<sup>cc</sup>, J.D. Hobbs<sup>bt</sup>, B. Hoeneisen<sup>l</sup>, H. Hoeth<sup>z</sup>, M. Hohlfeld<sup>v</sup>, S.J. Hong<sup>ae</sup>, S. Hossain<sup>bw</sup>, P. Houben<sup>ah</sup>, Y. Hu<sup>bt</sup>, Z. Hubacek<sup>j</sup>, V. Hynek<sup>i</sup>, I. Iashvili<sup>bq</sup>, R. Illingworth<sup>ax</sup>, A.S. Ito<sup>ax</sup>, S. Jabeen<sup>bj</sup>, M. Jaffré<sup>p</sup>, S. Jain<sup>bw</sup>, K. Jakobs<sup>w</sup>, C. Jarvis<sup>bi</sup>, R. Jesik<sup>aq</sup>, K. Johns<sup>as</sup>, C. Johnson<sup>br</sup>, M. Johnson<sup>ax</sup>, A. Jonckheere<sup>ax</sup>, P. Jonsson<sup>aq</sup>, A. Juste<sup>ax</sup>, E. Kajfasz<sup>o</sup>, A.M. Kalinin<sup>aj</sup>, J.M. Kalk<sup>bh</sup>, S. Kappler<sup>u</sup>, D. Karmanov<sup>al</sup>, P.A. Kasper<sup>ax</sup>, I. Katsanos<sup>br</sup>, D. Kau<sup>aw</sup>, V. Kaushik<sup>bz</sup>, R. Kehoe<sup>ca</sup>, S. Kermiche<sup>o</sup>, N. Khalatyan<sup>ax</sup>, A. Khanov<sup>bx</sup>, A. Kharchilava<sup>bq</sup>, Y.M. Kharzhev<sup>aj</sup>, D. Khatidze<sup>br</sup>, T.J. Kim<sup>ae</sup>, M.H. Kirby<sup>ba</sup>, M. Kirsch<sup>u</sup>, B. Klima<sup>ax</sup>, J.M. Kohli<sup>aa</sup>, J.-P. Konrath<sup>w</sup>, V.M. Korablev<sup>am</sup>, A.V. Kozelov<sup>am</sup>, J. Kraus<sup>bm</sup>, D. Krop<sup>bb</sup>, T. Kuhl<sup>x</sup>, A. Kumar<sup>bq</sup>, A. Kupco<sup>k</sup>, T. Kurča<sup>t</sup>, J. Kvita<sup>i</sup>, F. Lacroix<sup>m</sup>, D. Lam<sup>bc</sup>, S. Lammers<sup>br</sup>, G. Landsberg<sup>by</sup>, P. Lebrun<sup>t</sup>, W.M. Lee<sup>ax</sup>, A. Leflat<sup>al</sup>, J. Lellouch<sup>q</sup>, J. Leveque<sup>as</sup>, J. Li<sup>bz</sup>, L. Li<sup>av</sup>, Q.Z. Li<sup>ax</sup>, S.M. Lietti<sup>e</sup>, J.G.R. Lima<sup>az</sup>, D. Lincoln<sup>ax</sup>, J. Linnemann<sup>bm</sup>, V.V. Lipaev<sup>am</sup>, R. Lipton<sup>ax</sup>, Y. Liu<sup>g</sup>, Z. Liu<sup>f</sup>, A. Lobodenko<sup>an</sup>, M. Lokajicek<sup>k</sup>, P. Love<sup>ap</sup>, H.J. Lubatti<sup>cd</sup>, R. Luna<sup>c</sup>, A.L. Lyon<sup>ax</sup>, A.K.A. Maciel<sup>b</sup>, D. Mackin<sup>cb</sup>, R.J. Madaras<sup>at</sup>, P. Mättig<sup>z</sup>, C. Magass<sup>u</sup>, A. Magerkurth<sup>bl</sup>, P.K. Mal<sup>cd</sup>, H.B. Malbouisson<sup>c</sup>, S. Malik<sup>bo</sup>, V.L. Malyshev<sup>aj</sup>, H.S. Mao<sup>ax</sup>, Y. Maravin<sup>bg</sup>, B. Martin<sup>n</sup>, R. McCarthy<sup>bt</sup>, A. Melnitchouk<sup>bn</sup>, L. Mendoza<sup>h</sup>, P.G. Mercadante<sup>e</sup>

M. Merkin<sup>al</sup>, K.W. Merritt<sup>ax</sup>, A. Meyer<sup>u</sup>, J. Meyer<sup>v</sup>, T. Millet<sup>t</sup>, J. Mitrevski<sup>br</sup>, J. Molina<sup>c</sup>, R.K. Mommsen<sup>ar</sup>, N.K. Mondal<sup>ac</sup>, R.W. Moore<sup>f</sup>, T. Moulik<sup>bf</sup>, G.S. Muanza<sup>t</sup>, M. Mulders<sup>ax</sup>, M. Mulhearn<sup>br</sup>, O. Mundal<sup>v</sup>, L. Mundim<sup>c</sup>, E. Nagy<sup>o</sup>, M. Naimuddin<sup>ax</sup>, M. Narain<sup>by</sup>, N.A. Naumann<sup>ai</sup>, H.A. Neal<sup>bl</sup>, J.P. Negret<sup>h</sup>, P. Neustroev<sup>an</sup>, H. Nilsen<sup>w</sup>, H. Nogima<sup>c</sup>, S.F. Novaes<sup>e</sup>, T. Nunnemann<sup>y</sup>, V. O'Dell<sup>ax</sup>, D.C. O'Neil<sup>f</sup>, G. Obrant<sup>an</sup>, C. Ochando<sup>p</sup>, D. Onoprienko<sup>bg</sup>, N. Oshima<sup>ax</sup>, N. Osman<sup>aq</sup>, J. Osta<sup>bc</sup>, R. Otec<sup>j</sup>, G.J. Otero y Garzón<sup>ax</sup>, M. Owen<sup>ar</sup>, P. Padley<sup>cb</sup>, M. Pangilinan<sup>by</sup>, N. Parashar<sup>bd</sup>, S.-J. Park<sup>bs</sup>, S.K. Park<sup>ae</sup>, J. Parsons<sup>br</sup>, R. Partridge<sup>by</sup>, N. Parua<sup>bb</sup>, A. Patwa<sup>bu</sup>, G. Pawloski<sup>cb</sup>, B. Penning<sup>w</sup>, M. Perfilov<sup>al</sup>, K. Peters<sup>ar</sup>, Y. Peters<sup>z</sup>, P. Pétroff<sup>p</sup>, M. Petteni<sup>aq</sup>, R. Piegaia<sup>a</sup>, J. Piper<sup>bm</sup>, M.-A. Pleier<sup>v</sup>, P.L.M. Podesta-Lerma<sup>ag,3</sup>, V.M. Podstavkov<sup>ax</sup>, Y. Pogorelov<sup>bc</sup>, M.-E. Pol<sup>b</sup>, P. Polozov<sup>ak</sup>, B.G. Pope<sup>bm</sup>, A.V. Popov<sup>am</sup>, C. Potter<sup>f</sup>, W.L. Prado da Silva<sup>c</sup>, H.B. Prosper<sup>aw</sup>, S. Protopopescu<sup>bu</sup>, J. Qian<sup>bl</sup>, A. Quadt<sup>v,4</sup>, B. Quinn<sup>bn</sup>, A. Rakitine<sup>ap</sup>, M.S. Rangel<sup>b</sup>, K. Ranjan<sup>ab</sup>, P.N. Ratoff<sup>ap</sup>, P. Renkel<sup>ca</sup>, S. Reucroft<sup>bk</sup>, P. Rich<sup>ar</sup>, J. Rieger<sup>bb</sup>, M. Rijssenbeek<sup>bt</sup>, I. Ripp-Baudot<sup>s</sup>, F. Rizatdinova<sup>bx</sup>, S. Robinson<sup>aq</sup>, R.F. Rodrigues<sup>c</sup>, M. Rominsky<sup>bw</sup>, C. Royon<sup>r</sup>, P. Rubinov<sup>ax</sup>, R. Ruchti<sup>bc</sup>, G. Safronov<sup>ak</sup>, G. Sajot<sup>n</sup>, A. Sánchez-Hernández<sup>ag</sup>, M.P. Sanders<sup>q</sup>, A. Santoro<sup>c</sup>, G. Savage<sup>ax</sup>, L. Sawyer<sup>bh</sup>, T. Scanlon<sup>aq</sup>, D. Schaile<sup>y</sup>, R.D. Schamberger<sup>bt</sup>, Y. Scheglov<sup>an</sup>, H. Schellman<sup>ba</sup>, T. Schliephake<sup>z</sup>, C. Schwanenberger<sup>ar</sup>, A. Schwartzman<sup>bp</sup>, R. Schwienhorst<sup>bm</sup>, J. Sekaric<sup>aw</sup>, H. Severini<sup>bw</sup>, E. Shabalina<sup>ay</sup>, M. Shamim<sup>bg,\*</sup>, V. Shary<sup>r</sup>, A.A. Shchukin<sup>am</sup>, R.K. Shivpuri<sup>ab</sup>, V. Siccaldi<sup>s</sup>, V. Simak<sup>j</sup>, V. Sirotenko<sup>ax</sup>, P. Skubic<sup>bw</sup>, P. Slattery<sup>bs</sup>, D. Smirnov<sup>bc</sup>, G.R. Snow<sup>bo</sup>, J. Snow<sup>bv</sup>, S. Snyder<sup>bu</sup>, S. Söldner-Rembold<sup>ar</sup>, L. Sonnenschein<sup>q</sup>, A. Sopczak<sup>ap</sup>, M. Sosebee<sup>bz</sup>, K. Soustruznik<sup>i</sup>, B. Spurlock<sup>bz</sup>, J. Stark<sup>n</sup>, J. Steele<sup>bh</sup>, V. Stolin<sup>ak</sup>, D.A. Stoyanova<sup>am</sup>, J. Strandberg<sup>bl</sup>, S. Strandberg<sup>ao</sup>, M.A. Strang<sup>bq</sup>, E. Strauss<sup>bt</sup>, M. Strauss<sup>bw</sup>, R. Ströhmer<sup>y</sup>, D. Strom<sup>ba</sup>, L. Stutte<sup>ax</sup>, S. Sumowidagdo<sup>aw</sup>, P. Svoisky<sup>bc</sup>, A. Sznajder<sup>c</sup>, P. Tamburello<sup>as</sup>, A. Tanasijczuk<sup>a</sup>, W. Taylor<sup>f</sup>, J. Temple<sup>as</sup>, B. Tiller<sup>y</sup>, F. Tissandier<sup>m</sup>, M. Titov<sup>r</sup>, V.V. Tokmenin<sup>aj</sup>, T. Toole<sup>bi</sup>, I. Torchiani<sup>w</sup>, T. Trefzger<sup>x</sup>, D. Tsybychev<sup>bt</sup>, B. Tuchming<sup>r</sup>, C. Tully<sup>bp</sup>, P.M. Tuts<sup>br</sup>, R. Unalan<sup>bm</sup>, L. Uvarov<sup>an</sup>, S. Uvarov<sup>an</sup>, S. Uzunyan<sup>az</sup>, B. Vachon<sup>f</sup>, P.J. van den Berg<sup>ah</sup>, R. Van Kooten<sup>bb</sup>, W.M. van Leeuwen<sup>ah</sup>, N. Varelas<sup>ay</sup>, E.W. Varnes<sup>as</sup>, I.A. Vasilyev<sup>am</sup>, M. Vaupel<sup>z</sup>, P. Verdier<sup>t</sup>, L.S. Vertogradov<sup>aj</sup>, M. Verzocchi<sup>ax</sup>, F. Villeneuve-Seguié<sup>aq</sup>, P. Vint<sup>aq</sup>, P. Vokac<sup>j</sup>, E. Von Toerne<sup>bg</sup>, M. Voutilainen<sup>bp,5</sup>, R. Wagner<sup>bp</sup>, H.D. Wahl<sup>aw</sup>, L. Wang<sup>bi</sup>, M.H.L.S. Wang<sup>ax</sup>, J. Warchol<sup>bc</sup>, G. Watts<sup>cd</sup>, M. Wayne<sup>bc</sup>, G. Weber<sup>x</sup>, M. Weber<sup>ax</sup>, L. Welty-Rieger<sup>bb</sup>, A. Wenger<sup>w,6</sup>, N. Wermes<sup>v</sup>, M. Wetstein<sup>bi</sup>, A. White<sup>bz</sup>, D. Wicke<sup>z</sup>, G.W. Wilson<sup>bf</sup>, S.J. Wimpenny<sup>av</sup>, M. Wobisch<sup>bh</sup>, D.R. Wood<sup>bk</sup>, T.R. Wyatt<sup>ar</sup>, Y. Xie<sup>by</sup>, S. Yacoub<sup>ba</sup>, R. Yamada<sup>ax</sup>, M. Yan<sup>bi</sup>, T. Yasuda<sup>ax</sup>, Y.A. Yatsunenko<sup>aj</sup>, K. Yip<sup>bu</sup>, H.D. Yoo<sup>by</sup>, S.W. Youn<sup>ba</sup>, J. Yu<sup>bz</sup>, A. Zatserklyaniy<sup>az</sup>, C. Zeitnitz<sup>z</sup>, T. Zhao<sup>cd</sup>, B. Zhou<sup>bl</sup>, J. Zhu<sup>bt</sup>, M. Zielinski<sup>bs</sup>, D. Zieminska<sup>bb</sup>, A. Zieminski<sup>bb,\*</sup>, L. Zivkovic<sup>br</sup>, V. Zutshi<sup>az</sup>, E.G. Zverev<sup>al</sup>

<sup>a</sup> Universidad de Buenos Aires, Buenos Aires, Argentina

<sup>b</sup> LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

<sup>c</sup> Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>d</sup> Universidade Federal do ABC, Santo André, Brazil

<sup>e</sup> Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

<sup>f</sup> University of Alberta, Edmonton, Alberta, and Simon Fraser University, Burnaby, British Columbia, and York University, Toronto, Ontario, and McGill University, Montreal, Quebec, Canada

<sup>g</sup> University of Science and Technology of China, Hefei, People's Republic of China

<sup>h</sup> Universidad de los Andes, Bogotá, Colombia

<sup>i</sup> Center for Particle Physics, Charles University, Prague, Czech Republic

<sup>j</sup> Czech Technical University, Prague, Czech Republic

<sup>k</sup> Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

<sup>l</sup> Universidad San Francisco de Quito, Quito, Ecuador

<sup>m</sup> LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France

<sup>n</sup> LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, France

<sup>o</sup> CPPM, IN2P3/CNRS, Université de la Méditerranée, Marseille, France

<sup>p</sup> LAL, Université Paris-Sud, IN2P3/CNRS, Orsay, France

<sup>q</sup> LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France

<sup>r</sup> DAPNIA/Service de Physique des Particules, CEA, Saclay, France

<sup>s</sup> IPHC, Université Louis Pasteur et Université de Haute Alsace, CNRS/IN2P3, Strasbourg, France

<sup>t</sup> IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France

<sup>u</sup> III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany

<sup>v</sup> Physikalisches Institut, Universität Bonn, Bonn, Germany

<sup>w</sup> Physikalisches Institut, Universität Freiburg, Freiburg, Germany

<sup>x</sup> Institut für Physik, Universität Mainz, Mainz, Germany

<sup>y</sup> Ludwig-Maximilians-Universität München, München, Germany

<sup>z</sup> Fachbereich Physik, University of Wuppertal, Wuppertal, Germany

<sup>aa</sup> Panjab University, Chandigarh, India

<sup>ab</sup> Delhi University, Delhi, India

<sup>ac</sup> Tata Institute of Fundamental Research, Mumbai, India

<sup>ad</sup> University College Dublin, Dublin, Ireland

<sup>ae</sup> Korea Detector Laboratory, Korea University, Seoul, Republic of Korea

<sup>af</sup> SungKyunKwan University, Suwon, Republic of Korea

<sup>ag</sup> CINVESTAV, Mexico City, Mexico

<sup>ah</sup> FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

<sup>ai</sup> Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands

<sup>aj</sup> Joint Institute for Nuclear Research, Dubna, Russia

<sup>ak</sup> Institute for Theoretical and Experimental Physics, Moscow, Russia

<sup>al</sup> Moscow State University, Moscow, Russia

- <sup>am</sup> Institute for High Energy Physics, Protvino, Russia  
<sup>an</sup> Petersburg Nuclear Physics Institute, St. Petersburg, Russia  
<sup>ao</sup> Lund University, Lund, and Royal Institute of Technology and Stockholm University, Stockholm, and Uppsala University, Uppsala, Sweden  
<sup>ap</sup> Lancaster University, Lancaster, United Kingdom  
<sup>aq</sup> Imperial College, London, United Kingdom  
<sup>ar</sup> University of Manchester, Manchester, United Kingdom  
<sup>as</sup> University of Arizona, Tucson, AZ 85721, USA  
<sup>at</sup> Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA  
<sup>au</sup> California State University, Fresno, CA 93740, USA  
<sup>av</sup> University of California, Riverside, CA 92521, USA  
<sup>aw</sup> Florida State University, Tallahassee, FL 32306, USA  
<sup>ax</sup> Fermi National Accelerator Laboratory, Batavia, IL 60510, USA  
<sup>ay</sup> University of Illinois at Chicago, Chicago, IL 60607, USA  
<sup>az</sup> Northern Illinois University, DeKalb, IL 60115, USA  
<sup>ba</sup> Northwestern University, Evanston, IL 60208, USA  
<sup>bb</sup> Indiana University, Bloomington, IN 47405, USA  
<sup>bc</sup> University of Notre Dame, Notre Dame, IN 46556, USA  
<sup>bd</sup> Purdue University Calumet, Hammond, IN 46323, USA  
<sup>be</sup> Iowa State University, Ames, IA 50011, USA  
<sup>bf</sup> University of Kansas, Lawrence, KS 66045, USA  
<sup>bg</sup> Kansas State University, Manhattan, KS 66506, USA  
<sup>bh</sup> Louisiana Tech University, Ruston, LA 71272, USA  
<sup>bi</sup> University of Maryland, College Park, MD 20742, USA  
<sup>bj</sup> Boston University, Boston, MA 02215, USA  
<sup>bk</sup> Northeastern University, Boston, MA 02115, USA  
<sup>bl</sup> University of Michigan, Ann Arbor, MI 48109, USA  
<sup>bm</sup> Michigan State University, East Lansing, MI 48824, USA  
<sup>bn</sup> University of Mississippi, University, MS 38677, USA  
<sup>bo</sup> University of Nebraska, Lincoln, NE 68588, USA  
<sup>bp</sup> Princeton University, Princeton, NJ 08544, USA  
<sup>bq</sup> State University of New York, Buffalo, NY 14260, USA  
<sup>br</sup> Columbia University, New York, NY 10027, USA  
<sup>bs</sup> University of Rochester, Rochester, NY 14627, USA  
<sup>bt</sup> State University of New York, Stony Brook, NY 11794, USA  
<sup>bu</sup> Brookhaven National Laboratory, Upton, NY 11973, USA  
<sup>bv</sup> Langston University, Langston, OK 73050, USA  
<sup>bw</sup> University of Oklahoma, Norman, OK 73019, USA  
<sup>bx</sup> Oklahoma State University, Stillwater, OK 74078, USA  
<sup>by</sup> Brown University, Providence, RI 02912, USA  
<sup>bz</sup> University of Texas, Arlington, TX 76019, USA  
<sup>ca</sup> Southern Methodist University, Dallas, TX 75275, USA  
<sup>cb</sup> Rice University, Houston, TX 77005, USA  
<sup>cc</sup> University of Virginia, Charlottesville, VA 22901, USA  
<sup>cd</sup> University of Washington, Seattle, WA 98195, USA

## ARTICLE INFO

## Article history:

Received 15 March 2008  
 Received in revised form 20 April 2008  
 Accepted 12 May 2008  
 Available online 21 May 2008  
 Editor: L. Rolandi

## PACS:

14.80.Ly  
 12.60.Jv

## ABSTRACT

We present a search for the pair production of scalar top quarks,  $\tilde{t}$ , using 995 pb<sup>-1</sup> of data collected in  $p\bar{p}$  collisions with the DØ detector at the Fermilab Tevatron Collider at  $\sqrt{s} = 1.96$  TeV. Both scalar top quarks are assumed to decay into a charm quark and a neutralino ( $\tilde{\chi}_1^0$ ), where  $\tilde{\chi}_1^0$  is the lightest supersymmetric particle. This leads to a final state with two acoplanar charm jets and missing transverse energy. We find the yield of such events to be consistent with the standard model expectation, and exclude sets of  $\tilde{t}$  and  $\tilde{\chi}_1^0$  masses at the 95% C.L. that substantially extend the domain excluded by previous searches.

Published by Elsevier B.V.

Supersymmetry (SUSY) may provide a solution to the hierarchy problem if the SUSY particles have masses less than 1 TeV, strongly motivating the searches for supersymmetric objects at the Fermilab Tevatron Collider. SUSY predicts the existence of part-

ners with identical quantum numbers to all Standard Model (SM) particles except for spin. There exist two spin zero SUSY partners of the top quark corresponding to the latter's left and right handed states. Several arguments exist in favor of a light scalar top quark ( $\tilde{t}$ ). The  $\tilde{t}$  mass  $m_{\tilde{t}}$  receives negative contributions proportional to the top quark Yukawa coupling in the renormalization group equations. This makes the  $\tilde{t}$  weak eigenstates lighter than other squarks [1]. Mixing between the left and right handed superpartners of the top quark, being proportional to the top quark mass  $m_t$ , leads to a large mass splitting between the two physical eigenstates. This makes one of the  $\tilde{t}$  considerably lighter than the other. Additionally, a light  $\tilde{t}$  that strongly couples to the Higgs boson could also generate a large enough CP violating phase to explain the mechanism for electroweak baryogenesis [2].

\* Corresponding author.

E-mail address: mansooram@fnal.gov (M. Shamim).

<sup>1</sup> Visitor from Augustana College, Sioux Falls, SD, USA.

<sup>2</sup> Visitor from The University of Liverpool, Liverpool, UK.

<sup>3</sup> Visitor from ICN-UNAM, Mexico City, Mexico.

<sup>4</sup> Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.

<sup>5</sup> Visitor from Helsinki Institute of Physics, Helsinki, Finland.

<sup>6</sup> Visitor from Universität Zürich, Zürich, Switzerland.

✘ Deceased.

In R-parity conserving models [3], the lightest supersymmetric particle (LSP) is stable, and cosmological constraints indicate that it should be neutral and colorless [4]. In the following we assume conservation of R-parity and take  $\tilde{\chi}_1^0$ , the lightest of four SUSY particles that result from the mixing between the SUSY partners of the SM neutral gauge and Higgs bosons, to be the LSP.

In the search reported in this Letter, we consider the range  $m_{\tilde{t}} < m_b + m_{\tilde{\chi}_1^+}$  and  $m_{\tilde{t}} < m_W + m_b + m_{\tilde{\chi}_1^0}$ , where  $m_b$  is the  $b$  quark mass,  $m_{\tilde{\chi}_1^0}$  is the  $\tilde{\chi}_1^0$  mass and  $m_{\tilde{\chi}_1^+}$  is the  $\tilde{\chi}_1^+$  mass, with  $\tilde{\chi}_1^+$  being the lighter of two mass eigenstates resulting from the mixing of the SUSY partners of charged gauge and Higgs bosons. The dominant  $\tilde{t}$  decay mode in this model is the flavor changing process  $\tilde{t} \rightarrow c\tilde{\chi}_1^0$  and is assumed to occur with 100% branching fraction. The  $\tilde{t} \rightarrow t\tilde{\chi}_1^0$  decay is kinematically forbidden over the  $\tilde{t}$  mass range accessible in this search, and the tree level four-body decays  $\tilde{t} \rightarrow bf\tilde{f}'\tilde{\chi}_1^0$  can be neglected [5].

In  $p\bar{p}$  collisions,  $\tilde{t}$  pairs are produced via quark–antiquark annihilation and gluon fusion. The  $\tilde{t}$  pair production cross section ( $\sigma_{\tilde{t}\tilde{t}}$ ) primarily depends on  $m_{\tilde{t}}$ , and a weak dependence on other SUSY parameters affects only the higher-order corrections. At  $\sqrt{s} = 1.96$  TeV which is the centre-of-mass energy available at the Fermilab Tevatron Collider,  $\sigma_{\tilde{t}\tilde{t}}$  at next-to-leading-order (NLO), calculated with PROSPINO [6], ranges from 15 pb to 1 pb for  $100 \leq m_{\tilde{t}} \leq 160$  GeV. These cross sections are calculated using CTREQ6.1M parton distribution functions (PDFs) [7] and equal renormalization and factorization scales  $\mu_{\text{rf}} = m_{\tilde{t}}$ . A theoretical uncertainty of  $\approx 20\%$  is estimated due to scale and PDF choices. The  $\tilde{t}\tilde{t}$  event topology consists of two acoplanar charm jets and missing transverse energy ( $\cancel{E}_T$ ) from the neutralinos that escape detection. Searches for  $\tilde{t}$  pair production in the jets plus missing transverse energy mode have been reported by collaborations working at the CERN LEP collider [8], and the CDF [9,10] and DØ [11,12] collaborations. The highest excluded mass to date is  $m_{\tilde{t}} < 134$  GeV (95% C.L.) for  $m_{\tilde{\chi}_1^0} = 48$  GeV [12].

The  $\tilde{t}$  search is performed in the data collected with the DØ detector during Run IIa of the Tevatron and corresponds to an integrated luminosity of  $995 \pm 61$  pb $^{-1}$  [13]. A detailed description of the DØ detector can be found in [14]. The central tracking system consists of a silicon microstrip tracker and a fiber tracker, both located within a 2 T superconducting solenoidal magnet. A liquid-argon and uranium calorimeter covers pseudorapidity  $|\eta| \lesssim 4.2$ , where  $\eta = -\ln[\tan(\theta/2)]$ , and  $\theta$  is the polar angle with respect to the proton beam direction. An outer muon system, covering  $|\eta| < 2$ , consists of layers of tracking detectors and scintillation counters on both sides of 1.8 T iron toroids.

The data sample collected from April 2003 to February 2006 with the jets +  $\cancel{E}_T$  triggers was analyzed for the  $\tilde{t}$  search. The trigger conditions require the  $\cancel{E}_T$  and its separation from all jets to be greater than 30 GeV and  $25^\circ$ , respectively, where  $\cancel{E}_T$  is the transverse energy computed only from jets. Jets are reconstructed using an iterative midpoint cone algorithm with radius  $\mathcal{R}_{\text{cone}} = 0.5$  [15]. The data set is reduced to a sample of 1.5 million events by requiring at least two jets with  $p_T > 15$  GeV and  $\cancel{E}_T > 40$  GeV.

Signal samples are simulated using PYTHIA 6.323 [16] for  $m_{\tilde{t}}$  ranging from 95 GeV to 165 GeV and  $\tilde{\chi}_1^0$  masses from 45 GeV to 90 GeV. The largest expected backgrounds for this search are  $W$  and  $Z$  bosons produced in association with jets, denoted as  $V + \text{jets}$ . The  $V + \text{jets}$  and  $t\bar{t}$  processes are simulated using ALPGEN 2.05 [17] interfaced with PYTHIA for the generation of initial and final state radiation and hadronization. The background samples for single top quark and diboson production are simulated using COMPHEP [18] and PYTHIA, respectively. The PDF set CTREQ6L1 is used for both signal and background samples, and all generated events are subjected to full GEANT-based [19] simulation of the detec-

tor response. Simulated signal and background events are overlaid with recorded unbiased beam crossings to incorporate the effect of multiple interactions that occur in a single beam crossing. After reconstruction, simulated events are weighted properly to ensure that the instantaneous luminosity distribution is the same in data and the simulated Monte Carlo (MC) samples. A parametrization of the trigger efficiency measured from the data is applied to simulated MC events in order to fold in trigger effects. The multijet background, not included in the MC samples, is directly estimated from data.

A large data sample of  $Z/\gamma^*(\rightarrow ee) + \text{jets}$  events, corresponding to an integrated luminosity of  $1067 \pm 65$  pb $^{-1}$ , from the same data period as the  $\tilde{t}$  search, is used to improve the prediction of  $V + \text{jets}$  backgrounds. For this study,  $Z$  boson candidates are selected using two high transverse energy ( $E_T > 15$  GeV) clusters that deposit more than 90% of their energy in the electromagnetic calorimeter, have shower shapes consistent with expectations for electrons, are matched with tracks reconstructed in the central tracker, and form an invariant mass between 65 GeV and 115 GeV. At least two jets with  $p_T > 15$  GeV and  $|\eta_{\text{det}}| < 2.5$  are required, where  $|\eta_{\text{det}}|$  is the jet pseudorapidity calculated using the assumption that the jet originates from the detector center. The predicted number of  $Z/\gamma^*(\rightarrow ee) + \geq 2$  jets events is calculated using ALPGEN after correcting for differences in electron and jet reconstruction efficiencies between data and MC and normalizing the MC to the inclusive number of  $Z$  boson events in data. The ALPGEN prediction is corrected in each jet multiplicity bin by a reweighting function that depends on the transverse momentum of the  $Z$  boson to obtain better agreement between the model and data. The reweighting function is derived by fitting the ratio of the transverse momentum distribution of  $Z$  boson data to that from the MC prediction. After reweighting, all other kinematical variables in the  $Z/\gamma^*(\rightarrow ee) + \geq 2$  jets sample applicable to the  $\tilde{t}$  search are well described by MC.

The multijet background in  $Z/\gamma^*(\rightarrow ee) + \text{jets}$  events is estimated from a fit to the dielectron invariant mass distribution. The ratio of the number of  $ee$  events produced by  $\gamma^*$  to the combined number of events produced by  $Z$  boson and  $\gamma^*$  is determined from MC. This ratio is used to extract the multijet contribution by fitting the dielectron invariant mass in data with an exponential function for the combined contribution from multijet processes and  $\gamma^*$  and a Breit–Wigner convolved with a Gaussian for total number of  $Z$  boson events.

For the  $\tilde{t}$  search, the predicted SM background from  $V + \text{jets}$  sources is normalized to the number of  $Z/\gamma^*(\rightarrow ee) + 2$  jets events after subtracting the multijet background,  $N_{Z(\rightarrow ee)+2}^{\text{data}}$ . As an example, the normalization weight assigned to simulated  $Z(\rightarrow \nu\bar{\nu})$  events with  $n$  light partons is

$$w_{Z(\nu\bar{\nu})+n}^{\text{MC}} = f \frac{N_{Z(\rightarrow ee)+2}^{\text{data}} \sigma_{Z(\nu\bar{\nu})+n}^{\text{ALP}} \epsilon_{Z(\nu\bar{\nu})+n}}{N_{Z(\nu\bar{\nu})+n}^{\text{MC}} \sigma_{Z(\rightarrow ee)+2\text{lp}}^{\text{ALP}} \epsilon_{Z(\rightarrow ee)+2\text{lp}}}. \quad (1)$$

Here  $N_{Z(\nu\bar{\nu})+n}^{\text{MC}}$  is the number of simulated  $Z(\rightarrow \nu\bar{\nu}) + n$  light parton jets events;  $\sigma_{Z(\nu\bar{\nu})+n}^{\text{ALP}}$  and  $\sigma_{Z(\rightarrow ee)+2\text{lp}}^{\text{ALP}}$  are the cross sections predicted by ALPGEN for  $Z(\rightarrow \nu\bar{\nu}) + n$  and  $Z/\gamma^*(\rightarrow ee) + 2$  light parton jets, respectively; and  $\epsilon_{Z(\nu\bar{\nu})+n}$  and  $\epsilon_{Z(\rightarrow ee)+2\text{lp}}$  are the corresponding detection efficiencies. The factor  $f = 0.89 \pm 0.02$  is applied to correct for three effects: the absence of  $\gamma^*$  contribution to  $Z(\nu\bar{\nu}) + \text{jets}$  events, the normalization of MC light jets to a data sample that contains all flavors of jets, and the difference in the luminosities of the data set used for the  $\tilde{t}$  search ( $995$  pb $^{-1}$ ) and the  $Z/\gamma^*(\rightarrow ee) + 2$  jets data set ( $1067$  pb $^{-1}$ ).

The normalization weight assigned to simulated  $W(\rightarrow \ell\nu) + n$  light partons is

**Table 1**

Numbers of data events and cumulative signal efficiency for  $m_{\tilde{t}} = 150$  and  $m_{\tilde{\chi}_1^0} = 70$  GeV after each event selection

Selection	Events left	Signal eff. (%)
Initial selection and trigger	$1.5 \times 10^6$	55.9
C1: exactly two jets	464477	29.5
C2: $\cancel{E}_T > 40$ GeV	440161	27.5
C3: $\Delta\phi(\text{jet}_1, \text{jet}_2) < 165^\circ$	278505	26.5
C4: jet-1 $p_T > 40$ GeV	216382	24.7
C5: jet-1 $ \eta_{\text{det}}  < 1.5$	113591	24.6
C6: jet-2 $p_T > 20$ GeV	80987	22.0
C7: jet-2 $ \eta_{\text{det}}  < 1.5$	62910	20.1
C8: jet-1 jet-2 CPF $> 0.85$	49140	19.8
C9: isolated track veto	23832	13.4
C10: isolated electron veto	23194	13.3
C11: isolated muon veto	23081	13.3
C12: $\Delta\phi_{\text{max}} - \Delta\phi_{\text{min}} < 120^\circ$	9753	12.6
C13: $A > -0.05$	3733	12.0
C14: $\Delta\phi(\text{jet}, \cancel{E}_T) > 50^\circ$	3375	11.6
C15: $\cancel{E}_T > 60$ GeV	2288	10.0

$$w_{W(\ell\nu)+n}^{\text{MC}} = f \frac{N_{Z(ee)+2}^{\text{data}} \sigma_{W(\ell\nu)+n}^{\text{ALP}} \epsilon_{W(\ell\nu)+n}}{N_{W(\ell\nu)+n}^{\text{MC}} \sigma_{Z(ee)+2lp}^{\text{ALP}} \epsilon_{Z(ee)+2lp}} \alpha(p_T), \quad (2)$$

where

$$\alpha(p_T) = \frac{\left[ \frac{1}{\sigma_W^{\text{NLO}}} \frac{d\sigma_W^{\text{NLO}}}{dp_T} \right] \left[ \frac{1}{\sigma_Z^{\text{ALP}}} \frac{d\sigma_Z^{\text{ALP}}}{dp_T} \right]}{\left[ \frac{1}{\sigma_Z^{\text{NLO}}} \frac{d\sigma_Z^{\text{NLO}}}{dp_T} \right] \left[ \frac{1}{\sigma_W^{\text{ALP}}} \frac{d\sigma_W^{\text{ALP}}}{dp_T} \right]}, \quad (3)$$

is the product of the ratio of the normalized differential cross sections for  $W$  and  $Z$  bosons production at NLO [20] and predicted by ALPGEN, respectively.

The motivation behind using this technique is to lower the luminosity times cross section uncertainty ( $\approx 6.1\% \oplus 15\%$ ) on the predicted number of events towards the 5% statistical uncertainty of the  $Z/\gamma^*(\rightarrow ee) + 2$  jets normalization sample. The combined 15% uncertainty on the theoretical cross section for various background processes is mainly due to the choice of PDF and the renormalization and factorization scale. The signal and smaller backgrounds such as  $t\bar{t}$ , diboson, and single top quark production are normalized using the measured absolute luminosity. For these processes NLO cross sections were computed with MCFM 5.1 [21].

The search strategy for  $\tilde{t}$  involves three steps which include the application of the selection criteria on kinematical variables, heavy flavor (HF) tagging and optimization of the final selection depending on  $\tilde{t}$  and  $\tilde{\chi}_1^0$  masses. The data set for the  $\tilde{t}$  search is reduced to a sample of 2288 potential  $\tilde{t}\bar{\tilde{t}}$  candidates, by applying the 15 selection criteria denoted by C1–C15 and summarized in Table 1. The main motivation for C1 is to reduce the multijet background. Requirements C2 to C7 help in reducing the  $W +$  jets and multijet backgrounds. The condition on the charged particle fraction (CPF) in C8 requires that at least 85% of the jets' charged particle transverse momenta be associated with tracks originating from the selected primary vertex in the event. This track confirmation requirement removes events with spurious  $\cancel{E}_T$  due to the choice of an incorrect primary vertex. C9–C11 are applied to reject  $W +$  jets background with isolated leptons from  $W$  boson decay. For an electron to be isolated, the energy deposited in the calorimeter in a cone of radius 0.4 in  $\eta - \phi$  around the electron direction should not be more than 15% of the energy deposited in the electromagnetic layers inside a cone of radius 0.2. A muon is declared isolated if the sum of the energies of all tracks other than the muon in a cone of radius 0.5 is less than 2.5 GeV, and the calorimeter energy deposited in a hollow cone with inner and outer radii 0.1 and 0.4 around the muon direction is less than 5 GeV. A track with  $p_T > 5$  GeV is considered isolated if no other track with  $p_T > 1.5$  GeV is found in a hollow cone of inner and outer radii 0.1

and 0.4 around the track considered. This condition also helps suppress backgrounds with  $\tau$  leptons where the  $\tau$  decays hadronically. Remaining instrumental background is removed using a quantity defined by the angular separation between all jets and the  $\cancel{E}_T$  of the event,  $D = \Delta\phi_{\text{max}} - \Delta\phi_{\text{min}}$ , where  $\Delta\phi_{\text{max}}(\Delta\phi_{\text{min}})$  is the largest (smallest) azimuthal separation between a jet and  $\cancel{E}_T$ ; and an asymmetry variable defined as  $A = (\cancel{E}_T - \cancel{H}_T)/(\cancel{E}_T + \cancel{H}_T)$ . The requirements applied on these variables are given by C12 and C13. Fig. 1 shows that both of these variables are very effective in eliminating multijet background which dominates in data for large  $D$  and negative  $A$ .


The 2288 events selected in data can be compared to the  $2199 \pm 18^{+316}_{-321}$  events predicted from the simulation normalized to  $Z/\gamma^*(\rightarrow ee) + 2$  jets events or  $2292 \pm 19^{+527}_{-532}$  events predicted using absolute luminosity normalization, with the first quoted uncertainty due to finite MC statistics and the second due to systematic effects described in more detail below. The small remaining multijet background in the  $\tilde{t}$  search analysis is estimated after applying all analysis conditions shown in Table 1 except that on  $\cancel{E}_T$ . The SM  $V +$  jets background contribution, estimated from the MC, is subtracted from the data before fitting the  $\cancel{E}_T$  distribution with exponential and power law functions in the control region ( $40 < \cancel{E}_T < 60$  GeV). To estimate the multijet contribution in the signal region, we extrapolate the fit results to  $\cancel{E}_T > 60$  GeV. The average of the two fit results is taken as the multijet background estimate, while the difference between the two is taken as the systematic uncertainty. This amounts to  $14.4 \pm 10.7(\text{stat}) \pm 5.1(\text{sys})$  events contributed by multijet background before HF tagging and optimization of selection cuts.

After selecting candidate events on the basis of topology, HF tagging is used to identify charm jets in the final state. A neural network (NN) tagging tool [22] that combines information from three different D0 HF taggers to maximize the  $b$  quark tagging efficiency ( $\approx 73\%$ ) is used for this purpose.


The first tagger converts information from the impact parameter of the tracks identified in a jet into a probability that all tracks originate from the primary vertex, where the impact parameter is the distance of closest approach to the interaction point in a plane perpendicular to the beam axis. The second tagger identifies the presence of vertices that are significantly displaced from the primary vertex and associated with a jet. The third tagger makes use of the number of tracks with large impact parameter significance, where the significance is defined as the ratio of the impact parameter to its uncertainty. The result of the combination is a NN output. A requirement on the NN output is made that preserves high efficiency for detection of charm jets ( $\approx 30\%$ ) with a  $\approx 6\%$  probability for a light parton jet to be mistakenly tagged. The efficiency for  $c$  jet tagging is obtained by scaling the  $b$  jet tagging efficiency measured in the data by the  $c$ -tagging-to- $b$ -tagging efficiency ratio computed in the MC.

At the final stage of the analysis, additional selection criteria on three kinematic variables;  $\cancel{E}_T$ ,  $S = \Delta\phi_{\text{max}} + \Delta\phi_{\text{min}}$ , and  $H_T$ , with  $H_T$  defined as the scalar sum of the  $p_T$  of all jets, are optimized by maximizing the expected lower limit on the neutralino mass for a given  $m_{\tilde{t}}$ . The variable  $S$  after requiring at least one jet in the event to be HF tagged is shown in Fig. 2.


Minimum values of  $H_T$  are varied from 60 GeV to 140 GeV in steps of 20 GeV, while those for  $\cancel{E}_T$  are varied from 60 GeV to 100 GeV in steps of 10 GeV. Events having the values of these quantities above the minima are kept. Maximum values of  $S$  are tested between  $240^\circ$  and  $320^\circ$  in steps of  $20^\circ$ , and events having  $S$  below the minimum are retained. For each set of requirements, the expected value of the signal confidence level ( $CL_s$ ) [23] under the hypothesis that only background is present is evaluated using all  $\tilde{t}$  and  $\tilde{\chi}_1^0$  mass combinations, taking into account systematic uncertainties. The set of criteria that return  $\langle CL_s \rangle = 0.05$  for the



**Fig. 1.** Distributions of the asymmetry  $A = (\cancel{E}_T - \cancel{H}_T)/(\cancel{E}_T + \cancel{H}_T)$  with the requirement on  $D = \Delta\phi_{\max} - \Delta\phi_{\min}$  inverted (a) and applied (c). Distributions of  $D$  with the requirement on  $A$  inverted (b) and applied (d) for data (points with error bars), for SM backgrounds (histogram), and for a signal with  $m_{\tilde{t}} = 150$  GeV and  $m_{\tilde{\chi}_1^0} = 70$  GeV (hatched histogram). In all plots the signal contribution has been scaled up by five and  $\cancel{E}_T > 60$  GeV is required. The excess in data at  $A = 0$  and  $D = 0$ –10 degrees is consistent with the systematic uncertainties on the predicted background.



**Fig. 2.** Distributions of  $S = \Delta\phi_{\max} + \Delta\phi_{\min}$  for data (points with error bars), SM background (histogram), and a signal with  $m_{\tilde{t}} = 150$  GeV and  $m_{\tilde{\chi}_1^0} = 70$  GeV (hatched histogram) after requiring HF tagging but before optimization.



**Fig. 3.** Distributions of  $H_T$  after applying optimized requirements on  $\cancel{E}_T$  and  $S$  for data (points with error bars), SM background (histogram), and a signal with  $m_{\tilde{t}} = 150$  GeV and  $m_{\tilde{\chi}_1^0} = 70$  GeV (hatched histogram).

**Table 2**

Optimized values of selections, numbers of observed data and predicted background events. A requirement of  $\cancel{E}_T > 70$  GeV was chosen in all cases. The values of  $m_{\tilde{t}}$  and  $H_T$  are in GeV while those for  $S$  are in degrees

$m_{\tilde{t}}$	$H_T$	$S$	Observed	Predicted
95–130	$> 100$	$< 260$	83	$85.3 \pm 1.8^{+12.8}_{-13.0}$
135–145	$> 140$	$< 300$	57	$59.0 \pm 1.6^{+8.5}_{-8.8}$
150–160	$> 140$	$< 320$	66	$66.6 \pm 1.1^{+9.6}_{-10.0}$

**Table 3**

For three  $\tilde{t}$  and  $\tilde{\chi}_1^0$  mass combinations: signal efficiencies and the numbers of signal events expected. The first errors are statistical and second systematic. The nominal (NLO) signal cross section and upper limit at the 95% C.L. are also shown

$(m_{\tilde{t}}, m_{\tilde{\chi}_1^0})$ GeV	Efficiency (%)	Expected signal events	$\sigma_{\text{nom}}$ pb	$\sigma_{95}$ pb
(130, 55)	1.5	$51.9 \pm 2.7^{+7.2}_{-7.1}$	3.44	2.41
(140, 80)	0.9	$19.6 \pm 0.8^{+2.8}_{-2.5}$	2.24	2.87
(150, 70)	2.1	$30.8 \pm 1.2^{+4.2}_{-3.7}$	1.49	1.42

highest neutralino mass corresponding to a given  $m_{\tilde{t}}$  are chosen to be the optimal ones.


The optimized values of the selections for different  $m_{\tilde{t}}$  are given in Table 2 along with the number of events observed in data and expected SM background. In all cases a requirement of  $\cancel{E}_T \geq 70$  GeV is imposed. No contamination remains from multijet background at this point in the analysis; it is therefore neglected while setting the limit. Efficiencies for three signal mass points along with the expected numbers of events are shown in Table 3. The distribution of  $H_T$  after optimization but with the constraint on  $H_T$  removed is shown in Fig. 3. The final distribution of  $\cancel{E}_T$  is shown in Fig. 4. The detailed SM background composition is given in Table 4.

Systematic uncertainties are evaluated for each  $\tilde{t}$  and  $\tilde{\chi}_1^0$  mass combination for the optimized set of requirements. Sources of systematic uncertainty include jet energy scale, jet energy resolution, jet identification and reconstruction, the jet multiplicity requirement, trigger efficiency, data to MC scale factors, normalization of background, HF tagging, luminosity determination, choice of PDF, and  $W$  boson  $p_T$  reweighting. The effect of the jet multiplicity requirement on the background is studied using  $Z/\gamma^*(\rightarrow ee) + \text{jets}$  events. The spectrum of transverse momentum of the third jet in data events with three or more jets is observed to be very well described by the simulation generated with ALPGEN. The  $\approx 1\%$  statis-

**Table 4**

Numbers of predicted background events from different SM sources for a selection optimized for  $m_{\tilde{t}} \geq 150$  GeV. The uncertainties are due to the limited MC statistics

SM process	Number of events
$W(\rightarrow \ell\nu) + \text{jets}$	$20.0 \pm 0.7$
$Z(\rightarrow \nu\bar{\nu}) + \text{jets}$	$15.8 \pm 0.5$
$W(\rightarrow \ell\nu) + \text{HF} (b\bar{b}, c\bar{c})$	$12.6 \pm 0.5$
$Z(\rightarrow \nu\bar{\nu}) + \text{HF} (b\bar{b}, c\bar{c})$	$11.6 \pm 0.4$
$t\bar{t}$ and single top	$3.7 \pm 0.1$
$WW, WZ, ZZ$	$2.7 \pm 0.1$
$Z(\rightarrow \ell\ell) (e, \mu, \tau) + \text{jets}$	$0.1 \pm 0.01$
$Z(\rightarrow \ell\ell) (e, \mu, \tau) + \text{HF} (b\bar{b}, c\bar{c})$	$0.1 \pm 0.01$
Total	$66.6 \pm 1.1$



**Fig. 4.** Final distributions of  $E_T$  for data (points with error bars), SM background (histogram), and a signal with  $m_{\tilde{t}} = 150$  GeV and  $m_{\tilde{\chi}_1^0} = 70$  GeV (hatched histogram).

tical uncertainty of the lowest  $p_T$  bin, where the bulk of the events are, is taken as a systematic uncertainty introduced by the jet multiplicity requirement. To study the effect of the same requirement on the  $\tilde{t}$  signal, where a third jet enters an event primarily through initial or final state radiation, the  $p_T$  spectrum of the leading jet in simulated  $Z/\gamma^*(\rightarrow ee)$  events generated with PYTHIA is examined. Comparison between data and simulation shows a slight excess in data in the low  $p_T$  bin; this discrepancy is used to estimate a systematic uncertainty of  $\pm 1.5\%$  on the signal acceptance attributable to the jet multiplicity requirement. The uncertainty on the signal acceptance and background estimation due to the PDF choice was determined using the CTEQ6.1M PDF set.


The combined 10% uncertainty on the background normalization includes: 5% uncertainty from  $Z/\gamma^*(\rightarrow ee) + \text{jets}$  statistics assigned to all  $V + \text{jets}$  samples; 50% uncertainty on the NLO cross section assigned to the  $V + \text{HF}$  background; 6.1% luminosity uncertainty assigned to  $t\bar{t}$ , diboson, and single top quark background; and 8%, 6% and 15% uncertainties on NLO cross sections for  $t\bar{t}$ , diboson, and single top quark production, respectively. The uncertainty on the background estimation due to the  $W$  boson  $p_T$  reweighting is estimated using two different methods to estimate the  $W + \text{jets}$  background. In the first method, the  $W + \text{jets}$  background is estimated using the expression given in Eq. (2). In the second method, the same reweighting function as applied to the  $Z$  boson was used to reweight the  $W$  boson  $p_T$  which is equivalent to setting  $\alpha(p_T) = 1$  in Eq. (2). Detailed estimates of all systematic uncertainties are given in Table 5.

Using the assumption that  $\tilde{t}$  decays into a charm quark and a neutralino with 100% branching fraction and the nominal  $\tilde{t}$  pair production cross section, the largest  $m_{\tilde{t}}$  excluded by this analysis is 155 GeV, for a neutralino mass of 70 GeV at the 95% C.L. With the theoretical uncertainty on the  $\tilde{t}$  pair production cross section taken

**Table 5**

Breakdown of systematic uncertainties on the SM background and for a signal point with  $m_{\tilde{t}} = 150$  GeV and  $m_{\tilde{\chi}_1^0} = 70$  GeV

Source	SM background	Signal
Jet energy	+1.7%	+2%
Scale	-2.5%	-4%
Jet resolution	$\pm 1\%$	$\pm 1\%$
Jet reconstruction and identification	$\pm 0.8\%$	$\pm 0.1\%$
Trigger	$\pm 6\%$	$\pm 6\%$
Scale factor	$\pm 5\%$	$\pm 5\%$
Normalization	$\pm 10\%$	-
Luminosity	-	$\pm 6.1\%$
HF tagging	$\pm 4.1\%$	$\pm 3.5\%$
PDF choice	$\pm 4\%$	$\pm 8.7\%$
Two jet cut	-	-5.5%
$W$ boson $p_T$ reweighting	$\pm 3\%$	$\pm 1.5\%$



**Fig. 5.** Region in the  $\tilde{t}-\tilde{\chi}_1^0$  mass plane excluded at the 95% C.L. by the present search. The observed (expected) exclusion contour is shown as the brown solid (dashed) line. The yellow band represents the theoretical uncertainty on the scalar top quark pair production cross section due to PDF and renormalization and factorization scale choice. Results from previous searches [8,10,12] are also shown. The results from [10,12] correspond to the observed limits obtained with the one sigma lower bound of the predicted NLO cross section for stop pair production. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this Letter.)

into account, the largest limit on  $m_{\tilde{t}}$  is 150 GeV, for  $m_{\tilde{\chi}_1^0} = 65$  GeV. These results are shown in Fig. 5.

In summary, DØ has searched for scalar top quarks in jets plus missing transverse energy final states using  $1 \text{ fb}^{-1}$  of data. No evidence for  $\tilde{t}$  production has been found. This analysis substantially extends the excluded region of the  $\tilde{t}-\tilde{\chi}_1^0$  mass plane over the searches carried out previously.

## Acknowledgements

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Republic of Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation.

## References

- [1] H. Baer, et al., Phys. Rev. D 50 (1994) 4517, and references therein.



- [2] D. Delepine, et al., Phys. Lett. B 386 (1996) 183.
- [3] P. Fayet, Phys. Lett. B 238 (1977) 489.
- [4] J. Ellis, et al., Nucl. Phys. B 238 (1984) 453.
- [5] C. Boehm, A. Djouadi, Y. Mambrini, Phys. Rev. D 61 (2000) 095006.
- [6] W. Beenakker, et al., Nucl. Phys. B 515 (1998) 3.
- [7] J. Pumplin, et al., JHEP 0207 (2002) 012;  
D. Stump, et al., JHEP 0310 (2003) 046.
- [8] LEPSUSYWG Collaboration, ALEPH Collaboration, DELPHI Collaboration, L3 Collaboration, OPAL Collaboration, Note LEPSUSYWG/04-02.1, [http://lepsusy.web.cern.ch/lepsusy/www/squarks\\_summer04/stop\\_combi\\_208\\_final.html](http://lepsusy.web.cern.ch/lepsusy/www/squarks_summer04/stop_combi_208_final.html).
- [9] T. Aafolder, et al., CDF Collaboration, Phys. Rev. Lett. 84 (2000) 5704.
- [10] T. Aaltonen, et al., CDF Collaboration, Phys. Rev. D 76 (2007) 072010.
- [11] V.M. Abazov, et al., DØ Collaboration, Phys. Rev. Lett. 93 (2004) 011801.
- [12] V.M. Abazov, et al., DØ Collaboration, Phys. Lett. B 645 (2007) 119.
- [13] T. Andeen, et al., FERMILAB-TM-2365, 2007.
- [14] V.M. Abazov, et al., DØ Collaboration, Nucl. Instrum. Methods Phys. Res. A 565 (2006) 463.
- [15] G.C. Blazey, et al., in: U. Baur, R.K. Ellis, D. Zeppenfeld, (Eds.), Proceedings of the Workshop: QCD and Weak Boson Physics in Run II, Fermilab-Pub-00/297, 2000.
- [16] T. Sjöstrand, et al., Comput. Phys. Commun. 135 (2001) 238.
- [17] M.L. Mangano, et al., JHEP 0307 (2003) 001.
- [18] E. Boos, et al., CompHEP Collaboration, Nucl. Instrum. Methods Phys. Res. A 534 (2004) 250.
- [19] R. Brun, F. Carminati, CERN Program Library Long Writeup W5013, 1993, unpublished.
- [20] K. Melnikov, F. Petriello, Phys. Rev. D 74 (2006) 114017.
- [21] J.M. Campbell, R.K. Ellis, Phys. Rev. D 60 (1999) 113006.
- [22] T. Scanlon, Ph.D. thesis, Imperial College London, FERMILAB-THESIS-2006-43, 2006.
- [23] T. Junk, Nucl. Instrum. Methods Phys. Res. A 434 (1999) 435;  
A. Read, in: 1st Workshop on Confidence Limits, CERN Report No. CERN-2000-005, 2000.